



Lawrence Berkeley Laboratory

UNIVERSITY OF CALIFORNIA


Presented at the Conference on the Intersections
between Particle and Nuclear Physics,
Steamboat Springs, CO, May 23-30, 1984

HEAVY ION PHYSICS - INTRODUCTION TO THE HEAVY
ION PARALLEL SESSIONS AT CIPANP

L.S. Schroeder and T.W. Ludlam

June 1984

RECEIVED
LAWRENCE
BERKELEY LABORATORY
JUL 23 1984
LIBRARY AND
DOCUMENTS SECTION



LBL-17961
c.2

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

HEAVY ION PHYSICS - INTRODUCTION TO THE
HEAVY ION PARALLEL SESSIONS AT CIPANP*

L. S. Schroeder
Lawrence Berkeley Laboratory
University of California, Berkeley, CA 94720

T. W. Ludlam
Brookhaven National Laboratory
Upton, NY 11973

ABSTRACT

A brief introduction into some of the physics probed by heavy ion projectiles at low, intermediate and high energies is given. Emphasis is placed on nuclear matter under extreme conditions in this discussion, which should provide a common area of interest to both particle and nuclear physics.

INTRODUCTION

In putting together the six parallel sessions on heavy ion physics we have emphasized the area of "extreme conditions" in nuclear matter rather than the more conventional aspects such as nuclear structure. This area of heavy ion studies is getting more attention these days, particularly with the push to higher energies and therefore provides a relatively natural bridge to couple interests in both particle and nuclear physics.

*This work was supported by the Director, Office of Energy Research, Division of Nuclear Physics of the Office of High Energy and Nuclear Physics of the U. S. Department of Energy under contracts DE-AC03-76SF00098 and DE-AC02-76CH00016.

PHYSICS WITH HEAVY IONS

As a starting point it is convenient to discuss what happens in heavy ion collisions as a function of bombarding energy. This is done below by dividing the energy scale into: low, intermediate and high energies. The rough energy ranges listed are for fixed target experiments.

- 1) Low Energy: This covers the range of bombarding energies below ~ 30 - 50 MeV/nucleon. At the lower end of this scale, the nucleus is often thought of as a classical liquid; but with increasing energy the nucleonic composition starts to manifest itself and we enter a region where the nucleus can be roughly treated as a gas. The large numbers of nucleons present in these collisions allow extreme conditions in nuclei to be studied, including:
 - o superheavy element production/searches
 - o introduction of large angular momenta into the nuclear system (study of angular momentum limits reached, relaxation of system, . . .)
 - o local deposition of energy (hot spot formation and propagation)
 - o ability to study collective aspects of nuclei by using pion production near the absolute kinematic limits as a probe.

- 2) Intermediate Energy: This corresponds roughly to the interval from 50 to 1000 MeV/nucleon and spans the range from a simple "pure" nucleon gas treatment to the on-set of the hadron gas regime (i.e., excitation of the nucleonic degrees of freedom yielding nuclear matter with N^* , Δ , π , η , ρ , ω , . . . content). In this range the incident nuclear projectiles have the ability of delivering a large amount of energy over a large volume, allowing experiments to:
 - o probe extreme conditions of temperature (T) and baryon density (ρ) in nuclear matter with the goal of obtaining information on the equation of state $W(\rho, T)$
 - o search for unusual or exotic nuclear states or phenomena such as abnormal nuclei (Lee-Wick), pion condensation, Δ -matter, anomalous, . . .
 - o use the property of kinematic focussing (in peripheral collisions) at higher energies as a powerful tool for studying production of nuclei far from stability and in producing beams of secondary neutron-rich nuclei for conventional nuclear physics studies.

3) High Energy: Covering the region from a few GeV/nucleon to the highest energies available in the cosmic radiation. At these higher energies one expects to move from the arena of confined hadronic matter to one of deconfined quark matter (quark-gluon plasma). Heavy nuclei at high energy provide us with the opportunity of creating both high energy and baryon density in central nucleus-nucleus collisions with the potential of:

- o creating the quark-gluon plasma in both a baryon-rich (large ρ) and baryon-free (high T) region
- o forming in a controlled environment a new state of matter which is coupled to conditions in the early universe
- o probe the long-range aspects of QCD, and
- o extend our present studies of hot, dense hadronic matter to higher T , ρ .

Figure 1 shows the phase diagram of hadronic matter in the temperature-relative baryon density plane. It is only within the last decade that we have started to probe nuclei beyond the region of $T \approx 0$, $\rho/\rho_0 = 1$. Nucleus-nucleus collisions offer us a unique opportunity of probing a much larger region of this diagram, including the region of deconfined quark matter--the quark-gluon plasma.




Figure 1: Phase diagram for hadronic matter.

PROGRAM FOR HEAVY ION SESSIONS

Listed below is the program for the six parallel heavy ion sessions.

- Session A - J. Kogut (Univ. of Ill.): "Recent Results from Lattice Gauge Calculations"
(May 23)
- P. Carruthers (LANL): "Hydrodynamic Flow Structures in Quantum Field Theory"
 - A. Goldhaber (Stony Brook): "Nuclear Stopping Power"
- Session B - G. Yodh (Univ. of Maryland): "High Energy Nucleus-Nucleus Collisions: A Summary of Cosmic Ray Data"
(May 24)
- Y. Takahashi (Marshall Space Flight): "Recent Results from the JACEE Experiment"
 - G. Young (ORNL): "Subthreshold Production of Pions"
- Session C - D. Boal (Simon Fraser/MSU): "Low Energy Phase Transitions in Nuclear Matter"
(May 28)
- A. Hirsch (Purdue): "Fragment Production and the Liquid-Gas Phase Transition"
 - S. Fung (UC Riverside): "Source Size Measurements (HBT) in High Energy Nucleus-Nucleus Collisions and First Results on Uranium in Streamer Chamber at the Bevalac"
 - K. Crowe (UC Berkeley): " $\pi\pi$ Correlation Measurements in Fe + Fe Collisions"
- Session D - H.-G. Ritter (GSI/LBL): "Collective Flow and High Density Nuclear Matter"
(May 28)
- A. Sandoval (GSI/LBL): "Nuclear Matter Equation of State"
 - K. Nakai (Univ. Tokyo/KEK): "Pions, Protons and Excited Nucleons from Hadron-Nucleus Reactions at Several GeV"
- Session E - M. Tanaka (BNL): "Study of the Neutral Energy Spectrum in the Central Region of $\alpha\alpha$ Interactions at $\sqrt{s} = 124$ GeV at the CERN ISR"
(May 29)
- W. Zajc (Univ. of Penn.): "Particle Production in $\alpha\alpha$ Collisions at the CERN-ISR"
- Session F - C. M. Ko (Texas A&M): "Strange Particle Production in Relativistic Nuclear Collisions"
(May 29)
- D. Sivers (ANL): "Classical Chromodynamics and Extended Nuclear Matter"
 - P. J. S. Watson (Carleton Univ.): "Hadrons in Nuclear Matter"

This report was done with support from the Department of Energy. Any conclusions or opinions expressed in this report represent solely those of the author(s) and not necessarily those of The Regents of the University of California, the Lawrence Berkeley Laboratory or the Department of Energy.

Reference to a company or product name does not imply approval or recommendation of the product by the University of California or the U.S. Department of Energy to the exclusion of others that may be suitable.

TECHNICAL INFORMATION DEPARTMENT
LAWRENCE BERKELEY LABORATORY
UNIVERSITY OF CALIFORNIA
BERKELEY, CALIFORNIA 94720