

UC Davis

UC Davis Previously Published Works

Title

Evidence implicating blood-brain barrier impairment in the pathogenesis of acquired epilepsy following acute organophosphate intoxication

Permalink

<https://escholarship.org/uc/item/7j17w7ct>

Journal

Journal of Pharmacology and Experimental Therapeutics, 388(2)

ISSN

0022-3565

Authors

Bernardino, Pedro N

Luo, Audrey S

Andrew, Peter M

et al.

Publication Date

2024-02-01

DOI

10.1124/jpet.123.001836

Copyright Information

This work is made available under the terms of a Creative Commons Attribution-NoDerivatives License, available at <https://creativecommons.org/licenses/by-nd/4.0/>

Peer reviewed

Evidence implicating blood-brain barrier impairment in the pathogenesis of acquired epilepsy following acute organophosphate intoxication

Authors: Pedro N. Bernardino¹; Audrey S. Luo¹; Peter M. Andrew¹; Chelsea M. Unkel¹; Marco I. Gonzalez²; Angie Gelli³; Pamela J. Lein¹

¹ Department of Molecular Biosciences, University of California, Davis, School of Veterinary Medicine, Davis, CA, 95616, USA

² Department of Neurology, University of California, Davis, School of Medicine, Sacramento, CA, 95817, USA

³ Department of Pharmacology, University of California, Davis, School of Medicine, Davis, CA 95616, USA

Running title (max 60 characters): BBB impairment and epilepsy following acute OP intoxication

Corresponding author: Pamela J. Lein, PhD

Molecular Biosciences, UC Davis School of Veterinary Medicine
1089 Veterinary Research Drive
Davis, California, 95616, USA
e-mail: pjlein@ucdavis.edu; phone: 530-752-1970

Number of text pages: 36

Number of figures: 2

Number of words in Abstract: 197

Number of words in Introduction: 443

Number of words in Main text (not counting introduction): 12,706

Recommended section: Medical countermeasures (Special Section)

Non-standard abbreviation list:

AChE – acetylcholinesterase; AD – Alzheimer’s disease; BBB – blood-brain barrier; BM – basement membrane; CAM – cell adhesion molecule; CNS – central nervous system; DFP – diisopropylfluorophosphate; DPE – days-post exposure; EC – endothelial cell; IL – interleukin; Kir4.1 – inward-rectifying potassium channels; MCP – monocyte chemoattractant protein; MIP – macrophage inflammatory protein; MMP – matrix metalloproteinase; MRI – magnetic resonance imaging; nAChR – nicotinic acetylcholine receptor; NMDA – N-methyl-D-aspartate; NVU – neurovascular unit; OP - organophosphate; PDGFr- β - soluble platelet derived growth factor receptor β ; PG – prostaglandins; PTE – post-traumatic epilepsy; ROS/RNS – reactive oxygen and nitrogen species; SE – *status epilepticus*; SOC – standard of care; SRS – spontaneous recurrent seizures; TBI – traumatic brain injury; TGF- β – transforming growth factor beta; TJ – tight junction; TLE – temporal lobe epilepsy; TNF – tumor necrosis factor; VEGF – vascular endothelial growth factor; ZO-1 – zonula occludens-1

Abstract

Organophosphate (OP) poisoning can trigger cholinergic crisis, a life-threatening toxidrome that includes seizures and *status epilepticus* (SE). These acute toxic responses are associated with persistent neuroinflammation and spontaneous recurrent seizures (SRS), also known as acquired epilepsy. Blood-brain barrier (BBB) impairment has recently been proposed as a pathogenic mechanism linking acute OP intoxication to chronic adverse neurological outcomes. In this review, we briefly describe the cellular and molecular components of the BBB, review evidence of altered BBB integrity following acute OP intoxication, and discuss potential mechanisms by which acute OP intoxication may promote BBB dysfunction. We highlight the complex interplay between neuroinflammation and BBB dysfunction that suggests a positive feedforward interaction. Lastly, we examine research from diverse models and disease states that suggest mechanisms by which loss of BBB integrity may contribute to epileptogenic processes. Collectively, the literature identifies BBB impairment as a convergent mechanism of neurological disease and justifies further mechanistic research into how acute OP intoxication causes BBB impairment and its role in the pathogenesis of SRS, and potentially other long-term neurological sequelae. Such research is critical for evaluating BBB stabilization as a neuroprotective strategy for mitigating OP-induced epilepsy and possibly seizure disorders of other etiologies.

Significance Statement: Clinical and preclinical studies support a link between BBB dysfunction and epileptogenesis; however, a causal relationship has been difficult to prove. Mechanistic studies to delineate relationships between BBB dysfunction and epilepsy may provide novel insights into BBB stabilization as a neuroprotective strategy for mitigating epilepsy resulting from acute OP intoxication and non-OP causes, and potentially other adverse neurological conditions associated with acute OP intoxication, such as cognitive impairment.

1. Introduction

Organophosphate (OP) cholinesterase inhibitors represent a global public health threat with accidental and intentional OP poisonings impacting millions of people annually (Bird et al., 2014; Mew et al., 2017; Jett et al., 2020). OPs inhibit the enzyme acetylcholinesterase (AChE), which normally functions to limit cholinergic neurotransmission by hydrolyzing synaptic acetylcholine. Acute inhibition of AChE activity by >60-70% produces cholinergic crisis, a life-threatening toxidrome characterized by parasympathetic activation, seizures, *status epilepticus* (SE), and respiratory failure (Eddleston et al., 2008). The current standard of care (SOC) for treatment of acute OP intoxication includes supportive maintenance of cardiorespiratory parameters, and treatment with atropine to antagonize cholinergic muscarinic receptors, an oxime to reactivate AChE, and a benzodiazepine to control seizures (Eddleston et al., 2008; Glauser et al., 2016). While SOC improves survival following acute OP intoxication, it does not prevent long-term adverse neurologic consequences of acute OP intoxication, which include spontaneous recurrent seizures (SRS), also referred to as acquired epilepsy (Waheed et al., 2014; Talabani et al., 2018; Guignet et al., 2020; Gage et al., 2022). The development of more effective therapeutic strategies for mitigating epileptic and other chronic neurological sequelae is urgently needed to improve the quality of life of individuals who survive the acute toxic effects of OP poisoning.

The epileptogenic period after acute OP intoxication has been characterized in experimental models (de Araujo Furtado et al., 2010; de Araujo Furtado et al., 2012; Todorovic et al., 2012; Shrot et al., 2014; Deshpande and DeLorenzo, 2020; Guignet et al., 2020; Maupu et al., 2021), but pathogenic mechanisms underlying OP-induced epilepsy remain speculative. Studies of epilepsy caused by etiologies other than OP poisoning increasingly implicate blood-brain barrier (BBB) impairment in epileptogenesis (Marchi et al., 2012; van Vliet et al., 2015; van Lanen et al., 2021; Li et al., 2023). Critically, biomarkers of BBB dysfunction correlate with the development and progression of epilepsy in humans and in diverse preclinical models (Tenreiro et al., 2016; Bar-Klein et al., 2017; Dadas and Janigro, 2019; Mendes et al., 2019; Nation et al., 2019), suggesting a potential anti-epileptogenic benefit of protecting BBB function. While this hypothesis has yet to be tested in models of acute OP intoxication, recent studies have reported

BBB impairment following acute OP intoxication (Abdel-Rahman et al., 2002; Song et al., 2004; Bar-Klein et al., 2017; Rojas et al., 2021; Rojas et al., 2022).

In this review, we discuss the physiological and morphological properties of the BBB and summarize the current understanding and data gaps regarding BBB impairment in acute OP intoxication. We further elaborate on potential mechanisms underlying BBB dysfunction following OP exposure, with a focus on neuroinflammation, and consider the evidence implicating BBB disruption in epileptogenesis.

2. Structure and function of the BBB

The BBB plays an essential role in homeostatic control of the brain's microenvironment by restricting the movement of ions and other small hydrophilic molecules, macromolecules, and cells between the blood and brain parenchyma. Its physiological roles include: 1) controlling ion homeostasis in the extracellular milieu of the brain; 2) limiting movement of blood-borne neurotoxic molecules into the brain; 3) regulating passage of nutrients into the brain; and 4) restricting peripheral immune protein/cell infiltration (Saunders et al., 2008; Abbott et al., 2010; Profaci et al., 2020; Segura-Collar et al., 2022). These diverse functions are made possible by an assembly of specialized cellular and non-cellular structures that form the neurovascular unit (NVU), which is considered the functional unit of the BBB.

The NVU consists of endothelial cells (EC), adherence junction and tight junction (TJ) proteins, basement membrane (BM), pericytes and astrocytic end-feet (Fig. 1), and both the composition and function of the NVU are modulated by neuronal and microglial input (Hawkins and Davis, 2005; Obermeier et al., 2013; Obermeier et al., 2016). ECs and TJs establish a physical barrier unique to BBB vasculature. EC-TJ pairing, which forms a continuous structure throughout the vascular lining devoid of fenestrae, functions to restrict paracellular transport. Relative to blood vessel ECs in other organ systems, BBB ECs have reduced expression of leukocyte adhesion molecules, which limits diapedesis of white blood cells into the brain. BBB ECs express efflux transporters to facilitate transcellular movement of nutrients and promote removal of harmful hydrophilic molecules from the brain parenchyma (Wolburg and Lippoldt, 2002; Begley and Brightman, 2003; Hawkins and Davis, 2005; Abbott et al., 2006; Ransohoff and Engelhardt, 2012; Obermeier et al., 2013). Pericytes, a less-well characterized cell type in

the NVU, are more abundant around blood vessels in the brain than in other organs (Armulik et al., 2011). While there are gaps in knowledge about the function and activity of neurovascular pericytes, their absence is associated with embryonic lethality, increased BBB permeability, and atypical TJ distribution (Lindahl et al., 1997; Hellström et al., 2001; Obermeier et al., 2013). The BM, which comprises the NVU extracellular matrix, functions to stabilize the cellular assembly, enable crosstalk between the cellular components of the NVU, and augment the physical barrier separating the blood from the brain parenchyma (Del Zoppo et al., 2006; Profaci et al., 2020). The BM is comprised of diverse structural and adherence proteins produced by cells of the NVU. Inflammation and various disease states can alter BM composition; conversely, variation in BM assembly influences cell migration and signaling, which can modulate inflammation and disease progression (Sorokin, 2010; Obermeier et al., 2013; Profaci et al., 2020). The outermost parenchymal-adjacent NVU structure is astrocytic end-feet, which play a vital role in the development and maintenance of BBB integrity (Tao-Cheng et al., 1987; Díaz-Castro et al., 2023). Critically, their dysfunction results in BBB compromise (Menezes et al., 2014; Schreiner et al., 2015) and significant BBB leakage (Heithoff et al., 2021) in mouse models. Current evidence also supports a role for astrocytes in regulating local blood flow, osmolality balance mediated by specialized channels (e.g., aquaporin 4 and inward-rectifying potassium channels, Kir4.1), and production of growth factors responsible for BBB maturation and maintenance, such as angiopoetin-1, glial-derived neurotrophic factor (GDNF), basic fibroblast growth factor (bFGF) and transforming growth factor- β (TGF- β) (Zonta et al., 2003; Abbott et al., 2006; Nag, 2011; Heinemann et al., 2012; Wong et al., 2013; Cabezas et al., 2014).

Neurons communicate with the NVU through release of neurotransmitters that modulate local blood flow and BBB permeability, with the magnitude of these effects directly proportional to the level of neuronal activity (Vazana et al., 2016; Kaplan et al., 2020). Microglial activation and phenotypic state also influence BBB function. Pro-inflammatory microglia (M1-like) increase BBB permeability via release of reactive oxygen and nitrogen species and pro-inflammatory cytokines. In contrast, anti-inflammatory M2-like microglia protect BBB integrity via release of interleukin (IL)-10, which prevents decreased expression and/or restores physiologic expression levels of TJ proteins and other NVU components (Ronaldson and Davis, 2020; Knopp et al., 2022). Conversely, there is substantial evidence that loss of BBB integrity or

activation of NVU cells can activate microglia and promote neuroinflammation (Barkauskas et al., 2015; Bowyer et al., 2016; Thurgur and Pinteaux, 2019), indicating a complex, bidirectional relationship between these processes.

Proper BBB function requires coordination between multiple structural components and signaling pathways internal and external to the NVU, and these elements can be dysregulated by diverse pathological conditions, including chemical intoxication and inflammation. Establishing causal relationships between these processes has been challenging, in part because of the complex interactions between the BBB, neuronal activity and inflammation, but also because of the paucity of data regarding the temporal relationships between these events (Friedman, 2011). However, defining these relationships is critical to understanding the pathogenesis of diverse neurovascular, neurodegenerative, and epileptogenic conditions, and will provide insights regarding therapeutic strategies for preventing and/or treating these diseases.

3. Acute OP intoxication increases BBB permeability

Increased BBB permeability has been documented following acute OP intoxication in preclinical models. OP-induced BBB leakiness was first reported by Abdel-Rahman et al. (2002) who demonstrated that at 1 day post-exposure (DPE), acute sarin intoxication increased radioactivity in the brain parenchyma of rats following intravenous administration of [³H]-hexamethonium, a physiologically non-BBB penetrant molecule. Subsequent studies revealed that rats acutely intoxicated with sarin experienced BBB leakage at 2 DPE, 7 DPE and 1 month post-exposure as detected using contrast-enhanced magnetic resonance imaging (MRI), immunohistochemical analysis of serum albumin, and extravasation of Evan's blue into the brain (Bar-Klein et al., 2017). Western blot analyses of endogenous serum albumin in rat brains similarly indicated that acute intoxication with soman or diisopropylfluorophosphate (DFP) increased BBB permeability at 4 DPE (Rojas et al., 2021). Further analyses in the same DFP model revealed increased extravasation of albumin in the brain parenchyma at 3 h post-exposure, but not at 1 and 2 DPE (Rojas et al., 2022). There is also one report suggesting that exposure to paraoxon at a dose that did not cause SE increased leakage of horseradish peroxidase from capillaries within 10 min of exposure in juvenile, but not adult, rats (Song et al., 2004).

While the observation that acute OP intoxication increased BBB permeability is consistent across all published studies in which this outcome has been measured, the temporal progression and magnitude of BBB dysfunction varies between these reports, likely due to differences in experimental models and methodologies used to assess BBB integrity. It is well known that the spatiotemporal profile of neuropathology varies between OPs despite comparable cholinesterase inhibition (Pope, 1999; Bushnell and Moser, 2006; Costa, 2006; Rojas et al., 2021). Some of this variability is likely due to non-cholinergic mechanisms of OP toxicity, which are dependent on the structure of the OP in question. In the context of OP effects on BBB integrity, differing results likely also reflect variable seizure duration across studies. Specifically, in one study (Song et al., 2004), seizure activity was not reported and the dose of parathion used was unlikely to cause SE before brains were harvested for analysis; in other studies, SE was terminated at 30 or 60 min post-exposure (Bar-Klein et al., 2017; Rojas et al., 2021), while in the two remaining studies, seizures were not pharmacologically controlled and likely persisted for hours (Abdel-Rahman et al., 2002; Rojas et al., 2022). Methodological differences in assessing BBB integrity likely also contribute to variable results, e.g., the sensitivity and accuracy of detecting focal BBB leakage is significantly greater with histological evaluation compared to western blot analysis of albumin extravasation. Additionally, the spatiotemporal progression of BBB dysfunction has yet to be rigorously evaluated. Each published study to date assessed BBB leakage at a single time point or did not compare outcomes between time points, and unfortunately, the lone study that examined spatial characterization of BBB impairment (Bar-Klein et al., 2017) is confounded by the use of *in vivo* imaging parameters with low signal-to-noise ratio.

In summary, while there is consistent evidence of BBB impairment following acute OP intoxication, disparities in the published literature confound mechanistic or causal interpretations and complicate discussions concerning the therapeutic potential of BBB stabilization after OP exposures.

3.1 Putative mechanisms by which acute OP intoxication may cause BBB damage

Mechanistic characterization of neuropathological outcomes observed following acute OP intoxication provides clues as to how these exposures may cause BBB damage (Fig. 2). The canonical mechanism of acute OP toxicity is AChE inhibition (Tsai and Lein, 2021), resulting in

hyperstimulation of peripheral and central cholinergic synapses. BBB ECs express nicotinic acetylcholine receptors (nAChRs) that function to control vasodilation (Uchida and Hotta, 2009), and possibly also BBB permeability. In support of this possibility, in primary rat cerebral EC culture under oxidative stress, pharmacological activation of nAChRs dysregulated expression of TJs and adherence junction proteins coincident with increased trans-endothelial permeability (Hutamekalin et al., 2008). These data are consistent with *in vivo* observations of elevated BBB permeability in rats dosed with nicotine (Hawkins et al., 2005). Critically, nicotine-induced BBB permeability is attenuated by co-administration of nAChR antagonists (Hawkins et al., 2005), confirming the direct involvement of nAChR stimulation in BBB impairment under conditions of hypercholinergic stimulation. Cholinergic hyperstimulation also activates angiogenic processes to promote blood vessel growth, a process linked to greater vascular permeability, in a nAChR-dependent manner (Arias et al., 2009). Conversely, there are data suggesting that cholinergic hyperstimulation may protect BBB integrity via activation of cholinergic anti-inflammatory pathways and attenuation of OP-associated neuroinflammation (Pavlov et al., 2003; Dash et al., 2016; Colás et al., 2018; Martín et al., 2018).

Several processes downstream of the OP-induced hypercholinergic state likely also influence BBB health and function. Seizures and SE are known to increase brain vascular permeability (van Vliet et al., 2015; Löscher and Friedman, 2020), suggesting their role as primary drivers of BBB impairment following acute OP intoxication. A cross-sectional study of epileptic patients revealed significantly increased serological levels of BBB damage-associated markers hours after the onset of a convulsive episode (Cudna et al., 2023). While this study measured biomarkers in serum and not in brain tissue, these data are consistent with BBB dysregulation following a seizure event. Additionally, cellular and molecular changes associated with BBB leakiness are observed in epilepsy patients and animal models. Angiogenesis and decreased expression of TJ proteins has been reported in patients with temporal lobe epilepsy (TLE) (Rigau et al., 2007; van Vliet et al., 2007; Marchi and Lerner-Natoli, 2013). In patients with drug-resistant TLE, angiogenesis is associated with elevated expression of vascular endothelial growth factor (VEGF) and VEGF receptor, as well as altered TJ protein expression, specifically increased claudin-5 but decreased zonula occludens 1 (ZO-1) and occludin (Castañeda-Cabral et al., 2020).

Seizures induced by inhibition of voltage-gated potassium channel with 4-aminopyridine were reported to cause pericyte-associated changes in vascular diameter and BBB permeability in rat hippocampal slice cultures and in cortical arterioles of rats perfused with 4-aminopyridine containing cerebrospinal fluid (Prager et al., 2019). Similarly, in the hippocampus of mice with kainic acid-induced SE, pericyte proliferation and pericyte-microglia clustering was identified at 3 and 7 DPE (Klement et al., 2018), suggesting the involvement of pericytes in seizure-induced BBB dysfunction. However, additional research is needed to confirm this relationship. Reactive astrogliosis, which is often observed following seizures (Wilcox et al., 2015), may dysregulate physiological processes integral to BBB permeability, including potassium and glutamate homeostasis (David et al., 2009; Coulter and Steinhäuser, 2015). Following seizures, elevated matrix metalloproteinase (MMP) activity is observed, which correlates with lower TJ protein expression (Rempe et al., 2018). Importantly, SE-induced downregulation of TJ expression is attenuated by upstream MMP inhibition (Lischper et al., 2010; Rempe et al., 2018), suggesting MMPs are critically involved in regulating BBB integrity.

Oxidative stress is a widely recognized consequence of acute OP intoxication (Lukaszewicz-Hussain, 2010; Pearson and Patel, 2016; Vanova et al., 2018; Farkhondeh et al., 2020) triggered by dysregulated mitochondrial activity, glutamate-induced Ca^{2+} influx, lipid peroxidation and neuroinflammation (Patel et al., 1996; Middlemore-Risher et al., 2011; Farkhondeh et al., 2020; Song et al., 2020). Oxidative stress promotes BBB permeability directly by damaging cellular components of the NVU and indirectly by degrading TJs and BM (Song et al., 2020).

4. The role of neuroinflammation in OP-induced BBB impairment

Neuroinflammation is a physiologic response to central nervous system (CNS) insult characterized by an orchestrated activation and communication of glial cells resident to the brain and, in some cases, white blood cells from the periphery. This coordinated response involves secretion of inflammatory signaling factors to control injury and heal damaged tissue (Kraft and Harry, 2011; DiSabato et al., 2016; Shabab et al., 2017). However, excessive and persistent neuroinflammatory responses can be detrimental to nervous system function (Lyman et al., 2014; DiSabato et al., 2016), including dysregulation of the BBB (Takata et al., 2021).

Following acute OP intoxication, neuroinflammation is observed clinically and preclinically with the response varying by agent, exposure condition, and time of assessment post-intoxication (Guignet and Lein, 2019; Kanthasamy et al., 2019). Characterization of the spatiotemporal profile of OP-induced neuroinflammatory responses in preclinical models using *ex vivo* (Angoa-Pérez et al., 2010; Sisó et al., 2017) and/or *in vivo* (Flannery et al., 2016; Hobson et al., 2019) methods has demonstrated robust activation of astrocytes and microglia in multiple brain regions that vary temporally in a region-dependent manner. Multiple pro-inflammatory factors are upregulated following acute OP intoxication, including tumor necrosis factor (TNF)- α , IL-1 β , IL-6, monocyte chemoattractant protein (MCP)-1, and macrophage inflammatory protein (MIP)-1 α (Dillman et al., 2009; Johnson and Kan, 2010; Johnson et al., 2011; Hobson et al., 2019). In parallel, levels of reactive oxygen and nitrogen species (ROS/RNS) are increased (Puttachary et al., 2015; Pearson and Patel, 2016; Vanova et al., 2018; Guignet and Lein, 2019; Gage et al., 2021; Meyer et al., 2023). Interestingly, the spatiotemporal distribution of neuroinflammation observed in preclinical models of acute OP intoxication during the first week post-exposure (Flannery et al., 2016; Bar-Klein et al., 2017; Sisó et al., 2017) is very similar to that reported in a recent study of the regional characterization of BBB impairment (Bernardino et al., under review). However, no mechanistic or causal associations have yet been investigated between these two processes in the context of acute OP intoxication.

4.1. Inflammation-associated activation of CNS cells influences BBB function

Observations from preclinical models of seizures triggered by factors other than OPs suggest several pathways by which neuroinflammation could influence BBB integrity. One is activation of microglia. Microglia respond to diverse central nervous system insults by releasing pro-inflammatory and anti-inflammatory cytokines and chemokines, primarily IL-1 β , IL-6, TNF- α , MCP-1 CCL5, and CXCL1, as well as prostaglandins (PGs) and ROS (DiSabato et al., 2016). While these signaling molecules initiate and coordinate immune responses to CNS injury, they also alter the activity of NVU cells to disrupt homeostatic BBB function (Takata et al., 2021). For example, *in vitro* studies using rat and mice EC cultures demonstrated that LPS-activated microglia enhanced BBB permeability to sodium fluorescein and decreased the activity of the P-glycoprotein efflux pump, and these effects were blocked by inhibiting TNF- α or nicotinamide

adenine dinucleotide phosphate (NADPH) oxidase (Nishioku et al., 2010; Sumi et al., 2010; Matsumoto et al., 2012). IL-1 β , TNF- α , IL-6 and oncostatin M directly dysregulated expression and organization of TJ proteins in primary EC cultures derived from mouse or rat brains (Takata et al., 2008; Voirin et al., 2020), while other lipid mediators (e.g. PGE₂, PGF_{2 α} and sphingosine-1-phosphate) increased paracellular permeability in numerous *in vitro* and *in vivo* models (Dalvi et al., 2015; Kim et al., 2015; Mohan et al., 2018; Rojas et al., 2019; Nakagawa and Aruga, 2020; Takata et al., 2021). Collectively, these data suggest that microglia and microglia-derived inflammatory factors may promote BBB leakiness following acute OP intoxication.

Activated astrocytes can also impair the BBB by modulating NVU function. Astrocytic secretion of inflammatory signaling molecules, including VEGF-A, MMP-9, MCP-1, and IL-6, has been observed in an *in vitro* mouse model of ischemic stroke and *in vivo* rat model of brain ischemia (Shan et al., 2019). Clinical studies of human samples and *in vitro* studies with rat cell cultures indicated that VEGF and MMP-9 induced TJ rearrangement and cleavage (Machida et al., 2017; Castañeda-Cabral et al., 2020). Downregulation of VEGF and MMP-9 expression by glucagon-like peptide-1 receptor agonism preserved TJ organization and expression in a rat stroke model (Shan et al., 2019). Astrocytic MCP-1 is a chemoattractant that promotes trans-BBB migration of monocytes from the blood into brain parenchyma (Weiss et al., 1998) to augment neuroinflammatory responses.

There are limited data to support a role for pericyte-mediated inflammatory responses in BBB dysfunction. When stimulated by TNF- α *in vitro*, pericytes secreted a unique profile of pro-inflammatory regulators, including IL-6, MIP-1 α , IL-1 β and inducible nitric oxide synthase (iNOS), which promoted the release of MMP-9 (Takata et al., 2011; Matsumoto et al., 2014; Segura-Collar et al., 2022). *In vitro* pro-inflammatory activation of pericytes and ECs increased production of IL-8 coincident with trans-endothelial migration of neutrophils in a MMP-2/-9-dependent manner (Pieper et al., 2013), suggesting that pericytes enhance recruitment of immune cells to sites of CNS insult (Matsumoto et al., 2014).

Cytokines modulate neuronal excitability and synaptic neurotransmission (Vezzani and Viviani, 2015), and while these effects have not yet been causally linked to increased BBB leakage, there are studies hinting at such a relationship. ECs express N-methyl-D-aspartate (NMDA) receptors (Lu et al., 2019) and GABA receptors (Won et al., 2013), suggesting

receptiveness to modulation by neuronal input. *In vitro* exposure of ECs to glutamate disrupted normal patterns of occludin expression and promoted BBB dysfunction (András et al., 2007). Additionally, intra-cortical infusion of glutamate into mice enhanced extravasation of sodium fluorescein into the brain parenchyma independent of any effects on neuronal activation (Vazana et al., 2016). These observations support the hypothesis that cytokine-mediated shifts in neuron excitability and neurotransmission may indirectly influence BBB function.

4.2. Peripheral inflammation induces neuroinflammation in the CNS

There is experimental evidence that acute OP intoxication can also stimulate peripheral inflammatory responses (Lallement et al., 1993; Collombet et al., 2005; Collombet, 2011; Meyer et al., 2023). *In vitro* exposure of human EC to paraoxon enhanced transmigration of peripheral immune cells, increased expression of cell adhesion molecules (CAMs), IL-8, and caspase-1, and decreased VE-cadherin (Israelov et al., 2020). Such observations are complemented by *in vivo* data pointing to increased CAMs and caspase-1 transcription following paraoxon exposure in the mouse (Israelov et al., 2020). In other contexts (e.g., not in models of acute OP intoxication), peripheral inflammation enhanced BBB permeability through increased expression of CAMs on ECs and direct dysregulation of TJ and BM proteins (Bohatschek et al., 2001; Huang et al., 2021). Conversely, a growing body of evidence indicates that BBB impairment may drive neuroinflammation. For example, protection of BBB integrity significantly attenuated neuroinflammation induced by peripheral blood components (Aragon et al., 2017), suggesting that BBB leakiness promotes and sustains inflammatory responses in the CNS. The most obvious manifestation of BBB impairment triggering neuroinflammation is the migration of peripheral immune cells into the brain as a consequence of BBB dysfunction. Peripheral monocyte infiltration was associated with BBB impairment in drug-resistant TLE patients and a TLE rat model (Broekaart et al., 2018). Numerous studies in multiple sclerosis patients and preclinical models indicated that changes in BBB architecture, such as deterioration of TJ structure and increased EC expression of CAMs, facilitate and promote peripheral immune cell infiltration, which was linked to the initiation and maintenance of neuroinflammatory responses (Larochelle et al., 2011).

Accumulation of soluble blood-borne components in the brain parenchyma not only promotes the recruitment of peripheral leukocytes but also directly stimulates inflammatory responses (Sulhan et al., 2020; Suleymanova, 2021). Observations from preclinical models of neurodegenerative disease with concurrent peripheral inflammation suggest that cytokines released following peripheral immune activation can cross a leaky BBB to promote and/or amplify neuroinflammation (Herrera et al., 2015). There is also evidence suggesting that blood-borne components can initiate neuroinflammatory responses even in the absence of peripheral immune activation. For example, serum albumin, the most abundant protein in circulation, can enter the brain parenchyma following seizures induced by acute OP intoxication (Rojas et al., 2021; Rojas et al., 2022). In isolated guinea pig brain, perfusion with albumin exacerbated astrogliosis and microgliosis induced by seizure-like events (Vila Verde et al., 2021). Other data suggest that albumin itself may promote neuroinflammation independently of enhanced seizure activity. In mouse and rat preclinical models, albumin activated glial TGF- β signaling, which enhanced cytokine production (Ivens et al., 2007; Frigerio et al., 2012; Weissberg et al., 2015; Webster et al., 2017). Similar to albumin, the soluble blood-borne molecules fibrin and fibrinogen can promote immune activation through TGF- β signaling (Peng and Lv, 2022). Fibrin/fibrinogen can also form an insoluble matrix that directly activates microglia and macrophages by binding to CD11b/CD18 (Adams et al., 2007; Davalos et al., 2012). Antibody-mediated block of the fibrin/fibrinogen-CD11b/CD18 interaction attenuated pro-inflammatory cell activation and neurodegeneration in mouse models of Alzheimer's disease (AD) (Ryu et al., 2018).

Maintenance of CNS iron homeostasis is a critical function of the BBB, and disturbances in this balance that result in iron accretion are associated with neuroinflammatory responses (Ward et al., 2022). In a rat model of Parkinson's disease, BBB impairment preceded iron extravasation, which was followed by microglia activation (Olmedo-Díaz et al., 2017), defining the temporal relationship between these processes. While the mechanisms by which iron promotes neuroinflammation are unknown, CNS iron accumulation is associated with enhanced oxidative stress and pro-inflammatory glial activation (Urrutia et al., 2021; Ward et al., 2022).

In summary, based on the available clinical and preclinical data, the connection between BBB impairment and neuroinflammation appears to be a complex positive feed-forward

relationship. However, an understanding of the relative contributions of BBB impairment and neuroinflammation in epileptogenesis remains a significant data gap given extensive evidence implicating neuroinflammation itself as an epileptogenic process (Andrew and Lein, 2021). Therefore, in the context of acute OP intoxication, it is likely that both BBB impairment and neuroinflammation contribute to the development of chronic sequelae, including epilepsy/SRS.

5. Evidence that BBB impairment is involved in the pathogenesis of SRS/epilepsy

While clinical and preclinical evidence indicate that seizures of diverse etiology cause BBB impairment (see section 1), a growing body of research points to a role for BBB impairment in the development and progression of epilepsy (Oby and Janigro, 2006; Marchi et al., 2007; Friedman et al., 2009; Marchi et al., 2010; Bar-Klein et al., 2017). Traumatic brain injury (TBI) (Herman, 2002; Tomkins et al., 2008; Shlosberg et al., 2010), cerebrovascular disease (Pitkänen et al., 2016; Yang et al., 2018) and AD (Vossel et al., 2017), as well as multiple acute and long-term neuropathologies characterized by NVU dysfunction are associated with increased risk for epileptogenic disorders (Gorter et al., 2019; van Vliet and Marchi, 2022).

Preclinical studies of TBI reveal a biphasic BBB breakdown following the initial brain insult characterized by rapid onset of enhanced BBB permeability that peaks hours post-injury followed by secondary inflammation-mediated BBB disruption within days of the insult (Shlosberg et al., 2010). Clinical data confirm chronic BBB dysfunction in patients with TBI (Hay et al., 2015). The frequency, severity and persistence of BBB dysfunction is greater in patients with post-concussion syndrome and post-traumatic epilepsy (PTE) (Korn et al., 2005; Tomkins et al., 2008), suggesting a role for BBB disruption in neuronal hyperexcitability and seizures. BBB permeability following TBI is likely the result of physical tissue disruption, and secondary processes, including altered pericyte-endothelium crosstalk (Bhowmick et al., 2019), decreased TJ and adhesion protein expression (Obermeier et al., 2016), and VEGF-mediated angiogenesis (Castañeda-Cabral et al., 2020). Although the specific pathways by which TBI gives rise to seizures remain unclear, changes in BBB permeability are known to contribute to PTE development (Dadas and Janigro, 2019).

Some studies have identified cerebrovascular disease, particularly stroke (Stefanidou et al., 2022), as a risk factor for epilepsy. Indeed, post-stroke epilepsy, like PTE, is a significant

complication (Stefanidou et al., 2022), accounting for approximately 40% of new seizure onset in elderly stroke survivors (Tanaka and Ihara, 2017). Downstream consequences of BBB dysfunction following stroke include vascular edema, ion gradient disruption, and aberrant mitochondrial function, all of which are thought to promote epileptogenesis (Tanaka and Ihara, 2017).

Similarly, neurodegenerative diseases are associated with altered cerebral vasculature. Growing evidence implicates BBB breakdown in AD progression. Expression of soluble platelet derived growth factor receptor β (PDGFr- β) is elevated in patients with cognitive impairment while MMP-9 activity is increased in individuals with high genetic risk of AD (Zenaro et al., 2017; Uprety et al., 2021). Increased levels of soluble PDGFr- β , an indicator of pericyte damage, were correlated with MRI findings of BBB impairment, while MMP-9 has been implicated in the pathogenesis of BBB breakdown through dysregulation of TJs (Lischper et al., 2010; Rempe et al., 2018; Uprety et al., 2021). While no direct connection has yet been established between neurodegeneration-mediated BBB dysfunction and seizure generation, loss of BBB integrity is a well-documented characteristic of the aging brain (Senatorov et al., 2019).

5.1. Pathways associated with SRS/epilepsy are altered during BBB disruption

Ultimately, despite mechanistic and etiological differences between TBI, PTE, and AD, all are linked to BBB disruption and increased seizure risk. While a causal relationship between BBB dysfunction and SRS development has yet to be determined in these neurological disorders, these observations support the hypothesis that BBB impairment may influence epileptogenic processes, and have led to the identification of pro-epileptic signaling pathways triggered by BBB impairment. The most well-defined is the leakage of seizure-promoting serum molecules into the brain parenchyma, including albumin, potassium, and glutamate (Marchi et al., 2012). Several studies report serum albumin leakage and accumulation in the cortex of both animals and humans following BBB breakdown (Ivens et al., 2007; van Vliet et al., 2007; Marchi et al., 2010; Frigerio et al., 2012). Focal extravasation of albumin promotes hyperexcitability through binding to TGF- β receptors to activate TGF- β signaling pathways (Ivens et al., 2007; Heinemann et al., 2012). Additional serum proteins, like fibrinogen, are similarly known to activate TGF- β signaling (Peng and Lv, 2022).

TGF- β signaling results in phosphorylation and nuclear translocation of the Smad family of transcription factors that regulate expression of TGF- β -responsive genes. Consequences of TGF- β signaling thought to increase seizure susceptibility and neuronal excitability include downregulation of inward-rectifying potassium channels (Kir4.1) and aquaporin 4, upregulation of IL-1 β and other pro-inflammatory cytokines, impaired astrocytic glutamate metabolism, and excitatory synaptogenesis (Heinemann et al., 2012; Swissa et al., 2019). There is evidence that while albumin-mediated TGF- β activation enhances excitability, this alone is insufficient to induce epileptogenesis. For example, Frigerio et al. (2012) showed that intracerebral application of exogenous albumin increased seizure susceptibility but did not promote development of SRS. Consistent with this, in patients with cerebral vascular malformation, BBB dysfunction and consequent albumin accumulation in the brain increased susceptibility to seizures (Raabe et al., 2012). Thus, it seems likely that TGF- β signaling induced by blood-borne molecules synergizes with additional pathways to promote epileptogenic processes.

Glutamate dyshomeostasis is another process contributing to seizures and SRS development. It is proposed that the functionally intact BBB maintains physiologic glutamate balance between the brain milieu and serum, preventing glutamate extravasation (Gruenbaum et al., 2022). As a result, a loss of BBB function contributes to increased extracellular glutamate levels in the brain. While glutamate in ECs promotes BBB permeability via NMDA-mediated increases in intracellular calcium and subsequent oxidative stress (Sharp et al., 2005; Swissa et al., 2019; Deshpande and DeLorenzo, 2020), calcium accumulation is also observed in neurons, contributing to a lower seizure threshold and greater susceptibility to epileptic seizures (Silverman et al., 2002). Contributing to the cyclical events of BBB impairment leading to seizures and vice-versa, calpains, a family of calcium-dependent proteases, are activated by elevated intracellular calcium (Suzuki and Sorimachi, 1998), resulting in cleavage of intracellular and extracellular substrates, and higher BBB permeability. In preclinical studies, BBB dysfunction is prevented when calpain inhibitors are administered to rodents (Alluri et al., 2016; Tao et al., 2017). Although calpain directly compromises BBB function, it also indirectly enhances BBB permeability by modulating neurodegenerative and neuroinflammatory processes (Camins et al., 2006; Alluri et al., 2016; Metwally et al., 2021). Collectively, these data suggest a

potential for calpain inhibitors to stabilize the BBB and prevent downstream pathological consequences of BBB dysfunction.

Chronic disruption of TJ proteins is another key driver of epilepsy associated with BBB impairment. Claudin family proteins, specifically claudin-5, are unique structural components of EC TJs and critically important in regulating BBB permeability (Hashimoto et al., 2023). A mutation in claudin-5 in human patients and transgenic zebrafish expressing the same mutation caused BBB impairment and was associated with epileptogenesis (Deshwar et al., 2022). Additionally, Greene et al. (2022) demonstrated increased kainic acid seizure-induced sodium fluorescein extravasation in claudin-5 heterozygous mice (CLDN^{-/-}/CLDN^{+/+}) compared to those with intact claudin-5 expression. They also demonstrated that knockdown of claudin-5 contributed to the development of SRS and enhanced EC activation and migration, and that BBB leakage and seizure activity were attenuated in a mouse model of kainic acid epilepsy treated with the claudin-5 regulator, RepSox. These data suggest that claudin-5 is vital for normal BBB functioning and dysregulation of this critical TJ protein induces seizures/SRS via BBB breakdown. There are several mechanisms by which TJ protein expression can change, including induced MMP-2/-9 activity through diverse stimuli (Jin et al., 2010; Mehra et al., 2016). MMP-2/-9 activation decreases TJ proteins and subsequently increases BBB leakage (Ralay Ranaivo et al., 2012), resulting in persistent changes in NVU composition and, consequently, the brain milieu (Löscher and Friedman, 2020). Studies in stroke models show tissue plasminogen activator (tPA) as a key molecule inducing upregulation and activation of MMPs (Harston et al., 2010; Balami et al., 2013; Lakhan et al., 2013).

The therapeutic efficacy of stabilizing the BBB to prevent SRS development has yet to be comprehensively evaluated, but there is evidence to support this as a promising approach (Reiss et al., 2023). Treatment with small molecule inhibitors of TGF- β prior to kainic acid administration stabilized the BBB via upregulation of claudin-5 and attenuated seizure activity and neuroinflammation (Greene et al., 2022). Similarly, blocking peripheral albumin-induced TGF- β signaling with an angiotensin II type 1 receptor antagonist decreased BBB permeability and blocked development of SRS in a rat model of acute OP intoxication (Bar-Klein et al., 2014). Consistent with these observations, preclinical data suggest that TGF- β inhibition to suppress albumin-induced TGF- β signaling in astrocytes conserves claudin-5 expression and

mitigates abnormal epileptic stimulation (Hashimoto et al., 2023). Another possible approach for stabilizing the BBB is MMP-2/-9 inhibition. Treatment with IPR-179, an MMP-2/-9 inhibitor, conferred anti-seizure and anti-epileptogenic effects in a rat model of rapid kindling and a mouse model of kainic acid-induced epilepsy (Broekaart et al., 2021). VEGF-mediated microvascular alterations are also known to influence BBB integrity, suggesting VEGF modulation as another therapeutic target for restoration of BBB function and seizure prevention (van Lanen et al., 2021). Lastly, caspase-1 has emerged as a key contributor to BBB dysfunction via inflammatory processes under a variety of pathological conditions, including OP exposures (Rand and Cooper, 2021). Initial human *in vitro* and mouse *in vivo* experiments with a caspase-1 selective inhibitor, VX-765, preserved BBB function via inhibition of the caspase-1-mediated canonical inflammasome pathway and coincident mitigation of peripheral immune transmigration (Israelov et al., 2020)

While data support a causal relationship between BBB dysfunction and epilepsy, drawing a definitive conclusion is stymied by the fact that loss of BBB integrity occurs in unison with neuroinflammation, oxidative stress and other factors known to influence epileptogenesis. Currently, there is a lack of therapeutics that specifically target BBB-associated functions, rather, stabilization of the BBB is often the consequence of therapeutic effects on other pathologic processes. Regardless, drugs that stabilize the BBB are promising candidates to avert development of SRS, not only in the case of acute OP intoxication, but also for diverse disease states characterized by BBB dysfunction.

6. Key data gaps and conclusions

BBB dysfunction is a pathological process associated with numerous neurological disorders, suggesting its potential as a novel target to mitigate adverse neurological outcomes (Shlosberg et al., 2010; Thal and Neuhaus, 2014; Gorter et al., 2019; Greene et al., 2022). However, research efforts to understand the pathophysiology of BBB impairment and its role in epileptogenesis are in their infancy and there remain more questions than answers, particularly in the context of acute OP intoxication.

Compromised BBB integrity has been documented following acute OP intoxication (Abdel-Rahman et al., 2002; Song et al., 2004; Bar-Klein et al., 2017; Rojas et al., 2021; Rojas et al.,

2022); however, higher spatial and temporal resolution studies are needed to rigorously map the spatiotemporal profile of BBB impairment. A detailed natural history of BBB impairment post-OP intoxication is necessary to understand the dynamic interplay of BBB function and other well-documented pathologic processes, such as neuroinflammation, neurodegeneration and, importantly, SRS (Collombet et al., 2008; Aroniadou-Anderjaska et al., 2016; Sisó et al., 2017; Figueiredo et al., 2018; Reddy et al., 2020). Additionally, identifying which cellular and/or molecular components of the NVU are impacted by acute OP intoxication is essential for identifying candidate targets and therapeutic windows for intervention.

Mechanistic research of OP-induced BBB dysfunction is necessary to evaluate the relationship between BBB impairment and epilepsy/SRS. We propose numerous mechanisms of OP-induced BBB impairment; however, interventional studies are critical to demonstrate causal relationships and to confirm the anti-epileptogenic potential of BBB stabilization. Only with a better understanding of the processes both up- and down-stream of BBB impairment can progress be made in identifying candidate therapeutic intervention to maintain BBB function and, ultimately, prevent long-term consequences of BBB dysregulation, including SRS. Addressing these gaps in knowledge will prove critical in guiding the direction of future research. Given the growing evidence linking BBB impairment to seizures, both as a consequence and as a cause, across multiple disease models, it seems devoting research efforts to this issue is well-justified.

Acknowledgments:

We thank Suzette Smiley-Jewell (UC Davis) for providing feedback on early versions of this manuscript.

Data availability:

This article contains no datasets generated or analyzed during the current study.

Authors' contributions:

Participated in research design: N/A (review paper)

Conducted experiments: N/A (review paper)

Contributed new reagents or analytic tools: N/A (review paper)

Performed data analysis: N/A (review paper)

Wrote or contributed to the writing of the manuscript: Bernardino, Luo, Andrew, Unkel, Gonzalez, Gelli, Lein

References:

- Abbott NJ, Patabendige AA, Dolman DE, Yusof SR and Begley DJ (2010) Structure and function of the blood–brain barrier. *Neurobiology of disease* **37**:13-25.
- Abbott NJ, Rönnbäck L and Hansson E (2006) Astrocyte–endothelial interactions at the blood–brain barrier. *Nature reviews neuroscience* **7**:41-53.
- Abdel-Rahman A, Shetty A and Abou-Donia M (2002) Acute exposure to sarin increases blood brain barrier permeability and induces neuropathological changes in the rat brain: dose–response relationships. *Neuroscience* **113**:721-741.
- Adams RA, Bauer J, Flick MJ, Sikorski SL, Nuriel T, Lassmann H, Degen JL and Akassoglou K (2007) The fibrin-derived gamma377-395 peptide inhibits microglia activation and suppresses relapsing paralysis in central nervous system autoimmune disease. *The Journal of experimental medicine* **204**:571-582.
- Alluri H, Grimsley M, Anasooya Shaji C, Varghese KP, Zhang SL, Peddaboina C, Robinson B, Beeram MR, Huang JH and Tharakan B (2016) Attenuation of Blood-Brain Barrier Breakdown and Hyperpermeability by Calpain Inhibition. *The Journal of biological chemistry* **291**:26958-26969.
- András IE, Deli MA, Veszelka S, Hayashi K, Hennig B and Toborek M (2007) The NMDA and AMPA/KA receptors are involved in glutamate-induced alterations of occludin expression and phosphorylation in brain endothelial cells. *Journal of cerebral blood flow and metabolism : official journal of the International Society of Cerebral Blood Flow and Metabolism* **27**:1431-1443.
- Andrew PM and Lein PJ (2021) Neuroinflammation as a Therapeutic Target for Mitigating the Long-Term Consequences of Acute Organophosphate Intoxication. *Frontiers in pharmacology* **12**:674325.
- Angoa-Pérez M, Kreipke CW, Thomas DM, Van Shura KE, Lyman M, McDonough JH and Kuhn DM (2010) Soman increases neuronal COX-2 levels: possible link between seizures and protracted neuronal damage. *Neurotoxicology* **31**:738-746.
- Aragon MJ, Topper L, Tyler CR, Sanchez B, Zychowski K, Young T, Herbert G, Hall P, Erdely A, Eye T, Bishop L, Saunders SA, Muldoon PP, Ottens AK and Campen MJ (2017) Serum-borne bioactivity caused by pulmonary multiwalled carbon nanotubes induces neuroinflammation via blood-brain barrier impairment. *Proceedings of the National Academy of Sciences of the United States of America* **114**:E1968-e1976.
- Arias HR, Richards VE, Ng D, Ghafoori ME, Le V and Mousa SA (2009) Role of non-neuronal nicotinic acetylcholine receptors in angiogenesis. *The international journal of biochemistry & cell biology* **41**:1441-1451.
- Armulik A, Genové G and Betsholtz C (2011) Pericytes: developmental, physiological, and pathological perspectives, problems, and promises. *Developmental cell* **21**:193-215.
- Aroniadou-Anderjaska V, Figueiredo TH, Apland JP, Prager EM, Pidoplichko VI, Miller SL and Braga MF (2016) Long-term neuropathological and behavioral impairments after exposure to nerve agents. *Annals of the New York Academy of Sciences* **1374**:17-28.
- Balami JS, Sutherland BA and Buchan AM (2013) Complications associated with recombinant tissue plasminogen activator therapy for acute ischaemic stroke. *CNS & Neurological Disorders-Drug Targets (Formerly Current Drug Targets-CNS & Neurological Disorders)* **12**:155-169.

- Bar-Klein G, Cacheaux LP, Kamintsky L, Prager O, Weissberg I, Schoknecht K, Cheng P, Kim SY, Wood L, Heinemann U, Kaufer D and Friedman A (2014) Losartan prevents acquired epilepsy via TGF- β signaling suppression. *Annals of neurology* **75**:864-875.
- Bar-Klein G, Lublinsky S, Kamintsky L, Noyman I, Veksler R, Dalipaj H, Senatorov Jr VV, Swissa E, Rosenbach D and Elazary N (2017) Imaging blood-brain barrier dysfunction as a biomarker for epileptogenesis. *Brain* **140**:1692-1705.
- Barkauskas DS, Dorand RD, Myers JT, Evans TA, Barkauskas KJ, Askew D, Purgert R and Huang AY (2015) Focal transient CNS vessel leak provides a tissue niche for sequential immune cell accumulation during the asymptomatic phase of EAE induction. *Experimental neurology* **266**:74-85.
- Begley DJ and Brightman MW (2003) Structural and functional aspects of the blood-brain barrier. *Peptide transport and delivery into the central nervous system*:39-78.
- Bhowmick S, D'Mello V, Caruso D, Wallerstein A and Abdul-Muneer PM (2019) Impairment of pericyte-endothelium crosstalk leads to blood-brain barrier dysfunction following traumatic brain injury. *Exp Neurol* **317**:260-270.
- Bird S, Traub S and Grayzel J (2014) Organophosphate and carbamate poisoning. *UpToDate* **14**:339.
- Bohatschek M, Werner A and Raivich G (2001) Systemic LPS injection leads to granulocyte influx into normal and injured brain: effects of ICAM-1 deficiency. *Exp Neurol* **172**:137-152.
- Bowyer JF, Sarkar S, Tranter KM, Hanig JP, Miller DB and O'Callaghan JP (2016) Vascular-directed responses of microglia produced by methamphetamine exposure: indirect evidence that microglia are involved in vascular repair? *Journal of neuroinflammation* **13**:1-15.
- Broekaart DW, Bertran A, Jia S, Korotkov A, Senkov O, Bongaarts A, Mills JD, Anink JJ, Seco J, Baayen JC, Idema S, Chabrol E, Becker AJ, Wadman WJ, Tarragó T, Gorter JA, Aronica E, Prades R, Dityatev A and van Vliet EA (2021) The matrix metalloproteinase inhibitor IPR-179 has antiseizure and antiepileptogenic effects. *The Journal of clinical investigation* **131**.
- Broekaart DWM, Anink JJ, Baayen JC, Idema S, de Vries HE, Aronica E, Gorter JA and van Vliet EA (2018) Activation of the innate immune system is evident throughout epileptogenesis and is associated with blood-brain barrier dysfunction and seizure progression. *Epilepsia* **59**:1931-1944.
- Bushnell PJ and Moser VC (2006) Behavioral toxicity of cholinesterase inhibitors. *Toxicology of organophosphate & carbamate compounds*:347-360.
- Cabezas R, Ávila M, Gonzalez J, El-Bachá RS, Báez E, García-Segura LM, Jurado Coronel JC, Capani F, Cardona-Gomez GP and Barreto GE (2014) Astrocytic modulation of blood brain barrier: perspectives on Parkinson's disease. *Frontiers in cellular neuroscience* **8**:211.
- Camins A, Verdaguer E, Folch J and Pallàs M (2006) Involvement of calpain activation in neurodegenerative processes. *CNS drug reviews* **12**:135-148.
- Castañeda-Cabral JL, Colunga-Durán A, Ureña-Guerrero ME, Beas-Zárate C, Nuñez-Lumbreras MLA, Orozco-Suárez S, Alonso-Vanegas M, Guevara-Guzmán R, Deli MA, Valle-Dorado MG, Sánchez-Valle V and Rocha L (2020) Expression of VEGF- and tight

- junction-related proteins in the neocortical microvasculature of patients with drug-resistant temporal lobe epilepsy. *Microvascular research* **132**:104059.
- Colás L, Domercq M, Ramos-Cabrer P, Palma A, Gómez-Vallejo V, Padro D, Plaza-García S, Pulagam KR, Higuchi M, Matute C, Llop J and Martín A (2018) In vivo imaging of A7 nicotinic receptors as a novel method to monitor neuroinflammation after cerebral ischemia. *Glia* **66**:1611-1624.
- Collombet J-M (2011) Nerve agent intoxication: recent neuropathophysiological findings and subsequent impact on medical management prospects. *Toxicology and applied pharmacology* **255**:229-241.
- Collombet J-M, Mourcin F, Grenier N, Four E, Masqueliez C, Baubichon D, Lallement G and Hérodin F (2005) Effect of soman poisoning on populations of bone marrow and peripheral blood cells in mice. *Neurotoxicology* **26**:89-98.
- Collombet J-M, Piérard C, Béracochéa D, Coubard S, Burckhart M-F, Four E, Masqueliez C, Baubichon D and Lallement G (2008) Long-term consequences of soman poisoning in mice: Part 1. Neuropathology and neuronal regeneration in the amygdala. *Behavioural brain research* **191**:88-94.
- Costa LG (2006) Current issues in organophosphate toxicology. *Clinica chimica acta* **366**:1-13.
- Coulter DA and Steinhäuser C (2015) Role of astrocytes in epilepsy. *Cold Spring Harbor perspectives in medicine* **5**:a022434.
- Cudna A, Bronisz E, Jopowicz A and Kurkowska-Jastrzębska I (2023) Changes in serum blood-brain barrier markers after bilateral tonic-clonic seizures. *Seizure* **106**:129-137.
- Dadas A and Janigro D (2019) Breakdown of blood brain barrier as a mechanism of post-traumatic epilepsy. *Neurobiol Dis* **123**:20-26.
- Dalvi S, Nguyen HH, On N, Mitchell RW, Aukema HM, Miller DW and Hatch GM (2015) Exogenous arachidonic acid mediates permeability of human brain microvessel endothelial cells through prostaglandin E2 activation of EP 3 and EP 4 receptors. *Journal of neurochemistry* **135**:867-879.
- Dash PK, Zhao J, Kobori N, Redell JB, Hylin MJ, Hood KN and Moore AN (2016) Activation of Alpha 7 Cholinergic Nicotinic Receptors Reduce Blood-Brain Barrier Permeability following Experimental Traumatic Brain Injury. *The Journal of neuroscience : the official journal of the Society for Neuroscience* **36**:2809-2818.
- Davalos D, Ryu JK, Merlini M, Baeten KM, Le Moan N, Petersen MA, Deerinck TJ, Smirnov DS, Bedard C, Hakozaki H, Gonias Murray S, Ling JB, Lassmann H, Degen JL, Ellisman MH and Akassoglou K (2012) Fibrinogen-induced perivascular microglial clustering is required for the development of axonal damage in neuroinflammation. *Nature communications* **3**:1227.
- David Y, Cacheaux LP, Ivens S, Lapilover E, Heinemann U, Kaufer D and Friedman A (2009) Astrocytic dysfunction in epileptogenesis: consequence of altered potassium and glutamate homeostasis? *The Journal of neuroscience : the official journal of the Society for Neuroscience* **29**:10588-10599.
- de Araujo Furtado M, Lumley LA, Robison C, Tong LC, Lichtenstein S and Yourick DL (2010) Spontaneous recurrent seizures after status epilepticus induced by soman in Sprague-Dawley rats. *Epilepsia* **51**:1503-1510.

- de Araujo Furtado M, Rossetti F, Chanda S and Yourick D (2012) Exposure to nerve agents: from status epilepticus to neuroinflammation, brain damage, neurogenesis and epilepsy. *Neurotoxicology* **33**:1476-1490.
- Del Zoppo G, Milner R, Mabuchi T, Hung S, Wang X and Koziol J (2006) Vascular matrix adhesion and the blood–brain barrier. *Biochemical Society Transactions* **34**:1261-1266.
- Deshpande LS and DeLorenzo RJ (2020) Novel therapeutics for treating organophosphate-induced status epilepticus co-morbidities, based on changes in calcium homeostasis. *Neurobiology of disease* **133**:104418.
- Deshwar AR, Cytrynbaum C, Murthy H, Zon J, Chitayat D, Volpatti J, Newbury-Ecob R, Ellard S, Lango Allen H, Yu EP, Noche R, Walker S, Scherer SW, Mahida S, Elitt CM, Nicolas G, Goldenberg A, Saugier-Veber P, Lecoquierre F, Dabaj I, Meddaugh H, Marble M, Keppler-Noreuil KM, Drayson L, Baranano KW, Chassevent A, Agre K, Letard P, Bilan F, Le Guyader G, Laquerriere A, Ramsey K, Henderson L, Brady L, Tarnopolsky M, Bainbridge M, Friedman J, Capri Y, Athayde L, Kok F, Gurgel-Giannetti J, Ramos LLP, Blaser S, Dowling JJ and Weksberg R (2022) Variants in CLDN5 cause a syndrome characterized by seizures, microcephaly and brain calcifications. *Brain*.
- Díaz-Castro B, Robel S and Mishra A (2023) Astrocyte Endfeet in brain function and pathology: Open questions. *Annual Review of Neuroscience* **46**.
- Dillman JF, 3rd, Phillips CS, Kniffin DM, Tompkins CP, Hamilton TA and Kan RK (2009) Gene expression profiling of rat hippocampus following exposure to the acetylcholinesterase inhibitor soman. *Chemical research in toxicology* **22**:633-638.
- DiSabato DJ, Quan N and Godbout JP (2016) Neuroinflammation: the devil is in the details. *Journal of neurochemistry* **139 Suppl 2**:136-153.
- Eddleston M, Buckley NA, Eyer P and Dawson AH (2008) Management of acute organophosphorus pesticide poisoning. *The Lancet* **371**:597-607.
- Farkhondeh T, Mehrpour O, Forouzanfar F, Roshanravan B and Samarghandian S (2020) Oxidative stress and mitochondrial dysfunction in organophosphate pesticide-induced neurotoxicity and its amelioration: a review. *Environmental science and pollution research international* **27**:24799-24814.
- Figueiredo TH, Apland JP, Braga MF and Marini AM (2018) Acute and long-term consequences of exposure to organophosphate nerve agents in humans. *Epilepsia* **59**:92-99.
- Flannery BM, Bruun DA, Rowland DJ, Banks CN, Austin AT, Kukis DL, Li Y, Ford BD, Tancredi DJ, Silverman JL, Cherry SR and Lein PJ (2016) Persistent neuroinflammation and cognitive impairment in a rat model of acute diisopropylfluorophosphate intoxication. *J Neuroinflammation* **13**:267.
- Friedman A (2011) Blood–brain barrier dysfunction, status epilepticus, seizures, and epilepsy: a puzzle of a chicken and egg? *Epilepsia* **52**:19-20.
- Friedman A, Kaufer D and Heinemann U (2009) Blood-brain barrier breakdown-inducing astrocytic transformation: novel targets for the prevention of epilepsy. *Epilepsy research* **85**:142-149.
- Frigerio F, Frasca A, Weissberg I, Parrella S, Friedman A, Vezzani A and Noé FM (2012) Long-lasting pro-ictogenic effects induced in vivo by rat brain exposure to serum albumin in the absence of concomitant pathology. *Epilepsia* **53**:1887-1897.

- Gage M, Putra M, Wachter L, Dishman K, Gard M, Gomez-Estrada C and Thippeswamy T (2021) Saracatinib, a Src tyrosine kinase inhibitor, as a disease modifier in the rat DFP model: sex differences, Neurobehavior, gliosis, neurodegeneration, and nitro-oxidative stress. *Antioxidants* **11**:61.
- Gage M, Rao NS, Samidurai M, Putra M, Vasanthi SS, Meyer C, Wang C and Thippeswamy T (2022) Soman (GD) Rat Model to Mimic Civilian Exposure to Nerve Agent: Mortality, Video-EEG Based Status Epilepticus Severity, Sex Differences, Spontaneously Recurring Seizures, and Brain Pathology. *Frontiers in Cellular Neuroscience* **15**:573.
- Glauser T, Shinnar S, Gloss D, Alldredge B, Arya R, Bainbridge J, Bare M, Bleck T, Dodson WE, Garrity L, Jagoda A, Lowenstein D, Pellock J, Riviello J, Sloan E and Treiman DM (2016) Evidence-Based Guideline: Treatment of Convulsive Status Epilepticus in Children and Adults: Report of the Guideline Committee of the American Epilepsy Society. *Epilepsy currents* **16**:48-61.
- Gorter JA, Aronica E and van Vliet EA (2019) The Roof is Leaking and a Storm is Raging: Repairing the Blood-Brain Barrier in the Fight Against Epilepsy. *Epilepsy currents* **19**:177-181.
- Greene C, Hanley N, Reschke CR, Reddy A, Mäe MA, Connolly R, Behan C, O'Keeffe E, Bolger I, Hudson N, Delaney C, Farrell MA, O'Brien DF, Cryan J, Brett FM, Beausang A, Betsholtz C, Henshall DC, Doherty CP and Campbell M (2022) Microvascular stabilization via blood-brain barrier regulation prevents seizure activity. *Nature communications* **13**:2003.
- Gruenbaum BF, Zlotnik A, Fleidervish I, Frenkel A and Boyko M (2022) Glutamate neurotoxicity and destruction of the blood–brain barrier: key pathways for the development of neuropsychiatric consequences of TBI and their potential treatment strategies. *International journal of molecular sciences* **23**:9628.
- Guignet M, Dhakal K, Flannery BM, Hobson BA, Zolkowska D, Dhir A, Bruun DA, Li S, Wahab A, Harvey DJ, Silverman JL, Rogawski MA and Lein PJ (2020) Persistent behavior deficits, neuroinflammation, and oxidative stress in a rat model of acute organophosphate intoxication. *Neurobiol Dis* **133**:104431.
- Guignet M and Lein PJ (2019) Chapter Two - Neuroinflammation in organophosphate-induced neurotoxicity, in *Advances in Neurotoxicology* (Aschner M and Costa LG eds) pp 35-79, Academic Press.
- Harston GW, Sutherland BA, Kennedy J and Buchan AM (2010) The contribution of L-arginine to the neurotoxicity of recombinant tissue plasminogen activator following cerebral ischemia: a review of rtPA neurotoxicity. *Journal of Cerebral Blood Flow & Metabolism* **30**:1804-1816.
- Hashimoto Y, Greene C, Munnich A and Campbell M (2023) The CLDN5 gene at the blood-brain barrier in health and disease. *Fluids and barriers of the CNS* **20**:22.
- Hawkins BT and Davis TP (2005) The blood-brain barrier/neurovascular unit in health and disease. *Pharmacological reviews* **57**:173-185.
- Hawkins BT, Egleton RD and Davis TP (2005) Modulation of cerebral microvascular permeability by endothelial nicotinic acetylcholine receptors. *American journal of physiology Heart and circulatory physiology* **289**:H212-219.

- Hay JR, Johnson VE, Young AM, Smith DH and Stewart W (2015) Blood-Brain Barrier Disruption Is an Early Event That May Persist for Many Years After Traumatic Brain Injury in Humans. *Journal of neuropathology and experimental neurology* **74**:1147-1157.
- Heinemann U, Kaufer D and Friedman A (2012) Blood-brain barrier dysfunction, TGF β signaling, and astrocyte dysfunction in epilepsy. *Glia* **60**:1251-1257.
- Heithoff BP, George KK, Phares AN, Zuidhoek IA, Munoz-Ballester C and Robel S (2021) Astrocytes are necessary for blood-brain barrier maintenance in the adult mouse brain. *Glia* **69**:436-472.
- Hellström M, Gerhardt H, Kalén M, Li X, Eriksson U, Wolburg H and Betsholtz C (2001) Lack of pericytes leads to endothelial hyperplasia and abnormal vascular morphogenesis. *The Journal of cell biology* **153**:543-554.
- Herman ST (2002) Epilepsy after brain insult: targeting epileptogenesis. *Neurology* **59**:S21-26.
- Herrera AJ, Espinosa-Oliva AM, Oliva-Martin MJ, Carrillo-Jimenez A, Venero JL and de Pablos RM (2015) Collateral Damage: Contribution of Peripheral Inflammation to Neurodegenerative Diseases. *Current topics in medicinal chemistry* **15**:2193-2210.
- Hobson BA, Rowland DJ, Sisó S, Guignet MA, Harmany ZT, Bandara SB, Saito N, Harvey DJ, Bruun DA and Garbow JR (2019) TSPO PET using [18F] PBR111 reveals persistent neuroinflammation following acute diisopropylfluorophosphate intoxication in the rat. *Toxicological Sciences* **170**:330-344.
- Huang X, Hussain B and Chang J (2021) Peripheral inflammation and blood-brain barrier disruption: effects and mechanisms. *CNS neuroscience & therapeutics* **27**:36-47.
- Hutamekalin P, Farkas AE, Orbók A, Wilhelm I, Nagyoszi P, Veszélka S, Deli MA, Buzás K, Hunyadi-Gulyás E, Medzihradzky KF, Meksuriyen D and Krizbai IA (2008) Effect of nicotine and polyaromatic hydrocarbons on cerebral endothelial cells. *Cell biology international* **32**:198-209.
- Israelov H, Ravid O, Atrakchi D, Rand D, Elhaik S, Bresler Y, Twitto-Greenberg R, Omesi L, Liraz-Zaltsman S and Gosselet F (2020) Caspase-1 has a critical role in blood-brain barrier injury and its inhibition contributes to multifaceted repair. *Journal of Neuroinflammation* **17**:1-19.
- Ivens S, Kaufer D, Flores LP, Bechmann I, Zumsteg D, Tomkins O, Seiffert E, Heinemann U and Friedman A (2007) TGF-beta receptor-mediated albumin uptake into astrocytes is involved in neocortical epileptogenesis. *Brain* **130**:535-547.
- Jett DA, Sibrizzi CA, Blain RB, Hartman PA, Lein PJ, Taylor KW and Rooney AA (2020) A national toxicology program systematic review of the evidence for long-term effects after acute exposure to sarin nerve agent. *Critical reviews in toxicology* **50**:474-490.
- Jin R, Yang G and Li G (2010) Molecular insights and therapeutic targets for blood-brain barrier disruption in ischemic stroke: critical role of matrix metalloproteinases and tissue-type plasminogen activator. *Neurobiol Dis* **38**:376-385.
- Johnson EA, Dao TL, Guignet MA, Geddes CE, Koemeter-Cox AI and Kan RK (2011) Increased expression of the chemokines CXCL1 and MIP-1 α by resident brain cells precedes neutrophil infiltration in the brain following prolonged soman-induced status epilepticus in rats. *J Neuroinflammation* **8**:41.

- Johnson EA and Kan RK (2010) The acute phase response and soman-induced status epilepticus: temporal, regional and cellular changes in rat brain cytokine concentrations. *J Neuroinflammation* **7**:40.
- Kanthsamy A, Jin H, Charli A, Vellareddy A and Kanthsamy A (2019) Environmental neurotoxicant-induced dopaminergic neurodegeneration: a potential link to impaired neuroinflammatory mechanisms. *Pharmacology & therapeutics* **197**:61-82.
- Kaplan L, Chow BW and Gu C (2020) Neuronal regulation of the blood–brain barrier and neurovascular coupling. *Nature Reviews Neuroscience* **21**:416-432.
- Kim GS, Yang L, Zhang G, Zhao H, Selim M, McCullough LD, Kluk MJ and Sanchez T (2015) Critical role of sphingosine-1-phosphate receptor-2 in the disruption of cerebrovascular integrity in experimental stroke. *Nature communications* **6**:7893.
- Klement W, Garbelli R, Zub E, Rossini L, Tassi L, Girard B, Blaquiere M, Bertaso F, Perroy J, de Bock F and Marchi N (2018) Seizure progression and inflammatory mediators promote pericytosis and pericyte-microglia clustering at the cerebrovasculature. *Neurobiol Dis* **113**:70-81.
- Knopp RC, Banks WA and Erickson MA (2022) Physical associations of microglia and the vascular blood-brain barrier and their importance in development, health, and disease. *Current Opinion in Neurobiology* **77**:102648.
- Korn A, Golan H, Melamed I, Pascual-Marqui R and Friedman A (2005) Focal cortical dysfunction and blood-brain barrier disruption in patients with Postconcussion syndrome. *Journal of clinical neurophysiology : official publication of the American Electroencephalographic Society* **22**:1-9.
- Kraft AD and Harry GJ (2011) Features of microglia and neuroinflammation relevant to environmental exposure and neurotoxicity. *International journal of environmental research and public health* **8**:2980-3018.
- Lakhan SE, Kirchgessner A, Tepper D and Leonard A (2013) Matrix metalloproteinases and blood-brain barrier disruption in acute ischemic stroke. *Frontiers in neurology* **4**:32.
- Lallement G, Delamanche IS, Pernet-Marino I, Baubichon D, Denoyer M, Carpentier P and Blanchet G (1993) Neuroprotective activity of glutamate receptor antagonists against soman-induced hippocampal damage: quantification with an ω 3 site ligand. *Brain research* **618**:227-237.
- Larochelle C, Alvarez JI and Prat A (2011) How do immune cells overcome the blood-brain barrier in multiple sclerosis? *FEBS letters* **585**:3770-3780.
- Li H, Liu X, Wang R, Lu A, Ma Z, Wu S, Lu H, Du Y, Deng K, Wang L and Yuan F (2023) Blood-brain barrier damage and new onset refractory status epilepticus: An exploratory study using dynamic contrast-enhanced magnetic resonance imaging. *Epilepsia*.
- Lindahl P, Johansson BR, Leveen P and Betsholtz C (1997) Pericyte loss and microaneurysm formation in PDGF-B-deficient mice. *Science* **277**:242-245.
- Lischper M, Beuck S, Thanabalasundaram G, Pieper C and Galla HJ (2010) Metalloproteinase mediated occludin cleavage in the cerebral microcapillary endothelium under pathological conditions. *Brain research* **1326**:114-127.
- Löscher W and Friedman A (2020) Structural, molecular, and functional alterations of the blood-brain barrier during epileptogenesis and epilepsy: a cause, consequence, or both? *International journal of molecular sciences* **21**:591.

- Lu L, Hogan-Cann AD, Globa AK, Lu P, Nagy JI, Bamji SX and Anderson CM (2019) Astrocytes drive cortical vasodilatory signaling by activating endothelial NMDA receptors. *Journal of cerebral blood flow and metabolism : official journal of the International Society of Cerebral Blood Flow and Metabolism* **39**:481-496.
- Lukaszewicz-Hussain A (2010) Role of oxidative stress in organophosphate insecticide toxicity – Short review. *Pesticide Biochemistry and Physiology* **98**:145-150.
- Lyman M, Lloyd DG, Ji X, Vizcaychipi MP and Ma D (2014) Neuroinflammation: the role and consequences. *Neuroscience research* **79**:1-12.
- Machida T, Dohgu S, Takata F, Matsumoto J, Kimura I, Koga M, Nakamoto K, Yamauchi A and Kataoka Y (2017) Role of thrombin-PAR1-PKC θ/δ axis in brain pericytes in thrombin-induced MMP-9 production and blood-brain barrier dysfunction in vitro. *Neuroscience* **350**:146-157.
- Marchi N, Angelov L, Masaryk T, Fazio V, Granata T, Hernandez N, Hallene K, Diglaw T, Franic L, Najm I and Janigro D (2007) Seizure-promoting effect of blood-brain barrier disruption. *Epilepsia* **48**:732-742.
- Marchi N, Granata T, Ghosh C and Janigro D (2012) Blood-brain barrier dysfunction and epilepsy: pathophysiologic role and therapeutic approaches. *Epilepsia* **53**:1877-1886.
- Marchi N and Lerner-Natoli M (2013) Cerebrovascular remodeling and epilepsy. *The Neuroscientist : a review journal bringing neurobiology, neurology and psychiatry* **19**:304-312.
- Marchi N, Teng Q, Ghosh C, Fan Q, Nguyen MT, Desai NK, Bawa H, Rasmussen P, Masaryk TK and Janigro D (2010) Blood-brain barrier damage, but not parenchymal white blood cells, is a hallmark of seizure activity. *Brain research* **1353**:176-186.
- Martín A, Domercq M and Matute C (2018) Inflammation in stroke: the role of cholinergic, purinergic and glutamatergic signaling. *Therapeutic advances in neurological disorders* **11**:1756286418774267.
- Matsumoto J, Dohgu S, Takata F, Nishioku T, Sumi N, Machida T, Takahashi H, Yamauchi A and Kataoka Y (2012) Lipopolysaccharide-activated microglia lower P-glycoprotein function in brain microvascular endothelial cells. *Neuroscience letters* **524**:45-48.
- Matsumoto J, Takata F, Machida T, Takahashi H, Soejima Y, Funakoshi M, Futagami K, Yamauchi A, Dohgu S and Kataoka Y (2014) Tumor necrosis factor- α -stimulated brain pericytes possess a unique cytokine and chemokine release profile and enhance microglial activation. *Neuroscience letters* **578**:133-138.
- Maupu C, Enderlin J, Iger A, Oger M, Auvin S, Hassan-Abdi R, Soussi-Yanicostas N, Brazzolotto X, Nachon F and Dal Bo G (2021) Diisopropylfluorophosphate-induced status epilepticus drives complex glial cell phenotypes in adult male mice. *Neurobiology of Disease* **152**:105276.
- Mehra A, Ali C, Parcq J, Vivien D and Docagne F (2016) The plasminogen activation system in neuroinflammation. *Biochimica et biophysica acta* **1862**:395-402.
- Mendes NF, Pansani AP, Carmanhães ERF, Tange P, Meireles JV, Ochikubo M, Chagas JR, da Silva AV, Monteiro de Castro G and Le Sueur-Maluf L (2019) The Blood-Brain Barrier Breakdown During Acute Phase of the Pilocarpine Model of Epilepsy Is Dynamic and Time-Dependent. *Frontiers in neurology* **10**:382.

- Menezes MJ, McClenahan FK, Leiton CV, Aranmolate A, Shan X and Colognato H (2014) The extracellular matrix protein laminin $\alpha 2$ regulates the maturation and function of the blood–brain barrier. *Journal of Neuroscience* **34**:15260-15280.
- Metwally E, Zhao G and Zhang YQ (2021) The calcium-dependent protease calpain in neuronal remodeling and neurodegeneration. *Trends Neurosci* **44**:741-752.
- Mew EJ, Padmanathan P, Konradsen F, Eddleston M, Chang SS, Phillips MR and Gunnell D (2017) The global burden of fatal self-poisoning with pesticides 2006-15: Systematic review. *Journal of affective disorders* **219**:93-104.
- Meyer C, Rao NS, Vasanthi SS, Pereira B, Gage M, Putra M, Holtkamp C, Huss J and Thippeswamy T (2023) Peripheral and central effects of NADPH oxidase inhibitor, mitoapocynin, in a rat model of diisopropylfluorophosphate (DFP) toxicity. *Frontiers in Cellular Neuroscience*.
- Middlemore-Risher ML, Adam BL, Lambert NA and Terry AV, Jr. (2011) Effects of chlorpyrifos and chlorpyrifos-oxon on the dynamics and movement of mitochondria in rat cortical neurons. *The Journal of pharmacology and experimental therapeutics* **339**:341-349.
- Mohan S, Koller EJ, Fazal JA, De Oliveria G, Pawlowicz AI and Doré S (2018) Genetic deletion of PGF2 α -FP receptor exacerbates brain injury following experimental intracerebral hemorrhage. *Frontiers in neuroscience* **12**:556.
- Nag S (2011) Morphology and properties of brain endothelial cells. *The Blood-Brain and Other Neural Barriers: Reviews and Protocols*:3-47.
- Nakagawa S and Aruga J (2020) Sphingosine 1-Phosphate Signaling Is Involved in Impaired Blood–Brain Barrier Function in Ischemia–Reperfusion Injury. *Molecular Neurobiology* **57**:1594-1606.
- Nation DA, Sweeney MD, Montagne A, Sagare AP, D'Orazio LM, Pachicano M, Seppehrband F, Nelson AR, Buennagel DP, Harrington MG, Benzinger TLS, Fagan AM, Ringman JM, Schneider LS, Morris JC, Chui HC, Law M, Toga AW and Zlokovic BV (2019) Blood-brain barrier breakdown is an early biomarker of human cognitive dysfunction. *Nat Med* **25**:270-276.
- Nishioku T, Matsumoto J, Dohgu S, Sumi N, Miyao K, Takata F, Shuto H, Yamauchi A and Kataoka Y (2010) Tumor necrosis factor- α mediates the blood-brain barrier dysfunction induced by activated microglia in mouse brain microvascular endothelial cells. *Journal of pharmacological sciences* **112**:251-254.
- Obermeier B, Daneman R and Ransohoff RM (2013) Development, maintenance and disruption of the blood-brain barrier. *Nature medicine* **19**:1584-1596.
- Obermeier B, Verma A and Ransohoff RM (2016) The blood–brain barrier. *Handbook of clinical neurology* **133**:39-59.
- Oby E and Janigro D (2006) The blood-brain barrier and epilepsy. *Epilepsia* **47**:1761-1774.
- Olmedo-Díaz S, Estévez-Silva H, Orádd G, Af Bjerken S, Marcellino D and Virel A (2017) An altered blood-brain barrier contributes to brain iron accumulation and neuroinflammation in the 6-OHDA rat model of Parkinson's disease. *Neuroscience* **362**:141-151.
- Patel M, Day BJ, Crapo JD, Fridovich I and McNamara JO (1996) Requirement for superoxide in excitotoxic cell death. *Neuron* **16**:345-355.

- Pavlov VA, Wang H, Czura CJ, Friedman SG and Tracey KJ (2003) The cholinergic anti-inflammatory pathway: a missing link in neuroimmunomodulation. *Molecular medicine (Cambridge, Mass)* **9**:125-134.
- Pearson JN and Patel M (2016) The role of oxidative stress in organophosphate and nerve agent toxicity. *Annals of the New York Academy of Sciences* **1378**:17-24.
- Peng S and Lv K (2022) The role of fibrinogen in traumatic brain injury: from molecular pathological mechanisms to clinical management. *European journal of trauma and emergency surgery : official publication of the European Trauma Society*.
- Pieper C, Pieloch P and Galla H-J (2013) Pericytes support neutrophil transmigration via interleukin-8 across a porcine co-culture model of the blood–brain barrier. *Brain research* **1524**:1-11.
- Pitkänen A, Roivainen R and Lukasiuk K (2016) Development of epilepsy after ischaemic stroke. *The Lancet Neurology* **15**:185-197.
- Pope CN (1999) Organophosphorus pesticides: do they all have the same mechanism of toxicity? *Journal of Toxicology and Environmental Health Part B: Critical Reviews* **2**:161-181.
- Prager O, Kamintsky L, Hasam-Henderson LA, Schoknecht K, Wuntke V, Papageorgiou I, Swolinsky J, Muoio V, Bar-Klein G, Vazana U, Heinemann U, Friedman A and Kovács R (2019) Seizure-induced microvascular injury is associated with impaired neurovascular coupling and blood-brain barrier dysfunction. *Epilepsia* **60**:322-336.
- Profaci CP, Munji RN, Pulido RS and Daneman R (2020) The blood–brain barrier in health and disease: Important unanswered questions. *Journal of Experimental Medicine* **217**.
- Puttachary S, Sharma S, Stark S and Thippeswamy T (2015) Seizure-induced oxidative stress in temporal lobe epilepsy. *BioMed research international* **2015**.
- Raabe A, Schmitz AK, Pernhorst K, Grote A, Von Der Brelie C, Urbach H, Friedman A, Becker AJ, Elger CE and Niehusmann P (2012) Cliniconeuropathologic correlations show astroglial albumin storage as a common factor in epileptogenic vascular lesions. *Epilepsia* **53**:539-548.
- Ralay Ranaivo H, Hodge JN, Choi N and Wainwright MS (2012) Albumin induces upregulation of matrix metalloproteinase-9 in astrocytes via MAPK and reactive oxygen species-dependent pathways. *J Neuroinflammation* **9**:68.
- Rand D and Cooper I (2021) Caspase-1: an important player and possible target for repair of the blood-brain barrier underlying neurodegeneration. *Neural Regeneration Research* **16**:2390.
- Ransohoff RM and Engelhardt B (2012) The anatomical and cellular basis of immune surveillance in the central nervous system. *Nature Reviews Immunology* **12**:623-635.
- Reddy SD, Wu X, Kuruba R, Sridhar V and Reddy DS (2020) Magnetic resonance imaging analysis of long-term neuropathology after exposure to the nerve agent soman: correlation with histopathology and neurological dysfunction. *Annals of the New York Academy of Sciences* **1480**:116-135.
- Reiss Y, Bauer S, David B, Devraj K, Fidan E, Hattingen E, Liebner S, Melzer N, Meuth SG, Rosenow F, Rüber T, Willems LM and Plate KH (2023) The neurovasculature as a target in temporal lobe epilepsy. *Brain pathology (Zurich, Switzerland)* **33**:e13147.
- Rempe RG, Hartz AMS, Soldner ELB, Sokola BS, Alluri SR, Abner EL, Kryscio RJ, Pekcec A, Schlichtiger J and Bauer B (2018) Matrix Metalloproteinase-Mediated Blood-Brain

- Barrier Dysfunction in Epilepsy. *The Journal of neuroscience : the official journal of the Society for Neuroscience* **38**:4301-4315.
- Rigau V, Morin M, Rousset MC, de Bock F, Lebrun A, Coubes P, Picot MC, Baldy-Moulinier M, Bockaert J, Crespel A and Lerner-Natoli M (2007) Angiogenesis is associated with blood-brain barrier permeability in temporal lobe epilepsy. *Brain* **130**:1942-1956.
- Rojas A, Abreu-Melon J, Wang S and Dingledine R (2022) Time-dependent neuropathology in rats following organophosphate-induced status epilepticus. *Neurotoxicology* **91**:45-59.
- Rojas A, Chen D, Ganesh T, Varvel NH and Dingledine R (2019) The COX-2/prostanoid signaling cascades in seizure disorders. *Expert opinion on therapeutic targets* **23**:1-13.
- Rojas A, McCarren HS, Wang J, Wang W, Abreu-Melon J, Wang S, McDonough JH and Dingledine R (2021) Comparison of neuropathology in rats following status epilepticus induced by diisopropylfluorophosphate and soman. *Neurotoxicology* **83**:14-27.
- Ronaldson PT and Davis TP (2020) Regulation of blood–brain barrier integrity by microglia in health and disease: a therapeutic opportunity. *Journal of Cerebral Blood Flow & Metabolism* **40**:S6-S24.
- Ryu JK, Rafalski VA, Meyer-Franke A, Adams RA, Poda SB, Rios Coronado PE, Pedersen L, Menon V, Baeten KM, Sikorski SL, Bedard C, Hanspers K, Bardehle S, Mendiola AS, Davalos D, Machado MR, Chan JP, Plastira I, Petersen MA, Pfaff SJ, Ang KK, Hallenbeck KK, Syme C, Hakozaki H, Ellisman MH, Swanson RA, Zamvil SS, Arkin MR, Zorn SH, Pico AR, Mucke L, Freedman SB, Stavenhagen JB, Nelson RB and Akassoglou K (2018) Fibrin-targeting immunotherapy protects against neuroinflammation and neurodegeneration. *Nature immunology* **19**:1212-1223.
- Saunders NR, Ek CJ, Habgood MD and Dziegielewska KM (2008) Barriers in the brain: a renaissance? *Trends in neurosciences* **31**:279-286.
- Schreiner B, Romanelli E, Liberski P, Ingold-Heppner B, Sobottka-Brillout B, Hartwig T, Chandrasekar V, Johannssen H, Zeilhofer HU and Aguzzi A (2015) Astrocyte depletion impairs redox homeostasis and triggers neuronal loss in the adult CNS. *Cell reports* **12**:1377-1384.
- Segura-Collar B, Mata-Martínez P, Hernández-Laín A, Sanchez-Gomez P and Gargini R (2022) Blood-brain barrier disruption: a common driver of central nervous system diseases. *The Neuroscientist* **28**:222-237.
- Senatorov VV, Jr., Friedman AR, Milikovsky DZ, Ofer J, Saar-Ashkenazy R, Charbash A, Jahan N, Chin G, Mihaly E, Lin JM, Ramsay HJ, Moghbel A, Preininger MK, Eddings CR, Harrison HV, Patel R, Shen Y, Ghanim H, Sheng H, Veksler R, Sudmant PH, Becker A, Hart B, Rogawski MA, Dillin A, Friedman A and Kaufer D (2019) Blood-brain barrier dysfunction in aging induces hyperactivation of TGF β signaling and chronic yet reversible neural dysfunction. *Science translational medicine* **11**.
- Shabab T, Khanabdali R, Moghadamtousi SZ, Kadir HA and Mohan G (2017) Neuroinflammation pathways: a general review. *The International journal of neuroscience* **127**:624-633.
- Shan Y, Tan S, Lin Y, Liao S, Zhang B, Chen X, Wang J, Deng Z, Zeng Q, Zhang L, Wang Y, Hu X, Qiu W, Peng L and Lu Z (2019) The glucagon-like peptide-1 receptor agonist reduces inflammation and blood-brain barrier breakdown in an astrocyte-dependent manner in experimental stroke. *J Neuroinflammation* **16**:242.

- Sharp CD, Houghton J, Elrod JW, Warren A, Jackson THt, Jawahar A, Nanda A, Minagar A and Alexander JS (2005) N-methyl-D-aspartate receptor activation in human cerebral endothelium promotes intracellular oxidant stress. *American journal of physiology Heart and circulatory physiology* **288**:H1893-1899.
- Shlosberg D, Benifla M, Kaufer D and Friedman A (2010) Blood-brain barrier breakdown as a therapeutic target in traumatic brain injury. *Nature reviews Neurology* **6**:393-403.
- Shrot S, Ramaty E, Biala Y, Bar-Klein G, Daninos M, Kamintsky L, Makarovsky I, Statlender L, Rosman Y, Krivoy A, Lavon O, Kassirer M, Friedman A and Yaari Y (2014) Prevention of organophosphate-induced chronic epilepsy by early benzodiazepine treatment. *Toxicology* **323**:19-25.
- Silverman IE, Restrepo L and Mathews GC (2002) Poststroke seizures. *Archives of neurology* **59**:195-201.
- Sisó S, Hobson BA, Harvey DJ, Bruun DA, Rowland DJ, Garbow JR and Lein PJ (2017) Editor's Highlight: Spatiotemporal Progression and Remission of Lesions in the Rat Brain Following Acute Intoxication With Diisopropylfluorophosphate. *Toxicological sciences : an official journal of the Society of Toxicology* **157**:330-341.
- Song K, Li Y, Zhang H, An N, Wei Y, Wang L, Tian C, Yuan M, Sun Y, Xing Y and Gao Y (2020) Oxidative Stress-Mediated Blood-Brain Barrier (BBB) Disruption in Neurological Diseases. *Oxidative Medicine and Cellular Longevity* **2020**:4356386.
- Song X, Pope C, Murthy R, Shaikh J, Lal B and Bressler JP (2004) Interactive effects of paraoxon and pyridostigmine on blood-brain barrier integrity and cholinergic toxicity. *Toxicological Sciences* **78**:241-247.
- Sorokin L (2010) The impact of the extracellular matrix on inflammation. *Nature Reviews Immunology* **10**:712-723.
- Stefanidou M, Himali JJ, Devinsky O, Romero JR, Ikram MA, Beiser AS, Seshadri S and Friedman D (2022) Vascular risk factors as predictors of epilepsy in older age: The Framingham Heart Study. *Epilepsia* **63**:237-243.
- Suleymanova EM (2021) Behavioral comorbidities of epilepsy and neuroinflammation: Evidence from experimental and clinical studies. *Epilepsy & behavior : E&B* **117**:107869.
- Sulhan S, Lyon KA, Shapiro LA and Huang JH (2020) Neuroinflammation and blood-brain barrier disruption following traumatic brain injury: Pathophysiology and potential therapeutic targets. *Journal of neuroscience research* **98**:19-28.
- Sumi N, Nishioku T, Takata F, Matsumoto J, Watanabe T, Shuto H, Yamauchi A, Dohgu S and Kataoka Y (2010) Lipopolysaccharide-activated microglia induce dysfunction of the blood-brain barrier in rat microvascular endothelial cells co-cultured with microglia. *Cellular and molecular neurobiology* **30**:247-253.
- Suzuki K and Sorimachi H (1998) A novel aspect of calpain activation. *FEBS letters* **433**:1-4.
- Swissa E, Serlin Y, Vazana U, Prager O and Friedman A (2019) Blood-brain barrier dysfunction in status epilepticus: Mechanisms and role in epileptogenesis. *Epilepsy & behavior : E&B* **101**:106285.
- Takata F, Dohgu S, Matsumoto J, Takahashi H, Machida T, Wakigawa T, Harada E, Miyaji H, Koga M, Nishioku T, Yamauchi A and Kataoka Y (2011) Brain pericytes among cells constituting the blood-brain barrier are highly sensitive to tumor necrosis factor- α , releasing matrix metalloproteinase-9 and migrating in vitro. *J Neuroinflammation* **8**:106.

- Takata F, Nakagawa S, Matsumoto J and Dohgu S (2021) Blood-Brain Barrier Dysfunction Amplifies the Development of Neuroinflammation: Understanding of Cellular Events in Brain Microvascular Endothelial Cells for Prevention and Treatment of BBB Dysfunction. *Front Cell Neurosci* **15**:661838.
- Takata F, Sumi N, Nishioku T, Harada E, Wakigawa T, Shuto H, Yamauchi A and Kataoka Y (2008) Oncostatin M induces functional and structural impairment of blood–brain barriers comprised of rat brain capillary endothelial cells. *Neuroscience letters* **441**:163-166.
- Talabani JM, Ali AI, Kadir AM, Rashid R, Samin F, Greenwood D and Hay A (2018) Long-term health effects of chemical warfare agents on children following a single heavy exposure. *Human & experimental toxicology* **37**:836-847.
- Tanaka T and Ihara M (2017) Post-stroke epilepsy. *Neurochemistry international* **107**:219-228.
- Tao-Cheng J-H, Nagy Z and Brightman MW (1987) Tight junctions of brain endothelium in vitro are enhanced by astroglia. *Journal of Neuroscience* **7**:3293-3299.
- Tao XG, Shi JH, Hao SY, Chen XT and Liu BY (2017) Protective Effects of Calpain Inhibition on Neurovascular Unit Injury through Downregulating Nuclear Factor- κ B-related Inflammation during Traumatic Brain Injury in Mice. *Chinese medical journal* **130**:187-198.
- Tenreiro MM, Ferreira R, Bernardino L and Brito MA (2016) Cellular response of the blood-brain barrier to injury: Potential biomarkers and therapeutic targets for brain regeneration. *Neurobiol Dis* **91**:262-273.
- Thal SC and Neuhaus W (2014) The blood-brain barrier as a target in traumatic brain injury treatment. *Archives of medical research* **45**:698-710.
- Thurgur H and Pinteaux E (2019) Microglia in the neurovascular unit: blood–brain barrier–microglia interactions after central nervous system disorders. *Neuroscience* **405**:55-67.
- Todorovic MS, Cowan ML, Balint CA, Sun C and Kapur J (2012) Characterization of status epilepticus induced by two organophosphates in rats. *Epilepsy research* **101**:268-276.
- Tomkins O, Shelef I, Kaizerman I, Eliushin A, Afawi Z, Misk A, Gidon M, Cohen A, Zumsteg D and Friedman A (2008) Blood-brain barrier disruption in post-traumatic epilepsy. *Journal of neurology, neurosurgery, and psychiatry* **79**:774-777.
- Tsai Y-H and Lein PJ (2021) Mechanisms of organophosphate neurotoxicity. *Current opinion in toxicology* **26**:49-60.
- Uchida S and Hotta H (2009) Cerebral cortical vasodilatation mediated by nicotinic cholinergic receptors: effects of old age and of chronic nicotine exposure. *Biological & pharmaceutical bulletin* **32**:341-344.
- Uprety A, Kang Y and Kim SY (2021) Blood-brain barrier dysfunction as a potential therapeutic target for neurodegenerative disorders. *Archives of pharmacal research* **44**:487-498.
- Urrutia PJ, Bórquez DA and Núñez MT (2021) Inflaming the Brain with Iron. *Antioxidants (Basel, Switzerland)* **10**.
- van Lanen RH, Melchers S, Hoogland G, Schijns OE, Zandvoort MAV, Haeren RH and Rijkers K (2021) Microvascular changes associated with epilepsy: A narrative review. *Journal of cerebral blood flow and metabolism : official journal of the International Society of Cerebral Blood Flow and Metabolism* **41**:2492-2509.

- van Vliet EA, Aronica E and Gorter JA (2015) Blood-brain barrier dysfunction, seizures and epilepsy. *Seminars in cell & developmental biology* **38**:26-34.
- van Vliet EA, da Costa Araújo S, Redeker S, van Schaik R, Aronica E and Gorter JA (2007) Blood-brain barrier leakage may lead to progression of temporal lobe epilepsy. *Brain* **130**:521-534.
- van Vliet EA and Marchi N (2022) Neurovascular unit dysfunction as a mechanism of seizures and epilepsy during aging. *Epilepsia* **63**:1297-1313.
- Vanova N, Pejchal J, Herman D, Dlabkova A and Jun D (2018) Oxidative stress in organophosphate poisoning: role of standard antidotal therapy. *Journal of applied toxicology : JAT* **38**:1058-1070.
- Vazana U, Veksler R, Pell GS, Prager O, Fassler M, Chassidim Y, Roth Y, Shahar H, Zangen A and Raccach R (2016) Glutamate-mediated blood–brain barrier opening: implications for neuroprotection and drug delivery. *Journal of Neuroscience* **36**:7727-7739.
- Vezzani A and Viviani B (2015) Neuromodulatory properties of inflammatory cytokines and their impact on neuronal excitability. *Neuropharmacology* **96**:70-82.
- Vila Verde D, de Curtis M and Librizzi L (2021) Seizure-Induced Acute Glial Activation in the in vitro Isolated Guinea Pig Brain. *Frontiers in neurology* **12**:607603.
- Voirin A-C, Perek N and Roche F (2020) Inflammatory stress induced by a combination of cytokines (IL-6, IL-17, TNF- α) leads to a loss of integrity on bEnd. 3 endothelial cells in vitro BBB model. *Brain research* **1730**:146647.
- Vossel KA, Tartaglia MC, Nygaard HB, Zeman AZ and Miller BL (2017) Epileptic activity in Alzheimer's disease: causes and clinical relevance. *The Lancet Neurology* **16**:311-322.
- Waheed S, Sabeen A and Ullah Khan N (2014) New onset refractory status epilepticus as an unusual presentation of a suspected organophosphate poisoning. *Case reports in emergency medicine* **2014**:676358.
- Ward RJ, Dexter DT and Crichton RR (2022) Iron, Neuroinflammation and Neurodegeneration. *Int J Mol Sci* **23**.
- Webster KM, Sun M, Crack P, O'Brien TJ, Shultz SR and Semple BD (2017) Inflammation in epileptogenesis after traumatic brain injury. *J Neuroinflammation* **14**:10.
- Weiss JM, Downie SA, Lyman WD and Berman JW (1998) Astrocyte-derived monocyte-chemoattractant protein-1 directs the transmigration of leukocytes across a model of the human blood-brain barrier. *Journal of immunology (Baltimore, Md : 1950)* **161**:6896-6903.
- Weissberg I, Wood L, Kamintsky L, Vazquez O, Milikovsky DZ, Alexander A, Oppenheim H, Ardizzone C, Becker A and Frigerio F (2015) Albumin induces excitatory synaptogenesis through astrocytic TGF- β /ALK5 signaling in a model of acquired epilepsy following blood–brain barrier dysfunction. *Neurobiology of disease* **78**:115-125.
- Wilcox KS, Gee JM, Gibbons MB, Tvrdik P and White JA (2015) Altered structure and function of astrocytes following status epilepticus. *Epilepsy & Behavior* **49**:17-19.
- Wolburg H and Lippoldt A (2002) Tight junctions of the blood–brain barrier: development, composition and regulation. *Vascular pharmacology* **38**:323-337.
- Won C, Lin Z, Kumar TP, Li S, Ding L, Elkhali A, Szabó G and Vasudevan A (2013) Autonomous vascular networks synchronize GABA neuron migration in the embryonic forebrain. *Nature communications* **4**:2149.

- Wong AD, Ye M, Levy AF, Rothstein JD, Bergles DE and Searson PC (2013) The blood-brain barrier: an engineering perspective. *Frontiers in neuroengineering* **6**:7.
- Yang H, Rajah G, Guo A, Wang Y and Wang Q (2018) Pathogenesis of epileptic seizures and epilepsy after stroke. *Neurological research* **40**:426-432.
- Zenaro E, Piacentino G and Constantin G (2017) The blood-brain barrier in Alzheimer's disease. *Neurobiol Dis* **107**:41-56.
- Zonta M, Angulo MC, Gobbo S, Rosengarten B, Hossmann K-A, Pozzan T and Carmignoto G (2003) Neuron-to-astrocyte signaling is central to the dynamic control of brain microcirculation. *Nature neuroscience* **6**:43-50.

Footnotes:

No author has an actual or perceived conflict of interest with the contents of this article.

This work was supported by the National Institute of Neurological Disorders and Stroke CounterACT program, grant numbers NS079202 and NS127758 (PJL); the National Institute of General Medical Sciences training program, grant number T32GM135741 (CMU); the 2022-2023 Lodric Maddox Graduate Fellowship from the UC Davis School of Veterinary Medicine (PNB); and the 2020-2023 ARCS Foundation Northern California Scholar Program (PMA).

Reprint request: Pamela J. Lein, Department of Molecular Biosciences, School of Veterinary Medicine, University of California, Davis, CA. E-mail: pjlein@ucdavis.edu

Figure Legends:

Figure 1. Structure of the blood-brain barrier (BBB). The brain vasculature forms a specialized selective barrier that prevents the passage of red blood cells (RBC), leukocytes (LK), and many blood-borne molecules from the blood into the brain parenchyma. A physical barrier is composed of endothelial cells (EC) with limited to no fenestrae that form tight junctions (TJ) between adjacent ECs. Pericytes (PC) line the abluminal side of the ECs, and the EC-TJ-PC complex is embedded in a basement membrane (BM). Astrocyte (AC) projections, known as astrocytic end-feet (ACF), are critically important in inducing EC to form a BBB, and they regulate BBB permeability via signaling with PCs and ECs. Neurons (N) and microglia (MG) also influence the expression and organization of essential proteins in the BBB.

Figure 2. Putative mechanisms by which acute OP intoxication impairs BBB integrity to cause acquired epilepsy/spontaneous recurrent seizures (SRS). Acute OP intoxication inhibits acetylcholinesterase, which in turn causes: (1) accumulation of acetylcholine in central and peripheral synaptic clefts that can trigger (2) seizures that rapidly progress to status

epilepticus (SE). Sustained neuron depolarization as a result of SE activates glial cells and triggers excitotoxicity. (3) Activated astrocytes and microglia secrete pro-inflammatory factors and reactive oxygen species (ROS) that promote neurodegeneration and sustain seizure activity. In addition, (4) dying neurons release intracellular metabolites that contribute to neuroinflammation and oxidative stress. All four phenomena (1-4) have direct roles in causing (5) BBB impairment and consequent leakage of blood-borne components into the brain parenchyma. Albumin and fibrinogen access the brain parenchyma, activating (6) TGF- β signaling that induces (7) transcriptional modifications that decrease glutamate metabolism and expression of potassium ion channels (Kir4.1) and aquaporin 4 in astrocytic end-feet, and increase the release of pro-inflammatory cytokines and excitatory synaptogenesis. (8) Glutamate accumulation opens calcium channels in neurons and endothelial cells. (9) Elevated intracellular calcium levels activate calpain, which cleaves cytosolic and nuclear proteins, and if released into the extracellular space, extracellular matrix proteins. (10) Matrix metalloproteinases (MMPs), specifically MMP-2 and MMP-9, are activated by tissue plasminogen activator (tPA) and neuroinflammatory/oxidative factors that results in (11) chronic alterations of the tight junction (TJ) proteins. A leaky BBB promotes further leakage (3-5) and reduces the neuron's depolarization threshold, thus decreasing seizure threshold and increasing susceptibility to epileptic activity. Orange arrows indicate mechanisms and processes leading to BBB impairment (1-4); blue arrows indicate mechanisms and processes that are consequences of BBB impairment and lead to epilepsy/SRS (6-11); blue/orange dashed arrows indicate mechanisms and processes that are consequences of BBB impairment and contribute, directly or indirectly, to further impairment of the BBB (5, 6, 9, 11).