

# Lawrence Berkeley National Laboratory

## Recent Work

### Title

SUPERDEF0RMATION IN [SUP]104,105 PD

### Permalink

<https://escholarship.org/uc/item/7jx112k4>

### Authors

Macchiavelli, A.O.

Burde, J.

Diamond, R.M.

### Publication Date

1988-07-01



# Lawrence Berkeley Laboratory

UNIVERSITY OF CALIFORNIA

RECEIVED  
LAWRENCE  
BERKELEY LABORATORY

AUG 6 1988

LIBRARY AND  
DOCUMENTS SECTION

Submitted to Physical Review C


## Superdeformation in $^{104,105}\text{Pd}$

A.O. Macchiavelli, J. Burde, R.M. Diamond, C.W. Beusang,  
M.A. Deleplanque, R.J. McDonald, F.S. Stephens, and J.E. Draper

July 1988

**TWO-WEEK LOAN COPY**

*This is a Library Circulating Copy  
which may be borrowed for two weeks.*



LBL-25597  
e. 2

## **DISCLAIMER**

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

**SUPERDEFORMATION IN  $^{104,105}\text{Pd}$** 

A.O.Macchiavelli\*, J.Burde\*\*, R.M.Diamond, C.W.Beausang, M.A.Deleplanque  
R.J.McDonald, F.S.Stephens and J.E.Draper\*\*\*

Nuclear Science Division  
Lawrence Berkeley Laboratory  
University of California  
Berkeley, California 94720

In the present communication we report the discovery of a rotational band in  $^{104}\text{Pd}$  and one in  $^{105}\text{Pd}$  that can be interpreted as arising from superdeformed shapes. The moments of inertia,  $J^{(1)}$  and  $J^{(2)}$ , of these bands are similar to those measured in the Ce-Nd region, once the  $A^{5/3}$  mass dependence is removed. This implies a deformation  $\epsilon \sim 0.35-0.4$ . This is the third mass region where superdeformed bands have been found at high spins.

This work was supported by the Director, Office of Energy Research, Division of Nuclear Physics of the Office of High Energy and Nuclear Physics of the U.S. Department of Energy under Contract DE-AC03-76SF00098.

The understanding of the nuclear shape requires a knowledge of macroscopic properties, determined by the interplay of Coulomb, surface and rotational energies, and of microscopic properties associated with the detailed motion of the nucleons near the Fermi surface. Of special interest are those shapes known as "superdeformed" (SD) where the nucleus acquires a very elongated shape that can be approximately represented by an ellipsoid where the ratio of the long to short axis is considerably larger than that of normal deformation  $\sim 1.3:1$ . Within the framework of the anisotropic harmonic oscillator model<sup>1</sup> one can expect the existence of favorable shell gaps that appear regularly as a function of deformation and nucleon number. They are predicted to occur for particular "superdeformed magic numbers" and at deformations corresponding to integer ratios of the lengths of the axes (e.g.,  $\epsilon = 0.6$  corresponds to a ratio of 2:1). More realistic calculations provide a better prediction of regions of superdeformation<sup>2</sup> and these simple regularities are largely lost. Nevertheless, superdeformed shapes are predicted to occur, and their observation and systematic study will result in a better understanding of the interplay between the macroscopic and microscopic aspects of the nuclear motion.

The first observation of SD nuclei goes back to the discovery of fission isomers<sup>3</sup> and the identification of the rotational bands built on them<sup>4</sup>. This interesting topic of nuclear structure had to wait 15 years until other discrete superdeformed bands were discovered at high spins in the 150 and 130 mass regions<sup>5</sup>. This was possible only when large arrays of Compton-suppressed Ge detectors became operational and were able to extract these bands from the complex background that occurs in the spin region where they are found. Not much is yet known about the actual properties of these bands, such as their single-particle configuration, pairing correlations and their population and decay mechanisms. It is important, at this stage, to determine experimentally the regions where superdeformation occurs. In the present communication we report the discovery of rotational bands in  $^{104}\text{Pd}$  and  $^{105}\text{Pd}$  that can be interpreted as originating from a second minimum in the potential energy surface of these nuclei at high spin and large deformation. This is a new mass region where discrete superdeformed bands have been

found at high spins, the other known regions being around  $^{152}\text{Dy}$  and in the light Ce-Nd nuclei.

The experiment was carried out at the Lawrence Berkeley Laboratory 88-inch Cyclotron. Two stacked  $^{64}\text{Ni}$ , self-supporting targets  $0.5 \text{ mg/cm}^2$  thick, were bombarded with a 200 MeV  $^{48}\text{Ca}$  beam. The emitted  $\gamma$ -rays were detected in 20 Compton-suppressed Ge detectors from the HERA array at an event rate for triple and higher fold coincidences of approximately  $2000 \text{ s}^{-1}$ . A total of 70 million events were recorded event-by-event on magnetic tapes for subsequent off-line analysis.

A "doubles" sort, in which the events were broken into independent pairs showed the existence of a very strong ridge structure in the  $\gamma$ - $\gamma$  correlation matrix <sup>6</sup>. The observed energy spacing is consistent with that from a strongly deformed shape. A more detailed inspection revealed the presence of two discrete bands that were associated with  $^{104}\text{Pd}$  and  $^{105}\text{Pd}$ , the  $\alpha 4n$  and the  $\alpha 3n$  products, respectively. Figure 1a shows a background-subtracted spectrum obtained by summing spectra having gates within the band assigned to  $^{105}\text{Pd}$ . These transitions are also clearly seen in the spectrum in Fig. 1b, which corresponds to a gate on the known 306 keV ground-state transition in  $^{105}\text{Pd}$  <sup>7</sup>. A summed spectrum derived from gates in the band assigned to  $^{104}\text{Pd}$  is shown in Fig. 1c. A partial decay scheme is shown in Fig. 2 for the two nuclei.

$^{105}\text{Pd}$  : In an attempt to find links between the SD band and the yrast states, a "triples" sort was performed, requiring two  $\gamma$ -ray gates in the SD band or one gate in the band and the other in the ground state ( $g_{7/2}$ ) band before incrementing a spectrum with the third  $\gamma$ . As in previous cases, no obvious links could be found. However, the  $\gamma$ -ray intensities deduced from the triples sort of the data, summarized in Table 1, show that the band deexcites both to the  $g_{7/2}$  and the  $h_{11/2}$  bands with roughly equal intensity. The feeding of the  $g_{7/2}$  band seems to take place at a spin of  $39/2$  and therefore the deexcitation pattern suggests a spin of  $43/2$  for the first observed level of the new rotational band, with an uncertainty of  $\pm 2$  units. Angular correlations with gates on the band members or on the 306 keV ( $\Delta I=1$ ) transition established that the transitions are

consistent with stretched quadrupole character<sup>8</sup>, thus extending the spin sequence up to 83/2 for the top transition. With the branching derived from the triples sort, the relative intensities from a gate in the 306 keV transition and its known intensity, the population of this band was determined to be  $\sim 30\%$  of the total events leading to  $^{105}\text{Pd}$ .

$^{104}\text{Pd}$ : In contrast with  $^{105}\text{Pd}$ , the data from the triples sort were insufficient to provide additional useful information, but consistent with the doubles sort analysis (perhaps due to the relatively lower intensity of this product and of its band). From the single-gated spectra it was possible to conclude that the band feeds the yrast states at around spin 22, and therefore a spin  $24 \pm 2$  could be assigned for the lowest state of the band. The population of this band represents  $\sim 15\%$  of the events in  $^{104}\text{Pd}$ , as obtained from gates in low-spin members of the yrast band<sup>9</sup>. Relative intensities are also given in Table 1.

It can be argued that these bands should be populated when they are in the yrast region, i.e., close to the highest spins observed. Therefore, due to their large moment of inertia they would deexcite at  $\sim 2\text{-}3\text{MeV}$  above the yrast line.

The moments of inertia  $J^{(1)}$  and  $J^{(2)}$ , shown in Fig. 3, are similar to those measured in the mass 130 region, once the  $A^{5/3}$  dependence is removed, suggesting a deformation of  $\epsilon \sim 0.35\text{-}0.4$  for these bands. The gradual decrease in  $J^{(2)}$  can be qualitatively understood by the gradual alignment of the valence nucleons' angular momentum with the rotation axis, as the rotational frequency increases. Therefore, the generation of angular momentum becomes energetically more expensive and the nucleus approaches the "band termination" that will occur when all the angular momentum available for that configuration is aligned. In comparison with  $^{152}\text{Dy}$ , it is expected<sup>10</sup> that in the spin range observed a bigger proportion of this maximum angular momentum is exhausted for these lighter systems, and therefore the reduction of  $J^{(2)}$  is faster.

Both  $^{104}\text{Pd}$  and  $^{105}\text{Pd}$  can be interpreted as soft, slightly deformed, prolate rotors in the normal (low frequency) regime<sup>7,9</sup>. The observation of SD bands indicates the existence of a second minimum in the potential-energy surface of these nuclei. As mentioned before,

these minima are likely at  $\epsilon \sim 0.35-0.4$  and become yrast (lowest in energy) at a spin of about 40. The increase in  $J^{(2)}$  in  $^{105}\text{Pd}$  at  $\hbar\omega \sim 1$  MeV is quite interesting and may be caused by a band crossing. It is the first time such a crossing has been seen in a SD band.

The structure of these bands can be discussed in terms of the cranked shell model<sup>11</sup>. We have calculated the single-particle routhians at a deformation of  $\epsilon=0.4$  and, at the frequencies where the SD bands are observed, a plausible configuration involves levels from the  $N=4$  and  $5$  shells for both neutrons and protons, i.e.  $\pi(N=4)^4(N=5)^2 \nu(N=4)^{14}(N=5)^4$ . The deformation-driving effect of the  $\pi$  and  $\nu$   $N=5(h_{11/2})$  high-j orbitals will favor the stabilization of the second minima in the potential-energy surface of these nuclei. Furthermore, the calculations suggests that the increase of  $J^{(2)}$  in  $^{105}\text{Pd}$  at the highest frequencies can be attributed to the alignment of a pair of neutrons in the  $N=6$   $i_{13/2}$  level. The crossing is predicted to occur at  $\hbar\omega \sim 1.1$  MeV, very close to the observed frequency.

A more general calculation, regarding the existence of SD gaps in a realistic Wood-Saxon potential has been presented by Dudek et al.<sup>2</sup>. An inspection of single-particle levels as a function of deformation reveals the presence of gaps at large quadrupole deformation for  $Z=46$  and  $N=58$ . These predictions are consistent with our observations, although a quantitative comparison is beyond the scope of this work. It is interesting to note that the neutron number of  $N=58,59$  for the present cases can be related to the proton number  $Z=58,60$  in the Ce-Nd region.

In conclusion, we have observed bands in  $^{104}\text{Pd}$  and  $^{105}\text{Pd}$  consisting of stretched E2 transitions. Their regularity and measured moments of inertia suggest that these bands are based on a superdeformed configuration having  $\epsilon \sim 0.35-0.4$ , similar to those observed in the Ce-Nd region. The sudden increase in  $J^{(2)}$  in the SD band of  $^{105}\text{Pd}$  at the highest spins may be due to the alignment of a pair of  $i_{13/2}$  neutrons.



## References

- \* Permanent Address: Comisión Nacional de Energía Atómica, Buenos Aires, Argentina.
- \*\* Permanent Address: Racah Institute of Physics, Hebrew University, Jerusalem, Israel.
- \*\*\* Permanent Address: University of California, Davis, California 95611
- <sup>1</sup> A.Bohr and B.Mottelson, Nuclear Structure Vol. 2, (W.A.Benjamin Inc.,1975).
- <sup>2</sup> V.M.Strutinsky, Nucl. Phys. **A95**, 420 (1967).
- V.M.Strutinsky, Nucl. Phys. **A122**, 1 (1968).
- K.Neergaard, V.V.Pashkevich and S.Frauendorf, Nucl. Phys. **262**, 61 (1976)
- I.Ragnarsson, T.Bengtsson, G.Leander and S.Aberg, Nucl. Phys. **A347**, 287 (1980).
- J.Dudek and W.Nazarewicz, Phys. Rev. **C31**, 298 (1985).
- J.Dudek et al., Phys. Rev. Lett. **59**, 1405 (1987).
- <sup>3</sup> S.M. Polikanov et al., Sov. Phys. JETP **15**, 1016 (1962).
- <sup>4</sup> H.J.Specht, J.Weber, E.Konecny and D.Heunemann, Phys. Lett. **41B**, 2182 (1972).
- <sup>5</sup> P.J.Twin et al., Phys. Rev. Lett. **57**, 811 (1986).
- P.J.Nolan et al., J. Phys. **G11**, L17 (1985).
- E.M.Beck et al., Phys. Rev. Lett. **58**, 2182 (1987), Phys. Lett. **B195**, 531 (1987).
- B.Hass et al., Phys. Rev. Lett. **60**, 503 (1988).
- M.A.Deleplanque et al., Phys. Rev. Lett. in press.
- <sup>6</sup> O.Andersen et al., Phys. Rev. Lett. **43**, 687 (1979).
- <sup>7</sup> F.A.Rickey, J.A.Grau, L.E.Samuels and P.C.Simms, Phys. Rev. **C15**, 1530 (1977).
- <sup>8</sup> J.E.Draper, Nucl. Inst. Meth. **A247**, 481 (1986).
- <sup>9</sup> J.A.Grau et al., Phys. Rev. **C15**, 2302 (1976).
- <sup>10</sup> I.Ragnarsson, Phys. Lett. **B199**, 317 (1987).
- <sup>11</sup> R.Bengtsson and S.Frauendorf, Nucl. Phys. **A327**, 139 (1979).

Table 1

Gamma-ray energies, relative intensities and initial spin of transitions seen in the decay of the SD band. Intensities were derived from "doubles" and "triples" sorts of the data (see text).

$^{104}\text{Pd}$			$^{105}\text{Pd}$		
$E_\gamma$ (keV)	Intensity $L_\gamma$ <sup>a</sup> %	$I^\pi$	$E_\gamma$ (keV)	Intensity $L_\gamma$ <sup>a</sup> %	$I^\pi$
556	100	2 <sup>+</sup>	306	50	7/2 <sup>+</sup>
768	96	4 <sup>+</sup>	706	50	11/2 <sup>+</sup>
926	142 <sup>b</sup>	6 <sup>+</sup> , 16 <sup>+</sup>	891	51	15/2 <sup>+</sup>
971	87	8 <sup>+</sup>	854	45	19/2 <sup>+</sup>
802	175 <sup>b</sup>	10 <sup>+</sup> , 14 <sup>+</sup>	539	41	23/2 <sup>+</sup>
612	89	12 <sup>+</sup>	578	41	27/2 <sup>+</sup>
1064	100	18 <sup>+</sup>	795	35	31/2 <sup>+</sup>
1194	100	20 <sup>+</sup>	1014	35	35/2 <sup>+</sup>
			1178	24	39/2 <sup>+</sup>
1263	127 <sup>c</sup>	22 <sup>+</sup> , (26)			
1381	78	(28)	481	50	15/2 <sup>-</sup>
1511	93	(30)	772	52	19/2 <sup>-</sup>
1638	79	(32)	958	53	23/2 <sup>-</sup>
1763	92	(34)			
1919	78	(36)	1208	51	(47/2)
2079	64	(38)	1282	47	(51/2)
			1379	96	(55/2)
			1488	105	(59/2)
			1597	110	(63/2)
			1720	100	(67/2)
			1846	60	(71/2)
			2007	55	(75/2)
			2141	30	(79/2)
			2240	12	(83/2)

a : Intensities have an uncertainty of  $\pm 10\%$  .

b : Sum of transitions in the yrast sequence.

c : Includes the 1257keV transition in the yrast band.

### Figure Captions


Fig.1. a) Spectrum in coincidence with gates within the members of the SD band assigned to  $^{105}\text{Pd}$  ;  $\gamma$  -ray transitions are labeled by their energy in keV.

b) Spectrum in coincidence with the 306keV transition in  $^{105}\text{Pd}$  . The lines show the position of SD transitions. Transitions in the yrast band are labeled by their initial spin (See Fig. 2).

c) Same as a) for the band assigned to  $^{104}\text{Pd}$  . Yrast transitions are labeled by their initial spin.


Fig. 2. Partial decay schemes of the SD bands in  $^{104}\text{Pd}$  and  $^{105}\text{Pd}$  .

Fig 3. Moments of inertia for the observed rotational bands. The dotted area represents  $J^{(1)}$  with an uncertainty of  $\pm 2$  in the spin assignment. Open circles are for  $^{104}\text{Pd}$  and full circles for  $^{105}\text{Pd}$  . For comparison, the dynamic moment of inertia (scaled by  $A^{5/3}$ ) in the SD band in  $^{132}\text{Ce}$  is shown as a dotted line. The right-hand scale is in units of the rigid sphere.




XBL 884-1204

Figure 1



XBL 884-1203

Figure 2



XBL 884-1205

Figure 3

*LAWRENCE BERKELEY LABORATORY  
TECHNICAL INFORMATION DEPARTMENT  
UNIVERSITY OF CALIFORNIA  
BERKELEY, CALIFORNIA 94720*