

UC San Diego

UC San Diego Previously Published Works

Title

CO2 Sensing and CO2 Regulation of Stomatal Conductance: Advances and Open Questions

Permalink

<https://escholarship.org/uc/item/7k45f83h>

Journal

Trends in Plant Science, 21(1)

ISSN

1360-1385

Authors

Engineer, Cawas B
Hashimoto-Sugimoto, Mimi
Negi, Juntaro
[et al.](#)

Publication Date

2016

DOI

10.1016/j.tplants.2015.08.014

Peer reviewed



Published in final edited form as:

Trends Plant Sci. 2016 January ; 21(1): 16–30. doi:10.1016/j.tplants.2015.08.014.

CO₂ sensing and CO₂ regulation of stomatal conductance: advances and open questions

Cawas Engineer¹, Mimi Hashimoto-Sugimoto², Juntaro Negi², Maria Israelsson-Nordstrom³, Tamar Azoulay-Shemer¹, Wouter-Jan Rappel¹, Koh Iba², and Julian Schroeder¹

¹Division of Biological Sciences, Cell and Developmental Biology Section, and Center for Molecular Genetics, University of California, San Diego, 9500 Gilman Drive, La Jolla, CA 92093-0116, USA

²Department of Biology, Faculty of Sciences, Kyushu University, 744 Motoooka, Nishi-ku, Fukuoka 819-0395, Japan

³Umeå Plant Science Center, Department of Forest Genetics and Plant Physiology, Swedish University of Agricultural Sciences, SE-901 83 Umeå, Sweden

Abstract

Guard cells form epidermal stomatal gas exchange valves in plants and regulate the aperture of stomatal pores in response to changes in the carbon dioxide (CO₂) concentration in leaves. Moreover, the development of stomata is repressed by elevated CO₂ in diverse plant species. Evidence suggests that plants can sense CO₂ concentration changes via guard cells and via mesophyll tissues in mediating stomatal movements. We review new discoveries and open questions on mechanisms mediating CO₂-regulated stomatal movements and CO₂ modulation of stomatal development, which together function in CO₂-regulation of stomatal conductance and gas exchange in plants. Research in this area is timely in light of the necessity of selecting and developing crop cultivars which perform better in a shifting climate.

Keywords

Atmospheric carbon dioxide; guard cell signaling; stomatal movements; stomatal development; climate; agriculture

Importance of CO₂ regulation of stomatal conductance

Higher than ambient CO₂ concentrations mediate a closure of stomatal pores in plants and conversely, low CO₂ concentrations trigger opening of stomatal pores. Respiration in plant

Corresponding author: Schroeder, J. (jischroeder@ucsd.edu).

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

The authors declare no conflicts of interest.

leaves in the night (dark) causes a rapid rise in the intercellular CO₂ concentration (C_i) in leaves and measurements indicate that CO₂ levels can exceed 600 ppm (Figure 1A) CO₂ [1]. Moreover intercellular CO₂ concentrations ([CO₂]) can rapidly drop to below 200 ppm in the light [165±58 ppm] [1]. In parallel to the diurnal oscillation in C_i, global CO₂ levels have risen exponentially [2, 3] since the advent of the industrial revolution (Figure 1B). April 2014 was the first month in recorded history to have consistently had CO₂ levels above 400 ppm. This increase in atmospheric CO₂ causes a rise in leaf C_i. Stomatal pore apertures respond to these changes in C_i [4, 5]. A longer term effect of the continuing [CO₂] rise is the down-regulation of stomatal development in the leaf epidermis [6]. This developmental response was first discovered almost 3 decades ago [6], and has been subsequently confirmed with evidence in the fossil record [7, 8]. While a preponderance of species exhibit this response, some species show either an opposite effect or are unresponsive to elevated CO₂ concentrations [9, 10]. Despite the prime importance of these responses, little has been known about the underlying genetic mechanisms.

The combined effect of the above two CO₂ responses negatively impacts overall stomatal conductance on a global scale by reducing both stomatal apertures and the total numbers of stomata per unit leaf area. While decreased stomatal conductance is beneficial for limiting water loss from plant leaves [11] through reduced evapotranspiration, there is also a potential cost to such water savings. Fewer and more closed stomata in response to elevated CO₂ levels also reduce the evapotranspirative cooling ability of leaves, which in turn adds to leaf heat stress [12–14] under water-limited growth regimens. Heat stress, combined with drought and rising temperatures can reduce plant health [15] and crop productivity globally, thus impacting agricultural practices and possibly nutrient content [16, 17] and supply [18]. Furthermore, a large stomatal conductance can correlate with improved crop yield [19–21], and thus the downregulation of stomatal conductance by CO₂ may contribute to suboptimal yields when sufficient water is available. Additionally, CO₂ concentrations also affect leaf development and leaf area in some species and these responses have a direct bearing on water conservation and improving biomass in plants. These varied responses to changes in atmospheric CO₂ levels complicate efforts to model and predict the effects of rising CO₂ levels on global gas exchange in forests and ecosystems. Future research and attempts to adapt crop production to climate change would benefit significantly from a molecular understanding of CO₂-regulated plant gas exchange. Here we summarize recent discoveries of molecular and cellular mechanisms mediating CO₂ regulation of stomatal movements. Interested readers are directed to reviews summarizing previous metabolic and physiological aspects of stomatal function [22–24]. We also review a recently uncovered initial framework [25] for mechanisms mediating elevated CO₂ repression of stomatal development in plants, and point to open questions for further inquiry into this pathway. Predicting plant performance and plant responses to the combination of rising CO₂ and other environmental stresses such as heat and drought is reviewed elsewhere [12–14, 26–29].

Convergence of CO₂ and ABA signaling

Pioneering research has shown that the plant hormone abscisic acid (ABA) enhances the stomatal CO₂ response [30]. However, the precise signaling convergence point remains to be resolved [31–33]. Among the first genetic stomatal CO₂ signaling components

discovered were the dominant abscisic acid (ABA) -insensitive PP2C protein phosphatase mutations, *abi1-1* and *abi2-1*, which cause conditional CO₂ insensitivity [31, 34]. Molecular genetic components of the ABA and stomatal movement regulatory signal transduction machinery have been shown to be conserved over evolutionary timescales [35–37]. Interestingly, ABA signaling through ABA receptors, named PYR/RCARs, partially affects stomatal responses to elevated CO₂ [33]. A present model proposes that these ABA receptors may not directly mediate guard cell CO₂ signaling, but disruption slows CO₂ responses. These results suggest that CO₂ and ABA responses are mediated by converging pathways, such that the CO₂ response is weakened in the complete absence of an ABA response. Thus two models could be considered: (i) ABA and ABA receptors function upstream of the convergence point of CO₂ and ABA signaling, while synergistically amplifying common downstream signaling mechanisms [32, 33]. Or (ii) CO₂ elevation may rapidly (within 1 to 3 minutes) cause an elevation in the guard cell [ABA] concentration and thus mediate stomatal closing. The recent development of real time ABA reporters enables time-dependent monitoring of ABA concentration changes in single cells including guard cells [38, 39]. Such Förster resonance energy transfer (FRET) reporters will enable future analyses which address the question of whether [CO₂] elevation causes a rapid [ABA] increase in guard cells.

Roles for carbonic anhydrase enzymes in CO₂ signaling

The cellular sensing mechanisms for changes in CO₂ and/or bicarbonate concentrations that mediate rapid high CO₂-induced stomatal movements are key to understanding the underlying response. Carbonic anhydrase enzymes functioning in guard cells have been shown to be important for the rapid wild type stomatal movement response to changes in CO₂ levels [40]. These enzymes are beta carbonic anhydrases [40] and the CO₂ response of the alpha and gamma classes of carbonic anhydrases [41] has not yet been determined. Interestingly, complementation of double mutant plants by expressing a structurally-unrelated mammalian carbonic anhydrase preferentially in guard cells restores the WT response [40]. These data indicate that the catalytic activity of the carbonic anhydrase enzymes is involved in “transponding” [42] the CO₂ stimulus in this pathway and point to bicarbonate and/or protons as possible second messengers involved in signal transduction. The importance of the role of bicarbonate in stomatal aperture regulation was found in patch clamp studies. Elevation in the cytoplasmic bicarbonate concentration activates anion channels in guard cells required for stomatal closing [32]. A recent study has suggested that a MATE transporter-like protein, RHC1, functions as a bicarbonate sensor in this pathway [43].

Two key questions for stomatal CO₂ signaling are the tissue and sub-cellular sites of CO₂ perception. Is CO₂ sensed in the mesophyll or in the guard cells or in both cell types? CO₂ perception in animals is currently believed to be a vestigial sensory mechanism [42] and studies in animals have linked carbonic anhydrases to the perception of carbonation [44]. Adenylyl cyclase enzymes have been shown to play important roles in CO₂ sensing in mammalian and fungal systems [45, 46], but plant orthologs for these proteins have not been identified. Another area of scientific inquiry is whether photosynthesis plays a direct role in rapid CO₂-regulated stomatal movements. Of the six *Arabidopsis thaliana* beta carbonic

anhydrases, CA4 is localized at the plasma membrane while CA1 is mainly targeted to chloroplasts upon transient expression in tobacco (*Nicotiana benthamiana*) cells [40, 47]. This expression pattern raises the question: how does this expression pattern of carbonic anhydrases function in mediating stomatal responses to CO₂?

In a recent study, the intracellular targeting of the β -carbonic anhydrases CA1 and CA4 in guard cells was characterized in relation to their roles in CO₂ control of stomatal movements. Expression of fluorescently-tagged carbonic anhydrases in guard cells of *calca4* double mutant plants showed that expression of CA4 at the plasma membrane or CA1 in guard cell chloroplasts can restore CO₂-induced stomatal responses [48]. Mathematical modeling and experiments have revealed that spatial gradients of inter-cellular second messengers can play a role in signaling events [49]. In addition, recent computational and experimental studies in *Xenopus laevis* oocytes have demonstrated that increases or decreases in the external CO₂ concentration can lead to significant and transient spatial intracellular gradients of CO₂ and its reaction products including bicarbonate [50]. In this mathematical model, the plasma membrane was assumed to be permeable to CO₂ and carbonic anhydrase were assumed to be in the cytoplasm. Although reaction-kinetic parameters of carbonic anhydrases may be similar in plants, the size of guard cells is significantly smaller than *Xenopus laevis* oocytes: $\approx 5 \mu\text{m}$ compared to $\approx 1000 \mu\text{m}$. This smaller dimension leads to a much smaller diffusional time scale, which can be estimated by taking the ratio of the square of the cell size and the diffusion constant. Taking the diffusion constant of CO₂ to be of the same order as in oocytes, $D \approx 1000 \mu\text{m}^2/\text{s}$, the diffusion time in guard cells is less than 1 s [48], which is faster than the typical time scale of stomatal responses. Thus, CO₂ gradients are unlikely to occur in guard cells and play a significant role in regulating stomatal opening and closing. Consistent with this analysis, recent mathematical modeling addressing the spatial localization of carbonic anhydrases showed virtually identical dynamics of HCO₃⁻ concentration changes at any depth into modeled guard cells [48]. This modeling study predicts that carbonic anhydrases that are localized to the plasma membrane and to the cytoplasm produce effective intracellular HCO₃⁻ concentration changes in guard cells, while the participation of the chloroplast localized CA, in this process, is less likely [48]. This simplified model, together with functional analyses of CA1 and CA4, predicted that intracellular HCO₃⁻ concentration in guard cells, driven by carbonic anhydrases at the plasma membrane or in the cytosol, can regulate stomatal CO₂ responses. In addition, this research also suggests that in guard cells, chloroplast-localized carbonic anhydrases contribute to CO₂ control of stomatal closing via an unknown plastidial mechanism that will require further investigation [48].

Based on the above model and findings, new gas exchange analyses of single carbonic anhydrase mutants support the model that the plasma membrane-localized CA4 plays an important role in CO₂ signal transduction as the *ca4* mutant can show a slightly altered sensitivity, even if not always significant, to the CO₂ stimulus (Figure 1C and D). The *calca4* double mutant shows a clearly more pronounced slowed CO₂ response [40, 48] (Figure 1E). One possible model to explain the enhanced phenotype in *calca4* could be that either mutation alone causes a largely sub-threshold impairment in the CO₂ response. However, overexpression of either CA is sufficient for recovery of a more wild type-like

CO₂ response by over-coming this threshold effect [40, 48]. This threshold model is also consistent with the finding that the catalytic activity of carbonic anhydrases mediates the stomatal CO₂ response [40], and that CAs accelerate the rate of CO₂/ HCO₃⁻ + H⁺ equilibration.

thaliana βCA1 is mainly targeted to the chloroplast stroma [47]. Furthermore, YFP-tagged βCA1 in guard cell chloroplasts restored stomatal responses to CO₂ [48]. These data raised the question, whether guard cell photosynthesis functions directly in the stomatal-CO₂ response. Transgenic plants in which chlorophyll was directly removed from guard cell chloroplasts, but not from mesophyll chloroplasts, showed that chlorophyll-lacking guard cells continue to show CO₂-induced stomatal closing [51]. Together with pharmacological studies using norflurazon [52], these findings suggest that the role of βCA1 in CO₂ signaling does not lie in photosynthesis, and implies an unknown function of guard cell chloroplasts in the CO₂ response, which needs to be identified.

Mesophyll and guard cell mechanisms of CO₂ signaling and a role for photosynthesis

Photosynthesis plays a well-documented role in CO₂ regulation of stomatal apertures by lowering the intercellular CO₂ concentration (C_i) in leaves [4] (Figure 1A). In addition to this indirect function of photosynthesis via lowering of C_i, present studies have led to differing models for the site of direct CO₂ sensing in the published literature (for reviews see [53, 54]). Some studies have shown a role for guard cells in CO₂ sensing [52, 55–58]. Yet other studies indicate that mesophyll cells play a key role by directly sensing CO₂ [59, 60]. Presently it cannot be excluded that two pathways (mesophyll and guard cell) converge. Genetic evidence for CO₂ insensitive mutants in both pathways is needed. We will review each pathway separately.

CO₂ sensing in the mesophyll

Stomatal conductance was demonstrated to respond to intercellular [CO₂] rather than external leaf surface [CO₂] [4]. Based on results from experiments which showed that stomata in isolated epidermal tissues of *Tradescantia pallida* and *Pisum sativum* exhibited a limited response to CO₂ and that mesophyll tissue enhanced the response, it has been proposed that the mesophyll tissue in leaves plays an important role in CO₂ sensing [60, 61]. Studies revealed that an enhanced and reversible stomatal response to CO₂ occurs when mesophyll tissues are placed upon an excised leaf epidermis [60]. Studies suggest that this CO₂ response involves a diffusible substance (cite here refs 60 and 62: Mott et al Fujita et al)., Adding a polyethylene (diffusion disruption) barrier impairs mesophyll amplification of the stomatal response [62]. This result was proposed to show that mesophyll-derived signals could be small diffusible molecules in the aqueous phase [62]. While the precise nature of the mesophyll to stomata signal is unknown, possible candidates include: sucrose [53] or malate [63]. Abscisic acid (ABA) should also be considered as a possible diffusible small molecule, as ABA plays a synergistic role in the CO₂ response [30, 33, 34, 64]. Notably, ABA was shown to amplify the CO₂ response in *Vicia faba* guard cells which typically show a much more sluggish response to CO₂ without the addition of ABA [65].

Further research is needed to determine whether the level of the unknown mesophyll-derived amplifying factor is itself regulated by CO₂ concentration changes or whether such a factor is constitutively present, independent of CO₂ and thus amplifies CO₂ sensing mechanisms in guard cells. Identification of the mesophyll-derived signaling molecules is needed to elucidate the predicted CO₂ signaling mechanisms. Isolation of genetic mutants that show CO₂ insensitivity in regulation of stomatal movements and for which the underlying proteins function in mesophyll cells would also help advance an understanding of the underlying mechanisms.

Photosynthesis and CO₂-controlled movements

Studies have proposed that leaf photosynthesis directly mediates stomatal function during CO₂ regulation of stomatal movements [59, 60]. However, other studies suggest that these CO₂ sensing mechanisms do not depend on leaf photosynthesis [52, 57, 66]. Altering the photosynthetic capacity via Rubisco antisense plant lines did not affect stomatal movement rates and conductance responses at ambient CO₂ compared to the wild type [57, 66]. Norflurazon treated chlorophyll-less “albino” leaves showed functional CO₂ regulation of stomatal movements [40, 52]. Studies with Phosphoenolpyruvate carboxylase (PEPC) antisense construct lines showed that in C₄ plants, PEPC activity is required for stomatal aperture opening under high light fluxes and low C_i [67].

Photosynthesis can partake in red light-induced stomatal opening by the photosynthesis-derived decrease in intercellular leaf space [CO₂] (C_i) [68–70]. Similarly, increased intercellular [CO₂] (C_i) in *Xanthium strumarium* leaves induces a decline of their stomatal conductance [59]. The slope of the stomatal conductance change was gradual until the transition of photosynthesis from Rubisco-limitation to electron transport-limitation, after which point the slope steeply declined with high values of C_i. A shift in the slope of stomatal conductance to intercellular [CO₂] in concert with this type of photosynthesis limitation, was observed under different [O₂] and light intensities thus implicating a role of photosynthesis in determining stomatal aperture [59]. However, when the calculated intercellular CO₂ concentration was kept stable by adjusting ambient CO₂, the majority of red light-induced stomatal opening was not dependent on mesophyll-induced reductions in intercellular CO₂. Therefore this study also indicated a C_i-independent component of red light-induced stomatal opening. In addition, leaves treated with the photosystem II inhibitor DCMU still responded to CO₂ [59]. Taken together, these studies point to the question whether red light-induced stomatal opening occurs via more than one pathway, with one pathway mediated by low C_i signaling. A recent genetic study of the High Temperature *ht1* mutant supports a model of a C_i-dependent and a C_i-independent component of red light-induced stomatal opening [70]. Another recent study, showed that carbonic anhydrase *calca4* double mutant plants did not affect whole leaf assimilation rates at a wide range of C_i concentrations from < 50 to > 1200 ppm CO₂ [51], further supporting the model that *calca4* mutant does not affect the CO₂ response via impairing whole leaf photosynthesis.

The function of guard-cell photosynthesis in CO₂-induced stomatal responses was genetically investigated by studying transgenic plants that lack chlorophyll specifically in guard cells [51]. Gas exchange analyses and stomatal movement analysis showed that CO₂-

induced stomatal closure is not directly mediated by guard-cell photosynthesis or electron transport. Interestingly, ~45% of the stomata in these lines showed a deflated “thin-shaped” stomatal morphology, suggesting a different but key role of guard cell photosynthesis in the energization and turgor production in guard cells. The role of photosynthesis in the direct response to CO₂-mediated stomatal conductance regulation remains a subject of debate.

CO₂ sensing in guard cells

Several studies have shown the ability of isolated guard cells to respond to environmental stimuli (and specifically [CO₂] changes) suggesting that sensing and signaling pathway components for these reside in guard cells [40, 55, 56, 71–73]. One of the first reported instances is the classical work by J.D.B. Weyers and colleagues which showed that in response to changes in [CO₂], isolated guard cell protoplasts swell and shrink [56, 73]. The presently identified genetic mutants that impair CO₂ control of stomatal movements encode proteins which exert their function in guard cells including the carbonic anhydrases CA1, CA4 [40], the HT1 protein kinase [74], the SLAC1 anion channel [75, 76], the OST1 protein kinase [32, 77], the PATROL1 Munc13-like protein [78], the AtALMT12/QUAC1 R-type anion channel [79] and the RHC1 MATE transporter [43].

Role for protein kinases

Thermal imaging of plant leaves is a convenient way to predict their transpiration and stomatal conductance [80, 81]. The *ht1* (*high leaf temperature 1*; *ht1-1* and *ht1-2*) mutant was a first CO₂ response mutant selected through a thermal imaging screen in *Arabidopsis thaliana* [74]. HT1 encodes a protein kinase, and *ht1-1* mutation leads to decreased HT1 kinase activity resulting in a reduced stomatal CO₂ response. The *ht1-2* mutant abolishes measurable HT1 kinase activity. Leaf stomata of *ht1-2* mutant plants show a constitutively “high” CO₂ response, resulting in a low stomatal conductance and thus elevated leaf temperature. Nonetheless, *ht1-2* leaf stomata retain responses to blue light, fusicoccin and ABA. To date, at least five *ht1* alleles have been isolated by the same strategy (Hashimoto-Sugimoto, unpublished). HT1 may be a central negative regulator of stomatal high CO₂ signaling [74] (Figure 2A). The open stomata 1 (OST1) protein kinase is required for CO₂-induced stomatal closing [32]. A recent study has proposed that the HT1 protein kinase phosphorylates the OST1 protein kinase, thereby deactivating OST1 [43]. However, another study has found HT1 to be epistatic to OST1 in a high CO₂-induced stomatal closing assay [70]. Further research is warranted to fully understand the roles of OST1 and HT1 in stomatal movement control.

Role for calcium

Calcium ions have been shown to act as guard cell second messengers in CO₂ signal transduction in *Commelina communis* [72, 82] and *Arabidopsis thaliana* [58, 83]. An ABA-insensitive mutant, *gca2* (*growth controlled by abscisic acid*) shows a similar [Ca²⁺]_{cyt} transient rate at low and elevated CO₂, and exhibits a strongly attenuated stomatal closure in response to elevated CO₂ in leaves [58]. The *gca2* mutant also exhibits an altered ABA-induced [Ca²⁺]_{cyt} pattern in guard cells (Allen et al. 2001), suggesting GCA2 functions downstream of, or at the convergence point of CO₂ and ABA signaling. Research suggests

that the increased Ca^{2+} spiking rate of *gca2* mutant guard cells results from a more negative plasma membrane voltage in guard cells due to CO_2 and ABA insensitivity [58]. Consistent with this model, more negative membrane potentials cause an increased Ca^{2+} spike rate in guard cells [84–86]. Cloning of the *GCA2* gene may help us better understand the synergistic mechanisms in CO_2 and ABA signal transduction.

Wildtype guard cells of the *Arabidopsis thaliana* accession *Landsberg erecta* show a dampening of the cytosolic Ca^{2+} transient rate in response to elevated CO_2 and ABA [58, 85], similar to *Vicia faba* and *Commelina communis* guard cells exposed to ABA [84, 87]. CO_2 induced dampening in the Ca^{2+} transient rate was not as clearly resolved in guard cells of the *Arabidopsis thaliana* Columbia accession when compared to the *Landsberg erecta* accession (Figure 2B). These findings may be linked to a weaker stomatal closing response in the *Arabidopsis thaliana* Columbia guard cells relative to those in the *Landsberg erecta* accession. The findings that guard cells show such “spontaneous” Ca^{2+} transients without ABA or CO_2 , but that CO_2 signaling requires cytosolic Ca^{2+} led to formulation of a “ Ca^{2+} sensitivity priming” hypothesis [58]. This model postulates that the stomatal closing stimuli, CO_2 and ABA, enhance (prime) the Ca^{2+} sensitivity of stomatal closing mechanisms, as has been found for abscisic acid [86, 88] and for $\text{CO}_2/\text{HCO}_3^-$ activation of S-type anion channels in guard cells [32]. Initial molecular, genetic, biochemical and cellular mechanisms mediating stimulus-induced Ca^{2+} sensitivity priming have been identified for ABA signal transduction showing that type 2C protein phosphatases control the ABA-mediated Ca^{2+} signal specificity [89].

Role for intracellular membrane trafficking

Stomatal opening is initiated by the activation of H^+ -ATPases in the guard cell plasma membrane. Low $[\text{CO}_2]$ triggers a more negative plasma membrane voltage (hyperpolarization), and elevated $[\text{CO}_2]$ is likely to inhibit proton efflux by plasma membrane H^+ -ATPases [55]. However, investigation of a direct link between CO_2 and H^+ -ATPase regulation remains to be characterized. The *Arabidopsis thaliana patroll* mutant isolated in a thermal imaging screen, is impaired in stomatal opening in response to low $[\text{CO}_2]$ and light [78]. The *PATROL1* gene encodes a protein with a MUN domain [90]. Munc13 has been reported to be involved in membrane trafficking of neurotransmitter release in animals, and the MUN domain is the minimal Munc13 region sufficient for the activity of Munc13-1 in synaptic vesicle priming [90]. This suggests a molecular role of *PATROL1* in intracellular membrane traffic. *PATROL1* is expressed throughout plant tissues including guard cells, and located in endosomes. Intriguingly, environmental conditions affect the intracellular distribution of the *PATROL1* protein. The *PATROL1* protein is detected in close proximity to the plasma membrane of guard cells under stomatal opening conditions (well-watered plants and light), and is observed inside guard cells as numerous punctate structures under stomatal closure conditions (dark or desiccation). Loss of function mutation of *patroll* disturbs the normal plasma membrane targeting of the *Arabidopsis thaliana* H^+ -ATPase *AHA1*, but does not affect that of the S-type anion channel *SLAC1*, and the inward rectifying potassium channel *KAT1* [78]. This indicates that *PATROL1* is target-selective and it has been proposed to function in targeting or tethering the *AHA1* H^+ -ATPase to the plasma membrane during stomatal opening. Over-expression

of *PATROL1* in plants (*PATROL1*-OX) leads to a rapid and enhanced rise in stomatal conductance in response to low $[\text{CO}_2]$ or light, eventually resulting in higher CO_2 assimilation and increased biomass production than wild type plants [78]. Furthermore, *PATROL1*-OX plants show normal stomatal closure in response to high $[\text{CO}_2]$, darkness and desiccation. Whether the increased biomass of *PATROL1*-OX plants is solely due to stomatal conductance changes or additional roles in enhancing photosynthesis as well as cellular expansion will be interesting for future analyses

Plasma membrane regulators

Increased $[\text{CO}_2]$ has been shown to enhance anion channel activity in guard cells [40, 69, 71, 91]. S-type anion channels in the plasma membrane of guard cells were proposed to provide a central control mechanism for stomatal closing [92]. Genetic evidence for this model was obtained in two studies. Thermography was used to isolate the *Arabidopsis thaliana* mutant *cdi3* (*carbon dioxide insensitive 3*) that is impaired in CO_2 -dependent leaf temperature change [75]. The CDI3 protein is a distant homologue of bacterial and fungal C4-dicarboxylate transporters, and is localized specifically in the plasma membrane of guard cells. *cdi3* guard cell protoplasts exhibited a higher content of K^+ , Cl^- and Malate^{2-} [75]. Parallel to this research, the characterization of the ozone signaling mutant *rcd3* (radical induced cell death) in *Arabidopsis thaliana* revealed that *rcd3Arabidopsis thaliana* leaves are more sensitive to the ozone radical because stomatal closing is impaired in response to ozone, ABA, Ca^{2+} and CO_2 [76]. The *RCD3* gene was independently mapped to the same C4-dicarboxylate transporter homologous gene [76]. Patch clamp analyses of *rcd3* mutant alleles showed that S-type anion channel currents were greatly impaired, whereas the R-type anion channel and Ca^{2+} channel currents were intact in mutant guard cells [76] providing genetic evidence for findings suggesting that S-type and R-type anion channels are distinct from one another [93]. These results supported the idea that CDI3/RCD3 encodes an essential anion transporting subunit of the Slow (S-type) anion channels in plants and was re-named SLAC1 [75, 76]. S-type anion channels in guard cells are activated by phosphorylation events [94, 95]. Electrophysiological experiments in *Xenopus laevis* oocytes demonstrated that in the presence of the protein kinase OST1 [77, 96], SLAC1 generates S-type anion channel activity [97, 98]. These studies provided evidence that SLAC1 encodes an S-type anion channel, which is activated by OST1-mediated phosphorylation. Recent research has focused on the regulatory events that trigger anion channel activation. The calcium-dependent protein kinases CPK21 and CPK23 phosphorylate and activate SLAC1 [99]. CPK6 with CPK3 was found to participate in the ABA- and Ca^{2+} -dependent regulation of guard cell S-type anion channels and stomatal closure [100]. The functional reconstitution of ABA activation of SLAC1 was shown using either the CPK6 or OST1 protein kinases in *Xenopus laevis* oocytes [101].

Several regulators involved in the ABA-induced activation of SLAC1 have been identified. An example is the receptor-like kinase GHR1, which is mainly localized in the guard cell plasma membrane, and activates SLAC1 anion currents in *Xenopus laevis* oocytes [102]. However, the relative functions of the Ca^{2+} -dependent and Ca^{2+} -independent pathways, which are key for models of guard cell signaling *in planta*, remain unknown. It remains to be genetically dissected whether these two pathways function independently of one another.

Furthermore, the molecular mechanism by which CO₂ activates S-type anion channel activity is a matter of present interest. OST1 is a positive regulator of CO₂-induced stomatal closing and activation of the S-type anion channels in guard cells [32]. Tian et al. recently reported [43] that a MATE-type transporter, RHC1, is activated by bicarbonate and functions upstream of HT1, and furthermore that HT1 directly phosphorylates OST1 and inhibits OST1-induced activation of SLAC1 (Figure 2A).

Various mechanisms have been suggested that couple a rise in [CO₂] to changes in the activity of plasma membrane ion channels. The apoplastic malate concentration rises from 1.00 ± 0.60 to 3.10 ± 2.30 mM in response to high [CO₂], which can activate R-type anion channels in guard cells [103]. This rise in malate may result from malate efflux from guard cells, as stomatal closing has been shown to be accompanied by malate efflux [104]. Furthermore, R-type anion channels mediate malate efflux [79, 105]. An alternative model has also been proposed in which CO₂ causes malate release from mesophyll cells [103], but direct evidence or mechanisms for this alternate model have not yet been reported. The *AtALMT12/QUAC1* gene was shown to encode the anion transporting activity of R-type anion channels in guard cells [79, 106]. *AtALMT12/QUAC1*, a member of the aluminum-activated malate transporter family in *Arabidopsis thaliana*, is highly expressed in guard cells and is targeted to the plasma membrane. Plants lacking *AtALMT12/QUAC1* are impaired in CO₂-induced stomatal closure, as well as in ABA responses. Electrophysiological studies of loss-of-function mutant guard cells and *Xenopus laevis* oocytes expressing the protein revealed that *AtALMT12/QUAC1* represents the malate-sensitive R-type anion channel [79].

The ABC transporter *AtABCB14*, was suggested to encode a malate uptake transporter in the guard cell plasma membrane [107]. Plants lacking the *AtABCB14* transporter were reported to show a slightly more rapid high CO₂-induced stomatal closure in comparison to wild type controls. However, in isolated epidermal strips that contained guard cells, no difference in stomatal CO₂ responses was observed between wild type and *atabcb14* mutant. The authors suggested that *AtABCB14* removes extracellular malate which is known to activate anion channels [103] and consequently, that part of the CO₂ response is mediated by malate secreted into the apoplast. Recent research has also proposed a role for *AtABCB14* in auxin transport [108]. Further research on *AtABCB14* functions is needed to determine the range of transported substrates.

CO₂ modulation of stomatal development

[CO₂] regulates the development of stomata [6, 109, 110] and wild type *Arabidopsis thaliana* plants of the Columbia ecotype show a slight repression in stomatal development at elevated CO₂ [111]. Responses of stomatal density to long-term CO₂ doubling are varied in different accessions of *Arabidopsis thaliana*, however, the majority of plant species exhibit reduced stomatal development [7, 109]. The mechanisms by which CO₂ levels can modulate stomatal development [112–118] are of interest given the continuing rise in atmospheric CO₂ levels.

Research has suggested that CO₂ control of stomatal development is mediated by Ci in leaves [119]. A cell wall wax biosynthesis mutant, *hic*, was shown to disrupt this response and exhibited an inverted developmental response, with more stomata being produced at elevated CO₂ in the *hic* mutant [120]. While the precise mechanism of action is unknown, *HIC* codes for a putative 3-keto acyl coenzyme A synthase which is involved in the synthesis of very-long-chain fatty acids. Interestingly, other genetic mutations which affect cell wall wax deposition in plants also show changes in stomatal development [120–123]. Presently unknown systemic signals which travel from mature to young leaves can function in CO₂-mediated changes of stomatal density in emerging young leaves [124], and thus cuticular waxes may serve as candidates which affect the movement of diffusible signals. Recent research found that carbonic anhydrase mutant plants exhibit an inversion in their stomatal development response to elevated CO₂, showing increased stomatal development at elevated [CO₂] (500 ppm or 1000 ppm) compared to low [CO₂] (150 ppm) grown plants [25]. These effects were observed in both cotyledons and mature rosette leaves [25]. The findings that elevated CO₂ causes enhanced stomatal development in cotyledons of *ca1ca4* double mutant plants provides evidence that CO₂ can also exert a developmental response without long distance leaf to leaf signaling [25]. The epidermal patterning factor gene EPF2 [125, 126] has been shown to bind the ERECTA receptor kinase [127–129]. The ERECTA receptor kinase [130, 131] and transgenic modulation of EPF2 expression [132] have also been shown to play roles in determining transpiration and water use efficiency of plants via regulation of stomatal index. Interestingly, *epf2* mutant alleles also show an inverted stomatal development phenotype at elevated CO₂ [25]. Activation of the EPF2-pro-peptide requires EPF2 cleavage. Cell wall proteome analyses led to the identification of the CRSP protease which can cleave EPF2 *in vitro* and which is involved in EPF2 function *in vivo*. Mutations in the CA1 and CA4 carbonic anhydrases, *EPF2* and *CRSP* genes show an inverted stomatal development response to elevated CO₂ [25]. CAs are required for high CO₂ upregulation of EPF2 and CRSP mRNAs based on qPCR analyses [25]. The mechanisms by which and whether carbonic anhydrases modulate EPF2 and CRSP activity remain to be determined. Several new questions emerge from this framework model (Figure 3); chief among them are: (i) Is CO₂ regulation of EPF2 and CRSP transcripts due to feedback modulation or is the CO₂ response transcriptional and what are the transcriptional regulators mediating the elevated CO₂ response? (ii) What are the downstream signaling components beyond the EPF2 peptide and ERECTA receptor? (iii) Analyses of CRSP protease insertion mutant alleles have led to the suggestion that additional proteases may act in the CO₂ response [25] and hence, are there other proteases which cleave and activate EPF2? (iv) How is excess EPF2 degraded or sequestered? And (v) Why do the described mutants show an increase in stomatal density and index at elevated CO₂ rather than a decrease?

While initial experiments have explored aspects of the effects of environmental stress on stomatal development [109, 118, 124, 133, 134], future genetic studies involving the specific effects of abiotic factors such as CO₂, light, temperature and humidity [124, 133] will reveal if and how these signaling cascades overlap and interact. A recent study showed that ABA plays an important role in stomatal cell lineage specification [135] by regulating the initiation of stomatal development and inducing the expansion of epidermal pavement cells.

Another area of future research which appears promising is the response of plants at lower than ambient C_i that occurs in daylight [1]. The carbonic anhydrase mutants exhibit WT-like stomatal development at low CO_2 [6, 25]. These findings point to the possibility of a separate mechanism regulating stomatal development at these lower than ambient CO_2 levels. One possibility is that other carbonic anhydrases may play key roles at these low CO_2 concentrations and remain to be identified. Alternatively, if the enzymatic activity of carbonic anhydrases transmits the developmental CO_2 signal, it is plausible that at low $[CO_2]$ of 150 ppm, non-accelerated catalysis of CO_2 to bicarbonate and protons may be sufficient for this CO_2 response. The specific mechanisms of EPF2 peptide stability and abundance modulation are open questions and lend themselves to several avenues of future research which could target the precise timing of CO_2 -mediated stomatal developmental repression and the temporal scale of repressive inhibition.

Concluding remarks and future directions

The observed transcriptional response to a CO_2 stimulus of hundreds of genes [25, 136], including photosynthesis-linked genes, indicates that unidentified mechanisms for transcriptional control exist in this pathway, as well as in CO_2 -regulated photosynthesis. Future research aimed at isolating these players could involve bioinformatic mining of published data on stomatal gene networks and guard cell responses to environmental triggers [25, 136–142]. Systems and network biology tools could be employed to probe transcriptomic and proteomic resources to identify not only putative transcription factors, but also identify candidate response elements of known transcriptional regulators of the stomatal development and movement pathways. Such candidates may also aid efforts in engineering plants that maximize stomatal conductance under well-watered conditions and more water use-efficient plants capable of adjusting to elevated CO_2 in a more desirable fashion than present cultivars. Another area of future research is the need for studying the stomatal and physiological responses of plants to the atmospheric CO_2 levels predicted to be present in the next decade [132, 143]. Results from such studies could assist breeders and biotechnological choices for plant and crop cultivars and germplasm better capable of adapting to those CO_2 levels. As an example, experiments exploring natural variation in stomatal density [144] which impact plant water use and productivity, could be conducted. Along these lines, a recent comprehensive study on 374 *Arabidopsis thaliana* accessions [145] has unveiled a wealth and diversity in stomatal responses to the stimuli light and CO_2 in *Arabidopsis thaliana*. As a starting point, these varied responses (see Outstanding Questions Box) to the same stimulus may indicate a plethora of underlying mechanisms which mediate responses of plants to abiotic stress and the environment.

Acknowledgments

Work in our laboratories was supported by the National Science Foundation (MCB1414339) and in part the National Institutes of Health (GM060396-ES010337) to J.I.S. and by Grants-in-Aid for Scientific Research on Priority Areas (21114002 and 26221103) to K.I. and (15K18556) to J.N. from the Ministry of Education, Culture, Sports, Science, and Technology of Japan. T.A.-S. received initial support from a postdoctoral fellowship from the US-Israel Binational Agricultural Research and Development Fund (BARD) and partial support from the UCSD FISP program.

References

1. Hanstein S, de Beer D, Felle HH. Miniaturised carbon dioxide sensor designed for measurements within plant leaves. *Sensors and Actuators B-Chemical*. 2001; 81(1):107–114.
2. Keeling CD, et al. Atmospheric Carbon-Dioxide Variations at Mauna-Loa Observatory, Hawaii. *Tellus*. 1976; 28(6):538–551.
3. Neftel A, et al. Evidence from Polar Ice Cores for the Increase in Atmospheric CO₂ in the Past 2 Centuries. *Nature*. 1985; 315(6014):45–47.
4. Mott KA. Do Stomata Respond to CO₂ Concentrations Other Than Intercellular. *Plant Physiology*. 1988; 86(1):200–203. [PubMed: 16665866]
5. Medlyn BE, et al. On the validation of models of forest CO₂ exchange using eddy covariance data: some perils and pitfalls. *Tree Physiol*. 2005; 25(7):839–857. [PubMed: 15870053]
6. Woodward FI. Stomatal numbers are sensitive to increases in CO₂ from pre-industrial levels. *Nature*. 1987; 327(6123):617–618.
7. McElwain JC, Chaloner WG. Stomatal Density and Index of Fossil Plants Track Atmospheric Carbon-Dioxide in the Paleozoic. *Annals of Botany*. 1995; 76(4):389–395.
8. Chater C, Gray JE, Beerling DJ. Early evolutionary acquisition of stomatal control and development gene signalling networks. *Current Opinion in Plant Biology*. 2013; 16(5):638–646. [PubMed: 23871687]
9. Ferris R, Taylor G. Stomatal Characteristics of Four Native Herbs Following Exposure to Elevated CO₂. *Annals of Botany*. 1994; 73(4):447–453.
10. Jordan GJ. A critical framework for the assessment of biological palaeoproxies: predicting past climate and levels of atmospheric CO₂ from fossil leaves. *New Phytologist*. 2011; 192(1):29–44. [PubMed: 21770947]
11. Keenan TF, et al. Increase in forest water-use efficiency as atmospheric carbon dioxide concentrations rise. *Nature*. 2013; 499(7458):324–327. [PubMed: 23842499]
12. Long SP, Ort DR. More than taking the heat: crops and global change. *Current Opinion in Plant Biology*. 2010; 13(3):241–248. [PubMed: 20494611]
13. Long SP, et al. Rising atmospheric carbon dioxide: Plants face the future. *Annual Review of Plant Biology*. 2004; 55:591–628.
14. Leakey AD, et al. Elevated CO₂ effects on plant carbon, nitrogen, and water relations: six important lessons from FACE. *J Exp Bot*. 2009; 60(10):2859–2876. [PubMed: 19401412]
15. Easlon HM, Bloom AJ. The effects of rising atmospheric carbon dioxide on shoot-root nitrogen and water signaling. *Front Plant Sci*. 2013; 4:304. [PubMed: 23983674]
16. Wroblewitz S, et al. Effect of rising atmospheric carbon dioxide concentration on the protein composition of cereal grain. *J Agric Food Chem*. 2014; 62(28):6616–6625. [PubMed: 24976461]
17. Weigel HJ. Plant quality declines as CO₂ levels rise. *Elife*. 2014; 3:e03233. [PubMed: 24872506]
18. Battisti DS, Naylor RL. Historical warnings of future food insecurity with unprecedented seasonal heat. *Science*. 2009; 323:240–244. [PubMed: 19131626]
19. Lu ZM, et al. Stomatal conductance predicts yields in irrigated Pima cotton and bread wheat grown at high temperatures. *Journal of Experimental Botany*. 1998; 49:453–460.
20. Bahar B, Yildirim M, Barutcular C. Relationships between Stomatal Conductance and Yield Components in Spring Durum Wheat under Mediterranean Conditions. *Notulae Botanicae Horti Agrobotanici Cluj-Napoca*. 2009; 37(2):45–48.
21. Prashar A, et al. Infra-red Thermography for High Throughput Field Phenotyping in *Solanum tuberosum*. *Plos One*. 2013. 8(6)
22. Mansfield TA, Hetherington AM, Atkinson CJ. Some Current Aspects of Stomatal Physiology. *Annual Review of Plant Physiology and Plant Molecular Biology*. 1990; 41(1):55–75.
23. Vavasseur A, Raghavendra AS. Guard cell metabolism and CO₂ sensing. *New Phytologist*. 2005; 165(3):665–682. [PubMed: 15720679]
24. Assmann SM. From proton pump to proteome. Twenty-five years of research on ion transport in higher plants. *Plant Physiol*. 2001; 125(1):139–141. [PubMed: 11154317]

25. Engineer CB, et al. Carbonic anhydrases, EPF2 and a novel protease mediate CO control of stomatal development. *Nature*. 2014
26. Bishop KA, et al. Is there potential to adapt soybean (*Glycine max* Merr.) to future [CO₂]? An analysis of the yield response of 18 genotypes in free-air CO₂ enrichment. *Plant, Cell & Environment*. 2014 n/a-n/a.
27. Bishop KA, Leakey ADB, Ainsworth EA. How seasonal temperature or water inputs affect the relative response of C₃ crops to elevated [CO₂]: a global analysis of open top chamber and free air CO₂ enrichment studies. *Food and Energy Security*. 2014; 3(1):33–45.
28. Dow GJ, Bergmann DC. Patterning and processes: how stomatal development defines physiological potential. *Current Opinion in Plant Biology*. 2014; 21:67–74. [PubMed: 25058395]
29. Franks PJ, et al. Sensitivity of plants to changing atmospheric CO₂ concentration: from the geological past to the next century. *New Phytologist*. 2013; 197(4):1077–1094. [PubMed: 23346950]
30. Raschke K. Simultaneous requirement of carbon dioxide and abscisic acid for stomatal closing in *Xanthium strumarium* L. *Planta*. 1975; 125(3):243–259. [PubMed: 24435438]
31. Leymarie J, Vavasseur A, Lascève G. CO₂ sensing in stomata of *abi1-1* and *abi2-1* mutants of *Arabidopsis thaliana*. *Plant Physiology and Biochemistry*. 1998; 36(7):539–543.
32. Xue S, et al. Central functions of bicarbonate in S-type anion channel activation and OST1 protein kinase in CO₂ signal transduction in guard cell. *EMBO J*. 2011; 30(8):1645–1658. [PubMed: 21423149]
33. Merilo E, et al. PYR/RCAR receptors contribute to ozone-, reduced air humidity-, darkness-, and CO₂-induced stomatal regulation. *Plant Physiol*. 2013; 162(3):1652–1668. [PubMed: 23703845]
34. Webb AA, Hetherington AM. Convergence of the abscisic acid, CO₂, and extracellular calcium signal transduction pathways in stomatal guard cells. *Plant Physiol*. 1997; 114(4):1557–1560. [PubMed: 9276963]
35. Chater C, et al. Regulatory Mechanism Controlling Stomatal Behavior Conserved across 400 Million Years of Land Plant Evolution. *Current Biology*. 2011; 21(12):1025–1029. [PubMed: 21658944]
36. Ruszala EM, et al. Land Plants Acquired Active Stomatal Control Early in Their Evolutionary History. *Current Biology*. 2011; 21(12):1030–1035. [PubMed: 21658945]
37. Creese C, et al. Are fern stomatal responses to different stimuli coordinated? Testing responses to light, vapor pressure deficit, and CO₂ for diverse species grown under contrasting irradiances. *New Phytologist*. 2014; 204(1):92–104. [PubMed: 25077933]
38. Waadt R, et al. FRET-based reporters for the direct visualization of abscisic acid concentration changes and distribution in *Arabidopsis*. *Elife*. 2014; 3:e01739. [PubMed: 24737861]
39. Jones AM, et al. Abscisic acid dynamics in roots detected with genetically encoded FRET sensors. *Elife*. 2014; 3:e01741. [PubMed: 24737862]
40. Hu H, et al. Carbonic anhydrases are upstream regulators of CO₂-controlled stomatal movements in guard cells. *Nat Cell Biol*. 2010; 12(1):87–93. [PubMed: 20010812]
41. Moroney JV, Bartlett SG, Samuelsson G. Carbonic anhydrases in plants and algae. *Plant Cell and Environment*. 2001; 24(2):141–153.
42. Frommer WB. CO₂ Common Sense. *Science*. 2010; 327(5963):275–276. [PubMed: 20075235]
43. Tian W, et al. A molecular pathway for CO₂ response in *Arabidopsis* guard cells. *Nat Commun*. 2015; 6:6057. [PubMed: 25599916]
44. Chandrashekar J, et al. The taste of carbonation. *Science*. 2009; 326(5951):443–445. [PubMed: 19833970]
45. Chen YQ, et al. Soluble adenylyl cyclase as an evolutionarily conserved bicarbonate sensor. *Science*. 2000; 289(5479):625–628. [PubMed: 10915626]
46. Klengel T, et al. Fungal adenylyl cyclase integrates CO₂ sensing with cAMP signaling and virulence. *Current Biology*. 2005; 15(22):2021–2026. [PubMed: 16303561]
47. Fabre N, et al. Characterization and expression analysis of genes encoding alpha and beta carbonic anhydrases in *Arabidopsis*. *Plant Cell Environ*. 2007; 30(5):617–629. [PubMed: 17407539]

48. Hu H, et al. Distinct Cellular Locations of Carbonic Anhydrases Mediate CO₂ Control of Stomatal Movements. *Plant Physiol.* 2015
49. Chen W, Levine H, Rappel WJ. A mathematical analysis of second messenger compartmentalization. *Phys Biol.* 2008; 5(4):046006. [PubMed: 19075354]
50. Somersalo E, et al. A reaction-diffusion model of CO₂ influx into an oocyte. *J Theor Biol.* 2012; 309:185–203. [PubMed: 22728674]
51. Azoulay-Shemer T, et al. Guard cell photosynthesis is critical for stomatal turgor production, yet does not directly mediate CO₂- and ABA-induced stomatal closing. *The Plant Journal.* 2015 n/a-n/a.
52. Roelfsema MR, et al. Guard cells in albino leaf patches do not respond to photosynthetically active radiation, but are sensitive to blue light, CO₂ and abscisic acid. *Plant Cell Environ.* 2006; 29(8): 1595–1605. [PubMed: 16898020]
53. Lawson T, et al. Mesophyll photosynthesis and guard cell metabolism impacts on stomatal behaviour. *New Phytol.* 2014; 203(4):1064–1081. [PubMed: 25077787]
54. Lawson T. Guard cell photosynthesis and stomatal function. *New Phytol.* 2009; 181(1):13–34. [PubMed: 19076715]
55. Edwards A, Bowling DJF. Evidence for a CO₂ inhibited proton extrusion pump in the stomatal cells of *Tradescantia virginiana*. *Journal of Experimental Botany.* 1985; 36(162):91–98.
56. Weyers JDB, et al. Guard-Cell Protoplasts - Aspects of Work with an Important New Research Tool. *Physiologia Plantarum.* 1983; 58(3):331–339.
57. von Caemmerer S, et al. Stomatal conductance does not correlate with photosynthetic capacity in transgenic tobacco with reduced amounts of Rubisco. *Journal of Experimental Botany.* 2004; 55(400):1157–1166. [PubMed: 15107451]
58. Young JJ, et al. CO₂ signaling in guard cells: calcium sensitivity response modulation, a Ca²⁺-independent phase, and CO₂ insensitivity of the *gca2* mutant. *Proc Natl Acad Sci U S A.* 2006; 103(19):7506–7511. [PubMed: 16651523]
59. Messinger SM, Buckley TN, Mott KA. Evidence for involvement of photosynthetic processes in the stomatal response to CO₂. *Plant Physiol.* 2006; 140(2):771–778. [PubMed: 16407445]
60. Mott KA, Sibbersen ED, Shope JC. The role of the mesophyll in stomatal responses to light and CO₂. *Plant Cell Environ.* 2008; 31(9):1299–1306. [PubMed: 18541006]
61. Mott KA. Opinion: stomatal responses to light and CO₂ depend on the mesophyll. *Plant Cell Environ.* 2009; 32(11):1479–1486. [PubMed: 19627565]
62. Fujita T, Noguchi K, Terashima I. Apoplastic mesophyll signals induce rapid stomatal responses to CO₂ in *Commelina communis*. *New Phytol.* 2013; 199(2):395–406. [PubMed: 23560389]
63. Hedrich R, Marten I. Malate-induced feedback regulation of plasma membrane anion channels could provide a CO₂ sensor to guard cells. *EMBO J.* 1993; 12(3):897–901. [PubMed: 7681395]
64. Negi J, et al. New approaches to the biology of stomatal guard cells. *Plant Cell Physiol.* 2014; 55(2):241–250. [PubMed: 24104052]
65. Raschke K, Dittlich P. [(14)C]Carbon-dioxide fixation by isolated leaf epidermes with stomata closed or open. *Planta.* 1977; 134(1):69–75. [PubMed: 24419582]
66. von Caemmerer S, et al. Reductions of Rubisco activase by antisense RNA in the C₄ plant *Flaveria bidentis* reduces Rubisco carbamylation and leaf photosynthesis. *Plant Physiol.* 2005; 137(2):747–755. [PubMed: 15665240]
67. Cousins AB, et al. The role of phosphoenolpyruvate carboxylase during C₄ photosynthetic isotope exchange and stomatal conductance. *Plant Physiol.* 2007; 145(3):1006–1017. [PubMed: 17827274]
68. Olsen JE, Junttila O. Far red end-of-day treatment restores wild type-like plant length in hybrid aspen overexpressing phytochrome A. *Physiol Plant.* 2002; 115(3):448–457. [PubMed: 12081538]
69. Roelfsema MR, et al. CO₂ provides an intermediate link in the red light response of guard cells. *Plant J.* 2002; 32(1):65–75. [PubMed: 12366801]
70. Matrosova A, et al. The HT1 protein kinase is essential for red light-induced stomatal opening and genetically interacts with OST1 in red light and CO₂-induced stomatal movement responses. *New Phytologist.* 2015

71. Brearley J, Venis MA, Blatt MR. The effect of elevated CO₂ concentrations on K⁺ and anion channels of *Vicia faba* L. guard cells. *Planta*. 1997; 203(2):145–154.
72. Webb AAA, et al. Carbon dioxide induces increases in guard cell cytosolic free calcium. *Plant J*. 1996; 9(3):297–304.
73. Fitzsimons PJ, Weyers JDB. Separation and Purification of Protoplast Types from *Commelina-Communis* L Leaf Epidermis. *Journal of Experimental Botany*. 1983; 34(138):55–66.
74. Hashimoto M, et al. Arabidopsis HT1 kinase controls stomatal movements in response to CO₂. *Nat Cell Biol*. 2006; 8(4):391–397. [PubMed: 16518390]
75. Negi J, et al. CO₂ regulator SLAC1 and its homologues are essential for anion homeostasis in plant cells. *Nature*. 2008; 452(7186):483–486. [PubMed: 18305482]
76. Vahisalu T, et al. SLAC1 is required for plant guard cell S-type anion channel function in stomatal signalling. *Nature*. 2008; 452(7186):487–491. [PubMed: 18305484]
77. Mustilli AC, et al. Arabidopsis OST1 protein kinase mediates the regulation of stomatal aperture by abscisic acid and acts upstream of reactive oxygen species production. *Plant Cell*. 2002; 14(12):3089–3099. [PubMed: 12468729]
78. Hashimoto-Sugimoto M, et al. A Munc13-like protein in Arabidopsis mediates H⁽⁺⁾-ATPase translocation that is essential for stomatal responses. *Nat Commun*. 2013; 4:2215. [PubMed: 23896897]
79. Meyer S, et al. AtALMT12 represents an R-type anion channel required for stomatal movement in Arabidopsis guard cells. *Plant J*. 2010; 63(6):1054–1062. [PubMed: 20626656]
80. Merlot S, et al. Use of infrared thermal imaging to isolate Arabidopsis mutants defective in stomatal regulation. *Plant J*. 2002; 30(5):601–609. [PubMed: 12047634]
81. Raskin I, Ladyman JA. Isolation and characterization of a barley mutant with abscisic-acid-insensitive stomata. *Planta*. 1988; 173(1):73–78. [PubMed: 24226182]
82. Schwartz A, Ilan N, Grantz DA. Calcium Effects on Stomatal Movement in *Commelina communis* L. : Use of EGTA to Modulate Stomatal Response to Light, KCl and CO₂. *Plant Physiol*. 1988; 87(3):583–587. [PubMed: 16666189]
83. Hubbard KE, et al. Abscisic acid and CO₂ signalling via calcium sensitivity priming in guard cells, new CDPK mutant phenotypes and a method for improved resolution of stomatal stimulus-response analyses. *Ann Bot*. 2012; 109(1):5–17. [PubMed: 21994053]
84. Grabov A, Blatt MR. Membrane voltage initiates Ca²⁺ waves and potentiates Ca²⁺ increases with abscisic acid in stomatal guard cells. *Proc Natl Acad Sci U S A*. 1998; 95(8):4778–4783. [PubMed: 9539815]
85. Klusener B, et al. Convergence of calcium signaling pathways of pathogenic elicitors and abscisic acid in Arabidopsis guard cells. *Plant Physiol*. 2002; 130(4):2152–2163. [PubMed: 12481099]
86. Siegel RS, et al. Calcium elevation-dependent and attenuated resting calcium-dependent abscisic acid induction of stomatal closure and abscisic acid-induced enhancement of calcium sensitivities of S-type anion and inward-rectifying K channels in Arabidopsis guard cells. *Plant J*. 2009; 59(2):207–220. [PubMed: 19302418]
87. Staxen I, et al. Abscisic acid induces oscillations in guard-cell cytosolic free calcium that involve phosphoinositide-specific phospholipase C. *Proc Natl Acad Sci U S A*. 1999; 96(4):1779–1784. [PubMed: 9990101]
88. Chen ZH, et al. Dynamic regulation of guard cell anion channels by cytosolic free Ca²⁺ concentration and protein phosphorylation. *Plant J*. 2010; 61(5):816–825. [PubMed: 20015065]
89. Brandt B, et al. Calcium specificity signaling mechanisms in abscisic acid signal transduction in Arabidopsis guard cells. *Elife*. 2015:4.
90. Basu J, et al. A minimal domain responsible for Munc13 activity. *Nature Structural & Molecular Biology*. 2005; 12(11):1017–1018.
91. Marten H, et al. Silencing of NtMPK4 impairs CO₂-induced stomatal closure, activation of anion channels and cytosolic Casignals in *Nicotiana tabacum* guard cells. *Plant J*. 2008; 55(4):698–708. [PubMed: 18452588]
92. Schroeder JI, Hagiwara S. Cytosolic Calcium Regulates Ion Channels in the Plasma-Membrane of *Vicia-Faba* Guard-Cells. *Nature*. 1989; 338(6214):427–430.

93. Schroeder JI, Keller BU. Two types of anion channel currents in guard cells with distinct voltage regulation. *Proc Natl Acad Sci U S A.* 1992; 89(11):5025–5029. [PubMed: 1375754]
94. Schmidt C, et al. Strong regulation of slow anion channels and abscisic acid signaling in guard cells by phosphorylation and dephosphorylation events. *Proc Natl Acad Sci U S A.* 1995; 92(21): 9535–9539. [PubMed: 11607582]
95. Li J, et al. Regulation of abscisic acid-induced stomatal closure and anion channels by guard cell AAPK kinase. *Science.* 2000; 287(5451):300–303. [PubMed: 10634783]
96. Yoshida R, et al. ABA-activated SnRK2 protein kinase is required for dehydration stress signaling in Arabidopsis. *Plant Cell Physiol.* 2002; 43(12):1473–1483. [PubMed: 12514244]
97. Geiger D, et al. Activity of guard cell anion channel SLAC1 is controlled by drought-stress signaling kinase-phosphatase pair. *Proc Natl Acad Sci U S A.* 2009; 106(50):21425–21430. [PubMed: 19955405]
98. Lee SC, et al. A protein kinase-phosphatase pair interacts with an ion channel to regulate ABA signaling in plant guard cells. *Proc Natl Acad Sci U S A.* 2009; 106(50):21419–21424. [PubMed: 19955427]
99. Geiger D, et al. Guard cell anion channel SLAC1 is regulated by CDPK protein kinases with distinct Ca²⁺ affinities. *Proc Natl Acad Sci U S A.* 2010; 107(17):8023–8028. [PubMed: 20385816]
100. Mori IC, et al. CDPKs CPK6 and CPK3 function in ABA regulation of guard cell S-type anion- and Ca²⁺-permeable channels and stomatal closure. *PLoS Biol.* 2006; 4(10):e327. [PubMed: 17032064]
101. Brandt B, et al. Reconstitution of abscisic acid activation of SLAC1 anion channel by CPK6 and OST1 kinases and branched ABI1 PP2C phosphatase action. *Proc Natl Acad Sci U S A.* 2012
102. Hua D, et al. A plasma membrane receptor kinase, GHR1, mediates abscisic acid- and hydrogen peroxide-regulated stomatal movement in Arabidopsis. *Plant Cell.* 2012; 24(6):2546–2561. [PubMed: 22730405]
103. Hedrich R, et al. Malate-Sensitive Anion Channels Enable Guard-Cells to Sense Changes in the Ambient CO₂ Concentration. *Plant Journal.* 1994; 6(5):741–748.
104. Van Kirk CA, Raschke K. Release of Malate from Epidermal Strips during Stomatal Closure. *Plant Physiol.* 1978; 61(3):474–475. [PubMed: 16660318]
105. Keller BU, Hedrich R, Raschke K. Voltage-Dependent Anion Channels in the Plasma-Membrane of Guard-Cells. *Nature.* 1989; 341(6241):450–453.
106. Sasaki T, et al. Closing plant stomata requires a homolog of an aluminum-activated malate transporter. *Plant Cell Physiol.* 2010; 51(3):354–365. [PubMed: 20154005]
107. Lee M, et al. The ABC transporter AtABC14 is a malate importer and modulates stomatal response to CO₂. *Nat Cell Biol.* 2008; 10(10):1217–1223. [PubMed: 18776898]
108. Kaneda M, et al. ABC transporters coordinately expressed during lignification of Arabidopsis stems include a set of ABCBs associated with auxin transport. *J Exp Bot.* 2011; 62(6):2063–2077. [PubMed: 21239383]
109. Woodward FI. Potential impacts of global elevated CO₂ concentrations on plants. *Curr Opin Plant Biol.* 2002; 5(3):207–211. [PubMed: 11960737]
110. Hetherington AM, Woodward FI. The role of stomata in sensing and driving environmental change. *Nature.* 2003; 424(6951):901–908. [PubMed: 12931178]
111. Woodward FI, Kelly CK. The Influence of CO₂ Concentration on Stomatal Density. *New Phytologist.* 1995; 131(3):311–327.
112. Abrash EB, Lampard GR. A view from the top: new ligands controlling stomatal development in Arabidopsis. *New Phytol.* 2010; 186(3):561–564. [PubMed: 20522161]
113. Pillitteri LJ, Torii KU. Mechanisms of stomatal development. *Annu Rev Plant Biol.* 2012; 63:591–614. [PubMed: 22404473]
114. Nadeau JA, Sack FD. Control of stomatal distribution on the Arabidopsis leaf surface. *Science.* 2002; 296(5573):1697–1700. [PubMed: 12040198]
115. Bergmann DC, Lukowitz W, Somerville CR. Stomatal development and pattern controlled by a MAPKK kinase. *Science.* 2004; 304(5676):1494–1497. [PubMed: 15178800]

116. Hara K, et al. The secretory peptide gene EPF1 enforces the stomatal one-cell-spacing rule. *Genes Dev.* 2007; 21(14):1720–1725. [PubMed: 17639078]
117. MacAlister CA, Ohashi-Ito K, Bergmann DC. Transcription factor control of asymmetric cell divisions that establish the stomatal lineage. *Nature.* 2007; 445(7127):537–540. [PubMed: 17183265]
118. Gray JE. Plant development: three steps for stomata. *Curr Biol.* 2007; 17(6):R213–R215. [PubMed: 17371761]
119. Santrucek J, et al. Stomatal and pavement cell density linked to leaf internal CO₂ concentration. *Annals of Botany.* 2014; 114(2):191–202. [PubMed: 24825295]
120. Gray JE, et al. The HIC signalling pathway links CO₂ perception to stomatal development. *Nature.* 2000; 408(6813):713–716. [PubMed: 11130071]
121. Aharoni A, et al. The SHINE clade of AP2 domain transcription factors activates wax biosynthesis, alters cuticle properties, and confers drought tolerance when overexpressed in *Arabidopsis*. *Plant Cell.* 2004; 16(9):2463–2480. [PubMed: 15319479]
122. Jenks MA, et al. Leaf Epicuticular Waxes of the *Eceriferum* Mutants in *Arabidopsis*. *Plant Physiology.* 1995; 108(1):369–377. [PubMed: 12228482]
123. Post-Beittenmiller D. The cloned *Eceriferum* genes of *Arabidopsis* and the corresponding Glossy genes in maize. *Plant Physiology and Biochemistry.* 1998; 36(1–2):157–166.
124. Lake JA, et al. Plant development. Signals from mature to new leaves. *Nature.* 2001; 411(6834):154. [PubMed: 11346781]
125. Hara K, et al. Epidermal cell density is autoregulated via a secretory peptide, EPIDERMAL PATTERNING FACTOR 2 in *Arabidopsis* leaves. *Plant Cell Physiol.* 2009; 50(6):1019–1031. [PubMed: 19435754]
126. Hunt L, Gray JE. The signaling peptide EPF2 controls asymmetric cell divisions during stomatal development. *Curr Biol.* 2009; 19(10):864–869. [PubMed: 19398336]
127. Torii KU, et al. The *Arabidopsis* ERECTA gene encodes a putative receptor protein kinase with extracellular leucine-rich repeats. *Plant Cell.* 1996; 8(4):735–746. [PubMed: 8624444]
128. Lee JS, et al. Direct interaction of ligand-receptor pairs specifying stomatal patterning. *Genes Dev.* 2012; 26(2):126–136. [PubMed: 22241782]
129. Lee JS, et al. Competitive binding of antagonistic peptides fine-tunes stomatal patterning. *Nature.* 2015; 522(7557) 435–+
130. Masle J, Gilmore SR, Farquhar GD. The ERECTA gene regulates plant transpiration efficiency in *Arabidopsis*. *Nature.* 2005; 436(7052):866–870. [PubMed: 16007076]
131. Shpak ED, et al. Stomatal patterning and differentiation by synergistic interactions of receptor kinases. *Science.* 2005; 309(5732):290–293. [PubMed: 16002616]
132. Doheny-Adams T, et al. Genetic manipulation of stomatal density influences stomatal size, plant growth and tolerance to restricted water supply across a growth carbon dioxide gradient. *Philos Trans R Soc Lond B Biol Sci.* 2012; 367(1588):547–555. [PubMed: 22232766]
133. Lake JA, Woodward FI. Response of stomatal numbers to CO₂ and humidity: control by transpiration rate and abscisic acid. *New Phytol.* 2008; 179(2):397–404. [PubMed: 19086289]
134. Dow GJ, Bergmann DC, Berry JA. An integrated model of stomatal development and leaf physiology. *New Phytologist.* 2014; 201(4):1218–1226. [PubMed: 24251982]
135. Tanaka Y, et al. ABA inhibits entry into stomatal-lineage development in *Arabidopsis* leaves. *Plant Journal.* 2013; 74(3):448–457. [PubMed: 23373882]
136. Coupe SA, et al. Systemic signalling of environmental cues in *Arabidopsis* leaves. *J Exp Bot.* 2006; 57(2):329–341. [PubMed: 16330523]
137. Leonhardt N, et al. Microarray expression analyses of *Arabidopsis* guard cells and isolation of a recessive abscisic acid hypersensitive protein phosphatase 2C mutant. *Plant Cell.* 2004; 16(3):596–615. [PubMed: 14973164]
138. Pillitteri LJ, et al. Molecular Profiling of Stomatal Meristemoids Reveals New Component of Asymmetric Cell Division and Commonalities among Stem Cell Populations in *Arabidopsis*. *Plant Cell.* 2011; 23(9):3260–3275. [PubMed: 21963668]

139. Bauer H, et al. The stomatal response to reduced relative humidity requires guard cell-autonomous ABA synthesis. *Curr Biol.* 2013; 23(1):53–57. [PubMed: 23219726]
140. Pandey S, et al. Boolean modeling of transcriptome data reveals novel modes of heterotrimeric G-protein action. *Molecular Systems Biology.* 2010:6.
141. Hachez C, et al. Differentiation of Arabidopsis guard cells: analysis of the networks incorporating the basic helix-loop-helix transcription factor, FAMA. *Plant Physiol.* 2011; 155(3):1458–1472. [PubMed: 21245191]
142. Lau OS, et al. Direct roles of SPEECHLESS in the specification of stomatal self-renewing cells. *Science.* 2014; 345(6204):1605–1609. [PubMed: 25190717]
143. Franks PJ, et al. Increasing water-use efficiency directly through genetic manipulation of stomatal density. *New Phytologist.* 2015 n/a-n/a.
144. Woodward FI, Lake JA, Quick WP. Stomatal development and CO₂: ecological consequences. *New Phytologist.* 2002; 153(3):477–484.
145. Takahashi S, et al. Natural variation in stomatal responses to environmental changes among Arabidopsis thaliana ecotypes. *PLoS One.* 2015; 10(2):e0117449. [PubMed: 25706630]
146. Meure CM, et al. Law Dome CO₂, CH₄ and N₂O ice core records extended to 2000 years BP. *Geophysical Research Letters.* 2006; 33(14)
147. Bergmann D. Stomatal development: from neighborly to global communication. *Curr Opin Plant Biol.* 2006; 9(5):478–483. [PubMed: 16890476]

<p>Outstanding questions box</p>	<ul style="list-style-type: none"> • Stomatal guard cells simultaneously perceive various cues from the environment, and are sensory organs that process this information in ways that most benefit the plant. Thus for sessile plants, stomata play key roles in avoiding and mitigating environmental and biotic stress. However, many aspects of the molecular mechanisms and network underlying stomatal function remain unknown. A few important questions in stomatal CO₂ responses follow: • How does the chloroplastic localization of carbonic anhydrase1 in guard cells function in transducing the CO₂ stimulus into stomatal movement responses? • These recent findings suggest an important role for guard cell chloroplasts in CO₂ signalling for stomatal movements, that however does not require guard cell photosynthesis. Hence, what are the underlying plastidial mechanisms which transmit the CO₂ signal in guard cells and how do these converge with the cytosolic bicarbonate signaling branch? • What are the guard cell bicarbonate sensors and what is the biochemical protein basis by which intracellular bicarbonate functions as a second messenger in guard cells? • What is the mechanism by which mesophyll cells contribute to stomatal CO₂ responses? • What is the major CO₂ responsive mechanism and what are the transcription factors involved in the CO₂-regulated stomatal development response? • When such necessary advanced information becomes available can we robustly manipulate stomatal gas exchange and plant responses to the continuing rise in atmospheric CO₂ levels?
<p>Trends box</p>	<ul style="list-style-type: none"> • Research on how plant guard cells, which form epidermal stomatal gas exchange valves, regulate the aperture of stomatal pores in response to changes in the carbon dioxide (CO₂) concentration is of current interest given the necessity of selecting and developing crop cultivars which perform better in a shifting global climate. Understanding of the underlying CO₂ response mechanisms is also needed for predictive modelling efforts to better understand plant responses to rising atmospheric CO₂ levels. • Recent discoveries in guard cell CO₂ and secondary messenger signalling, contributions of subcellular localization of the CO₂ binding carbonic anhydrases, interplay with the stress hormone abscisic acid and the role of photosynthesis in stomatal responses to the CO₂ stimulus point to new models and new open questions in CO₂ signal transduction. • Elucidation of the molecular mechanisms controlling stomatal development and identification of initial mechanisms mediating elevated CO₂ repression of stomatal development points to a signaling model and to new avenues for further research on this pathway.




Figure 1. Change of leaf CO₂ concentration (C_i) in response to light and darkness, atmospheric CO₂ rise and stomatal gas exchange response to ambient [CO₂] changes
 (A) Effects of light on [CO₂] (C_i) in the *Vicia faba* substomatal cavity at an external ambient [CO₂] of 350 ppm. Sub-stomatal cavity carbon dioxide concentrations were measured with a potentiometric CO₂-biosensor microprobe inserted into leaves through open stomata [1] (Reprinted with permission from Elsevier publishing group). (B) Carbon dioxide concentrations in the earth's atmosphere [2, 146]. (Reprinted with permission from: https://scripps.ucsd.edu/programs/keelingcurve/wp-content/plugins/sio-blueloon/graphs/co2_800k_zoom.png). (C–E) Effects of CO₂ on gas exchange in carbonic anhydrase mutant leaves. Raw (C and E) and normalized (D) stomatal conductance values for *ca1* and *ca4* single carbonic anhydrase mutants and wild type (WT). Individual ambient CO₂ treatments were each 30 minutes in duration. 1 leaf from each of 3 separate plants per genotype were analyzed over a period of 3 weeks and averaged. Errors represent s.e.m.




Figure 2. CO₂ signaling pathway in stomatal movement regulation and putative convergence points with ABA signaling

(A) A simplified model for carbon dioxide and abscisic acid (ABA) signal transduction pathways in guard cells that mediate stomatal closure. The current model involves the enzymatic function of the carbonic anhydrases CA1 and CA4 with bicarbonate ions as intermediary signaling molecule. Downstream calcium, protein kinases, and ion channels are required for the stomatal closure response to CO₂. Note that amplifying signals from the mesophyll [60, 62] are not shown here for simplicity (see text for details). Different pathway components are color-coded for ease of viewing: ABA genes = green; CO₂ genes = brown; kinases = purple; channels = blue. Abbreviations: ABA = Abscisic acid; PYR/RCAR = ABA receptors; CO₂ = carbon dioxide; CA = carbonic anhydrase; HCO₃⁻ = bicarbonate Ca = calcium; ABI1 = protein phosphatase 2C. (B) The rate of cytosolic calcium transient production in guard cells of the Arabidopsis thaliana Col-0 ecotype are not significantly modulated by the CO₂ concentration, in contrast to the *Ler* ecotype. Cytosolic calcium transients in guard cells pre-exposed to low CO₂, switched to high CO₂, then returned to low CO₂, at the indicated concentrations. Buffer composition was 10 mM KCl/50 mM CaCl₂/10 mM Mes-Tris, pH 6.15. Data are the means of n=27 guard cells ±SE.

CO₂ sensing and signaling during stomatal development




Figure 3. CO₂ control of stomatal development

A current framework [25, 120] includes EPF2, CRSP, the carbonic anhydrases CA1 and CA4 and the cell wall wax biosynthesis mutant *hic* [120]. The secreted extracellular protease CRSP cleaves and activates EPF2, which is involved in extracellular communication resulting in a modulatory repression of stomatal development in response to elevated CO₂. Experiments have shown that the ERECTA receptor kinase has specific EPF2 binding activity [128, 129]. Downstream signaling components include key stomatal fate transcriptional regulators (see: [113, 147]). Signaling components and mechanisms are indicated by question marks (see text for details). Abbreviations: CO₂ = carbon dioxide; CA = carbonic anhydrase; CRSP = CO₂-responsive secreted protease; EPF = epidermal patterning factor; HIC = high carbon dioxide mutant; HCO₃⁻ = bicarbonate.