

UC Irvine

UC Irvine Previously Published Works

Title

Fungus propagules in plastids: the mycosome hypothesis

Permalink

<https://escholarship.org/uc/item/7k5663t4>

Journal

International Microbiology, 6(1)

ISSN

1139-6709

Author

Atsatt, Peter R.

Publication Date

2003-03-13

DOI

10.1007/s10123-003-0103-6

Peer reviewed

Peter R. Atsatt

Fungus propagules in plastids: the mycosome hypothesis

Received: 11 October 2002 / Accepted: 25 November 2002 / Published online: 13 March 2003
© Springer-Verlag and SEM 2003

Abstract A stress-induced “mycosome” phase of *Aureobasidium pullulans* consisting of minute reproductive propagules that may revert directly to walled yeast cells is described. Mycosomes detected by light- and electron-microscopy reproduce within senescent plant plastids, and display three developmental pathways: wall-less cells (protoplasts), yeast cells, or membrane-bounded spherules that harbor plastids. Widespread in plant and algal cells, mycosomes are produced by both ascomycete and basidiomycete fungi.

Keywords *Aureobasidium pullulans* · Fungi · Plastids · Mycosomes · Someangia

Introduction

Minute fungus propagules that I call “mycosomes” are associated with algal and plant plastid development. I hypothesize that certain fungi have a cryptic life-history phase specialized to use the lipids and nitrogen of senescing plastids for their reproduction. Mycosomes are defined as wall-deficient reproductive propagules (diameter ca. 0.1–1.0 μm) that develop into larger forms: wall-less cells (protoplasts), yeast cells, or membrane-bounded thylakoid-containing spherules in which mycosomes reproduce. I discovered mycosomes when yeast cells cultured from the cytoplasm of diverse plants developed from minute precursors. Most of the enlarged unicells produced conidia and mycelia morphologically and physiologically similar to the pleiomorphic ascomycete *Aureobasidium pullulans* (ATCC strain 90393). Various pigmented (black, olive green, yellow brown,

rust brown or hyaline) on glucose-yeast nitrogen base agar [31], all isolates were fermentation-negative, urease-positive and utilized citrate [4] and ascorbate but usually not iso-ascorbate as a sole carbon source [3]. Well known as ubiquitous on aerial plant surfaces [2, 13], *A. pullulans* also has an endobiotic phase [23, 25, 26] that has not been described. Aureobasidin antibiotic production inhibits pathogenic fungi [14, 29]. A dothideaceous loculoascomycete, *A. pullulans* is either a species complex, or an unusually variable asexual state of *Discosphaerina fulvida* [32]. In this study I seek the developmental cycle of mycosomes, including their plastid relationship. If plant-derived mycosomes give rise to *A. pullulans* yeast, then axenic fungal cultures should regenerate mycosomes capable of reversion to *A. pullulans*. I describe *A. pullulans* mycosome life history in vitro, and also some in vivo stages. Mycosome distribution was sought in diverse photosynthetic organisms.

Materials and methods

Organisms and media

Plants

Albica sp., *Cycas revoluta*, *Ginkgo biloba*, *Ornithogalum caudatum*, *Polytrichum* sp., *Sequoia sempervirens* (UC Irvine Arboretum); *Ephedra* sp. (Mohave desert, Calif.); *Phoradendron californicum*, *Cuscuta subinclusa*, *Castilleja stanantha*, marine intertidal *Zostera* sp. (Orange County, Calif.); *Cassytha filiformis* (Oahu, Hawaii); *Pilostyles thurberi* (Texas); *Lycopodium* sp. (Wards Natural Science Biologicals, Rochester, N.Y.).

Algae and cyanobacteria

Culture Collection of Algae at the University of Texas (UTEX): *Anabaena flos-aquae* 1444, *Lyngbya kuetzingii* 1547, *Cylindrocapsa licheniforme* 2014, *Trebouxia erici* 910, *Glaucocystis nostochinearum* 64, *Chlorella kessleri* 2229, *Chlamydomonas reinhardtii* 2337, *Nanochloris eucaryotum* 2502, *Prototheca kruegeri* 329, *Trentepohlia* sp. 1227, *Klebsormidium flaccidum* 321, *Zygnema circumcarinatum* 1559, *Mesotaenium caldariorum* 41, *Spyridia filamentosa* 1508, and *Phaeodactylum tricorneratum* 640; UC Davis

P.R. Atsatt
Department of Ecology and Evolutionary Biology,
University of California, Irvine, CA 92697-2525, USA
E-mail: pratsatt@uci.edu
Tel.: +1-949-4992744
URL: www.mycosomes.info

research collection: *Coleochaete scutata* 262; Wards Natural Science Biologicals: *Mougeotia* sp. W 0280, *Zygnema circumcarinatum* W0900, *Dunaliella salina* W 0171, *Navicula pelliculosa* W 1210, *Tribonema aequale* W 1400; Connecticut Valley Teaching Collection: *Spirogyra* sp., *Chlamydomonas* sp.

Media

Media contained the following components in 100 ml sterile distilled water:

AC	0.1 g ammonium nitrate + C (0.02 g citric acid, pH ca. 4.5)
E	10 ⁻⁶ M ethylene, from 2-chloroethyl-phosphonic acid
ACE	AC + E
ACEG	ACE + G (2 ml glycerol)
ACPc	AC + 0.5% L- α -phosphatidylcholine
B	0.3 g bovine serum albumin (fraction V; Sigma, St. Louis, Mo.)
BG	B + G
BGE	BG + E
BMG	B + M (10 g mannitol) + G
BMGYM	BMG + 2 g Difco YM broth (Difco, Detroit, Mich.)
BAC	B + AC
BCP	B + C + 50 ml prune extract (five sliced prunes boiled in 500 ml water, extract filter-sterilized 0.2 μ m)
MSk	10% Murishige-Skoog minerals + B + 0.3 g CaCl ₂ + 0.6 g MgSO ₄
MSk-2	10% MSk minerals + E + B + 0.1 ml glycerol
YMA	Difco yeast-maltose broth + 2% agar

Media were autoclaved or filter-sterilized as necessary. Gelman cellulose filters of 0.2 μ m-pore diameter were used to attempt mycosome separation from plant and fungus cytoplasm. Media containing plant or fungus cytoplasm was filtered through tandem (5.0 μ m above, 0.2 μ m below), autoclaved 25 mm filter-holders attached to a 200 ml sterile plastic syringe.

Mycosome-phase from plant cytoplasm

Psilotum nudum, a leafless vascular plant, served as a primary experimental host. The mycosome phase develops in *Psilotum* plastid-rich cytoplasm (prc) expelled by glass rod maceration from surface sterilized, water-stressed, post-reproductive stem tissue, and transferred to water or other liquid media.

Experimental techniques

Tissue senescence

I conditioned plant tissue and algal cells in cold water, or acidic AC medium, sometimes containing ethylene (E or ACE), because exogenous ethylene, water stress and prolonged refrigeration promote a sudden rise in plant ethylene production and accompanying senescence symptoms, particularly plastid yellowing [30]. Tissue age and acidic hydration (ca. pH 4.5) are key mycosome-induction factors. Surface-infected young (dark green) *Psilotum* stems do not yield yeast cells, whereas most older (yellowish green, naturally dehiscent sporangia) samples are yeast-positive after acidic hydration. Prc samples from both young and older stem segments usually yield *A. pullulans* prior to surface sterilization, and are negative for fungi when cultured in YM medium immediately after sterilization. When sterilized cohorts of the two age classes are conditioned for 48 h in AC medium before prc samples are added to YM medium, young stems continue to be negative for yeast, whereas most older stem samples yield *A. pullulans* unicells.

Verification of sterility

All work was performed under a laminar-flow hood, with sterile technique. Surface and internal forms of *A. pullulans* are distinguished as follows. Surface organisms are killed by stem submersion for 10 min in 20% Chlorox bleach followed by three washes in sterile distilled water (dH₂O). Post-reproductive *Psilotum* stems with dehiscent sporangia are cut into 2 cm segments and mixed prior to selecting three for each prc sample. Surface sterilization efficacy is verified when prc samples cultured 48 h in liquid yeast-maltose medium (YM; Difco) are negative for fungi. The mycosome phase is induced by conditioning the remaining cohort of sterilized stem segments in dH₂O for 3–7 days in the refrigerator or 48 h in acidic AC medium at room temperature. After conditioning, stem segments are rewashed and macerated in a few drops of dH₂O, and the prc pipetted into a shallow layer of dH₂O (or other medium). Upon settling in Corning 50-mm plastic plates with tight-fitting lids, the chloroplasts form a thin covering over the container bottom. Culture samples monitored by light microscopy are yeast-negative for approximately 24 h. Between 36 and 48 h, a single sample may contain hundreds of small yeast cells, far in excess of numbers that could be generated by rare surface cells that may escape sterilization. Tissue age differences and mycosome mortality from excessive sterilization may produce inconsistent results.

Green plastids were cultured because they retain membrane integrity and respond differently than senescent forms. Prc from green post-reproductive stems refrigerated 7 days in water was cultured in BMGYM on YMA (Fig. 3k, l). To determine whether green plastids could be conditioned after cytoplasm release from stems, a single prc sample from a fresh post-reproductive green stem was subdivided into three treatments: (1) 96 h in BMGYM (non-conditioned control), (2) 96 h in acidic BCP, and (3) 48 h in BCP, followed by 48 h in BMGYM (Fig. 3d–f).

Mycosomes from *A. pullulans* pure cultures

Single-cell clones of *Psilotum*-derived *A. pullulans* were stressed in dH₂O, AC medium, ethylene, antibiotics, dye, or combinations of these, and macerated as per plant cells. To test mycosome longevity, *A. pullulans* cells were inoculated into flasks with dH₂O containing erythromycin, ampicillin, L-canavine or amphoterecin-B at 0.5 mg/ml. After 14 months at room temperature, water samples were observed (Fig. 1k, l) and transferred to ACEG on YMA.

Electron microscopy

Samples were fixed in 4% glutaraldehyde, post-fixed in 2% osmium tetroxide/0.1 M PO₄, and plastic embedded. Thin sections stained in 2% uranyl acetate followed by lead citrate were observed with a Zeiss EM 10 microscope. To stabilize mycosome and yeast cell dissociation from fragmenting senescent plastids during fixation for EM, I cultured prc in modified MSk medium (–CaCl₂/MgSO₄, +0.1 ml glycerol) on glass disks, fixed the material at 24 h, and embedded the surface with agar to prevent movement (Fig. 5g).

Light microscopy

Color images are unstained or stained with aniline blue (AB) in lactophenol for fungi, I₂KI for starch, sudan black (SB) for lipid, or fluorescent 4', 6-diamidino-2-phenylindole (DAPI)-DNA stain, and photographed in transmitted light, phase contrast or UV on a Zeiss Axiophot photomicroscope. Areas of concentrated AB stain appear red under phase contrast.

Mycosome-phase from algal cytoplasm

I sampled three cyanobacteria and 12 algal axenic cultures from UTEX, and a research culture of *Coleochaete scutata*. The UTEX vials were stressed for 6 weeks under Growlux lamps (16-h days);




Fig. 1 Mycosome (ms) developmental stages from pure cultures of *Aureobasidium pullulans* (a–l) and *Psilotum nudum* (m–p). **a** Budding ms in hyphae (left), “lipid bodies” (center) and developing conidia (right) [$\times 2,000$, aniline blue (AB) + Sudan black (SB)]. **b–j** ms and yeast recovered from $0.2\text{-}\mu\text{m}$ filtration: **b** two someangia with numerous ms ($\times 1,250$); **c** densely staining ms (red) that **d** form yeast cells ($\times 2,300$). **e–i** Yeast development within someangia ($\times 3,500$): **e** a yeast within the someangium; **f** a fragmented someangium that produced multiple yeast. Note transition from spherical to ovoid shape; **g, h** ms produce nonstaining forms within a membrane; **i** the central ms develops as a yeast cell, and the someangium constricts (arrows) around the peripheral ms. **j** “Empty” and nonstaining ms ($\times 2,200$). **k–m** Ms bud and enlarge as spherules containing one or more ms ($\times 8,000$). **n** Spherules produce wall-deficient hyphae ($\times 1,250$), or **o** may accumulate lipids ($\times 2,000$, SB+AB). **p** Ms released by narrow constricting filaments enlarge as protoplasts that form conidia ($\times 1,000$). Experimental conditions: **a** *A. pullulans* (Ap) 5 days in AC (for definition of media, see Materials and methods), + 10 days in AC over BMGYM agar; **b–f, j** Ap 14 days in AC, filtered $0.2\text{-}\mu\text{m}$ into ACE on YMA, transferred to fresh ACE on YMA at 5 days; **g–i** Ap 5 days in AC, medium cheesecloth-filtered into ACEG for 48 h; **m–o** *P. nudum* (Pn) plastid-rich cytoplasm (prc) in AC, filtered $0.2\text{-}\mu\text{m}$ onto YMA for 5 days (**m, n**) and 21 days (**o**); **p** Pn 48 h in dH₂O, prc 60 h in BMG on YMA

the cells were then conditioned for 48 h in 10^{-6} M ethylene, macerated with a sterile glass rod, plated in modified MSK medium on agar, and observed from 48 h to 16 days. A second sample of *T. erici* 910 obtained from UTEX 1 year later was macerated in ACPc medium and plated over YMA for nearly 6 months. Non-sterile algal cultures obtained from commercial teaching collections were also senesced, macerated and cultured. Fungal isolates were identified to genus using Yeast-IDent (API, Plainview, N.Y.).

Results

Mycosome-phase from *A. pullulans*

Conditioned in sterile distilled water, AC medium or antibiotics, *A. pullulans* produces internal conidia indistinguishable from external conidia, and minute AB-staining bodies hypothesized to be mycosomes. Mycosome-cycle stages also become prominent when AC-treated cells are transferred to fresh AC over rich (BMGYM) agar. These hyphae develop mycosomes and budding spheres (Fig. 1a, left arrows), apparent lipid bodies (center arrow), and developing conidia (right arrows). The mycosome latency test showed that yeast cells could be recovered from all four water + antibiotic cultures after 14 months at room temperature, and samples examined prior to plating revealed large numbers of mycosomes (Fig. 1k, l).

Mycosomes derived from fungus pure cultures rarely transform to yeast cells in conditioning media. Reversion of mycosomes to yeast may occur within 24 h after transfer of a few drops of these media to YM agar covered with fresh AC medium containing ethylene, phosphatidylcholine or prune extract. Mycosomes from *A. pullulans* (Fig. 1b–i) and plant prc (Fig. 1m–p) pass through $0.2\text{ }\mu\text{m}$ filters, but not in large numbers. Since detection relied on subsequent transformation to yeast (medium-dependent), frequency was not quantified. *A. pullulans* mycosomes filtered ($0.2\text{ }\mu\text{m}$) into ACE on YM agar enlarged to become structures I call

“someangia” (Fig. 1b–d). Someangia are defined as membrane-bounded acytoplasmic protoplasts that produce mycosomes within invaginating membranes. The mycosomes may stain densely (red) with AB (Fig. 1b, c). They become oval and less dense (blue) as they mature (Fig. 1d). A single mycosome may divide within an expanding protoplast membrane (Fig. 1g, h) and later enlarge as a yeast cell (Fig. 1i). The peripheral non-staining mycosomes may be released by constriction of the protoplast membrane (Fig. 1i, arrows). The peripheral mycosomes remain undeveloped (Fig. 1e), or enlarge as yeast cells that cause fragmentation of the someogenic protoplast (Fig. 1f).

Mycosomes also enlarge as distinctive membrane-bounded spherules that appear empty or contain material that does not stain with AB (Fig. 1j). Spherules usually contain a single mycosome (Fig. 1k)—in rare cases two to three—within a central area that becomes non-staining as the budding spherule enlarges (Fig. 1l). The parent spherule displays a densely staining ring-like peripheral cytoplasm (Fig. 1m) that narrows with age. I interpret spherules and someogenic protoplasts to be alternative forms of a common mycosome progenitor. Spherules may produce a single yeast cell (not shown), wall-deficient hyphae (Fig. 1n), or lipid-accumulating forms that bud mycosomes (Fig. 1o). Reticulate narrow filaments partition into linear arrays of independent mycosomes that may enlarge as vacuolate protoplasts that produce internal conidia (Fig. 1p).




Fig. 2a–d Mycosomes within *A. pullulans* cells. **a** Ms (arrows) enlarge as electron-dense bodies within double membrane-bounded organelles called someangia. Note ms are smaller than nuclei (N), ($\times 8,000$). **b** Double membrane (dm) with membrane lamellae and two ms, each with a prominent inclusion (arrow) ($\times 21,000$). **c** Mycosomes develop cell walls internal to the double membranes ($\times 18,000$). **d** Someangia accumulate lipid material and appear as osmiophilic bodies surrounded by an electron-lucent “line” (arrows), the space between the membranes ($\times 8,200$). Experimental conditions: Ap cells from Pn prc 48 h in BMGYM on YMA (**a**, **b**, **d**) and *Cuscuta* cytoplasm 40 h in dH₂O on YMA (**c**)

Mycosome ontogeny

A. pullulans cells from prc produce mycosomes within membrane-rich, double membrane-bounded organelles (Fig. 2a, b). I interpret these as homologs of the someangia I observed in culture (Fig. 1b). Small electron-opaque bodies (Fig. 2a, arrows) appear to originate within the double membrane envelope and to move into the membrane lamellae. They enlarge as spherical mycosomes; some show a heterogeneous electron-opaque inclusion (Fig. 2b, arrow). Single mycosomes bud from the someangium and develop a wall internal to its two bounding membranes (Fig. 2c). The someangia of yeast cultured in YM medium contain lipid and degenerating mycosomes, or appear as osmiophilic bodies (Fig. 2d).


Mycosomes within plastids and plastids within fungus protoplasts

Plastids cultured from prc contain mycosomes. Plastids also occur within fungus protoplasts. Each experimental outcome is sensitive to plant tissue age, conditioning treatment and the medium in which the cytoplasm is placed. Well-senesced plastids observed immediately after stem maceration produce abundant plastoglobuli (Fig. 3a, double arrows). I define two types of plastoglobuli: lipid bodies that develop in the plastid stroma and

contain mostly neutral lipids, and gray spherical bodies with a high glycolipid content that are enclosed by a membrane structurally similar to the outer monolayer of thylakoid membranes [22]. I postulate that the latter contain mycosomes that enlarge as spherules (Fig. 3a, arrow) similar to those from *A. pullulans* (Fig. 1n). Larger forms contain multiple mycosomes (Fig. 3b, arrows) and some of these produce yeast (Fig. 3c).

Although healthy green plastids are typically resistant to mycosome development, green plastids conditioned 48 h in BCP medium, followed by the addition of 50% rich medium for 48 h showed mycosome to yeast transformation ($n = 4$, Fig. 3f). Controls exposed only to acidic BCP (conditioning medium) for 96 h were negative for yeast, but the mature plastids contained many small dense bodies as well as larger spheroidal bodies I interpret to be growth-arrested yeast cells (Fig. 3d, e; arrows). Non-conditioned controls cultured 96 h in rich medium showed no signs of mycosome development, as

Fig. 3a–n Mycosomes develop within plastids that sometimes occur within a fungus membrane or protoplast. **a** Ms enlarge from the plastoglobuli (double arrows) of senescent plastids ($\times 1,000$). **b** Lipid bodies contain ms ($\times 1,000$). **c** A yeast cell from a lipid body that contains ms (arrow) ($\times 2,000$). **d–f** Mycosome-to-yeast transition in green plastids ($\times 3,000$). **g** A *Psilotum* plastid (arrow) within a fungus protoplast ($\times 1,000$). **h** A fungus protoplast extends from the plastid margin ($\times 800$). **i** A plastid enclosed by an AB-staining, budding membrane ($\times 1,200$); **j** A 4', 6-diamidino-2-phenylindole (DAPI)-stained plastid enclosed by blue-fluorescing fungus membranes containing ms-sized bodies with DNA-like fluorescence ($\times 1,500$). **k, l** Spherical or elongate protoplasts develop from the margin of green plastids ($\times 1,800$). **m, n** Yeast develop at the margin of plastid starch ($\times 1,250$). Experimental conditions: **a, b** Pn 24 h dry, 77 h in B, prc observed at maceration; **c** Pn prc 5 days in BGE; **d–f** see Methods; **g–j** as in Fig. 1p; **k** and **l** Pn 6 days cold in dH₂O, prc 48 h in BMGYM; **m** Pn 48 h in AC, prc 5 days in BMGYM; **n** Pn 7 days cold in dH₂O, prc 40 h in ACEG



expected. Rich medium is universally inhibitory prior to acidic conditioning. BCP was apparently a key stimulant for mycosome development within green plastids. In the absence of senescence and *in vivo* conditioning, no chloroplasts occurred within a fungus membrane or protoplast. Individual plastids may have been infected with mycosomes during *in vitro* conditioning. Filter-sterilized (0.2 μm) prune extract cannot be excluded as a mycosome source.

Chloroplasts also developed within AB-staining protoplasts cultured in BMG medium at pH > 6.5. For reasons unknown, these robust conidia-producing forms (Fig. 1p) were not observed after 1986. The protoplasts included 80 \times 130 μm diameter units with senescent lipid-rich plastids and 15–20 conidia, others with green

plastids (Fig. 3g), single plastids with protoplast extensions (Fig. 3h), or single plastids enveloped by a blue-staining membrane with small “buds” (Fig. 3i). Under UV light, plastid thylakoid membranes autofluoresce red or red-orange, and DAPI-stained fungus cells and protoplasts fluoresce blue and show bright spots that are presumably nuclei. Bright DAPI fluorescence was seen within blue membranes that enclose red-orange plastids (Fig. 3j). These figures graphically support my hypothesis that individual plastids develop within a mycosome-containing fungus membrane or protoplast (see also Fig. 5). Conditioned *in vivo* and cultured in rich medium, green plastids produce multiple globose or elongate protrusions (Fig. 3k, l) that may indicate aberrant mycosome development stages. In Fig. 3k, note that the




Fig. 4a–j Mycosomes contain plastid pigments and starch. **a** A fungus protoplast that contains plastids, with ms at its margin ($\times 2,000$, no stain). **b** A similar fungus protoplast whose plastids fluoresce yellow under UV illumination. External ms fluoresce orange. Starch grain at *arrow* ($\times 1,200$); ms stages seen with **c** phase contrast and **d** UV microscopy. Budding yeast precursors show variable fluorescence, whereas yeast do not fluoresce ($\times 2,500$). **e** Mycosomes are green in the center (also **g**), become yellow with increasing size ($\times 3,800$), and (**f**) autofluoresce ($\times 2,000$). **g** Some yeast contain green inclusions assumed to be plastids ($\times 8,000$). **h** Mycosomes containing starch, from macerated *A. pullulans* cultured in prc ($\times 300$). **i** Some starch-producing ms contain an AB-staining body ($\times 2,500$). **j** Starch-producing ms in the process of division, one produces a yeast cell ($\times 1,400$). Experimental conditions: **a–g** Pn 8 days cold in dH₂O, prc 48 h in dH₂O; **h** Pn prc filtered 1.0- μ m in B, plus Ap treated 9 days in B, 24 h development; **i** as in Fig. 3n; **j** Pn 48 h in MSk-2, prc 48 h in MSk-2

membrane associated with the protrusions (arrows) is separate from the chloroplast. Mycosomes were also detected within membranes that envelope iodine-stained starch [33]. Yeast cells develop primarily at starch grain margins, either within barely visible membranes

(Fig. 3m), or within narrow protoplasts that surround a central starch grain (Fig. 3n).

Mycosomes from plastids have plastid traits

Fungus protoplasts that contain senescent plastids (Fig. 4a) autofluoresce yellow under UV illumination (Fig. 4b). The peripheral mycosomes fluoresce orange, an indication of the presence of chlorophyll and/or carotenoids, presumably of plastid origin. Fungus membranes fail to autofluoresce; they appear greenish against the yellow-fluorescing plastids. Compared in phase contrast and UV (Fig. 4c, d), the orange-fluorescing mycosomes may produce fluorescing buds, show weak fluorescence and nonfluorescing buds (inset), or produce nonfluorescing yeast. Mycosomes from fully senesced plastids show green centers when small, they become yellow with increased size (Fig. 4e), and fluoresce orange under UV (Fig. 4f).

Viewed with phase contrast microscopy, the larger mycosomes may contain a single green body. Such green bodies are occasionally seen in yeast cells (Fig. 4g), suggesting that mycosomes that become yeast contain plastids that remain briefly photosynthetic within their walls. The plastid-within-mycosome hypothesis is consistent with the fact that mycosomes often produce reddish-staining starch (Fig. 4h) similar to the amylopectin component [24] of plant starch. Although some fungi do produce a purple-staining amylose-starch, fungus starch lacks amylopectin. Sectioned for EM, the starch-producing mycosomes show a wall-like boundary and an electron-dense center without defining characteristics [33]. Some double-stained mycosomes show a well-developed AB-staining body enclosed by iodine-staining starch (Fig. 4i). Similar mycosomes were recovered from *A. pullulans* pure cultures on two occasions. Yeast cells also develop as division products of starch-producing mycosomes (Fig. 4j, top), illustrating the dual developmental potential of this fungus propagule.

Plastids enclosed inside fungal protoplasts

Plastids cultured from *Psilotum* were often enclosed by a distinctive electron-opaque boundary considered to be a fungus protoplast fused with the outer plastid membrane (Fig. 5a). These contain electron-dense and larger opaque bodies that are indistinguishable from plastoglobuli at low magnification (Fig. 5a). Enlargement may show a membrane-like boundary, heterogeneous content, division-like images and clustered grouping (Fig. 5b, d). Mycosomes are typically associated with thylakoid membranes (Fig. 5c), and many show spherical structures (arrow) that are common elements within derived forms (Fig. 6a, b; arrows). Intermediate stages (Fig. 5e) were infrequent, yet several small walled cells, between 0.5–1.0 μ m in diameter (Fig. 5f), were found associated with fragmenting thylakoid membranes and osmiophilic




Fig. 5a–g Mycosomes and yeast from *Psilotum* plastids, transmission electron microscopy. **a** An electron-opaque protoplast surrounds a plastid that contains numerous ms ($\times 13,000$). **b** Enlargement of inset ms cluster from **a** ($\times 45,500$); **c** Mycosomes associated with plastid thylakoid membranes often show spherical structures (arrow, see also Fig. 6a, b) ($\times 34,600$). **d** Typical ms size range (0.2–0.5 μm) during their division ($\times 32,000$). **d–f** Postulated spherical to ovoid developmental sequence from ca. 0.2 μm (**d**) to 0.40 μm (**e**), to a walled yeast 0.8 μm in length (**f**), (**e**, $\times 80,000$) (**f**, $\times 47,000$). **g** An *A. pullulans* yeast attached to the thylakoid membranes of a senescent amyloplast immobilized during fixation ($\times 11,100$). Experimental conditions: **a–f** Pn 24 h dry, 24 h in dH_2O , prc 45 h in BMGYM; **g** see Methods

bodies. In an experiment designed specifically to prevent movement during fixation, agar embedding of prc cultured on glass disks produced the profile in Fig. 5g, one of several yeast cells found attached to the thylakoid membranes of degenerating plastids with starch.


Cultured plastids show predominantly electron-dense mycosomes (Fig. 6a, b) containing heterogeneous spheroid areas defined by lighter density, plus spherical structures (arrows). These mycosomes may divide to produce ring-like forms (Fig. 6b) I identify as plastids that contain mycosomes (pcm): pcm also enlarge from electron-opaque bodies $< 0.2 \mu\text{m}$ in diameter, situated within the parent plastid envelope or, more frequently, associated with thylakoid membranes; pcm show central fragments and strands that emanate from a narrow peripheral cytoplasm (Fig. 6b, c); pcm in the envelope may be enclosed by a fungus membrane fused with the outer plastid membrane (Fig. 6c, arrow); pcm are also produced within protoplast extensions (Fig. 6d, asterisks) that are continuous with the outer plastid membrane.

The pcm “envelope” contains numerous electron opaque units, one or more of which enlarge as plastoglobuli (Fig. 6d, arrows) that are indistinguishable from those within the parent plastid. Independent pcm released from *Psilotum* plastids contain thylakoid membranes and mycosomes (Fig. 6e, f), or may be packed exclusively with mycosomes that appear to divide inward from the envelope. The large mycosome within the pcm seen in Fig. 6f is indistinguishable from yeast that develop abnormally (Fig. 6g); pcm genesis from minute electron opaque bodies may also occur in small protoplasts that originate from plastid thylakoid membranes (Fig. 6h). The plasma membrane-associated bodies (Figs. 6h, i, arrows) enlarge as pcm, identified by their electron-opaque membrane-bounded bodies (Fig. 6i, double arrow). The pcm bodies (Fig. 6j, arrows) apparently divide inward and enlarge as plastoglobuli associated with thylakoid membranes (Fig. 6j, right). Small pcm about 1.0 μm in diameter may produce single starch grains as well as thylakoid membranes.

Mycosome phase from algal monocultures

Trebouxia erici, *Coleochaete scutata* and *Phaeodactylum tricoratum* extracts yielded basidiomycete yeast (Table 1). *Cryptococcus* and *Rhodotorula* species are known to live inside plants [16, 25]. The transition from mycosome to yeast was documented (light micrographs) in cultures of both *T. erici* and *C. scutata*. Although the remaining UTEX cultures were yeast-negative, it is significant that typical mycosome stages were present in most cases.

Fig. 6 *Psilotum* plastids produce plastids containing ms (pcm) (a–f) and fungus protoplasts that also produce pcm (h–j). **a** A ms with a spherical structure (arrow, also **b**) and lighter gray areas, attached to thylakoids ($\times 12,000$). **b** A dividing pcm that came from a thylakoid-associated mycosome ($\times 21,000$). **c** A pcm within a plastid envelope (st starch, $\times 32,000$). **d** pcm (asterisks) with plastoglobuli (arrows), within a protoplast extension continuous with the outer plastid membrane ($\times 14,000$). **e** A pcm with thylakoid membranes and a membrane-bounded ms ($\times 26,000$). **f** A pcm ($\times 11,000$) contains an asymmetric ms indistinguishable from a yeast cell (**g**, $\times 12,000$) that developed abnormally. **h** A fungal protoplast with plasma membrane-associated bodies (arrows), from a *Psilotum* plastid ($\times 56,000$). **i** A fungal protoplast with bodies (arrows) that enlarge as pcm (double arrows, $\times 24,000$); **j** Maturing pcm with membrane-bounded ms (arrows) that divide into the stroma and enlarge as plastoglobuli ($\times 21,000$). Experimental conditions: Pn 72 h in dH₂O, prc 46 h in BMGYM



Cyanobacteria (*Anabaena flos-aquae* and *Cylindrospermum licheniforme*) produced mycosomes and spherules at the margins of senesced cells after 5 months, similar in all respects to forms derived from plants. Mycosomes that stained with AB were abundant in the periplasm of senescent *Glaucozystis* cells and some plastids (cyanelles) budded similar bodies from narrowed extensions. *Prototheca*, a colorless alga, yielded typical mycosome-containing membranes, and large AB-staining spherules clustered along thin filaments. *Trentepohlia* plastids accumulate a bright yellow β -carotene that provides a distinctive plastid pigment marker. Extract from this lichen alga produced reticulate membranes with mycosomes enlarging as bright yellow, dark-centered spherules, consistent with plant observations that mycosomes contain pigmented lipids similar to the plastids of their host organism.

Extract from *T. erici* 910 in ACPc medium produced an unusually compact, slow-growing mound of black filaments apparent only after 22 weeks of culture. *T. erici* 910 is a lichen alga, isolated from *Cladonia cristatella* by V. Ahmadjian (Clark University, Worces-

ter, Mass.) and contributed to UTEX in 1975. Ahmadjian identified my fungus isolate as a variant of his original lichen fungus from *C. cristatella*.

In contrast to UTEX axenic cultures, yeast were readily isolated from several non-sterile algal cultures. Because algae cannot be surface sterilized, only yeast that associated with mycosomes are listed in Table 1. Mycosomes were observed within senescent algal cells in cases marked with an a.

Discussion

Plant-derived mycosomes transform to *A. pullulans* yeast capable of regenerating mycosomes that revert back to the typical walled phase. More complex than miniature yeast cells, mycosomes are membrane-bounded fungal propagules that develop as walled cells, mycosome-producing protoplasts (someangia), or spherules with variable content. In *A. pullulans* pure cultures, the spherule may remain empty or develop one or more mycosomes. In contrast, spherules from senescent

Table 1 Photosynthetic organisms that yielded fungi associated with one or more phases of mycosome life history. *Ap* *Aureobasidium pullulans*, *Ca* *Candida* sp., *Cr* *Cryptococcus* sp., –ID unidentified but not *Ap*, *Lf* lichen-fungus, *Rh* *Rhodotorula* sp., *Tr* *Trichosporon* sp.

Plants	Fungus	Algae	Fungus
<i>Cuscuta subinclusa</i>	Ap	<i>Coleochaete</i> UCD 262 ^a	Rh
<i>Castilleja stanantha</i>	Ap	<i>Trebouxia</i> UTEX 910 ^a	Cr
<i>Cassytha filiformis</i>	–ID	<i>Trebouxia</i> UTEX 910	Lf
<i>Phoradendron californicum</i>	Ap	<i>Phaeodactylum</i> UTEX 640	Tr
<i>Pilostyles thurberi</i>	–ID	<i>Mougeotia</i> sp. ^a	Ca
<i>Ornithogalum caudatum</i>	Ap	<i>Spirogyra</i> sp. ^a	–ID
<i>Albica</i> sp.	Ap	<i>Zygnema circumcarinatum</i> ^a	Tr
<i>Zostera</i> sp.	Ap	<i>Dunaliella salina</i>	Rh
<i>Sequoia sempervirens</i>	Ap	<i>Chlamydomonas</i> sp.	Rh
<i>Ephedra</i> sp.	Ap	<i>Navicula pelliculosa</i>	Ca
<i>Ginkgo biloba</i>	AP	<i>Tribonema aequale</i>	Rh
<i>Cycas revoluta</i>	Ap		
<i>Polytrichum</i> sp.	Ap		
<i>Psilotum nudum</i>	Ap		
<i>Psilotum nudum</i>	Rh		
<i>Lycopodium</i> sp.	Ap		

^aMycosomes were observed within algal cells

plastids and plastid-containing fungus protoplasts may contain a central green body, plastid pigments, or starch with high amylopectin content.

Mycosomes develop inside plant plastids that have been incorporated into a wall-less stage of the fungus. Proplastids may be enclosed inside fungus protoplasts, but it is not clear how these organelles acquire mycosomes. Mature chloroplasts that contain mycosomes produce yeast cells, protoplasts and pcm; pcm develop from small electron-dense bodies and may be homologous to spherules observed with light microscopy. The presence of plastids within mycosomes can be confirmed with molecular probes, as can the affinity of the DNA-containing mycosomes.

Mycosomes that develop into internal conidia are comparable to those that form without wall layers in continuity between the conidium and the parent cell (e.g., endogenous conidia). While endogenous conidia are unknown in ascomycetes, since meiotic ascospores that develop within the double membrane ascus vesicle [15] are typically endogenous [10], the genetic potential for the double membrane someangium is likely to be present in mycosome-forming fungi. Mycosome discovery has likely been obstructed by fungus culture under nutrient-rich (laboratory) conditions. Internal conidia are rarely reported for *A. pullulans* [11, 18], but “lipid bodies” are commonly seen. The *A. pullulans* someangium becomes densely osmiophilic on typical nutrient agar (Fig. 2d), masking the mycosomes within. Similar electron-opaque bodies noted in several dimorphic fungal pathogens [1, 6, 7, 8, 9] may be someangia.

Mycosome-producing fungi are probably widespread. The causative agent of Dutch elm disease, *Ophiostoma ulmi*, produces “microendospores” [19] that pass

through 0.45- and 0.22- μ m Millipore filters. Injected into trees, these clear filtrates produced typical disease symptoms, even several months later [20]. *Candida* species consistently produce viable units filterable through Seitz sterilizing pads, sintered glass filters, or Gelman cellulose filters of 0.2- μ m pore diameter [17]. The microhyphae (minimal diameter 0.2 μ m) of *Fusarium oxysporum* f. sp. *dianthi* [21] may be mycosome stages. Mycosome ontogeny to mature ascomycete and basidiomycete yeast should be induced in vivo, and described in detail by comparative transmission electron microscopy studies. In situ hybridization with rRNA targeted probes [13, 27], new imaging techniques [12] and DNA probes should identify mycosomes and their filamentous phase within plant cells and tissue. Fungus-specific nuclear markers [5, 28] are predicted to demonstrate nuclei in mycosomes and to track mycosome behavior in plastids with multiple nucleoids.

Acknowledgements I thank Steve Weller for use of his photomicroscope, Sue Fisher for fixing and sectioning TEM material, and V. Ahmadjian for identifying the *T. erici* fungus. Encouragement and editing by Lynn Margulis helped distill 10 years of research into this short version. Micah S. Dunthorn, Richard Mullins, Art Weis, Steve Frank, Robin Bush and anonymous reviewers also improved the manuscript. Questions underlying this work arose from parasitic plant studies supported by the National Science Foundation. I am grateful to the Whitehall Foundation for funding this exploratory research.

References

1. Chamberland H, Ouellette GB (1977) Formes d'inclusions osmiophiles dans les cellules de *Ceratocystis ulmi*. *Can J Bot* 55:695–710
2. Cooke WB (1959) An ecological life history of *Aureobasidium pullulans* (de Barry) Arnaud. *Mycopathol Mycol Appl* 12:1–45
3. Costamagna L, Rosi I, Garuccio I, Arrigoni O (1986) Ascorbic acid utilization by some yeasts. *Can J Microbiol* 32:756–758
4. De Hoog GS, Yurlova NA (1994) Conidiogenesis, nutritional physiology and taxonomy of *Aureobasidium* and *Hormonema*. *Antonie van Leeuwenhoek* 65:41–54
5. Fernandez-Abalos JM, Fox H, Pitt C, Wells B, Doonan JH (1998) Plant-adapted green fluorescent protein is a versatile vital reporter for gene expression, protein localization and mitosis in the filamentous fungus, *Aspergillus nidulans*. *Mol Microbiol* 27:121–130
6. Garrison RG (1985) Cytological and ultrastructural aspects of dimorphism. In: Szaniszló PJ, Harris JL (eds) *Fungal dimorphism, with emphasis on fungi pathogenic for humans*. Plenum Press, New York, pp 15–47
7. Garrison RG, Boyd KS (1978) Role of the conidium in dimorphism of *Blastomyces dermatitidis*. *Mycopathologia* 64:29–33
8. Groove SN, Oujecdsky KB, Szaniszló PJ (1973) Budding in the dimorphic fungus *Phialophora dermatitidis*. *J Bacteriol* 81:522–527
9. Gunasekaran M, Hess WM, Weber DJ (1972) Ultrastructure and lipid changes in *Pyrenochaeta terrestris* during aging. *Can J Microbiol* 19:491–496
10. Hennebert GL, Sutton BC (1994) Unitary parameters in conidiogenesis. In: Hawksworth DL (ed) *Ascomycete systematics, problems and perspectives in the nineties*. Plenum Press, London, pp 65–76
11. Hermanides-Nijhof EJ (1984) *Aureobasidium* and allied genera. *Stud in Mycol (Baarn)* 15:141–177

12. Howard RJ (2001) Cytology of fungal pathogens and plant-host interactions. *Curr Opin Microbiol* 4:365–373
13. Li S, Spear RN, Andrews JH (1997) Quantitative fluorescence in situ hybridization of *Aureobasidium pullulans* on microscope slides and leaf surfaces. *Appl Environ Microbiol* 63:3261–3267
14. Lima G, Arru S, De Curtis F, Arras G (1999) Influence of antagonist, host fruit and pathogen on the biological control of postharvest fungal diseases by yeasts. *J Ind Microbiol Biotechnol* 23:223–229
15. Madelin MF (1981) Ultrastructural morphogenesis in higher fungi. In: Turian G, Hohl HR (eds) *The fungal spore: morphogenetic controls*. Academic Press, London, pp 95–106
16. Marquarding I (2000) Endophytische Pilze aus *Trifolium repens* L. mit besonderer Berücksichtigung isolierter Hefen und deren taxonomischer Einordnung. *Bibl Mycol* 184:1–143
17. Mattman LH (1992) *Cell wall deficient forms*. CRC Press, Boca Raton, Fla., pp 239–248
18. Meyers SP, Ahearn DG, Cook WL (1970) Mycological studies of Lake Champlain. *Mycologia* 62:504–515
19. Ouellette GB, Gagnon C (1960) Formation of microendospores in *Ceratocystis ulmi* (Buism.) C. Moreau. *Can J Bot* 38:235–241
20. Ouellette GB, Methot N, Chamberland H, Cote C, Lafontaine J-G (1995) Cytology of irregular growth forms of *Ophiostoma ulmi* and *Ophiostoma novo-ulmi* growing through millipore filter membranes and sterilized elm wood sections. *Can J Microbiol* 41:1095–1110
21. Ouellette GB, Baayen RP, Simard M, Rioux D (1999) Ultrastructural and cytochemical study of colonization of xylem vessel elements of susceptible and resistant *Dianthus caryophyllus* by *Fusarium oxysporum* f.sp. *dianthi*. *Can J Bot* 77:644–663
22. Picher M, Grenier G, Purcell M, Proteau L, Beaumont G (1993) Isolation and purification of intralamellar vesicles from *Lemma minor* L. chloroplasts. *New Phytol* 123:657–663
23. Pugh GJF, Buckley NG (1971) *Aureobasidium pullulans*: an endophyte in sycamore and other trees. *Trans Br Mycol Soc* 57:227–231
24. Salehuzzaman SNIM, Jacobsen E, Visser RGF (1993) Isolation and characterization of a cDNA encoding granule-bound starch synthase in cassava (*Manihot esculenta* Crantz) and its antisense expression in potato. *Plant Mol Biol* 23:947–962
25. Schweigkofler W, Prillinger H (1997) Analysis of endophytic and latent pathogenic fungi isolated from wooden parts of the grapevine *Vitis vinifera* in Austria and the South Tyrol. *Mitt Klosterneuburg* 47:149–158
26. Smith H, Wingfield MJ, Petrini O (1996) *Botryosphaeria dothidea* in *Eucalyptus grandis* and *Eucalyptus nitens* in South Africa. *For Ecol Manage* 89:189–195
27. Sterflinger K, Hain M (1999) In situ hybridization with rRNA targeted probes as a new tool for the detection of black yeast and meristematic fungi. *Stud Mycol* 43:23–30
28. Suelmann R, Sievers N, Fischer R (1997) Nuclear traffic in fungal hyphae: in vivo study of nuclear migration and positioning in *Aspergillus nidulans*. *Mol Microbiol* 25:757–769
29. Takesako K, Kuroda H, Inoue T, Haruna F, Yoshikawa Y, Kato I, Uchida K, Hiratani T, Yamaguchi H (1993) Biological properties of aureobasidin A, a cyclic depsipeptide antifungal antibiotic. *J Antibiot* 46:1414–1420
30. Woltering EJ, Harren F (1989) Early changes in ethylene production during senescence of carnation and *Phalenopsis* flowers measured by laser photoacoustic detection. In: Clijsters H, De Proft M, Marcelle R, Van Poucke M (eds) *Biochemical and physiological aspects of ethylene production in lower and higher plants*. Kluwer, Dordrecht, pp 263–270
31. Wickerham LJ, Kurtzman CP (1975) Synergistic color variants of *Aureobasidium pullulans*. *Mycologia* 67:342–361
32. Yurlova NA, de Hoog GS, Gerrits van den Ende AHG (1999) Taxonomy of *Aureobasidium* and allied genera. *Stud Mycol* 43:63–69
33. Electronic supplementary material: <http://www.mycosomes.info>