

# UC Merced

## UC Merced Previously Published Works

### Title

Peatland Acidobacteria with a dissimilatory sulfur metabolism

### Permalink

<https://escholarship.org/uc/item/7k67g7ks>

### Journal

The ISME Journal: Multidisciplinary Journal of Microbial Ecology, 12(7)

### ISSN

1751-7362

### Authors

Hausmann, Bela

Pelikan, Claus

Herbold, Craig W

et al.

### Publication Date

2018-07-01

### DOI

10.1038/s41396-018-0077-1

Peer reviewed

1 **Peatland *Acidobacteria***  
2 **with a dissimilatory sulfur metabolism**

3 Bela Hausmann <sup>1,2</sup>, Claus Pelikan <sup>1</sup>, Craig W. Herbold <sup>1</sup>, Stephan Köstlbacher <sup>1</sup>,  
4 Mads Albertsen <sup>3</sup>, Stephanie A. Eichorst <sup>1</sup>, Tijana Glavina del Rio <sup>4</sup>, Martin Huemer <sup>1</sup>,  
5 Per H. Nielsen <sup>3</sup>, Thomas Ratten <sup>5</sup>, Ulrich Stingl <sup>6</sup>, Susannah G. Tringe <sup>4</sup>, Daniela Trojan <sup>1</sup>,  
6 Cecilia Wentrup <sup>1</sup>, Dagmar Wobken <sup>1</sup>, Michael Pester <sup>2,7</sup>, and Alexander Loy <sup>1</sup>

7 <sup>1</sup> University of Vienna, Research Network Chemistry meets Microbiology, Department of Microbiology and Ecosystem  
8 Science, Division of Microbial Ecology, Vienna, Austria

9 <sup>2</sup> University of Konstanz, Department of Biology, Konstanz, Germany

10 <sup>3</sup> Aalborg University, Department of Chemistry and Bioscience, Center for Microbial Communities, Aalborg, Denmark

11 <sup>4</sup> US Department of Energy Joint Genome Institute, Walnut Creek, CA, USA

12 <sup>5</sup> University of Vienna, Research Network Chemistry meets Microbiology, Department of Microbiology and Ecosystem  
13 Science, Division of Computational Systems Biology, Vienna, Austria

14 <sup>6</sup> University of Florida, Fort Lauderdale Research and Education Center, UF/IFAS, Department for Microbiology and Cell  
15 Science, Davie, FL, USA

16 <sup>7</sup> Leibniz Institute DSMZ, Braunschweig, Germany

17 Correspondence: Michael Pester, Leibniz Institute DSMZ, Inhoffenstraße 7B, 38124 Braunschweig, Germany. Phone:  
18 +49 531 2616 237. Fax: +49 531 2616 418. E-mail: [michael.pesther@dsmz.de](mailto:michael.pesther@dsmz.de)

19

20

## 21 **Abstract**

22 Sulfur-cycling microorganisms impact organic matter decomposition in wetlands and consequently greenhouse gas emissions  
23 from these globally relevant environments. However, their identities and physiological properties are largely unknown. By  
24 applying a functional metagenomics approach to an acidic peatland, we recovered draft genomes of seven novel *Acidobacteria*  
25 species with the potential for dissimilatory sulfite (*dsrAB*, *dsrC*, *dsrD*, *dsrN*, *dsrT*, *dsrMKJOP*) or sulfate respiration (*sat*,  
26 *aprBA*, *qmoABC* plus *dsr* genes). Surprisingly, the genomes also encoded DsrL, which so far was only found in sulfur-  
27 oxidizing microorganisms. Metatranscriptome analysis demonstrated expression of acidobacterial sulfur-metabolism genes in  
28 native peat soil and their upregulation in diverse anoxic microcosms. This indicated an active sulfate respiration pathway,  
29 which, however, might also operate in reverse for dissimilatory sulfur oxidation or disproportionation as proposed for the  
30 sulfur-oxidizing *Desulfurivibrio alkaliphilus*. *Acidobacteria* that only harbored genes for sulfite reduction additionally encoded  
31 enzymes that liberate sulfite from organosulfonates, which suggested organic sulfur compounds as complementary energy  
32 sources. Further metabolic potentials included polysaccharide hydrolysis and sugar utilization, aerobic respiration, several  
33 fermentative capabilities, and hydrogen oxidation. Our findings extend both, the known physiological and genetic properties of  
34 *Acidobacteria* and the known taxonomic diversity of microorganisms with a DsrAB-based sulfur metabolism, and highlight  
35 new fundamental niches for facultative anaerobic *Acidobacteria* in wetlands based on exploitation of inorganic and organic  
36 sulfur molecules for energy conservation.

37

38

## 39 Introduction

40 Specialized microorganisms oxidize, reduce, or disproportionate sulfur compounds of various oxidation states (–II to +VI) to  
41 generate energy for cellular activity and growth and thereby drive the global sulfur cycle. The capability for characteristic  
42 sulfur redox reactions such as dissimilatory sulfate reduction or sulfide oxidation is not confined to single taxa but distributed  
43 across different, often unrelated taxa. The true extent of the taxon-diversity within the different guilds of sulfur  
44 microorganisms is unknown (Wasmund *et al.*, 2017). However, ecological studies employing specific sulfur metabolism genes  
45 (e.g., dissimilatory adenylyl-sulfate reductase-encoding *aprBA*, dissimilatory sulfite reductase-encoding *dsrAB*, or *soxB* that  
46 codes for a part of the thiosulfate-oxidizing Sox enzyme machinery) as phylogenetic and functional markers have repeatedly  
47 demonstrated that only a minor fraction of the sulfur metabolism gene diversity in many environments can be linked to  
48 recognized taxa (Meyer *et al.*, 2007; Müller *et al.*, 2015; Watanabe *et al.*, 2016). A systematic review of *dsrAB* diversity has  
49 revealed that the reductive bacterial-type enzyme branch of the DsrAB tree contains at least thirteen family-level lineages  
50 without any cultivated representatives. This indicates that major taxa of sulfate-/sulfite-reducing microorganisms have not yet  
51 been identified (Müller *et al.*, 2015).

52 Wetlands are among those ecosystems that harbor a diverse community of microorganisms with reductive-type DsrAB, most  
53 of which cannot be identified because they are distant from taxonomically classified DsrAB sequences (Pester *et al.*, 2012).  
54 Sulfur-cycling microorganisms provide significant ecosystem services in natural and anthropogenic wetlands, which are major  
55 sources of the climate-warming greenhouse gas methane (Kirschke *et al.*, 2013; Saunio *et al.*, 2016). While inorganic sulfur  
56 compounds are often detected only at low concentration (lower  $\mu\text{M}$  range), fast sulfur cycling nevertheless ensures that  
57 oxidized sulfur compounds such as sulfate are rapidly replenished for anaerobic respiration. The activity of sulfate-reducing  
58 microorganisms (SRM) fluctuates with time and space, but at peak times can contribute considerably to the anaerobic  
59 mineralization of organic carbon in wetlands (Pester *et al.*, 2012). Simultaneously, SRM prevent methane production by  
60 rerouting carbon flow away from methanogenic archaea. Peat microorganisms that are affiliated to known SRM taxa, such as  
61 *Desulfosporosinus*, *Desulfomonile*, and *Syntrophobacter*, are typically found in low abundance (Loy *et al.*, 2004; Costello and  
62 Schmidt, 2006; Dedysh *et al.*, 2006; Kraigher *et al.*, 2006; Pester *et al.*, 2010; Steger *et al.*, 2011; Tveit *et al.*, 2013; Hausmann  
63 *et al.*, 2016). In contrast, some microorganisms that belong to novel, environmental *dsrAB* lineages can be considerably more  
64 abundant in wetlands than species-level *dsrAB* operational taxonomic units of known taxa (Steger *et al.*, 2011). However, the  
65 taxonomic identity of these novel *dsrAB*-containing microorganisms and their role in sulfur and carbon cycling has yet to be  
66 revealed.

67 To identify these unknown DsrAB-encoding organisms and further investigate their fundamental ecological niches, we  
68 recovered thirteen metagenome-assembled genomes (MAGs) encoding DsrAB from a peat soil through a targeted, functional  
69 metagenomics approach. We analyzed expression of predicted physiological capabilities of the MAGs by metatranscriptome  
70 analyses of anoxic peat soil microcosms that were periodically stimulated by small additions of individual fermentation  
71 products with or without supplemented sulfate (Hausmann *et al.*, 2016). Here, we show that facultatively anaerobic members  
72 of the diverse *Acidobacteria* community in wetlands employ one or more types of dissimilatory sulfur metabolism.

73

74

## 75 **Materials and methods**

### 76 **Anoxic microcosm experiments, stable isotope probing, and nucleic acids isolation**

77 DNA and RNA samples were retrieved from a previous peat soil microcosm experiment (Hausmann *et al.*, 2016). Briefly, soil  
78 from 10–20 cm depth was sampled from an acidic peatland (Schlöppnerbrunnen II, Germany) in September 2010, and stored at  
79 4 °C for one week prior to nucleic acids extractions and set-up of soil slurry incubations. Individual soil slurry microcosms  
80 were incubated anoxically (100% N<sub>2</sub> atmosphere) in the dark at 14 °C, and regularly amended with low amounts (<200 μM)  
81 of either formate, acetate, propionate, lactate, butyrate, or without any additional carbon sources (six replicates each). In  
82 addition, half of the microcosms for each substrate were periodically supplemented with low amounts of sulfate (initial  
83 amendment of 190–387 μM with periodic additions of 79–161 μM final concentrations). DNA and RNA were extracted from  
84 the native soil and RNA was additionally extracted from the soil slurries after 8 and 36 days of incubation.

85 Furthermore, DNA was obtained from the heavy, <sup>13</sup>C-enriched DNA fractions of a previous DNA-stable isotope probing  
86 (DNA-SIP) experiment with soil from the same site (Pester *et al.*, 2010). Analogous to the single-substrate incubations, anoxic  
87 soil slurries were incubated for two months with low-amounts of sulfate and a <sup>13</sup>C-labelled mixture of formate, acetate,  
88 propionate, and lactate. DNA was extracted, separated on eight replicated density gradients, and DNA from a total of 16 heavy  
89 fractions (density 1.715–1.726 g mL<sup>-1</sup>) was pooled for sequencing.

90 Additional DNA was obtained from soils that were sampled from different depths in the years 2004 and 2007 (Steger *et al.*,  
91 2011).

### 92 **Quantitative PCR and metagenome/-transcriptome sequencing**

93 Abundances of *Acidobacteria* subdivision 1, 2, and 3 in soil samples from different years and depths were determined by  
94 newly-developed 16S rRNA gene-targeted real-time quantitative PCR (qPCR) assays (Supplementary Methods). Native soil  
95 DNA (two libraries), heavy <sup>13</sup>C-enriched DNA (three libraries), and native soil RNA, and RNA samples from the microcosms  
96 were sequenced on an Illumina HiSeq2000 system (Supplementary Methods).

### 97 **Binning, phylogeny, and annotation of DsrAB-encoding genomes**

98 The differential coverage binning approach by Albertsen *et al.* (2013) was applied to extract MAGs of interest. The raw  
99 FASTQ paired-end reads were imported into the CLC Genomics Workbench 5.5.1 (CLC Bio) and trimmed using a minimum  
100 Phred quality score of 20 with no ambiguous nucleotides allowed. TruSeq adapters were removed and a minimum length filter  
101 of 50 nt was applied. This resulted in 214, 171, 233, 49, and 294 million reads after quality filtering and trimming for the two  
102 native soil and three SIP metagenomes, respectively (84–95% of the raw reads). All reads were co-assembled using CLCs *de*  
103 *novo* assembly algorithm (kmer size 63, bubble size 50, minimum scaffold size 1000 nt). The reads from all five metagenomes  
104 were independently mapped to the assembled scaffolds using CLCs map to reference function (minimum length 0.7, minimum  
105 similarity 0.95) to obtain the scaffold coverage. The SIP metagenomes were merged into one mapping. 137, 112, and 376  
106 million reads could be mapped to the two native soil metagenomes and the SIP metagenome, respectively (64–66% of quality  
107 filtered reads). Gene prediction of the complete assembly was performed using prodigal (Hyatt *et al.*, 2010). In addition to the  
108 detection and taxonomic classification of 105 essential marker genes (Albertsen *et al.*, 2013), *dsrA* and *dsrB* homologs were  
109 identified using TIGRFAM's hidden Markov model (HMM) profiles TIGR02064 and TIGR02066, respectively, with  
110 HMMER 3.1 (Eddy, 2011) and the provided trusted cut-offs. Additional *dsrAB*-containing scaffolds were identified by using  
111 tblastn with the published DsrAB database as a query against the assembly (Müller *et al.*, 2015). DsrAB sequences were

112 classified by phylogenetic analysis (Supplementary Methods; Müller *et al.*, 2015). Binning and decontamination was finalized  
113 utilizing the G+C content and tetramer frequencies of the scaffolds, as well as paired-end information, as described and  
114 recommended in Albertsen *et al.* (2013). Completeness, contamination, and strain heterogeneity was estimated using CheckM  
115 1.0.6 (Parks *et al.*, 2015) with lineage-specific marker sets selected at phylum rank (or class rank for *Proteobacteria*). MAGs  
116 were taxonomically classified by phylogenomic analysis of concatenated marker sequences and calculation of average nucleic  
117 and amino acid identities (ANI, AAI, Supplementary Methods). MAGs were annotated using the MicroScope annotation  
118 platform (Vallenet *et al.*, 2017) and eggNOG (Huerta-Cepas *et al.*, 2016). Genes of interest (Supplementary Table S2) were  
119 manually curated using the full range of tools integrated in MicroScope annotation platform (Supplementary Methods).

## 120 **Genome-centric activity analysis: iRep and metatranscriptomics**

121 The index of replication (iRep) was calculated for each MAG with the combined native soil metagenomes. Settings and  
122 thresholds were applied as recommended (Brown *et al.*, 2016) using bowtie2 (Langmead and Salzberg, 2012) and the iRep  
123 script with default settings. Quality-filtered metatranscriptome reads were mapped to all genomes using bowtie2 and counted  
124 with featureCounts (Liao *et al.*, 2014). To determine gene expression changes, we applied the DESeq2 pipeline with  
125 recommended settings (Love *et al.*, 2014) (Supplementary Methods).

## 126 **Data availability**

127 Metagenomic and -transcriptomic data were deposited under the BioProject accession numbers PRJNA412436 and  
128 PRJNA412438, respectively, and can also be obtained via the JGI's genome portal (JGI Proposal ID 605). MAGs are available  
129 at MicroScope (<https://www.genoscope.cns.fr/agc/microscope/>) and were deposited at the European Nucleotide Archive  
130 (Supplementary Table S1). DsrAB sequences were deposited at NCBI GenBank under the accession numbers MG182080–  
131 MG182141.

132

133

## 134 Results

### 135 Functional metagenomics: Recovery of *dsrAB*-containing genomes from native soil 136 and <sup>13</sup>C-DNA fraction metagenomes

137 This study was conducted with soil samples from the Schlöppnerbrunnen II peatland in Germany, which is a long-term study  
138 site with active sulfur cycling and harbors a large diversity of unknown microorganisms with divergent *dsrAB* genes (Steger *et al.*, 2011; Pester *et al.*, 2012). We initially generated co-assembled metagenomes from native peat soil DNA (53 Gb) and a  
139 pool of DNA extracts from the heavy fractions of a previous DNA-stable isotope probing (DNA-SIP) experiment with soil  
140 from the same peat (101 Gb). The heavy fractions, which were obtained from anoxic peat incubations with periodically  
141 supplemented sulfate and a mixture of <sup>13</sup>C-labelled formate, acetate, propionate, and lactate at low concentrations, were  
142 enriched in DNA from *Desulfosporosinus* and also harbored DNA from yet unidentified *dsrAB*-containing microorganisms  
143 (Pester *et al.*, 2010). Based on the metagenome data, the native peat was dominated by *Acidobacteria* (61%), but also had  
144 *Actinobacteria*, *Alphaproteobacteria*, and *Deltaproteobacteria* as abundant (>5%) phyla/classes (Figure 1). Dominance of  
145 *Acidobacteria*, *Alpha*- and *Deltaproteobacteria* is typical for peatlands (Dedysh, 2011). Quantitative PCR confirmed that  
146 *Acidobacteria* subdivisions 1, 2, and 3 persistently dominated the Schlöppnerbrunnen II peat microbiota in oxic and anoxic soil  
147 layers (Supplementary Methods, Figure 1), as observed in other peatlands (Serkebaeva *et al.*, 2013; Urbanová and Bárta, 2014;  
148 Ivanova *et al.*, 2016).

150 We identified 36 complete or partial *dsrAB* genes on scaffolds of the co-assembled metagenome and subsequently recovered  
151 thirteen MAGs of DsrAB-encoding bacteria by differential coverage binning (Supplementary Table S1, Albertsen *et al.*, 2013).  
152 Twenty-eight *dsrAB* sequences were part of the reductive bacterial-type DsrAB family branch and were closely related to  
153 previously recovered sequences from this and other wetlands (Supplementary Figure S1). These *dsrAB* sequences were  
154 affiliated to the known SRM genera *Desulfosporosinus* (*Firmicutes*, n=1, one MAG) and *Syntrophobacter*  
155 (*Deltaproteobacteria*, n=3, two MAGs), the *Desulfobacca acetoxidans* lineage (n=1), and the uncultured DsrAB family-level  
156 lineages 8 (n=19, seven MAGs) and 10 (n=4). Six sequences grouped with the oxidative bacterial-type DsrAB family and were  
157 distantly affiliated with *Sideroxydans lithotrophicus* (*Betaproteobacteria*, n=5, two MAGs) or *Rhodomicrobium vanniellii*  
158 (*Alphaproteobacteria*, n=1) (Supplementary Figure S2). Interestingly, two of our sequences (n=2, one MAG) and a DsrAB  
159 sequence from the candidate phylum *Rokubacteria* (Hug *et al.*, 2016) formed a completely novel basal lineage outside the four  
160 previously recognized DsrAB enzyme families (Supplementary Figure S2) (Müller *et al.*, 2015). The thirteen partial to near  
161 complete *dsrAB*-containing MAGs had moderate to no detectable contamination as assessed by CheckM and manual curation  
162 (Supplementary Table S1) (Parks *et al.*, 2015) and derived from *Acidobacteria* subdivisions 1 and 3 (SbA1–7),  
163 *Desulfosporosinus* (SbF1), *Syntrophobacter* (SbD1, SbD2), *Betaproteobacteria* (SbB1, SbB2), and *Verrucomicrobia* (SbV1),  
164 as inferred by phylogenetic analysis of DsrAB sequences (Supplementary Figures S1 and S2) and concatenated sequences of  
165 single-copy, phylogenetic marker genes (Supplementary Figure S3). Only the *Desulfosporosinus* and *Syntrophobacter* MAGs  
166 contained rRNA gene sequences.

167 Phylogenomic analysis showed that *Acidobacteria* MAGs SbA1, SbA5, and SbA7 are affiliated with subdivision 1, while  
168 SbA3, SbA4, and SbA6 are affiliated with subdivision 3 (Supplementary Figure S3). The partial MAG SbA2 lacked the marker  
169 genes used for phylogenomic treeing, but was unambiguously assigned to *Acidobacteria* using an extended marker gene set  
170 (Albertsen *et al.*, 2013) and DsrAB phylogeny. The two near complete (96%) MAGs SbA1 and SbA5 have a size of 5.4 and  
171 5.3 Mb, respectively. The G+C content of all acidobacterial MAGs ranges from 58% to 63% (Supplementary Table S1). This  
172 in accordance with genome characteristics of acidobacterial isolates, which have genome sizes of 4.1–10.0 Mb and G+C

173 contents of 57–62% (Ward *et al.*, 2009; Rawat *et al.*, 2012.). SbA1 and SbA7 form a monophyletic clade in the *Acidobacteria*  
174 subdivision 1 with an AAI (Rodríguez-R and Konstantinidis, 2014) of 63% (Supplementary Figure S3) and DsrAB identity of  
175 80% as was calculated with T-Coffee 11 (Notredame *et al.*, 2000) using the unfiltered reference alignment without the  
176 intergenic region (Müller *et al.*, 2015). They have 56% AAI to their closest relative, *Ca. Koribacter versatilis*, which is lower  
177 than AAIs among members of known acidobacterial genera (60–71%). The third MAG from subdivision 1, SbA5, is affiliated  
178 with *Terracidiphilus gabretensis* with an AAI of 61%. DsrAB identity of SbA5 to SbA1 and SbA7 is 79%. The three  
179 subdivision 3 MAGs form a monophyletic clade with *Ca. Solibacter usitatus* (Supplementary Figure S3). SbA3, SbA4, and  
180 SbA6 have AAIs of 59–73% amongst them and 61–62% to *Ca. S. usitatus*. DsrAB identity amongst the three MAGs is 80–  
181 94% and 74–79% to the subdivisions 1 MAGs.

182 The DsrAB sequences encoded on all seven MAGs belong to the uncultured DsrAB family-level lineage 8 (Supplementary  
183 Figure S1), which so far only consisted of environmental *dsrAB* sequences of unknown taxonomic identity (Müller *et al.*,  
184 2015). Based on these MAGs and metatranscriptome analyses of anoxic peat soil microcosms, we describe the putative  
185 metabolic capabilities of these novel DsrAB-encoding *Acidobacteria*. Details on the other MAGs will be described elsewhere  
186 (Hausmann *et al.*, unpublished; Anantharaman *et al.*, unpublished). Functional interpretations of the recovered MAGs are  
187 made under the premise that the genomes are not closed, and thus it is unknown if genes are absent in these organisms or are  
188 missing due to incomplete sequencing, assembly, or binning.

## 189 **Dissimilatory sulfur metabolism**

190 Although *Acidobacteria* are abundant in diverse environments with active sulfur cycling (Serkebaeva *et al.*, 2013; Urbanová  
191 and Bárta, 2014; Sánchez-Andrea *et al.*, 2011; Wang *et al.*, 2012), this is the first discovery of members of this phylum with a  
192 putative dissimilatory sulfur metabolism. SbA2, SbA3, and SbA7 encode the complete canonical pathway for dissimilatory  
193 sulfate reduction, including homologs for sulfate transport (*sulP* and/or *dass*, not in SbA7) and activation (*sat*, *ppa*, *hppA*),  
194 adenosine 5'-phosphosulfate (APS) reduction (*aprBA*, *qmoABC*), and sulfite reduction (*dsrAB*, *dsrC*, *dsrMKJOP*) (Figure 2,  
195 Supplementary Table S2a) (Santos *et al.*, 2015). In the AprBA tree, the acidobacterial sequences are part of a large cluster of  
196 yet uncultured organisms and *Deltaproteobacteria* and *Firmicutes* that respire sulfate, sulfite, or thiosulfate (Supplementary  
197 Figure S4) (Watanabe *et al.*, 2016). SbA1, SbA4, SbA5, and SBA6 have an incomplete sulfate reduction gene set but contain  
198 all *dsr* genes for sulfite reduction. Several other *dsr* genes were present on some of the MAGs. The *dsrD* and *dsrN* genes  
199 occurred in pairs. The acidobacterial DsrD sequences have the same conserved, hydrophobic residues as *Desulfovibrio vulgaris*  
200 DsrD (Supplementary Figure S5) (Mizuno *et al.*, 2003). Ubiquity of DsrD among SRM suggests an essential function in sulfate  
201 reduction, but the physiological role of this small protein is unresolved (Hittel and Voordouw, 2000). DsrN is a homolog of  
202 cobyrinate a,c-diamide synthase in cobalamin biosynthesis and may be involved in amidation of the siroheme prosthetic group  
203 of DsrAB (Lübbe *et al.*, 2006). DsrV, a homolog of precorrin-2 dehydrogenase, and DsrWa, a homolog of uroporphyrin-III C-  
204 methyltransferase, may also be involved in siroheme biosynthesis (Holkenbrink *et al.*, 2011). DsrT is required for sulfide  
205 oxidation in *Chlorobaculum tepidum*, but also found in SRM (Holkenbrink *et al.*, 2011). The presence of *dsrMK*-paralogs  
206 (*dsrM2*, *dsrK2*) upstream of *dsrAB* is not uncommon in SRM (Pereira *et al.*, 2011). DsrMK are present in all *dsrAB*-containing  
207 microorganisms and are a transmembrane module involved in reduction of cytoplasmic DsrC-trisulfide in SRM, the final step  
208 in sulfate reduction (Santos *et al.*, 2015). DsrC encoded on the MAGs have the two essential cysteine residues at the C-  
209 terminal end for full functionality (Venceslau *et al.*, 2014). Interestingly, *dsrC* forms a gene duo with *dsrL* downstream of  
210 *dsrAB* in all seven MAGs. This is surprising, because *dsrL* is not found in SRM but in sulfur oxidizers. DsrL is highly  
211 expressed and essential for sulfur oxidation by the purple sulfur bacterium *Allochromatium vinosum* (Lübbe *et al.*, 2006;



212 Weissgerber *et al.*, 2014). DsrL is a cytoplasmic iron-sulfur flavoprotein with proposed NAD(P)H:acceptor oxidoreductase  
213 activity and was copurified with DsrAB from the soluble fraction of *A. vinosum* (Dahl *et al.*, 2005). The acidobacterial DsrL  
214 sequences are shorter than their homolog in *A. vinosum* (Supplementary Table S2a), but have the same functional domains  
215 (Supplementary Figure S6). Given the possible role of DsrL in sulfur oxidation, we sought to detect additional genes indicative  
216 of oxidative sulfur metabolism in the acidobacterial MAGs. However, genes for Sox enzyme machinery (*soxABXYZ*),  
217 thiosulfate dehydrogenase (*tsdA*), sulfide:quinone reductase (*sqr*), adenylyl-sulfate reductase membrane anchor subunit  
218 (*aprM*), flavocytochrome c sulfide dehydrogenase (*fccAB*), sulfur reductase (*sreABC*), thiosulfate reductase (*phsABC*),  
219 polysulfide reductase (*psrABC*), membrane-bound sulfite oxidizing enzyme (*soeABC*), cytoplasmic sulfur trafficking enzymes  
220 (*tusA*, *dsrE2*, *dsrEFH*), or DsrQ/DsrU (unknown functions) were absent (Laska *et al.*, 2003; Holkenbrink *et al.*, 2011; Lenk *et*  
221 *al.*, 2012; Wasmund *et al.*, 2017). SbA1, SbA3, SbA4, and SbA6 contain genes that have only low homology to *soxCD/sorAB*,  
222 periplasmic sulfite-oxidizing enzymes (Supplementary Results) and, thus, might have another function (Ghosh and Dam,  
223 2009).

224 Despite ongoing sulfur cycling, concentrations of inorganic sulfur compounds such as sulfate are low (lower  $\mu\text{M}$  range) in the  
225 Schlössnerbrunnen II peatland (Schmalenberger *et al.*, 2007; Küsel *et al.*, 2008; Knorr and Blodau, 2009). Enzymatic release  
226 of inorganic sulfur compounds from organic matter might thus represent a significant resource for sulfur-dissimilating  
227 microorganisms. Therefore, we specifically searched for genes coding for known organosulfur reactions that yield sulfite  
228 (Wasmund *et al.*, 2017). Genes for cysteate sulfo-lyase (*cuyA*), methanesulfonate monooxygenase (*msmABCD*),  
229 sulfoacetaldehyde acetyltransferase (*xsc*), and taurine dioxygenase (*tauD*) were absent. However, *suyAB*, coding for the (*R*-  
230 sulfolactate sulfo-lyase complex that cleaves (*R*)-sulfolactate into pyruvate and sulfite (Denger and Cook, 2010), were present  
231 in SbA4 and SbA5 (Supplementary Table S2a). Intriguingly, SbA4 and SbA5 only have capability for sulfite reduction. SbA5  
232 also encodes the racemase machinery for (*S*)-sulfolactate to (*R*)-sulfolactate, (*S*)-sulfolactate dehydrogenase (*slcC*) and (*R*-  
233 sulfolactate dehydrogenase (*comC*); the regulator gene *suyR* or the putative importer SlcHFG were absent (Denger and Cook,  
234 2010). Pyruvate may be used as an energy and carbon source, while sulfite could be used as an electron acceptor for anaerobic  
235 respiration (Simon and Kroneck, 2013).

## 236 **Respiration**

237 Cultivated *Acidobacteria* of subdivisions 1 and 3 are strict aerobes or facultative anaerobes (e.g., Eichorst *et al.*, 2007;  
238 Kulichevskaya *et al.*, 2010, 2014; Pankratov and Dedysh, 2010; Dedysh *et al.*, 2012; Pankratov *et al.*, 2012). Accordingly, we  
239 found respiratory chains encoded in all acidobacterial MAGs (Figure 3, Supplementary Results), with (near) complete operons  
240 for NADH dehydrogenase 1, succinate dehydrogenase (lacking in SbA2), one or both types of quinol—cytochrome-c  
241 reductase, low-affinity terminal oxidases, and ATP synthase (lacking in SbA2) (Supplementary Tables S2b–h). High-affinity  
242 terminal oxidases, putatively involved in detoxification of oxygen (Ramel *et al.*, 2013; Giuffrè *et al.*, 2014), are limited to four  
243 MAGs (Supplementary Table S2g). Genes for dissimilatory nitrogen or iron metabolisms are absent, with the exception of a  
244 putative metal reductase in SbA2 of unclear physiological role (Supplementary Results).

## 245 **Hydrogen utilization and production**

246 We identified [NiFe] hydrogenases of groups 1, 3, and 4 (Greening *et al.*, 2016) in SbA1–7 (Supplementary Table S2j).  
247 Membrane-bound group 1 hydrogenases (SbA1, SbA3, SbA5) consume hydrogen from the periplasm as an electron donor to  
248 generate energy, possibly coupled to sulfate/sulfite reduction. In contrast to other *Acidobacteria*, no group 1h/5 hydrogenases,  
249 which are coupled to oxygen respiration, were identified (Greening *et al.*, 2015). Cytoplasmic group 3 hydrogenases (all

250 MAGs) are bidirectional and proposed to be involved in energy-generating hydrogen oxidation and/or fermentative hydrogen  
251 production. Membrane-bound group 4 hydrogenases (SbA1, SbA5, SbA4, SbA6) produce H<sub>2</sub> and are postulated to conserve  
252 energy by proton translocation by oxidizing substrates like formate (group 4a) or carbon monoxide (via ferredoxin, group 4c)  
253 (Figure 3).

## 254 **A versatile heterotrophic physiology**

255 *Acidobacteria* are known for their capability to degrade simple and polymeric carbohydrates (Kulichevskaya *et al.*, 2010,  
256 2014; Pankratov and Dedysh, 2010; Dedysh *et al.*, 2012; Eichorst *et al.*, 2011; Pankratov *et al.*, 2012; Rawat *et al.*, 2012;  
257 Huber *et al.*, 2016), supported by many diverse carbohydrate-active enzymes encoded on their genomes (Ward *et al.*, 2009;  
258 Rawat *et al.*, 2012). Accordingly, the MAGs recovered in our study also contain many genes encoding diverse carbohydrate-  
259 active enzymes (Supplementary Methods, Figure 4). These include glycoside hydrolases (GH, 1.0–4.0% of all genes),  
260 polysaccharide lyases (0.07–0.3%), and carbohydrate esterases (0.7–1.4%) that are generally involved in degradation of  
261 complex sugars, but also glycosyltransferases (0.9–1.4%) for biosynthesis of carbohydrates. Functional GH families (assigned  
262 by EC number) putatively involved in cellulose and hemicellulose degradation were most prevalent (Supplementary Table S4).  
263 Specifically, the most often encountered EC numbers encompassed by the different GH families represented cellulose (EC  
264 3.2.1.4, e.g., GH5, GH74), xyloglucan (EC 3.2.1.150, EC 3.2.1.151, e.g., GH5, GH74), or xylan (EC 3.2.1.8, EC 3.2.1.37, e.g.,  
265 GH5) degradation, which is similar to the situation found in other members of *Acidobacteria* subdivision 1 and 3 (Ward *et al.*,  
266 2009; Rawat *et al.*, 2012). Further EC numbers that were often encountered in the various detected GH families were  
267 associated with oligosaccharide degradation (EC 3.2.1.21, e.g., GH2) or  $\alpha$ -N-acetylgalactosaminidase activity (EC 3.2.1.49,  
268 e.g., GH109). Degradation of cellulose and hemicellulose yields glucose and all MAGs encode glycolysis and pentose  
269 phosphate pathways (Figure 3, Supplementary Results).  $\alpha$ -N-acetylgalactosaminidase releases N-acetylgalactosamine residues  
270 from glycoproteins that are commonly found in microbial cell walls and extracellular polysaccharides (Bodé *et al.*, 2013). N-  
271 acetylgalactosamine can not be directly utilized via glycolysis, however the additionally required enzymes are present (Figure  
272 3; Supplementary Results). Under oxic conditions, organic carbon could be completely oxidized to CO<sub>2</sub> via the citric acid  
273 cycle (Figure 3). Alternatively, we also identified fermentative pathways. SbA3 encodes the bifunctional aldehyde-alcohol  
274 dehydrogenase AdhE that yields ethanol (Figure 3). All MAGs encode additional aldehyde and alcohol dehydrogenases  
275 without clear substrate specificity that could also ferment acetyl-CoA to ethanol. SbA7 and SbA5 encode a L-lactate  
276 dehydrogenase (Ldh) yielding lactate from pyruvate, while six MAGs encode L-lactate dehydrogenases (LldD, GlcDEF,  
277 LutABC) that presumably perform the reverse reaction (Figure 3). Similarly, we identified pathways for acetate and/or  
278 propionate production or utilization in all MAGs (Figure 3; Supplementary Results). SbA1 and SbA3 potentially produce H<sub>2</sub>  
279 via formate C-acetyltransferase PflD, which cleaves pyruvate into acetyl-CoA and formate. SbA1 encodes for the membrane-  
280 bound formate hydrogenlyase complex (*fdhF*, *hyf* operon) that produces H<sub>2</sub> and might also translocate protons. SbA3  
281 harbours an uncharacterized, cytoplasmic, monomeric FDH (*fdhA*) to transform formate to H<sub>2</sub>. SbA1, SbA3, SbA4, and SbA6  
282 also encode membrane-bound, periplasmic FDH (*fdo* operon) that transfers electrons into the membrane quinol pool, as a non-  
283 fermentative alternative of formate oxidation (Figure 3, Supplementary Table S2j).

## 284 **DsrAB-encoding *Acidobacteria* are metabolically active under anoxic conditions**

285 We calculated the index of replication (iRep) based on the native peat soil metagenomes to assess whether DsrAB-encoding  
286 *Acidobacteria* were active *in situ* (Brown *et al.*, 2016). SbA1 and SbA5, which were sufficiently complete ( $\geq 75\%$ ) for a  
287 reliable estimate, had iRep values of 1.21 and 1.19, respectively. This shows that a fraction of each population was  
288 metabolically active, i.e., on average 21% of SbA1 and 19% of SbA5 cells were actively replicating at the time of sampling.

289 Concordantly, SbA1–7 were also transcriptionally active in the same native soil samples. 35–46% of the SbA1–7 genes were  
290 expressed in at least one replicate. SbA1 and SbA5 contributed a considerable fraction (0.4% and 1.8%, respectively,  
291 Supplementary Table S1) of the total mRNA reads in the native soil metatranscriptome. These data likely underestimate the  
292 metabolic activity of SbA1–7 *in situ* because freshly sampled soil was stored at 4 °C for one week prior to nucleic acids  
293 extraction.

294 We further analyzed metatranscriptome data from a series of anoxic incubations of the peat soil with or without individual  
295 substrates (formate, acetate, propionate, lactate, or butyrate) and with or without supplemental sulfate (Hausmann *et al.*, 2016).  
296 While the incubations were not designed to specifically test for the MAG-inferred metabolic properties, they still allowed us to  
297 evaluate transcriptional response of the DsrAB-encoding *Acidobacteria* under various anoxic conditions (Supplementary  
298 Methods and Results). All treatments triggered shifts in genome-wide gene expression; more genes were significantly ( $p < 0.05$ )  
299 upregulated (73–933) than downregulated (14–81) as compared to the native soil. Upregulated genes included sulfur  
300 metabolism enzymes, high-affinity terminal oxidases, group 1 and 3 hydrogenases, aldehyde-alcohol dehydrogenase AdhE,  
301 glycoside hydrolases, and other carbon metabolism enzymes (Supplementary Table S3, Supplementary Figure S7).  
302 Significantly upregulated glycoside hydrolase genes belonged to GH family 2, 3, 5, 9, 10, 18, 20, 23, 26, 28, 29, 30, 33, 35, 36,  
303 38, 43, 44, 50, 51, 55, 74, 76, 78, 79, 88, 95, 97, 105, 106, 109, and 129 in MAGs SbA1–6. None of the GH genes were  
304 significantly downregulated in the incubations. Noteworthy genes that were significantly downregulated were superoxide  
305 dismutases (*sodA*) in SbA2 and SbA4 (Supplementary Table S3a).

306

307

## 308 Discussion

309 Diverse members of the phylum *Acidobacteria* are abundant in various ecosystems, particularly in soils and sediments with  
310 relative abundances typically ranging from 20–40% (Janssen, 2006). *Acidobacteria* are currently classified in 26 subdivisions  
311 based on their 16S rRNA phylogeny (Barns *et al.*, 2007). Given their phylogenetic breadth, comparably few isolates and  
312 genomes are available to explore their metabolic capabilities. Currently isolated species of subdivisions 1, 3, 4, and 6 are  
313 aerobic chemoorganotrophs that grow optimally at neutral or low pH (Dedysh, 2011; Eichorst *et al.*, 2011; Huber *et al.*, 2014,  
314 2016). Furthermore, subdivision 4 contains an anoxygenic phototroph (Garcia Costas *et al.*, 2012; Tank and Bryant, 2015),  
315 subdivisions 8 and 23 contain anaerobes (Liesack *et al.*, 1994; Coates *et al.*, 1999; Losey *et al.*, 2013), subdivisions 1, 3, and  
316 23 fermenters (Pankratov *et al.*, 2012; Kulichevskaya *et al.*, 2014; Losey *et al.*, 2013; Myers and King, 2016) and subdivision  
317 4, 8, 10 and 23 thermophiles (Izumi *et al.*, 2012; Losey *et al.*, 2013; Crowe *et al.*, 2014; Tank and Bryant, 2015).

318 *Acidobacteria* are known as dominant inhabitants of wetlands worldwide, in particular members of subdivision 1, 3, 4, and 8  
319 (Dedysh, 2011). Strains in the genera *Granulicella* (Pankratov and Dedysh, 2010), *Telmatobacter* (Pankratov *et al.*, 2012),  
320 *Bryocella* (Dedysh *et al.*, 2012) and *Bryobacter* (Kulichevskaya *et al.*, 2010) have been isolated from acidic wetlands and are  
321 presumably active in plant-derived polymer degradation (such as cellulose) (Dedysh, 2011; Pankratov *et al.*, 2011; Schmidt *et al.*  
322 *et al.*, 2015; Juottonen *et al.*, 2017), and in nitrogen and iron cycling (Küsel *et al.*, 2008; Kulichevskaya *et al.*, 2014).

323 Here, we provide metagenomic and metatranscriptomic evidence that the newly discovered species represent at least three  
324 novel genera in *Acidobacteria* subdivision 1 and 3 (Supplementary Figure S3) and possess a dissimilatory sulfur metabolism.  
325 The seven acidobacterial MAGs from the Schlöppnerbrunnen II peatland encode the complete canonical pathway for  
326 dissimilatory sulfite or sulfate reduction. The sulfate reduction pathway, however, could also operate in reverse as proposed for  
327 a sulfur-oxidizing deltaproteobacterium (Thorup *et al.*, 2017). The phylogenetic separation into two subdivisions as based on  
328 the concatenated marker gene tree is also apparent in the DsrAB phylogeny (Supplementary Figure S1). The acidobacterial  
329 DsrAB sequences are distributed among two monophyletic clades within the uncultured family-level lineage 8, which is part of  
330 the reductive, bacterial-type DsrAB branch (Müller *et al.*, 2015). The phylogenetic breadth of the acidobacterial DsrAB  
331 sequences is representative for the intra-lineage sequence divergence within uncultured DsrAB lineage 8, which suggests that  
332 this entire lineage represents yet uncultivated bacteria of the phylum *Acidobacteria*. Members of this uncultured DsrAB  
333 lineage are widespread in freshwater wetlands (Supplementary Figure S1) (Pester *et al.*, 2012). In particular, they represented  
334 an abundant fraction of the DsrAB diversity and were permanent autochthonous inhabitants of oxic and anoxic soil layers in  
335 the analyzed Schlöppnerbrunnen II peatland (Steger *et al.*, 2011; Pester *et al.*, 2010).

336 Presence of the complete gene set for canonical dissimilatory sulfate reduction suggests that the pathway is functional, as the  
337 genetic capability for sulfate reduction can be rapidly lost by adaptive evolution if unused (Hillesland *et al.*, 2014). Except for  
338 a truncated *aprB* on SbA6, we found no indications of pseudogenes, i.e., unexpected internal stop codons or reading frame  
339 shifts, for any of the sulfate/sulfite reduction genes on the acidobacterial MAGs (Müller *et al.*, 2015). In addition, sulfur  
340 metabolism genes of each MAG were expressed in the native soil and the anoxic microcosms (Supplementary Table S3a).  
341 Many sulfur metabolism genes were even significantly upregulated in the anoxic microcosms, with *dsrC* and *aprBA* among the  
342 top 10 most expressed genes in SbA7 (Supplementary Table S3a). These findings further support full functionality of the  
343 acidobacterial dissimilatory sulfur pathways under anoxic condition.

344 Known SRM typically couple sulfate respiration to oxidation of fermentation products such as volatile fatty acids, alcohols, or  
345 hydrogen (Rabus *et al.*, 2013). While other microorganisms in the Schlöppnerbrunnen II soil, such as *Desulfosporosinus*,

346 showed sulfate- and substrate-specific responses in our microcosms, hundreds of acidobacterial 16S rRNA phylotypes did not  
347 (with the exception of two) (Hausmann *et al.*, 2016). Gene expression patterns of DsrAB-encoding *Acidobacteria* in the  
348 individual anoxic microcosms as analyzed in the present study were ambiguous. Genes for putative oxidation of the  
349 supplemented substrates (formate, acetate, propionate, lactate, butyrate) were not specifically upregulated, neither without nor  
350 with supplemental sulfate. However, sulfur metabolism genes were upregulated in several incubations as compared to no-  
351 substrate-controls, suggesting indirect stimulation of a sulfur-based metabolism (Supplementary Results, Supplementary Table  
352 S3a). Indirect changes in microbial activity after the addition of fresh organic matter is often observed in soils (priming effects,  
353 Blagodatskaya and Kuzyakov, 2008). One explanation for this priming effect is the co-metabolism theory stating that easily  
354 available substrates provide the energy for microorganisms to produce extracellular enzymes to make immobile carbon  
355 accessible, which is then also available to other microorganisms. The DsrAB-encoding *Acidobacteria* have a large genetic  
356 repertoire to utilize carbohydrates and monomeric sugars (Figure 3). This is in accordance with the carbohydrate utilization  
357 potential previously described for subdivision 1 and 3 *Acidobacteria* (Ward *et al.*, 2009; Rawat *et al.*, 2012). Yet utilization of  
358 monomeric sugars is a rare feature of known SRM (Cord-Ruwisch *et al.*, 1986; Stetter, 1988) and utilization of poly- or  
359 oligosaccharides by sulfate-reducing bacteria was not yet reported. While the studied *Acidobacteria* expressed many of their  
360 glycoside hydrolase genes in our anoxic peat soil microcosms, further experiments are required to confirm if DsrAB-encoding  
361 *Acidobacteria* couple degradation of carbohydrate polymers or monomers to sulfate reduction.

362 It is intriguing to propose that MAGs SbA2, SbA3, and SbA7 derive from acidobacterial SRM as they lack known sulfur  
363 oxidation genes, except *dsrL* (Supplementary Figure S6), and express the complete dissimilatory sulfate reduction pathway  
364 (Supplementary Table S2a), including reductive, bacterial-type *dsrAB*, and *dsrD* that may be exclusive to SRM (Hittel and  
365 Voordouw, 2000; Dahl and Friedrich, 2008; Ghosh and Dam, 2009; Rabus *et al.*, 2015). However, the functions of DsrL and  
366 DsrD are yet unresolved, which prevents functional predictions based only on these genes. The proposal of an alternative  
367 hypothesis that these novel *Acidobacteria* reverse the sulfate reduction pathway for dissimilatory sulfur oxidation or sulfur  
368 disproportionation, bases on findings with the deltaproteobacterium *Desulfurivibrio alkaliphilus* (Thorup *et al.*, 2017). *D.*  
369 *alkaliphilus* also lacks known sulfur oxidation genes (including *dsrL*), except for *sqr*, and is proposed to gain energy by  
370 coupling sulfide oxidation via a reversed sulfate reduction pathway (with a reductive-type DsrAB) to the dissimilatory  
371 reduction of nitrate/nitrite to ammonium. Sulfide oxidation in acidobacterial MAGs SbA2, SbA3, and SbA7 could proceed  
372 analogous to the pathway models proposed by Thorup *et al.* (2017) and Christiane Dahl (Dahl, 2017). Briefly, hydrogen  
373 sulfide might react with DsrC either spontaneously (Ijssennagger *et al.*, 2015) or via an unknown sulfur transfer mechanism to  
374 form persulfated DsrC. Persulfated DsrC is then oxidized by DsrMKJOP, thereby transferring electrons into the membrane  
375 quinone pool, and releasing a DsrC-trisulfide, which is the substrate for DsrAB (Santos *et al.*, 2015; Dahl, 2017). It was  
376 hypothesized that electrons released during DsrC-trisulfide oxidation to sulfite and DsrC are transferred to DsrL (Dahl, 2017).  
377 Further sulfite oxidation to sulfate would be catalyzed by AprBA-QmoABC and Sat.

378 The acidobacterial MAGs have the genomic potential to use oxygen as terminal electron acceptor and might thus couple  
379 sulfide oxidation to aerobic respiration. Alternative electron acceptors for biological sulfur oxidation in wetlands could include  
380 nitrate/nitrite and metals such as Fe(III) (Küsel *et al.*, 2008). However, known genes for dissimilatory nitrate reduction and  
381 metal reduction (Weber *et al.*, 2006) were absent from these acidobacterial MAGs. Only SbA2 encodes a putative metal  
382 reduction complex that was recently characterized in *Desulfotomaculum reducens* (Otwell *et al.*, 2015). At this time, it is  
383 unclear whether DsrAB-encoding *Acidobacteria* are capable of Fe(III) respiration, as seen in *Geothrix fermentans* (Coates *et al.*  
384 *et al.*, 1999) and certain isolates in subdivision 1 (Blöthe *et al.*, 2008; Kulichevskaya *et al.*, 2014).

385

386

387 **Proposal of the acidobacterial *Candidatus* genera**  
388 ***Sulfotelmatobacter*, *Sulfotelmatomonas*, and *Sulfopaludibacter***

389 Based on combined interpretation of phylogeny (concatenated phylogenetic marker genes, DsrAB), genomic (ANI, AAI) and  
390 genetic (DsrAB) distances, and characteristic genomic features of dissimilatory sulfur metabolism (Figure 3), in accordance  
391 with Konstantinidis *et al.* (2017), we classify MAGs SbA1, SbA7, SbA5, SbA3, SbA4, and SbA6 into three new acidobacterial  
392 *Candidatus* genera, including *Candidatus* species names for the >95% complete MAGs SbA1 and SbA5. In-depth  
393 phylogenomic analysis of SbA2 was not possible and therefore it is tentatively assigned to *Acidobacteria* subdivision 3.

394 *Acidobacteria* subdivision 1

- 395 • *Ca.* genus *Sulfotelmatobacter* (Sul.fo.tel.ma.to.bac'ter. L. n. *sulfur*, sulfur; Gr. n. *telma*, -tos, swamp, wetland; N.L.  
396 masc. n. *bacter*, bacterium; N.L. masc. n. *Sulfotelmatobacter*, a bacterium from a swamp metabolizing sulfur) with  
397 *Ca.* *Sulfotelmatobacter kueseliae* MAG SbA1 (kue.se'li.ae. N.L. gen. n. *kueseliae*, of Kuesel, honouring Kirsten  
398 Küsel, for her work on the geomicrobiology of wetlands) and *Ca.* *Sulfotelmatobacter* sp. MAG SbA7.
- 399 • *Ca.* *Sulfotelmatomonas gaucii* MAG SbA5 (Sul.fo.tel.ma.to.mo.nas. L. n. *sulfur*, sulfur; Gr. n. *telma*, -tos, swamp,  
400 wetland; N.L. fem. n. *monas*, a unicellular organism; N.L. fem. n. *Sulfotelmatomonas*, a bacterium from a swamp  
401 metabolizing sulfur; gau'.ci.i. N.L. gen. n. *gaucii*, of Gauci, in honour of Vincent Gauci, for his pioneering work on  
402 the interplay of wetland sulfate reduction and global methane emission).

403 *Acidobacteria* subdivision 3

- 404 • *Ca.* genus *Sulfopaludibacter* (Sul.fo.pa.lu.di.bac'ter. L. n. *sulfur*, sulfur; L. n. *palus*, -udis, L. swamp; N.L. masc. n.  
405 *bacter*, bacterium; N.L. masc. n. *Sulfopaludibacter*, a bacterium from a swamp metabolizing sulfur) with *Ca.*  
406 *Sulfopaludibacter* sp. MAG SbA3, *Ca.* *Sulfopaludibacter* sp. MAG SbA4, and *Ca.* *Sulfopaludibacter* sp. MAG SbA6.
- 407 • *Acidobacteria* bacterium sp. MAG SbA2.

408

409

410

411



## 412 **Conclusion**

413 Sulfur cycling exerts important control on organic carbon degradation and greenhouse gas production in wetlands, but  
414 knowledge about sulfur microorganisms in these globally important ecosystems is scarce (Pester *et al.*, 2012). Here, we show  
415 by genome-centric metagenomics and metatranscriptomics that members of the phylum *Acidobacteria* have a putative role in  
416 peatland sulfur cycling. The genomic repertoire of these novel *Acidobacteria* species encompassed recognized acidobacterial  
417 physiologies, such as a facultative anaerobic metabolism, oxygen respiration, fermentation, carbohydrate degradation, and  
418 hydrogen metabolism, but was additionally augmented with a DsrAB-based dissimilatory sulfur metabolism (Figure 5). Based  
419 on their genetic repertoire and previous findings on reversibility of the dissimilatory sulfate reduction pathway (Dannenberg *et*  
420 *al.*, 1992; Fuseler and Cypionka, 1995; Fuseler *et al.*, 1996; Thorup *et al.*, 2017) , it is intriguing to speculate that the described  
421 peatland *Acidobacteria* could use the same pathway for both sulfate reduction and sulfide oxidation. The described DsrAB-  
422 carrying *Acidobacteria* that only encoded the pathway for dissimilatory sulfite reduction had additional genes for sulfite-  
423 producing enzymes, which suggests that organosulfonates might be their primary substrate for sulfur respiration. Our results  
424 not only extend the current understanding of the genetic versatility and distribution of dissimilatory sulfur metabolism among  
425 recognized microbial phyla, but also underpin the challenge to unambiguously differentiate between reductive or oxidative  
426 sulfur metabolism solely based on (meta-)genome/transcriptome data (Thorup *et al.*, 2017).

427

428



## 429 **Conflict of Interest**

430 The authors declare no conflict of interest.

431

## 432 **Acknowledgements**

433 We are grateful to Norbert Bittner for support during field sampling, Doris Steger and Pinsurang Deevong for their  
434 contributions to qPCR analysis, and Florian Goldenberg for maintaining the Life Science Computer Cluster at the Division of  
435 Computational Systems Biology (University of Vienna). We thank the staff of the Joint Genome Institute (JGI) for  
436 metagenome and metatranscriptome library preparation, sequencing, and standard bioinformatics support, Bernhard Schink for  
437 help in naming of bacterial taxa, and Christiane Dahl, Petra Pjevac, Marc Mußmann, and Kenneth Wasmund for valuable  
438 discussions and feedback. We acknowledge the LABGeM (CEA/IG/Genoscope & CNRS UMR8030) and the France  
439 Génomique National infrastructure (funded as part of Investissement d'avenir program managed by Agence Nationale pour la  
440 Recherche, contract ANR-10-INBS-09) for support with the MicroScope annotation platform. The work conducted by the JGI  
441 was supported by the Office of Science of the U.S. Department of Energy under Contract No. DE-AC02-05CH11231. This  
442 research was supported by the Austrian Science Fund (FWF, P23117-B17, P25111-B22, P26392-B20, and I1628-B22), the JGI  
443 (CSP 605), the German Research Foundation (DFG, PE 2147/1-1), the European Union (FP7-People-2013-CIG, Grant No  
444 PCIG14-GA-2013-630188), and the VILLUM FONDEN (research grant 15510).

445

446

## 447 **References**

- 448 Albertsen M, Hugenholtz P, Skarshewski A, Nielsen KL, Tyson GW, Nielsen PH. (2013). Genome sequences of rare,  
449 uncultured bacteria obtained by differential coverage binning of multiple metagenomes. *Nat Biotechnol* **31**: 533–538.
- 450 Barns SM, Cain EC, Sommerville L, Kuske CR. (2007). *Acidobacteria* phylum sequences in uranium-contaminated subsurface  
451 sediments greatly expand the known diversity within the phylum. *Appl Environ Microbiol* **73**: 3113–3116.
- 452 Blagodatskaya E, Kuzyakov Y. (2008). Mechanisms of real and apparent priming effects and their dependence on soil  
453 microbial biomass and community structure: critical review. *Biol Fertil Soils* **45**: 115–131.
- 454 Blöthe M, Akob DM, Kostka JE, Göschel K, Drake HL, Küsel K. (2008). pH gradient-induced heterogeneity of Fe(III)-  
455 reducing microorganisms in coal mining-associated lake sediments. *Appl Environ Microbiol* **74**: 1019–1029.
- 456 Bodé S, Fancy R, Boeckx P. (2013). Stable isotope probing of amino sugars – a promising tool to assess microbial interactions  
457 in soils. *Rapid Commun Mass Spectrom* **27**: 1367–1379.
- 458 Brown CT, Olm MR, Thomas BC, Banfield JF. (2016). Measurement of bacterial replication rates in microbial communities.  
459 *Nat Biotechnol* **34**: 1256–1263.
- 460 Coates JD, Ellis DJ, Gaw CV, Lovley DR. (1999). *Geothrix fermentans* gen. nov., sp. nov., a novel Fe(III)-reducing bacterium  
461 from a hydrocarbon-contaminated aquifer. *Int J Syst Bacteriol* **49**: 1615–1622.
- 462 Cord-Ruwisch R, Ollivier B, Garcia J-L. (1986). Fructose degradation by *Desulfovibrio* sp. in pure culture and in coculture  
463 with *Methanospirillum hungatei*. *Curr Microbiol* **13**: 285–289.
- 464 Costello EK, Schmidt SK. (2006). Microbial diversity in alpine tundra wet meadow soil: novel Chloroflexi from a cold, water-  
465 saturated environment. *Environ Microbiol* **8**: 1471–1486.
- 466 Crowe MA, Power JF, Morgan XC, Dunfield PF, Lagutin K, Rijpstra IC *et al.* (2014). *Pyrinomonas methylaliphatogenes* gen.  
467 nov., sp. nov., a novel group 4 thermophilic member of the phylum *Acidobacteria* from geothermal soils. *Int J Syst Evol*  
468 *Microbiol* **64**: 220–227.
- 469 Dahl C. (2017). Sulfur metabolism in phototrophic bacteria. In: Hallenbeck PC (ed). *Modern topics in the phototrophic*  
470 *prokaryotes*. Springer, pp 27–66.
- 471 Dahl C, Engels S, Pott-Sperling AS, Schulte A, Sander J, Lübke Y *et al.* (2005). Novel genes of the *dsr* gene cluster and  
472 evidence for close interaction of Dsr proteins during sulfur oxidation in the phototrophic sulfur bacterium *Allochromatium*  
473 *vinosum*. *J Bacteriol* **187**: 1392–1404.
- 474 Dahl C, Friedrich CG. (2008). *Microbial Sulfur Metabolism*. Springer, Berlin Heidelberg.
- 475 Dannenberg S, Kroder M, Dilling W, Cypionka H. (1992). Oxidation of H<sub>2</sub>, organic compounds and inorganic sulfur  
476 compounds coupled to reduction of O<sub>2</sub> or nitrate by sulfate-reducing bacteria. *Arch Microbiol* **158**: 93–99.

- 477 Dedysh SN. (2011). Cultivating uncultured bacteria from northern wetlands: knowledge gained and remaining gaps. *Front*  
478 *Microbiol* **2**: 184.
- 479 Dedysh SN, Kulichevskaya IS, Serkebaeva YM, Mityaeva MA, Sorokin VV, Suzina NE *et al.* (2012). *Bryocella elongata* gen.  
480 nov., sp. nov., a member of subdivision 1 of the *Acidobacteria* isolated from a methanotrophic enrichment culture, and  
481 emended description of *Edaphobacter aggregans* Koch *et al.* 2008. *Int J Syst Evol Microbiol* **62**: 654–664.
- 482 Dedysh SN, Pankratov TA, Belova SE, Kulichevskaya IS, Liesack W. (2006). Phylogenetic analysis and in situ identification  
483 of *Bacteria* community composition in an acidic *Sphagnum* peat bog. *Appl Environ Microbiol* **72**: 2110–2117.
- 484 Denger K, Cook AM. (2010). Racemase activity effected by two dehydrogenases in sulfolactate degradation by  
485 *Chromohalobacter salexigens*: purification of (*S*)-sulfolactate dehydrogenase. *Microbiology* **156**: 967–974.
- 486 Eddy SR. (2011). Accelerated profile HMM searches. *PLoS Comput Biol* **7**: e1002195.
- 487 Eichorst SA, Breznak JA, Schmidt TM. (2007). Isolation and characterization of soil bacteria that define *Terriglobus* gen. nov.,  
488 in the phylum *Acidobacteria*. *Appl Environ Microbiol* **73**: 2708–2717.
- 489 Eichorst SA, Kuske CR, Schmidt TM. (2011). Influence of plant polymers on the distribution and cultivation of bacteria in the  
490 phylum *Acidobacteria*. *Appl Environ Microbiol* **77**: 586–596.
- 491 Fuseler K, Cypionka H. (1995). Elemental sulfur as an intermediate of sulfide oxidation with oxygen by *Desulfobulbus*  
492 *propionicus*. *Arch Microbiol* **164**: 104–109.
- 493 Fuseler K, Krekeler D, Sydow U, Cypionka H. (1996). A common pathway of sulfide oxidation by sulfate-reducing bacteria.  
494 *FEMS Microbiol Lett* **144**: 129–134.
- 495 Garcia Costas AM, Liu Z, Tomsho LP, Schuster SC, Ward DM, Bryant DA. (2012). Complete genome of *Candidatus*  
496 *Chloracidobacterium thermophilum*, a chlorophyll-based photoheterotroph belonging to the phylum *Acidobacteria*. *Environ*  
497 *Microbiol* **14**: 177–190.
- 498 Ghosh W, Dam B. (2009). Biochemistry and molecular biology of lithotrophic sulfur oxidation by taxonomically and  
499 ecologically diverse bacteria and archaea. *FEMS Microbiol Rev* **33**: 999–1043.
- 500 Giuffrè A, Borisov VB, Arese M, Sarti P, Forte E. (2014). Cytochrome *bd* oxidase and bacterial tolerance to oxidative and  
501 nitrosative stress. *Biochim Biophys Acta* **1837**: 1178–1187.
- 502 Greening C, Biswas A, Carere CR, Jackson CJ, Taylor MC, Stott MB *et al.* (2016). Genomic and metagenomic surveys of  
503 hydrogenase distribution indicate H<sub>2</sub> is a widely utilised energy source for microbial growth and survival. *ISME J* **10**: 761–  
504 777.
- 505 Greening C, Carere CR, Rushton-Green R, Harold LK, Hards K, Taylor MC *et al.* (2015). Persistence of the dominant soil  
506 phylum *Acidobacteria* by trace gas scavenging. *Proc Natl Acad Sci USA* **112**: 10497–10502.

507 Hausmann B, Knorr K-H, Schreck K, Tringe SG, Glavina del Rio T, Loy A *et al.* (2016). Consortia of low-abundance bacteria  
508 drive sulfate reduction-dependent degradation of fermentation products in peat soil microcosms. *The ISME Journal* **10**: 2365–  
509 2375.

510 Hillesland KL, Lim S, Flowers JJ, Turkarslan S, Pinel N, Zane GM *et al.* (2014). Erosion of functional independence early in  
511 the evolution of a microbial mutualism. *Proc Natl Acad Sci USA* **111**: 14822–14827.

512 Hittel DS, Voordouw G. (2000). Overexpression, purification and immunodetection of DsrD from *Desulfovibrio vulgaris*  
513 Hildenborough. *Antonie Van Leeuwenhoek* **77**: 271–280.

514 Holkenbrink C, Barbas SO, Møllerup A, Otaki H, Frigaard N-U. (2011). Sulfur globule oxidation in green sulfur bacteria is  
515 dependent on the dissimilatory sulfite reductase system. *Microbiology* **157**: 1229–1239.

516 Huber KJ, Geppert AM, Wanner G, Fösel BU, Wüst PK, Overmann J. (2016). The first representative of the globally  
517 widespread subdivision 6 *Acidobacteria*, *Vicinamibacter silvestris* gen. nov., sp. nov., isolated from subtropical savannah soil.  
518 *Int J Syst Evol Microbiol* **66**: 2971–2979.

519 Huber KJ, Wüst PK, Rohde M, Overmann J, Foesel BU. (2014). *Aridibacter famidurans* gen. nov., sp. nov. and *Aridibacter*  
520 *kavangonensis* sp. nov., two novel members of subdivision 4 of the *Acidobacteria* isolated from semiarid savannah soil. *Int J*  
521 *Syst Evol Microbiol* **64**: 1866–1875.

522 Huerta-Cepas J, Szklarczyk D, Forslund K, Cook H, Heller D, Walter MC *et al.* (2016). eggNOG 4.5: a hierarchical orthology  
523 framework with improved functional annotations for eukaryotic, prokaryotic and viral sequences. *Nucleic Acids Res* **44**: D286–  
524 D293.

525 Hug La, Thomas BC, Sharon I, Brown CT, Sharma R, Hettich RL *et al.* (2016). Critical biogeochemical functions in the  
526 subsurface are associated with bacteria from new phyla and little studied lineages. *Environ Microbiol* **18**: 159–173.

527 Hyatt D, Chen G-L, Locascio PF, Land ML, Larimer FW, Hauser LJ. (2010). Prodigal: prokaryotic gene recognition and  
528 translation initiation site identification. *BMC Bioinformatics* **11**: 119.

529 Ijssennagger N, Belzer C, Hooiveld GJ, Dekker J, van Mil SWC, Müller M *et al.* (2015). Gut microbiota facilitates dietary  
530 heme-induced epithelial hyperproliferation by opening the mucus barrier in colon. *Proc Natl Acad Sci USA* **112**: 10038–10043.

531 Ivanova AA, Wegner C-E, Kim Y, Liesack W, Dedysh SN. (2016). Identification of microbial populations driving biopolymer  
532 degradation in acidic peatlands by metatranscriptomic analysis. *Mol Ecol* **25**: 4818–4835.

533 Izumi H, Nunoura T, Miyazaki M, Mino S, Toki T, Takai K *et al.* (2012). *Thermotomaculum hydrothermale* gen. nov., sp.  
534 nov., a novel heterotrophic thermophile within the phylum *Acidobacteria* from a deep-sea hydrothermal vent chimney in the  
535 Southern Okinawa Trough. *Extremophiles* **16**: 245–253.

536 Janssen PH. (2006). Identifying the dominant soil bacterial taxa in libraries of 16S rRNA and 16S rRNA genes. *Appl Environ*  
537 *Microbiol* **72**: 1719–1728.

- 538 Juottonen H, Eiler A, Biasi C, Tuittila E-S, Yrjälä K, Fritze H. (2017). Distinct anaerobic bacterial consumers of cellobiose-  
539 derived carbon in boreal fens with different CO<sub>2</sub>/CH<sub>4</sub> production ratios. *Appl Environ Microbiol* **83**: e02533–16.
- 540 Kirschke S, Bousquet P, Ciais P, Saunio M, Canadell JG, Dlugokencky EJ *et al.* (2013). Three decades of global methane  
541 sources and sinks. *Nat Geosci* **6**: 813–823.
- 542 Knorr K-H, Blodau C. (2009). Impact of experimental drought and rewetting on redox transformations and methanogenesis in  
543 mesocosms of a northern fen soil. *Soil Biol Biochem* **41**: 1187–1198.
- 544 Konstantinidis KT, Rosselló-Móra R, Amann R. (2017). Uncultivated microbes in need of their own taxonomy. *ISME J* **11**:  
545 2399–2406.
- 546 Kraigher B, Stres B, Hacin J, Ausec L, Mahne I, van Elsas JD *et al.* (2006). Microbial activity and community structure in two  
547 drained fen soils in the Ljubljana Marsh. *Soil Biol Biochem* **38**: 2762–2771.
- 548 Kulichevskaya IS, Suzina NE, Liesack W, Dedysh SN. (2010). *Bryobacter aggregatus* gen. nov., sp. nov., a peat-inhabiting,  
549 aerobic chemo-organotroph from subdivision 3 of the *Acidobacteria*. *Int J Syst Evol Microbiol* **60**: 301–306.
- 550 Kulichevskaya IS, Suzina NE, Rijpstra WIC, Sinninghe Damsté JS, Dedysh SN. (2014). *Paludibaculum fermentans* gen. nov.,  
551 sp. nov., a facultative anaerobe capable of dissimilatory iron reduction from subdivision 3 of the *Acidobacteria*. *Int J Syst Evol*  
552 *Microbiol* **64**: 2857–2864.
- 553 Küsel K, Blöthe M, Schulz D, Reiche M, Drake HL. (2008). Microbial reduction of iron and porewater biogeochemistry in  
554 acidic peatlands. *Biogeosciences* **5**: 1537–1549.
- 555 Langmead B, Salzberg SL. (2012). Fast gapped-read alignment with Bowtie 2. *Nat Methods* **9**: 357–359.
- 556 Laska S, Lottspeich F, Kletzin A. (2003). Membrane-bound hydrogenase and sulfur reductase of the hyperthermophilic and  
557 acidophilic archaeon *Acidianus ambivalens*. *Microbiology* **149**: 2357–2371.
- 558 Lenk S, Moraru C, Hahnke S, Arnds J, Richter M, Kube M *et al.* (2012). *Roseobacter* clade bacteria are abundant in coastal  
559 sediments and encode a novel combination of sulfur oxidation genes. *ISME J* **6**: 2178–2187.
- 560 Liao Y, Smyth GK, Shi W. (2014). featureCounts: an efficient general purpose program for assigning sequence reads to  
561 genomic features. *Bioinformatics* **30**: 923–930.
- 562 Liesack W, Bak F, Kreft J-U, Stackebrandt E. (1994). *Holophaga foetida* gen. nov., sp. nov., a new, homoacetogenic  
563 bacterium degrading methoxylated aromatic compounds. *Arch Microbiol* **162**: 85–90.
- 564 Losey NA, Stevenson BS, Busse H-J, Sinninghe Damsté JS, Rijpstra WIC, Rudd S *et al.* (2013). *Thermoanaerobaculum*  
565 *aquaticum* gen. nov., sp. nov., the first cultivated member of *Acidobacteria* subdivision 23, isolated from a hot spring. *Int J*  
566 *Syst Evol Microbiol* **63**: 4149–4157.
- 567 Love MI, Huber W, Anders S. (2014). Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2.

- 568 *Genome Biol* **15**: 550.
- 569 Loy A, Küsel K, Lehner A, Drake HL, Wagner M. (2004). Microarray and functional gene analyses of sulfate-reducing  
570 prokaryotes in low-sulfate, acidic fens reveal cooccurrence of recognized genera and novel lineages. *Appl Environ Microbiol*  
571 **70**: 6998–7009.
- 572 Lübke YJ, Youn H-S, Timkovich R, Dahl C. (2006). Siro(haem)amide in *Allochromatium vinosum* and relevance of DsrL and  
573 DsrN, a homolog of cobyrinic acid *a,c*-diamide synthase, for sulphur oxidation. *FEMS Microbiol Lett* **261**: 194–202.
- 574 Meyer B, Imhoff JF, Kuever J. (2007). Molecular analysis of the distribution and phylogeny of the *soxB* gene among sulfur-  
575 oxidizing bacteria: evolution of the Sox sulfur oxidation enzyme system. *Environ Microbiol* **9**: 2957–2977.
- 576 Mizuno N, Voordouw G, Miki K, Sarai A, Higuchi Y. (2003). Crystal structure of dissimilatory sulfite reductase D (DsrD)  
577 protein—possible interaction with B- and Z-DNA by its winged-helix motif. *Structure* **11**: 1133–1140.
- 578 Müller AL, Kjeldsen KU, Rattei T, Pester M, Loy A. (2015). Phylogenetic and environmental diversity of DsrAB-type  
579 dissimilatory (bi)sulfite reductases. *ISME J* **9**: 1152–1165.
- 580 Myers MR, King GM. (2016). Isolation and characterization of *Acidobacterium ailaui* sp. nov., a novel member of  
581 *Acidobacteria* subdivision 1, from a geothermally heated Hawaiian microbial mat. *Int J Syst Evol Microbiol* **66**: 5328–5335.
- 582 Notredame C, Higgins DG, Heringa J. (2000). T-Coffee: a novel method for fast and accurate multiple sequence alignment. *J*  
583 *Mol Biol* **302**: 205–217.
- 584 Otwell AE, Sherwood RW, Zhang S, Nelson OD, Li Z, Lin H *et al.* (2015). Identification of proteins capable of metal  
585 reduction from the proteome of the Gram-positive bacterium *Desulfotomaculum reducens* MI-1 using an NADH-based activity  
586 assay. *Environ Microbiol* **17**: 1977–1990.
- 587 Pankratov TA, Dedysh SN. (2010). *Granulicella paludicola* gen. nov., sp. nov., *Granulicella pectinivorans* sp. nov.,  
588 *Granulicella aggregans* sp. nov. and *Granulicella rosea* sp. nov., acidophilic, polymer-degrading acidobacteria from  
589 *Sphagnum* peat bogs. *Int J Syst Evol Microbiol* **60**: 2951–2959.
- 590 Pankratov TA, Ivanova AO, Dedysh SN, Liesack W. (2011). Bacterial populations and environmental factors controlling  
591 cellulose degradation in an acidic *Sphagnum* peat. *Environ Microbiol* **13**: 1800–1814.
- 592 Pankratov TA, Kirsanova LA, Kaparullina EN, Kevbrin VV, Dedysh SN. (2012). *Telmatobacter bradus* gen. nov., sp. nov., a  
593 cellulolytic facultative anaerobe from subdivision 1 of the *Acidobacteria*, and emended description of *Acidobacterium*  
594 *capsulatum* Kishimoto *et al.* 1991. *Int J Syst Evol Microbiol* **62**: 430–437.
- 595 Parks DH, Imelfort M, Skennerton CT, Hugenholtz P, Tyson GW. (2015). CheckM: assessing the quality of microbial  
596 genomes recovered from isolates, single cells, and metagenomes. *Genome Res* **25**: 1043–1055.
- 597 Pereira IAC, Ramos AR, Grein F, Marques MC, Marques da Silva S, Venceslau SS. (2011). A comparative genomic analysis  
598 of energy metabolism in sulfate reducing bacteria and archaea. *Front Microbiol* **2**: 69.

- 599 Pester M, Bittner N, Deevong P, Wagner M, Loy A. (2010). A ‘rare biosphere’ microorganism contributes to sulfate reduction  
600 in a peatland. *ISME J* **4**: 1591–1602.
- 601 Pester M, Knorr K-H, Friedrich MW, Wagner M, Loy A. (2012). Sulfate-reducing microorganisms in wetlands – fameless  
602 actors in carbon cycling and climate change. *Front Microbiol* **3**: 72.
- 603 Rabus R, Hansen TA, Widdel F. (2013). Dissimilatory sulfate- and sulfur-reducing prokaryotes. In: Rosenberg E, DeLong EF,  
604 Lory S, Stackebrandt E, Thompson F (eds). *The Prokaryotes – prokaryotic physiology and biochemistry*. Springer Berlin  
605 Heidelberg, pp 309–404.
- 606 Rabus R, Venceslau SS, Wöhlbrand L, Voordouw G, Wall JD, Pereira IAC. (2015). A post-genomic view of the  
607 ecophysiology, catabolism and biotechnological relevance of sulphate-reducing prokaryotes. *Adv Microb Physiol* **66**: 55–321.
- 608 Ramel F, Amrani A, Pieulle L, Lamrabet O, Voordouw G, Seddiki N *et al.* (2013). Membrane-bound oxygen reductases of the  
609 anaerobic sulfate-reducing *Desulfovibrio vulgaris* Hildenborough: roles in oxygen defence and electron link with periplasmic  
610 hydrogen oxidation. *Microbiology* **159**: 2663–2673.
- 611 Rawat SR, Männistö MK, Bromberg Y, Häggblom MM. (2012). Comparative genomic and physiological analysis provides  
612 insights into the role of *Acidobacteria* in organic carbon utilization in Arctic tundra soils. *FEMS Microbiol Ecol* **82**: 341–355.
- 613 Rodriguez-R LM, Konstantinidis KT. (2014). Bypassing cultivation to identify bacterial species. *Microbe Mag* **9**: 111–118.
- 614 Santos AA, Venceslau SS, Grein F, Leavitt WD, Dahl C, Johnston DT *et al.* (2015). A protein trisulfide couples dissimilatory  
615 sulfate reduction to energy conservation. *Science* **350**: 1541–1545.
- 616 Saunio M, Bousquet P, Poulter B, Peregon A, Ciais P, Canadell JG *et al.* (2016). The global methane budget 2000–2012.  
617 *Earth Syst Sci Data* **8**: 697–751.
- 618 Sánchez-Andrea I, Rodríguez N, Amils R, Sanz JL. (2011). Microbial diversity in anaerobic sediments at Río Tinto, a naturally  
619 acidic environment with a high heavy metal content. *Appl Environ Microbiol* **77**: 6085–6093.
- 620 Schmalenberger A, Drake HL, Küsel K. (2007). High unique diversity of sulfate-reducing prokaryotes characterized in a depth  
621 gradient in an acidic fen. *Environ Microbiol* **9**: 1317–1328.
- 622 Schmidt O, Horn MA, Kolb S, Drake HL. (2015). Temperature impacts differentially on the methanogenic food web of  
623 cellulose-supplemented peatland soil. *Environ Microbiol* **17**: 720–734.
- 624 Serkebaeva YM, Kim Y, Liesack W, Dedysh SN. (2013). Pyrosequencing-based assessment of the *Bacteria* diversity in  
625 surface and subsurface peat layers of a northern wetland, with focus on poorly studied phyla and candidate divisions. *PLoS*  
626 *One* **8**: e63994.
- 627 Simon J, Kroneck PMH. (2013). Microbial sulfite respiration.
- 628 Steger D, Wentrup C, Brauneegger C, Deevong P, Hofer M, Richter A *et al.* (2011). Microorganisms with novel dissimilatory



- 629 (bi)sulfite reductase genes are widespread and part of the core microbiota in low-sulfate peatlands. *Appl Environ Microbiol* **77**:  
630 1231–1242.
- 631 Stetter KO. (1988). *Archaeoglobus fulgidus* gen. nov., sp. nov.: a new taxon of extremely thermophilic archaeobacteria. *Syst*  
632 *Appl Microbiol* **10**: 172–173.
- 633 Tank M, Bryant DA. (2015). *Chloracidobacterium thermophilum* gen. nov., sp. nov.: an anoxygenic microaerophilic  
634 chlorophotoheterotrophic acidobacterium. *Int J Syst Evol Microbiol* **65**: 1426–1430.
- 635 Thorup C, Schramm A, Findlay AJ, Finster KW, Schreiber L. (2017). Disguised as a sulfate reducer: growth of the  
636 deltaproteobacterium *Desulfurivibrio alkaliphilus* by sulfide oxidation with nitrate. *MBio* **8**: e00671–17.
- 637 Tveit A, Schwacke R, Svenning MM, Urich T. (2013). Organic carbon transformations in high-Arctic peat soils: key functions  
638 and microorganisms. *ISME J* **7**: 299–311.
- 639 Urbanová Z, Bárta J. (2014). Microbial community composition and *in silico* predicted metabolic potential reflect  
640 biogeochemical gradients between distinct peatland types. *FEMS Microbiol Ecol* **90**: 633–646.
- 641 Vallenet D, Calteau A, Cruveiller S, Gachet M, Lajus A, Josso A *et al.* (2017). MicroScope in 2017: an expanding and  
642 evolving integrated resource for community expertise of microbial genomes. *Nucleic Acids Res* **45**: D517–D528.
- 643 Venceslau SS, Stockdreher Y, Dahl C, Pereira IAC. (2014). The ‘bacterial heterodisulfide’ DsrC is a key protein in  
644 dissimilatory sulfur metabolism. *Biochim Biophys Acta* **1837**: 1148–1164.
- 645 Wang Y, Sheng H-F, He Y, Wu J-Y, Jiang Y-X, Tam NF-Y *et al.* (2012). Comparison of the levels of bacterial diversity in  
646 freshwater, intertidal wetland, and marine sediments by using millions of illumina tags. *Appl Environ Microbiol* **78**: 8264–  
647 8271.
- 648 Ward NL, Challacombe JF, Janssen PH, Henrissat B, Coutinho PM, Wu M *et al.* (2009). Three genomes from the phylum  
649 *Acidobacteria* provide insight into the lifestyles of these microorganisms in soils. *Appl Environ Microbiol* **75**: 2046–2056.
- 650 Wasmund K, Mußmann M, Loy A. (2017). The life sulfuric: microbial ecology of sulfur cycling in marine sediments. *Environ*  
651 *Microbiol Rep* **9**: 323–344.
- 652 Watanabe T, Kojima H, Fukui M. (2016). Identity of major sulfur-cycle prokaryotes in freshwater lake ecosystems revealed by  
653 a comprehensive phylogenetic study of the dissimilatory adenylylsulfate reductase. *Sci Rep* **6**: 36262.
- 654 Weber KA, Achenbach LA, Coates JD. (2006). Microorganisms pumping iron: anaerobic microbial iron oxidation and  
655 reduction. *Nat Rev Microbiol* **4**: 752–764.
- 656 Weissgerber T, Sylvester M, Kröninger L, Dahl C. (2014). A comparative quantitative proteomic study identifies new proteins  
657 relevant for sulfur oxidation in the purple sulfur bacterium *Allochromatium vinosum*. *Appl Environ Microbiol* **80**: 2279–2292.
- 658





## 660 **Figure legends**

### 661 **Figure 1**

662 Microbial community composition in Schlössnerbrunnen II peatland in samples from different years and soil depths. (a)  
663 Abundance of phyla and proteobacterial classes in the native soil (relative to all classified reads/amplicons). Taxa less  
664 abundant than 1% are grouped in grey. Coverage abundance is based on metagenomic reads mapped to classified scaffolds.  
665 Amplicon abundance is based on *rrn* operon-copy number-corrected abundance of 16S rRNA gene operational taxonomic  
666 units (Hausmann *et al.*, 2016). (b) Relative abundance of acidobacterial subdivisions (SD) in the native soil samples as  
667 determined by 16S rRNA gene qPCR assays. In addition, all subdivisions more abundant than 1% in a 16S rRNA gene  
668 amplicon dataset are shown (Hausmann *et al.*, 2016).

### 669 **Figure 2**

670 Organization of dissimilatory sulfur metabolism genes on acidobacterial MAGs SbA1–7. Red: *sat*; orange: *aprBA*, *qmoABC*;  
671 green: *dsrABCMKJOPM2K2*; blue: *dsrD*; turquoise: *dsrL*; violet: *dsrNVWa*; pink: *suyAB*, *comC*, *slcC*; 1–4 (grey): syntenic  
672 genes encoding for conserved proteins of unknown function; white: genes of unknown function or not involved in sulfur  
673 metabolism. In SbA2 all genes are on one scaffold (scaffold 0lkb). Gene fragments at contig borders are indicated by an  
674 asterisk. *aprB* in SbA6, indicated by two asterisks, is truncated, which indicates a pseudogene or is due to an assembly error.  
675 Scaffolds are separated by two slashes.

### 676 **Figure 3**

677 Metabolic model as inferred from analysis of acidobacterial MAGs SbA1–7. Sulfur metabolism is highlighted in yellow.  
678 Enzymes and transporters are shown in blue font. Glycoside hydrolases are shown in pink font (Supplementary Table S2).  
679 Extracellular compounds are in parentheses. A slash (/) indicates isozymes, i.e., enzymes that perform the same function, but  
680 are distinctly different or have more than one established name. AcdA+B, MaeB+Pta, MeaB+Mce, Tal+Pgi: bifunctional  
681 fusion genes/proteins. Otherwise the plus sign (+) indicates protein complexes. TCA: tricarboxylic acid cycle. FDH: formate  
682 dehydrogenase. Hase: hydrogenase. NDH: NADH dehydrogenase. HCO: haem-copper oxidase. TO: terminal oxidase. KDG:  
683 2-dehydro-3-deoxy-D-gluconate. KDGP: 2-dehydro-3-deoxy-D-gluconate 6-phosphate. Expression of at least one copy of  
684 every enzyme and transporter was observed in the incubation samples.

### 685 **Figure 4**

686 Glycoside hydrolase genes are enriched in acidobacterial genomes/MAGs compared to genomes from other taxa that encode  
687 DsrA/DsrB. DsrAB-containing MAGs SbA1–7 are shown as solid symbols and numbered accordingly. X-axis shows the total  
688 number of predicted CDS per genome/MAG.

### 689 **Figure 5**

690 Putative lifestyles of DsrAB-encoding *Acidobacteria*.

691  
692  
693  
694

**Fig. 1.**







Fig. 2.



**Fig. 3.**



**Fig. 4.**



**Fig. 5.**

