

UC Irvine

UC Irvine Previously Published Works

Title

Health effects of acid aerosols formed by atmospheric mixtures.

Permalink

<https://escholarship.org/uc/item/7kj4s0hx>

Journal

Environmental Health Perspectives, 79

Authors

Kleinman, M T

Phalen, R F

Mautz, W J

et al.

Publication Date

1989-02-01

Peer reviewed

Health Effects of Acid Aerosols Formed by Atmospheric Mixtures

by Michael T. Kleinman,* Robert F. Phalen,*
William J. Mautz,* Richard C. Mannix,*
Thomas R. McClure,* and T. Timothy Crocker*

Under ambient conditions, sulfur and nitrogen oxides can react with photochemical products and airborne particles to form acidic vapors and aerosols. Inhalation toxicological studies were conducted, exposing laboratory animals, at rest and during exercise, to multicomponent atmospheric mixtures under conditions favorable to the formation of acidic reaction products. Effects of acid and ozone mixtures on early and late clearance of insoluble radioactive particles in the lungs of rats appeared to be dominated by the oxidant component (i.e., the mixture did cause effects that were significantly different from those of ozone alone). Histopathological evaluations showed that sulfuric acid particles alone did not cause inflammatory responses in centriacinar units of rat lung parenchyma (expressed in terms of percent lesion area) but did cause significant damage (cell killing followed by a wave of cell replication) in nasal respiratory epithelium, as measured by uptake of tritiated thymidine in the DNA of replicating cells. Mixtures of ozone and nitrogen dioxide, which form nitric acid, caused significant inflammatory responses in lung parenchyma (in excess of effects seen in rats exposed to ozone alone), but did not damage nasal epithelium. Mixtures containing acidic sulfate particles, ozone, and nitrogen dioxide damaged both lung parenchyma and nasal epithelia. In rats exposed at rest, the response of the lung appeared to be dominated by the oxidant gas-phase components, while responses in the nose were dominated by the acidic particles. In rats exposed at exercise, however, mixtures of ozone and sulfuric acid particles significantly (2.5-fold) elevated the degree of lung lesion formation over that seen in rats exposed to ozone alone under an identical exercise protocol.

Introduction

Data obtained after severe air pollution episodes, such as those that occurred in London in the 1950s, suggest that acid particles, especially sulfuric acid (H_2SO_4), were present in high concentrations and may have contributed to increased morbidity and mortality among London residents (1,2). In fact, many of the so-called killer smog episodes observed during the past 60 years have occurred under conditions that were favorable to the formation of airborne acids (3). The air during these episodes contained mixtures of gases, vapors and particles, including sulfur dioxide (SO_2), nitrogen dioxide (NO_2), nitric oxide (NO), substantial quantities of carbonaceous and noncarbonaceous particles, trace metals, and water in vapor and droplet forms. Under ambient conditions these

substances can react to yield sulfate- and nitrate-containing acidic particles and vapors and, when sunlight is present, ozone (O_3) (4-8).

The amount of sulfate directly emitted to the global atmosphere by natural or human processes as sulfur trioxide of sulfuric acid is small. The major acid sulfate formation pathway is aqueous phase oxidation of SO_2 in droplet aerosols at high relative humidity (9). The rate of conversion of SO_2 to sulfate in coal-fired source plumes is variable but is generally estimated to be less than 15% per hour (and frequently is less than 5% per hour) (10,11). Oxides of nitrogen, emitted from anthropogenic stationary and mobile combustion sources, are converted to nitric acid (HNO_3) vapor and particulate nitrates by photochemical processes involving hydrocarbons and O_3 during daylight periods and via reactions with O_3 at night (7,12) and indoors (13). In fogs, the rate of conversion of SO_2 to sulfate is greatly accelerated (14) and, in addition, if aldehydes or ketones are present, strongly acidic hydroxymethanesulfonic acids can be formed (15). From

*Air Pollution Health Effects Laboratory, Department of Community and Environmental Medicine, University of California, Irvine, CA 92717.

this brief review, it is apparent that concentrations of acidic and oxidant compounds in ambient air could be strongly correlated. Such associations have, in fact, been reported in field epidemiological studies (16).

Animal toxicological studies have been performed to examine the expected range and mechanisms of health effects attributable to oxidant-acid mixtures. Some of these studies have indicated that synergistic interactions occur (17–19), but others have not (20,21). The aforementioned studies, however, were not designed to specifically examine the health effect potentials for reacting mixtures that lead to formation of airborne acidity. This paper will report on recent inhalation toxicologic studies using laboratory rats in which the atmospheres contained acid precursors and the exposure conditions permitted formation of acids by gas-phase and gas and particle reactions. The studies described involve complex atmospheric mixtures with up to seven intentionally added components, plus their reaction products. The end points that were selected as indices of relevant (to human populations) biological effects include: a) changes in clearance rates of inhaled insoluble radioactive tracer particles, both short term (presumably from nasal and tracheobronchial airways) and long term (presumably from alveolar regions of the lung); b) histopathologic focal lesions in the deep lung; and c) cell replacement (which is presumably stimulated by cell killing) in upper respiratory tract epithelium.

Methods

Atmosphere Generation and Characterization

The atmospheres for study contained acidic and nonacidic sulfates, SO_2 , O_3 , NO_2 , catalytic metal ions (Fe^{+3} , Mn^{+2}), and insoluble particles. Sulfate [H_2SO_4 or ammonium sulfate ($(\text{NH}_4)_2\text{SO}_4$)] aerosols containing catalytic metal ions were generated by nebulizing a solution containing H_2SO_4 or $(\text{NH}_4)_2\text{SO}_4$, ferric sulfate, and manganese (II) sulfate into highly purified dilution of air using a Collison nebulizer (22). Aerosols were passed through an ^{85}Kr discharger to reduce static charges to the Boltzmann equilibrium level. NO_2 and SO_2 were precisely metered into purified air from compressed gas cylinders. O_3 was generated from medical-grade oxygen using a commercial ozonizer (Sander Ozonizer, Type III). Insoluble particles of ferric oxide (Fe_2O_3) were generated by nebulization of dilute colloidal suspensions of Fe_2O_3 , yielding spherical aggregate particles having an aerodynamic diameter of about $0.3\text{ }\mu\text{m}$. The aerosols were dried at 50°C in a flow-through furnace, diluted with dry air, passed through an ^{85}Kr discharger, and introduced into the multicomponent atmosphere mixture.

NO_2 and NO were monitored using a chemiluminescent detector (Monitor Labs 8080). SO_2 was monitored using a TECO Series 43 pulsed fluorescence detector. O_3 was measured using a UV absorption mon-

itor (Dasibi Model 1003-AH). During exposure the atmosphere was monitored using real-time detectors: Climet 208 and GCA RAM-1 optical particle detectors for aerosols with diameters from 0.3 to $10\text{ }\mu\text{m}$ and a TSI Model 3030 electrical aerosol size analyzer for particles with diameters ranging from 0.05 to $0.5\text{ }\mu\text{m}$. Cascade impactor samples (Sierra Model 212 8-stage impactor) were collected to provide size-classified particles for gravimetric and chemical analyses. Aerosols were collected on precleaned quartz fiber or fluorocarbon filters for gravimetric determinations and for chemical analyses of atmospheric components and reaction products. Nylon backup filters were used to collect HNO_3 vapor (from the reaction of NO_2 and O_3). A point-to-plane electrostatic precipitator (23) was used to obtain aerosol samples for electron microscopic size characterization. Ion exchange chromatography was used to analyze for sulfate, sulfite, nitrate, and nitrite ions in filter extracts. Samples extracted in dilute acid were analyzed for manganese and iron by atomic absorption spectrometry.

Animals and Exposures

Six-week-old, barrier-reared, male Sprague-Dawley rats (Hilltop Lab Animals, Inc.) were delivered to the laboratory in filter-equipped shipping containers. The rats were housed in a laminar-flow air barrier system (with chemical and particulate scrubbing) for 1 week prior to the start of each experiment. Microbiological assays supplied by the breeder and histopathological examinations performed at our laboratory were used to determine that the rats were free of major respiratory infections. Rats were randomly assigned to groups (6–12 per group); control groups were exposed to purified air, and matched groups were exposed to test atmospheres. In early experiments, rats were exposed whole-body in a chamber. This resulted in significant neutralization of acids by ammonia from excreta, hence the recent studies were performed using a nose-only exposure system, shown schematically in Figure 1A. In the earlier whole-body exposure studies with H_2SO_4 , to attain a titratable acid concentration equivalent to 3.5 mg/m^3 , it was necessary to maintain the total aerosol sulfate mass at 5 mg/m^3 (24).

Exposures of exercising rats were performed using a modified Quinton 10-rodent treadmill. The treadmill (Fig. 1B) was enclosed and sealed against significant leaks (25), and a stainless-steel manifold equally distributed the flow to the 10 treadmill runways. The inner surfaces of the 10 treadmill runways were lined with a stainless-steel sheet to minimize reactions with corrosive gases. Ports in each runway permitted sampling of exposure atmospheres in the rat breathing zone. Exhaust lines at the rear of the treadmill conducted the atmosphere and expired respiratory gases to a mixing chamber for metabolic measurements. Measurements of O_2 , CO_2 , and flow rates of inspired and mixed expired gases permitted the determination of average metabolic rate of exercising animals. Rats




FIGURE 1. Systems used for exposing rats to acid-containing pollutant mixtures. (A) The nose-only system prevents neutralization of acids by ammonia liberated by excreta. (B) The treadmill systems allows for simultaneous exposures of up to 10 rats.

were exposed to pollutant atmospheres or to clean air (sham exposures) for 4 hr at rest, 3 hr at light (8 meters/min) continuous exercise, or 1.5 hr at intermittent moderate exercise (24 meters/min for 6 min, rest for 1 min).

Particle Clearance

Clearance of inhaled radioactively labeled insoluble particles (26) by mucociliary and macrophage-mediated processes was quantitatively measured to determine the degree to which the ability of the respiratory tract to eliminate deposited particulate material (and by implication, toxic or infectious agents) was affected

by exposure to pollutant atmospheres. Tracer particles ($1.5 \mu\text{m}$ mass median aerodynamic diameter, MMAD) labeled with ^{51}Cr (200 mCi/g particles) were suspended in distilled water, nebulized into dry dilution air, and the resulting dried particles passed through an ^{85}Kr discharger and routed to a nose-only exposure chamber in which up to 20 rats could be simultaneously exposed (24,27). About 20 min was sufficient to deliver sufficient radioactivity to the rats' lungs. Initial chest counts were obtained on all of the rats, following which they were exposed for 4 hr to either purified air or to a pollutant atmosphere. The chest count data were used as baselines for the subsequent clearance measurements. Early clearance, presumably representing material removed from the upper

respiratory tract, was characterized by collecting and counting the rat's feces at 10 fixed times during the first 50 hr after the deposition of the tracer particles. Late, presumably deep lung, clearance was characterized by 5 chest counts during the subsequent 400 hr (27).

Autoradiography and Histopathology

Uptake of Radiolabeled DNA Precursors by Cells. When inhaled pollutants kill respiratory tract cells, the damaged tissue is repaired by the replication of cells adjacent to the dead cells. In preparation for cell division, DNA precursors are taken up by the replicating cells and, by introducing tritiated thymidine (^3HT), a radiolabeled nucleotide, it is possible to radioactively tag these cells and subsequently map their locations and numbers in the airways by autoradiography (28,29). These locations, then, represent epithelia in which cell death has been followed by a wave of new cell production. The number of cells that take up ^3HT is an index of the magnitude of the tissue damage.

Seventeen hours after the end of a pollutant or clean air exposure, the animals were injected IP with ^3HT (1 $\mu\text{Ci/g}$ of body weight), and, 1 hr later, were deeply anesthetized and killed. The lungs were excised and fixed by airway perfusion with 10% neutral-buffered formalin at 30 cm H_2O pressure for 72 hr (30). The trachea and larynx of each rat were dissected out and immersed in formalin. The heads were removed and the portions of the heads containing the intact nasal cavity fixed in formalin and decalcified with neutral TIS-buffered EDTA for 3 weeks. The trachea was split longitudinally, embedded in glycol methacrylate, and sectioned at 2 μm . Specimens of the nasal cavity, 2 to 3 mm thick, were cut vertically through the hard palate and nasal septa at level II (31), embedded in glycol methacrylate, and sectioned at 2 μm . Rates of cell turnover in tracheal and nasal sections were evaluated using autoradiography. Slides were dipped in Kodak NTB-2 nuclear track emulsion, and exposed for 30 days in a light-tight refrigerated box. After exposure, the slides were developed photographically, stained with toluidine blue and examined by light microscopy. Four distinct epithelial types were differentiated in the nasal sections (Fig. 2): squamous, olfactory, and two types of respiratory epithelia. Of the respiratory epithelia, one type, designated R, is a pseudostratified high columnar epithelium with ciliated and goblet cells and is located over the medial surfaces of the turbinates and the nasal septum. The second type, designated R1, is simple columnar epithelium with ciliated cells, but without goblet cells, and is found in the maxillary fossae lateral to the turbinates. Epithelial cells were quantitated by aligning a microscope eyepiece grid with the basement membrane and counting all cells, differentiating by cell type, within each of 10 grid spaces. The number of cells labeled with ^3HT per unit grid space was expressed as the percent of labeled to total cells, by cell type.

Lesions in Lung Parenchymal Tissue. A single 2 to 4 hr inhalation exposure to O_3 causes death of alveolar epithelial cells. Macrophages and some other leukocytes infiltrate the injured tissue to remove dead cells. The process of macrophage and leukocyte accumulation is called inflammation, and the locus of the inflammatory response is referred to as a lesion. Oxidant-related inflammatory responses occur in scattered centriacinar units of the alveolar parenchymal zone of the lung. Each centriacinar unit consists of the smallest airway, the terminal bronchiole, and the alveolar duct and alveolar sacs to which it leads. These units comprise most of the alveolar (parenchymal) zone of a mammalian lung, but only a fraction of these centriacinar units are injured after exposure and develop an infiltrate of inflammatory cells by which they are identified as lesions. Since most inflamed centriacinar units are surrounded by units of normal appearance, the lesions are isolated and are referred to as focal lesions. At exposure concentrations of about 0.6 to 0.8 ppm of O_3 , focal lesions appear by 18 hr postexposure, reach maximum intensity at about 48 hr postexposure, and subsequently resolve. At higher concentrations, O_3 exposure causes more widespread destruction of alveolar epithelial cells, greater numbers of lesions, and pulmonary edema, which can lead to death of the animal (29).

To assess parenchymal lung damage, groups of six or more rats were killed at 48 hr postexposure, and their lungs were fixed in the inflated state, as described previously. The volume of the right, middle lobe was measured by displacement. Longitudinal sections of this lobe, after paraffin embedding, were cut at 6 μm thickness and stained with hematoxylin and eosin. In preliminary studies, we compared rat lung lobes for the distribution and volume fraction of inflammatory foci in the alveolar zone. There were no significant differences between lobes, hence the middle lobe of the right lung has been used routinely. Histologic sections were analyzed quantitatively by light microscopy using a morphometric process described by Elias and Hyde (32). Each lung lobe section was scanned systematically at low microscopic magnification to record the number of times the scanning point overlaid predominantly alveolar areas made up of centriacinar units as opposed to nonalveolar areas that contained structures including bronchi, blood vessels, and connective tissue. This process defines the alveolar and nonalveolar lung volume fractions. The alveolar volume density was recorded and the systematic scan was repeated at higher magnification to record the frequency with which the scanning point overlaid inflammation in a centriacinar unit in an alveolar zone. This frequency divided by the alveolar volume density was used to calculate the extent of lung injury expressed as the volume density of centriacinar units with inflammation within the total volume of the alveolar zone. This datum was multiplied by 100 and recorded as the percent lesion area.




FIGURE 2. Inflammatory responses in centricinar units of lung parenchyma, expressed as percent lesion area, in rats exposed to acid-oxidant mixtures. The points on the graph represent the group mean, and the bars represent 1 SE of the mean. For details of exposure conditions and atmosphere composition, see text.

Data Analysis

Clearance. Fecal clearance curves, which were representative of early clearance (0–50 hr), were fit to log-normal functions, and the time at which 50% of the total activity was excreted was determined for each rat (T_{50}). For late clearance (2–17 days), a biological halftime (T_L) for the radioactivity in the rat's chest (decay corrected to the time of deposition) was determined. Group mean values for control and pollutant atmosphere-exposed rats were compared using two-tailed t -tests, with $p \leq 0.10$.

Histopathology and Autoradiography. Group mean values of lung injury expressed as percent lesion area were contrasted among control and exposed groups using one-tailed analyses of variance. If there

were significant ($p \leq 0.05$) mean differences between the atmosphere groups, pairwise comparisons were examined using the Tukey multiple comparison test.

Results

Effects of Acid Atmospheres on Clearance

Groups of rats were exposed to acidic and nonacidic sulfate aerosols alone and in combination with O₃. Purified air-exposed rats were used as negative controls; rats exposed to O₃ (0.6–0.8 ppm) were used as positive controls.

H₂SO₄ and non-acidic sulfate aerosols alone at 3.5

mg/m³ produced relatively small effects on early and late clearance after whole body exposure (24). The greatest effects observed were by H₂SO₄ and ferric sulfate at low (40%) relative humidity. The effects were of about the same order of magnitude, altering late clearance significantly when compared to the purified air-exposed group of rats.

Combinations of 0.8 ppm O₃ with 3.5 mg/m³ of the acidic and nonacidic sulfate aerosols resulted in effects that were similar to those of O₃ alone. In general, the effects of the aerosols with O₃ were qualitatively similar to those of the O₃ alone, and in most cases the effects of the combination were less than those of O₃ alone at both high and low relative humidity did produce greater effects on early and late clearance than did the O₃ alone at high relative humidity. These effects were of greater magnitude than those produced by the nonacidic aerosols in combination with O₃.

An atmosphere containing 0.8 ppm O₃, 5 ppm SO₂, and 1 mg/m³ H₂SO₄ at high relative humidity significantly delayed early clearance when compared with the purified air control group, but did not affect late clearance. However, the same atmosphere, but with (NH₄)₂SO₄ substituted for H₂SO₄, affected early clearance to the same extent and produced a significant acceleration in late clearance; the effects of this mixture were slightly, but not statistically significantly, greater than those of 0.8 ppm O₃ alone.

Histopathological Effects of Acid Atmospheres

Quantitations of inflammatory changes in centriacinar units in lung parenchyma and of rates of cell turnover in nasal and tracheal epithelia were undertaken in groups of rats exposed to a series of atmospheres ranging in complexity from clean air to a seven-component aerosol and gas mixture. Rats were exposed either at rest, nose-only, or while exercising on the rodent treadmill (Fig. 1B). The data shown in Figure 2 are a composite of independent experiments conducted over several years. Each experiment included its own control group and formed a self-consistent data set.

Effects of Acid Atmospheres on Lung Parenchyma

The inflammatory changes in centriacinar units expressed as percent lesion area in the lung sections of each rat was averaged for groups of 6 to 10 rats. Data from rats exposed in four separate experiments are summarized in Figure 2. Within each experiment the group mean averages were contrasted between exposure conditions using analysis of variance. Tukey's multiple comparison test was used to test whether differences in effects on tissues between the experimental

conditions, on a pairwise basis, were significant at the $\alpha \leq 0.05$ level.

Figure 2A examines a comparison of clean air, O₃ (0.6 ppm), and an acid-oxidant mixture containing H₂SO₄ particles laced with Fe⁺³ and Mn⁺² catalytic ions (1 mg/m³, 0.3 μ m MMAD), SO₂ (5 ppm), and O₃ (0.6 ppm) in rats exposed for 4 hr at rest. The acid-oxidant mixture was significantly more effective than O₃ alone in producing lung lesions. None of the clean air groups in this experiment, or in any of the other experiments, exhibited measurable inflammatory responses.

Figure 2B demonstrates the potentiating effect of 3 hr of low-level exercise. The control group was exposed to the acid mixture without O₃, to O₃ alone, and to the acid-O₃ mixture, at the same concentrations as in the previous experiment. The acid mixture alone did not effect the parenchyma; the oxidant-acid mixture was about three times more potent than O₃ alone.

The potentiating effect of exercise is further explored in Figure 2C, where two different exercise levels are contrasted. The light exercise exposure was sufficient to approximately double the rat's resting oxygen uptake over a 3-hr period. The moderate, intermittent exercise rate increased oxygen uptake rate, but the animals were exposed for only 1.5 hr, and the quantities of pollutants inhaled under the two exercise regimes were approximately equivalent. While the light exercise group showed about the same level of lung damage as in the previous experiment, the moderate intermittent exercise protocol caused a significantly greater response.

Figure 2D examines the interaction of O₃, NO₂ (2.5 ppm), and the SA aerosol mixture. In addition, one group of rats was exposed to an oxidant-aerosol mixture in which (NH₄)₂SO₄ was substituted for H₂SO₄ in the particle mixture. These exposures, which were conducted in resting rats exposed nose-only, demonstrated a striking synergism between O₃ and NO₂, but the addition of either the acid or the nonacid sulfate mixture did not increase the lesion area compared to O₃ and NO₂ alone. One notes that significant concentrations of HNO₃ vapor (2 mg/m³; 0.8 ppm) were observed in atmospheric mixtures which contained NO₂ (2.5 ppm) and O₃ (0.6 ppm) together, both alone, or with other components.

Cell Killing and Turnover in Nasal and Tracheal Tissue

Tissues from the same rats whose lung lesion data were summarized in Figures 2B and 2D were examined for uptake of ³HT. Analyses of variance showed that there were no significant differences in cell labeling index values attributable to exposure atmospheres in the olfactory, respiratory, or tracheal epithelium. The group mean labeling index for squamous epithelium showed that there was a slight, but statistically significant ($p \leq 0.05$), reduction in cell turnover in pollutant-exposed animals relative to the controls.

Analyses of variance demonstrated striking and significant ($p \leq 0.01$) effects on cell killing and turnover in the R1 epithelium. As shown in Figure 3, the animals exposed while exercising to acid-containing atmospheres (SA and SA + O₃) exhibited significantly greater rates of cell turnover than rats at rest exposed to either O₃ alone, clean air, or to the acid-oxidant mixture. As shown in Figure 4, rats exposed at rest to either O₃ or O₃ + NO₂ showed little or no increase in the cell labeling index compared to clean air. However, the addition of acidic or nonacidic sulfate aerosol to the oxidant gas mixture resulted in a 10-fold increase in the fraction of labeled cells. It should be noted that the nonacidic sulfate aerosol was in fact acidified by adsorption of HNO₃ from the vapor phase, which was generated by the reaction of O₃ and NO₂. Reactions between the oxidant gases produced about 2 mg/m³ (0.8 ppm) of HNO₃ vapor. The AS particles contained 370 µg/m³ of nitrate ion (presumably from adsorption of HNO₃ vapor), which would provide acidity equivalent to about 290 µg/m³ H₂SO₄ on a mole-to-mole basis, which is about one-half of the acidity of the SA aerosol.

Discussion and Conclusions

In general, our data do not indicate that the acidity of the aerosol is a major determinant of the effects upon particle clearance by pollutant atmospheres. In aerosol-only atmospheres, the acidic aerosol did not produce effects of greater magnitude than nonacid aerosols. In multicomponent atmospheres containing O₃, the results support the hypothesis that H₂SO₄ plays a minor role in altering respiratory tract clearance.




FIGURE 3. Damage to nasal respiratory epithelium in rats exposed to O₃, acid aerosols (SAM), O₃-acid aerosol mixtures.




FIGURE 4. Damage to nasal respiratory epithelium in rats exposed to combined O₃ plus NO₂ with and without particulate sulfates (SA, AS). Note that in these studies all exposures were conducted with rats at rest.

O₃, which has a relatively low solubility in water and which is not efficiently adsorbed in the upper respiratory tract, did not cause any significant increase in cell turnover rates, as measured by cell labeling with ³HT. The type R1 respiratory epithelium in the noses of rats exposed while exercising was markedly affected by the atmospheres that contained the H₂SO₄ mixture. When O₃ was added to the mixture, the labeling index was not significantly different. The addition, however, of NO₂ to the O₃-particle mixtures produced a striking increase in the impact of the inhaled pollutants on the nasal epithelia. The component to which this effect is most likely attributable is the acid aerosol, either particles that were acidic intrinsically (H₂SO₄) or particles that were not originally acidic [e.g., (NH₄)₂SO₄] but which adsorbed HNO₃ formed by the reaction of O₃ and NO₂ and became acidic.

Some hypotheses suggested by these results are that the augmentation in centriacinar injury is due to (a) an increase in penetration and deposition of O₃ in the parenchyma when acidic pollutants are also present, because of changes in breathing pattern; (b) O₃ sensitization of tissues which were subsequently (or simultaneously) affected by acidic pollutants; or (c) the formation of intermediate chemical products which were toxic.

Alarie (30) reported that inhalation of sensory irritants caused reflex-driven changes in breathing patterns. Animals that inhaled substances such as H₂SO₄ or SO₂ shifted to a pattern of slow, deep breathing. This type of pattern might result in enhanced deposition of gases such as SO₂ in the upper respiratory tract, but

could, in fact, allow greater numbers of small particles as well as gases of low water solubility such as O_3 to penetrate to the parenchyma more readily. The observations of damage to lung parenchyma in this study are consistent with these mechanisms.

With respect to formation of new toxic species in the oxidant/acid mixtures, we have observed significant levels of HNO_3 in oxidant mixtures containing O_3 and NO_2 and from chemical kinetics and stoichiometry we know that there must also be reactive intermediates such as nitrate radicals and nitrogen pentoxide present (the toxicology of these substances is not well understood). It is possible that more than one of the proposed mechanisms are active simultaneously. Current data are insufficient to identify or apportion the relative importance of each mechanism. The results discussed in this paper have only concerned short-term acute exposures. Data from current experiments of longer exposure duration are being evaluated and may shed additional light on the mechanisms by which acids and oxidants interact to cause damage to respiratory tract epithelium.

In summary, the effect of acid-oxidant mixtures on both early and late clearance appears to be dominated by the oxidant component, in short-term (4 hr) exposures at relatively high acid concentrations (3.5 mg/m^3). Acid- O_3 mixtures are significantly more potent than O_3 alone, with respect to causing inflammatory changes in centriacinar units of the lung parenchyma in exercising animals, but the effects in animals exposed at rest again appear to be mainly attributable to O_3 . O_3 - NO_2 mixtures with associated HNO_3 vapor were much more potent than O_3 alone in terms of lung toxicity. The addition of acidic or nonacidic sulfate particles to the exposure atmospheres containing O_3 and NO_2 did not result in augmentation of the oxidant-related effects, suggesting that effects of vapor phase acidity were not augmented by the presence of particle acidity. Cell killing in the nasal respiratory epithelium is caused by H_2SO_4 in exercising rats but neither O_3 nor O_3 plus NO_2 alone caused damage to nasal epithelia. Animals exposed at rest to mixtures of O_3 , NO_2 , and sulfate aerosols, under conditions favorable to the acidification of the particles by HNO_3 formed by reactions between the two oxidant gases, showed labeling in about 8% of the cells in the R1 nasal epithelium. Thus it appears that exposure atmospheres in which HNO_3 , and other reactive nitrogen compounds, are formed, i.e., those containing O_3 and NO_2 , are capable of causing health effects in excess of those attributable to either of the two oxidant gases alone.

An interesting interaction between O_3 and H_2SO_4 was seen only in exercising rats. Exercising rats exposed to relatively high concentrations of an atmospheric mixture containing H_2SO_4 aerosol (1 mg/m^3) and SO_2 (5 ppm) did not exhibit parenchymal lesions. Exercising rats exposed to 0.6 ppm O_3 typically did show parenchymal lesions. When, however, both O_3 and the O_3 mixed with H_2SO_4 were present, the degree of lesion formation was significantly elevated (2.5

times that observed in rats exposed to O_3 alone, at the same exercise level). Exercise was apparently an important factor in this process; rats exposed to an identical acid/ O_3 mixture while at rest did not exhibit parenchymal lesions.

The authors thank Mary Azizizn, Charles Bufalino, Debra Daniels, Richard Lejniaks, Tuan Nguyen, Robert Walters, and Rosario Zamora for their technical assistance. These studies were supported in part by programs funded by the Electric Power Research Institute, Southern California Edison Company, and the California Air Resources Board.

REFERENCES

1. Waller, R. E., and Lawther, P. J. Further observations on London fog. *Br. Med. J.* 2: 1473-1475 (1957).
2. Lawther, P. J., MacFarlane, A. J., Waller, R. E., and Brooks, A. G. E. Pulmonary function and sulphur dioxide: Some preliminary findings. *Environ. Res.* 10: 355-367 (1975).
3. Lippmann, M. Airborne acidity: Estimates of exposure and human health effects. *Environ. Health Perspec.* 63: 63-70 (1985).
4. Jacob, D. J., Waldman, J. M., Munger, J. W., and Hoffmann, M. R. Chemical composition of fogwater collected along the California coast. *Environ. Sci. Technol.* 19: 730-736 (1985).
5. Jacob, D. J., Munger, J. W., Waldman, J. M., and Hoffmann, M. R. The H_2SO_4 - HNO_3 - NH_3 system at high humidities and in fogs. 1. Spatial and temporal patterns in the San Joaquin Valley in California. *J. Geophys. Res.* 91: 1073-1088.
6. Luria, M., Sharf, G. The influence of light intensity, SO_2 , NO_x and C_3H_6 on sulfate aerosol formation. *J. Atmos. Chem.* 2: 321-329 (1985).
7. Russel, A. G., McRae, G. J., and Cass, G. R. The dynamics of nitric acid production and the fate of nitrogen oxides. *Atmos. Environ.* 19: 893-903 (1985).
8. Kleinman, M. T., Phalen, R. F., and Crocker, T. T. Generation and characterization of complex gas and particle mixtures for inhalation toxicologic studies. In: *Short-term Bioassays in the Analysis of Complex Environmental Mixtures IV.* (M. D. Waters, S. S. Sandhu, J. Lewtas, L. Claxton, G. Stauss, and S. Nesnow, Eds.), Plenum Publishing, New York, 1985, pp. 181-191.
9. Oppenheimer, M. The relationship of sulfur emissions to sulfate in precipitation. *Atmos. Environ.* 17: 451-460 (1983).
10. Forrest, J., and Newman, L. Further studies on the oxidation of sulfur dioxide in coal-fired power plant plumes. *Atmos. Environ.* 11: 465-474 (1977).
11. Forrest, J., Garber, R., and Newman, L. Formation of sulfate, ammonium and nitrate in the oil-fired power plant plume. *Atmos. Environ.* 13: 1287-1297 (1979).
12. Platt, U. F., Winer, A. M., Buermann, H. W., Atkinson, R., and Pitts, J. N., Jr. Measurement of nitrate radical concentrations in continental air. *Environ. Sci. Technol.* 18: 365-369 (1984).
13. Nazaroff, W. W., and Cass, G. R. Mathematical modeling of chemically reactive pollutants in indoor air. *Environ. Sci. Technol.* 20: 924-934 (1986).
14. Eatough, D. J., Arthur, R. J., Eatough, N. L., Hill, M. W., Mangelson, N. F., Richter, B. E., and Cooper, J. A. Rapid conversion of SO_2 (g) to sulfate in a fog bank. *Environ. Sci. Technol.* 18: 855-859 (1984).
15. Munger, J. W., Tiller, C., and Hoffmann, M. R. Identification of hydroxymethanesulfonate in fog water. *Science* 231: 247-249 (1986).
16. Lioy, P. J., Spektor, D., Thurston, G., Citak, K., Lippmann, M., Bock, N., Speizer, F. E., and Hayes, C. The design considerations for ozone and acid aerosol exposure and health investigations: The Fairview Lake Summer Camp—Photochemical Smog case study. *Environ. Int.* 13: 271-283 (1987).
17. Last, J. A., Hyde, D. M., and Chang, D. P. Y. A mechanism of synergistic lung damage by ozone and a respirable aerosol. *Exper. Lung Res.* 7: 223-235 (1984).
18. Last, J. A., Hyde, D. M., Guth, D. J., and Warren, D. L. Synergistic interaction of ozone and respirable aerosols on rat lungs. I. Importance of aerosol acidity. *Toxicology* 39: 247-257 (1986).

19. Schlesinger, R. B. Effects of intermittent inhalation exposures to mixed atmospheres of NO₂ and H₂SO₄ on rabbit alveolar macrophages. *J. Toxicol. Environ. Health* 22: 301-312 (1987).
20. Cavender, F. L., Steinhagen, W. H., Ulrich, C. E., Busey, W. M., Cockrell, B. Y., Haseman, J. K., Hogan, M.D., Drew, R. T. Effects in rats and guinea pigs of short-term exposures to sulfuric acid mist, ozone and their combination. *J. Toxicol. Environ. Health* 2: 521-533 (1977).
21. Silbaugh, S. A., and Mauderly, J. L. Effects of ozone and sulfuric acid on gas trapping in the guinea pig lung. *J. Toxicol. Environ. Health* 18: 133-141 (1986).
22. May, K. R. The Collision nebulizer: Description, performance and application. *J. Aerosol Sci.* 4: 235-243 (1973).
23. Morrow, P. E., and Mercer, T. T. A point-to-plane electrostatic precipitator for particle size sampling. *Am. Ind. Hyg. Assoc. J.* 25: 8-14 (1973).
24. Phalen, R. F., Kenoyer, J. L., Crocker, T. T., and McClure, T. R. Effects of sulfate aerosols in combination with ozone on elimination of tracer particles inhaled by rats. *J. Toxicol. Environ. Health* 6: 797-810 (1980).
25. Mautz, W. J., Phalen, R. F., McClure, T. R., and Bufalino, C. A rodent treadmill for inhalation toxicological studies and respirometry. *J. Appl. Physiol. Respir. Environ. Exercise Physiol.* 58: 673-679 (1985).
26. Hinrichs, R. J., Kenoyer, J. L., Phalen, R. F., and Crocker, T. T. Labeling of monodisperse polystyrene microspheres with tightly bound ⁵¹Cr. *Am. Ind. Hyg. Assoc. J.* 22: 570-575 (1978).
27. Mannix, R. C., Phalen, R. F., Kenoyer, J. L., and Crocker, T. T. Effect of sulfur dioxide-sulfate exposure on rat respiratory clearance. *Am. Ind. Hyg. Assoc. J.* 43: 679-685 (1982).
28. Shorter, R. G., Titus, J. L., and Divertie, M. B. Cell turnover in the respiratory tract. *Dis. Chest* 46: 138-142 (1938).
29. Evans, M. J., and Bils, R. F. Identification of cells labeled with tritiated thymidine in the pulmonary alveolar walls of the mouse. *Am. Rev. Respir. Dis.* 100: 372-378 (1969).
30. McClure, T. R., Diller, P. J., and Phalen, R. F. An airway perfusion apparatus for whole lung fixation. *Lab. Animal Sci.* 32: 195-196 (1982).
31. Young, J. T. Histopathologic examination of the rat nasal cavity. *Fundam. Appl. Toxicol.* 1: 309-312 (1981).
32. Elias, H., and Hyde, D. M. An elementary introduction to stereology. Quantitative microscopy. *Am. J. Anat.* 159: 411-446 (1980).
33. Alarie, Y. Sensory irritation by airborne chemicals. *CRC Crit. Rev. Toxicol.* 2: 299-363 (1973).