

c.2



Lawrence Berkeley Laboratory

UNIVERSITY OF CALIFORNIA

Materials & Chemical Sciences Division

RECEIVED
LAWRENCE
BERKELEY LABORATORY

APR 22 1987

LIBRARY
DOCUMENTS SECTION

Submitted to Chemical Physics Letters


PHOTOELECTRON ASYMMETRY PARAMETER FOR THE S 2p LEVEL OF SO₂: A ONE-DIMENSIONAL BARRIER CASE

T.A. Ferrett, P.A. Heimann, H.G. Kerkhoff,
U. Becker, D.W. Lindle, and D.A. Shirley

March 1987

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy
which may be borrowed for two weeks.*



LBL-22400
c.2

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

Photoelectron Asymmetry Parameter for the S 2p Level
of SO₂: a One-dimensional Barrier Case

T.A. Ferrett,* P.A. Heimann,** H.G. Kerckhoff,†
U. Becker,† D.W. Lindle,* and D.A. Shirley

Materials and Chemical Sciences Division
Lawrence Berkeley Laboratory
and
Department of Chemistry
University of California
Berkeley, California 94720

Abstract

We present a measurement of the asymmetry parameter (β) for S 2p core-level photoionization of SO₂ in the vicinity of several continuum resonances. Results are contrasted with $\beta(2p)$ for the central atoms of the molecules SF₆ and SiF₄. Atomic calculations for $\beta(S\ 2p)$ appear to reproduce qualitatively the molecular curve for SO₂ aside from a shift to lower β -values for the molecular case.

* Present address: National Bureau of Standards, Gaithersburg, MD 20899.

** Present address: Technische Universität München, Physik-Department E20, 8046 Garching b. München, Federal Republic of Germany.

† Permanent address: Technische Universität Berlin, Institut für Strahlungs- und Kernphysik, Sekr. PN 3-2, Hardenbergstr. 36, D-1000 Berlin 12, Federal Republic of Germany.

The series of core-level absorption spectra including H_2S , SO_2 , SiF_4 , and SF_6 have been used to illustrate the wide extent and variety of discrete and continuum resonances in molecular photoemission [1,2]. The SO_2 $L_{2,3}$ photoabsorption spectrum [1], shown in Fig. 1, displays a number of resonances. Features a through d have been ascribed to discrete molecular-orbital excitations and features f and g to Rydberg transitions. High resolution measurements have revealed even more Rydberg structure just below the S 2p edge (175 eV) [3].

In this work, we concentrate on the S 2p continuum, where oscillations over the broad resonances h and i have been related to analogous features in SF_6 (t_{2g} , e_g) and SiF_4 (e , t_2) caused by potential-barrier effects [1,2,4]. A centrifugal barrier on the perimeter of the molecule can be created by the sensitive interplay of attractive (electrostatic) and repulsive (centrifugal) forces. Quasibound states are formed just below the barrier and can be generally associated with excitations to unoccupied molecular orbitals. In the one-electron approximation, the decay of shape resonances leads to an enhancement only in the corresponding main-line continuum channel in the photoelectron spectrum (S 2p for a resonance above the $L_{2,3}$ edge).

These shape resonances are especially pronounced above the S 2p edge of SF_6 . The electronegative F atoms in this high symmetry molecule serve to enhance both the magnitude and the spatial extent of the barrier relative to other molecules. In addition, strong

multi-electron effects are exhibited in the decay of the e_g resonance in SF_6 to both the S 2p main line and a S 2p shake-up satellite [5]. In contrast, recent experiments [6,7] on Si 2p photoionization of SiF_4 showed that the fairly intense continuum features in this molecule do conform to the one-electron description [2,8] of quasibound continuum states, with observed decay only to the Si 2p main line.

Despite these differences in the resonant decay characteristics, both SF_6 [5] and SiF_4 [9] exhibit S 2p and Si 2p asymmetry parameters which are nearly identical as a function of kinetic energy. This similarity in $\beta(2p)$ is puzzling, because it seems to imply that the asymmetry parameter, normally sensitive to details of the potential field, is unaffected by the substantial difference between the SiF_4 and SF_6 potential barriers. To explore this question further, we have measured $\beta(S\ 2p)$ for SO_2 , where the analogous continuum features are much weaker. The SO_2 molecular potential lies between the atomic and the three-dimensional high-symmetry molecular case. In "one" dimension, along the S-O molecular axis, the barrier should more resemble that in SiF_4 and SF_6 . In perpendicular directions the potential is more atomic, and the photoelectron should escape more easily. The object of this study was to ascertain whether $\beta(2p)$ would reflect this intermediate character, or whether it would simply follow the atomic or molecular curves.

Photons emitted from the new 4° "grasshopper" monochromator at

the Stanford Synchrotron Radiation Laboratory (SSRL) ionized an effusive jet of SO_2 . The β -parameter was obtained from the ratio of intensities in two time-of-flight electron detectors oriented at 0° and 54.7° with respect to the photon polarization vector, and using Yang's theorem [10]. The experimental methods have been described in detail elsewhere [11-13]. Calibration of β as a function of kinetic energy was based on the known values of β for Xe(4d) [14,15], Ne(2s), and Ne(2p) [16] photoemission. Systematic errors contribute ± 0.10 to the error in β beyond the statistical errors shown in the figure. The photon-energy calibration was obtained to within ± 1.0 eV from the time-to-energy converted spectra using known binding and kinetic energies of the following photoemission and Auger peaks: Ne(2s), Ne(2p) [17], Xe(4d) [15], Xe 4d Auger [12,18], and SO_2 (S 2p) [3].

In Fig. 2 we show β (S 2p) for SO_2 along with a solid curve to represent the experimental results [5] for β (S 2p) of SF_6 . Although not shown in Fig. 2, β (Si 2p) for SiF_4 [9] is nearly identical to the curve for SF_6 as a function of kinetic energy. For SO_2 , β (S 2p) increases smoothly from threshold, in contrast to a sharp change in β (S 2p) for SF_6 near 15 eV kinetic energy associated with the e_g shape resonance. In comparison with β (S 2p) for SF_6 and SiF_4 , there is no pronounced minimum for SO_2 . From Fig. 1 we see that the SO_2 continuum feature i ($h\nu=196$ eV) is within the range of our measurement, but no apparent oscillation is present in β (S 2p).

We also compare to Hartree-Slater central field calculations [19] of β (2p) for atomic sulfur (dashed curve, Fig. 2). The atomic β

displays a minimum near threshold caused by the energy-dependent interaction of the ϵ_s and ϵ_d photoelectron waves. The $\beta(S\ 2p)$ for SO_2 shows the same qualitative trend as the atomic curve, aside from a shift to a lower absolute value of β . A previous comparison has been made between $\beta(Si\ 2p)$ for SiF_4 and atomic calculations for silicon with a similar result [9]. It was found that variation of the charge on Si to more positive values (relative to the neutral potential) tended to lower the β curves [9]. The atomic curve thus simulates the energy dependence of the molecular β reasonably well for SO_2 ; the lower value of β in molecules is partly due to effective charge differences on the central atom. There is a somewhat larger deviation from the atomic curve at lower kinetic energies; there is no minimum apparent in $\beta(S\ 2p)$ for SO_2 . The minimum might be very close to threshold, as in the S atom, and thus outside the range of our results.

In conclusion, SO_2 demonstrates resonant effects in the $L_{2,3}$ absorption spectrum (Fig. 1), but overall $\beta(S\ 2p)$ behaves much like its atomic sulfur counterpart. We note that photoemission experiments on the valence ionization of SO_2 found no evidence for potential-barrier effects [20]. Another example to study is H_2S , for which there are no detectable continuum resonances above the S 2p threshold.¹ We believe that $\beta(S\ 2p)$ for H_2S should conform even more closely to atomic predictions.

Acknowledgements

We gratefully acknowledge S.T. Manson for performing the atomic calculation for the sulfur 2p asymmetry parameter. This work was supported by the Director, Office of Energy Research, Office of Basic Energy Sciences, Chemical Sciences Division of the U.S. Department of Energy under Contract No. DE-AC03-76SF00098. It was performed at the Stanford Synchrotron Radiation Laboratory, which is supported by the Department of Energy's Office of Basic Energy Sciences.

References


1. T.M. Zimkina and A.S. Vinogradov, J. Phys. (Paris) 32, C4-3 (1971).
2. J.L. Dehmer, J. Chem. Phys. 56, 4496 (1972).
3. N.A. Shklyueva, L.N. Mazalov, and V.V. Murakhtanov, Zh. Strukt. Khim. 20, 733 (1979).
4. A.V. Kondratenko, L.N. Mazalov, and K.M. Neiman, Opt. Spec. 49, 266 (1980).
5. T.A. Ferrett, D.W. Lindle, P.A. Heimann, M.N. Piancastelli, P.H. Kobrin, H.G. Kerkhoff, U. Becker, W.D. Brewer, and D.A. Shirley (unpublished results).
6. G.M. Bancroft, S. Aksela, H. Aksela, K.H. Tan, B.W. Yates, L.L. Coatsworth, and J.S. Tse, J. Chem. Phys. 84, 5 (1986).
7. T.A. Ferrett, M.N. Piancastelli, D.W. Lindle, P.A. Heimann, and D.A. Shirley (unpublished results).
8. J.L. Dehmer in Resonances in Electron-Molecule Scattering, van der Waals Complexes, and Reactive Chemical Dynamics, edited by D.G. Truhlar (American Chemical Society, Washington, D.C., 1984); J.L. Dehmer, D. Dill, and A.C. Parr in Photophysics and Photochemistry in the Vacuum Ultraviolet, edited by S.P. McGlynn, G. Findley, and R. Huebner (Reidel, Dordrecht, Holland, 1985); J.L. Dehmer, A.C. Parr, and S.H. Southworth in Handbook on Synchrotron Radiation, Vol. II, edited by G.V. Marr (North Holland, Amsterdam, 1986).

9. P.R. Keller, J.W. Taylor, F.A. Grimm, P. Senn, T.A. Carlson, and M.O. Krause, Chem. Phys. 74, 247 (1983).
10. C.N. Yang, Phys. Rev. 74, 764 (1948).
11. M.G. White, R.A. Rosenberg, G. Gabor, E.D. Poliakoff, G. Thornton, S. Southworth, and D.A. Shirley, Rev. Sci. Instrum. 50, 1288 (1979).
12. S. Southworth, C.M. Truesdale, P.H. Kobrin, D.W. Lindle, W.D. Brewer, and D.A. Shirley, J. Chem. Phys. 76, 143 (1982).
13. S. Southworth, U. Becker, C.M. Truesdale, P.H. Kobrin, D.W. Lindle, S. Owaki, and D.A. Shirley, Phys. Rev. A 28, 261 (1983).
14. S.H. Southworth, Ph.D. Thesis, University of California, Berkeley (1981).
15. S.H. Southworth, P.H. Kobrin, C.M. Truesdale, D. Lindle, S. Owaki, and D.A. Shirley, Phys. Rev. A 24, 2257 (1981).
16. F. Wuilleumier and M.O. Krause, J. Electron Spectrosc. 15, 15 (1979).
17. W. Perssen, Phys. Scr. 3, 133 (1971); C.E. Moore, Atomic Energy Levels, NBS Circular 467 (U.S. GPO, Washington, D.C., 1949), Vol. 1.
18. L.O. Werme, T. Bergmark, and K. Siegbahn, Phys. Scr. 16, 141 (1972).
19. S.T. Manson (private communication).
20. D.M.P. Holland, A.C. Parr, and J.L. Dehmer, J. Electron Spectrosc. 32, 237 (1983).

Figure Captions


9
u
Fig. 1 The photoabsorption coefficient μ (arbitrary scale) for SO_2 in the vicinity of the $L_{2,3}$ edge. Notation for features a through i is taken from Ref. 1.

Fig. 2 The S 2p asymmetry parameter (β) for SO_2 as a function of photon energy (bottom scale) and kinetic energy (top scale). The solid curve represents experimental data for $\beta(\text{S } 2p)$ of SF_6 [5], plotted as a function of kinetic energy. Hartree-Slater theory curve [19] for the atomic $\beta(2p)$ of sulfur is also shown (dashed), as a function of kinetic energy.



XBL 8611-4291

Figure 1



XBL 8611-4572

Figure 2

This report was done with support from the Department of Energy. Any conclusions or opinions expressed in this report represent solely those of the author(s) and not necessarily those of The Regents of the University of California, the Lawrence Berkeley Laboratory or the Department of Energy.

Reference to a company or product name does not imply approval or recommendation of the product by the University of California or the U.S. Department of Energy to the exclusion of others that may be suitable.

*LAWRENCE BERKELEY LABORATORY
TECHNICAL INFORMATION DEPARTMENT
UNIVERSITY OF CALIFORNIA
BERKELEY, CALIFORNIA 94720*