

UNIVERSITY OF  
CALIFORNIA

*Radiation  
Laboratory*

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy  
which may be borrowed for two weeks.  
For a personal retention copy, call  
Tech. Info. Division, Ext. 5545*

BERKELEY, CALIFORNIA

## **DISCLAIMER**

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

Copy 2

UCRL-1033  
Unclassified-Chemistry Distribution

UNIVERSITY OF CALIFORNIA

Radiation Laboratory

Contract No. W-7405-eng-48

**UNCLASSIFIED**

CARRIER-FREE RADIOISOTOPES FROM CYCLOTRON TARGETS

XV. PREPARATION AND ISOLATION OF  $Mn^{32}$  FROM CHROMIUM\*

Herman R. Haymond, Warren M. Garrison and Joseph G. Hamilton

December 4, 1950

Berkeley, California

CARRIER-FREE RADIOISOTOPES FROM CYCLOTRON TARGETS  
XV. PREPARATION AND ISOLATION OF  $Mn^{52}$  FROM CHROMIUM\*

Herman R. Haymond, Warren M. Garrison and Joseph G. Hamilton

Crocker Laboratory, Radiation Laboratory, and Divisions of Medical Physics, Experimental Medicine, and Radiology; University of California, Berkeley and San Francisco, California.

December 4, 1950

Deuteron bombardment of chromium produces<sup>1</sup> three radioisotopes of manganese (46-min  $Mn^{51}$ , 6.5-day and 21-min  $Mn^{52}$ , and 310-day  $Mn^{54}$ ) by (d,n) and (d,2n) reaction. This paper describes a radiochemical procedure used in isolating carrier-free radio-manganese (6.5-day  $Mn^{52}$ ) from a chromium target which had been bombarded with 19 Mev deuterons in the 60-inch cyclotron at Crocker Laboratory. The  $Mn^{51}$  and  $Mn^{52m}$  had decayed out prior to chemical separation. The 310-day period of  $Mn^{54}$  was not observed, presumably due to the low abundance of  $Cr^{53,54}$ . Other possible concurrent reactions include the formation of radio-vanadium by (d,c) reaction and radio-titanium by (d,αp) reaction. A quantitative separation from vanadium is obtained with the procedure described here. Radio-titanium,  $Ti^{51}$ , was not detected.

Spectrographically-pure<sup>2</sup> chromium powder, supported on a water-cooled aluminum target plate by 0.25 mil platinum foil, was bombarded for 20 μa-hr. at a beam intensity of 10 μa. After bombardment, the chromium powder (approx. 1 gm.) was dissolved in a minimum volume of 12N hydrochloric acid and the resultant solution was evaporated to incipient dryness. Twenty milligrams of ferric chloride were added and the chromic chloride containing the

---

\*This document is based on work performed under Contract No. W-7405-eng-48-A for the Atomic Energy Commission.

<sup>1</sup> G. T. Saaborg and I. Perlman, Rev. Mod. Phys., 20, 585 (1948)

<sup>2</sup> Manganese was not detected by spectrographic analysis.

carrier-free radio-manganese was re-dissolved in 50 ml of 1N hydrochloric acid. This solution was then added with stirring to 200 ml of 2.5N sodium hydroxide solution previously saturated with bromine. Under these conditions, the carrier-free radio-manganese, presumably in the tetrapositive state, is quantitatively carried on the ferric hydroxide precipitate<sup>3</sup>. Chromium and vanadium are retained in the supernatant. Two additional reprecipitations were made to insure complete separation. The ferric hydroxide was washed, dissolved in 6N hydrochloric acid and iron was extracted with ethyl ether. The solution of carrier-free radio-manganese in 6N hydrochloric acid was evaporated to dryness after the addition of 5 ml of isotonic saline solution. The activity dissolved quantitatively with the addition of 5 ml of water.

The decay curve was followed for eight half-lives and showed a single period of 6.2 days which agrees closely with the reported<sup>4</sup> value for Mn<sup>52</sup>. Absorption measurements in aluminum showed the assigned<sup>5</sup> 0.6 Mev positron. The activity was further identified by chemical separations using manganese, chromium, vanadium and titanium carriers. Over 99 percent of a sample of the activity was recovered in the manganese fraction.

We wish to thank Mr. T. Putnam, Mr. G. B. Rossi and the crew of the 60-inch cyclotron for bombardments and Professor G. T. Seaborg for his continued interest in this work.

---

<sup>3</sup> Maxwell, Gile, Garrison and Hamilton, J. Chem. Phys. 17 1340 (1949)

<sup>4</sup> J. J. Livingood and G. T. Seaborg, Phys. Rev. 54 391 (1938)

<sup>5</sup> W. C. Peacock and M. Deutsch, Phys. Rev. 69 306 (1946)