

Lawrence Berkeley National Laboratory

LBL Publications

Title

CO 2 conversion to phenyl isocyanates by uranium(vi) bis(imido) complexes

Permalink

<https://escholarship.org/uc/item/7m45d5wm>

Journal

Chemical Communications, 56(3)

ISSN

1359-7345

Authors

Maria, Leonor
Bandeira, Nuno AG
Marçalo, Joaquim
et al.

Publication Date

2020-01-02

DOI

10.1039/c9cc07411b

Peer reviewed

CO₂ conversion to phenyl isocyanates by uranium(vi) bis(imido) complexes†‡

Leonor Maria,^{*ab} Nuno A. G. Bandeira,^{id*cd} Joaquim Marçalo, Isabel C. Santos^a and John K. Gibson^d^{idab}

ABSTRACT:

Uranium(vi) *trans*-bis(imido) complexes [U(κ^4 -{(tBu₂ArO)₂Me₂-cyclam})-(NPh)(NPh^R)] react with CO₂ to eliminate phenyl isocyanates and afford uranium(vi) *trans*-[O=U=NR]²⁺ complexes, including [U(κ^4 -{(tBu₂ArO)₂Me₂-cyclam})-(NPh)(O)] that was crystallographically characterized. DFT studies indicate that the reaction proceeds by endergonic formation of a cycloaddition intermediate; the secondary reaction to form a dioxo uranyl complex is both thermodynamically and kinetically hindered.

Uranium complexes have emerged as attractive candidates for activation of the very stable C=O bonds of carbon dioxide,^{1–3} which must occur during CO₂ functionalization. U(III) with appropriate supporting ligands can effectively reduce carbon dioxide, giving rise to various products that include an end-on-bound CO₂^{•–} U(IV) complex,⁴ uranium oxo species,^{5,6} carbon monoxide,^{5,6} carbonates,^{7,8} oxalates,⁹ and isocyanate.¹⁰ It was also demonstrated that U(IV) complexes with suitable ancillary ligands and functionalities can facilitate CO₂ insertion into U–E bonds (E = C, N, O, S).^{11–14} Meyer and co-workers showed that U(V) terminal oxo complexes [(^RArO)₃tacn}U=O] are formed upon treatment of U(V) mono-imidos with CO₂, probably *via* [2+2] cycloaddition with elimination of isocyanates.^{13,15} Similar CO₂ transformations have been observed with high oxidation state early transition metals, such as titanium imido complexes.^{16–19}

Activation of CO₂ by hexavalent uranium imido complexes has not yet been achieved, and in general CO₂ activation by U(VI) complexes is rare.²⁰ In such a rare case, the terminal nitride [U(Tren^{TIPS})(N)] reacted with CO₂ to yield a U(VI) oxo-cyanate complex by bond metathesis and cleavage of a C=O bond, with rapid decomposition of the product to [U^V(Tren^{TIPS})(O)].²⁰

Several structurally characterized uranium(VI) imido complexes have recently been reported,^{21–23} but reactivity studies of the U=NR bonds remain scarce,²⁴ particularly for U(VI) *trans*-bis(imido) complexes. Such *trans*-[RN=U=NR]²⁺ systems notably are isoelectronic with thermodynamically stable uranyl, {O=U=O}²⁺, but exhibit greater bond covalency.²⁵ It has been proposed that the inverse *trans*-influence (ITI)²⁶ of the two uranium–nitrogen multiple bonds stabilizes and reduces reactivity of the {RN=U=NR}²⁺ moiety. Accordingly, Boncella and co-workers showed that treating [U(=N^tBu)₂I₂(OPPh₃)₂] with PhN=C=O does not result in a uranium oxo complex but rather an imido transfer takes place to form [U(=N^tBu)(=NPh)I₂(OPPh₃)₂].²⁷ We here demonstrate that U(VI) *trans*-bis(imido) complexes supported by a bis(phenolate) cyclam ligand react with CO₂ to produce stable *trans*-[O=U=NR]²⁺ with elimination of aryl isocyanate.

Based on previous results for azobenzene activation with U(III) complex [U(κ^6 -{(tBu₂ArO)₂Me₂-cyclam})]I (1), with formation of [U(κ^4 -{(tBu₂ArO)₂Me₂-cyclam})-(NPh)₂] (2),²⁸ the initial focus studies was on the synthesis of the new U(VI) bis(imido) [U(κ^4 -{(tBu₂ArO)₂Me₂-cyclam})-(NPh)(NTol)] (3), with a goal of assessing reactivity with CO₂. The reaction of 4-methylazobenzene (TolNNPh) with 2 equiv. of U(III) complex 1 in toluene at room temperature resulted in four-electron cleavage of the N=N double bond, affording the soluble bis(imido) [U(κ^4 -{(tBu₂ArO)₂Me₂-cyclam})-(NPh)(NTol)] (3) and the insoluble U(IV) compound [U(κ^6 -{(tBu₂ArO)₂Me₂-cyclam})]I₂, in a 1 : 1 ratio (Scheme 1).

Quality single crystals for X-ray diffraction were obtained by slow evaporation of a *n*-hexane/thf solution for two weeks. Refinement of the XRD data confirmed the expected U(VI) complex, with the phenyl imido and *p*-methyl phenyl imido ligands in a *trans* arrangement (N5–U1–N6 = 173.7(4)°) with an

^a Centro de Ciências e Tecnologias Nucleares (C²TN), Instituto Superior Técnico, Universidade de Lisboa, 2695-066 Bobadela, Portugal.
E-mail: leonorm@ctn.tecnico.ulisboa.pt


^b Centro de Química Estrutural (CQE), Instituto Superior Técnico, Universidade de Lisboa, 2695-066 Bobadela, Portugal

^c BioISI and Centro de Química e Bioquímica, Faculdade de Ciências, Universidade de Lisboa, 1749-016 Lisboa, Portugal. E-mail: nuno.bandeira@ciencias.ulisboa.pt

^d Chemical Sciences Division, Lawrence Berkeley National Laboratory, Berkeley, CA 94720, USA

† A data set collection of computational results is available in the ioChem-BD repository³⁸ and can be accessed *via* <https://doi.org/10.19061/iochem-bd-6-21>.

‡ Electronic supplementary information (ESI) available: Computational and experimental details, NMR spectra, and detailed X-ray data. CCDC 1954718 and 1954719. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/c9cc07411b



Scheme 1 Synthesis of uranium(vi) bis(imido) complexes **2** and **3**.


overall distorted octahedral geometry (Fig. S2 in ESI[†]). Structural parameters of **3**, including short U–N_{imido} distances of 1.909(6) and 1.911(7) Å, are similar to those in the symmetric bis(imido) complex **2** (1.895(2) and 1.907(2) Å)²⁸ (structural parameters are in ESI[†] Table S2).

The ¹H NMR spectrum of **3** is characteristic of a diamagnetic compound, consistent with the asymmetric structure identified in the solid state. ¹H and ¹³C NMR (see Fig. S3 and S4 in ESI[†]) revealed two sets of resonances for the phenolate arms of $\{({}^t\text{Bu}_2\text{ArO})_2\text{Me}_2\text{-cyclam}\}^{2-}$ and two distinct sets of resonances for the phenyl imido and *p*-methyl phenyl imido ligands.

Frozen brown solutions of **2** and **3** in benzene-*d*₆ were exposed to an excess of CO₂ (*ca.* 30 equiv.). Upon warming to room temperature, the colour of the solutions changed slowly from brown to dark cherry red. Following the reaction by ¹H NMR revealed that bis(imido) U(vi) complex **2** was completely consumed within 7 hours, with ingrowth of new *ortho*, *meta*, and *para* phenyl proton resonances at 5.19, 6.90 and 5.75 ppm, and aromatic resonances between 6.86 and 6.58 ppm, consistent with formation of a new uranium phenyl imido complex and elimination of phenyl isocyanate (Fig. S6 and S7 in ESI[†]).

The ¹H NMR spectrum of the new complex exhibited integrated intensities for the $\{({}^t\text{Bu}_2\text{ArO})_2\text{Me}_2\text{-cyclam}\}^{2-}$ and phenyl imido protons of 1 : 1, and revealed loss of the C₂ symmetry of the bis(phenolate) cyclam ligand. The benzylic protons gave rise to two AB systems with *J*_{AB} coupling of 12.1 and 12.3 Hz, and the four ^tBu phenolate substituents appeared as 4 singlets, indicating that the two phenolate arms of $\{({}^t\text{Bu}_2\text{ArO})_2\text{Me}_2\text{-cyclam}\}^{2-}$ have different chemical environments. In addition, two resonances for the NCH₃ protons were observed at 2.01 and 1.99 ppm and the phenyl imido protons showed only one set of resonances, which indicates free rotation about the N–C_{ipso} bond. The ¹³C NMR data (Fig. S8–S10 in ESI[†]) corroborated the ¹H NMR results, consistent with formation of U(vi) oxo imido complex $[\text{U}(\kappa^4\text{-}\{({}^t\text{Bu}_2\text{ArO})_2\text{Me}_2\text{-cyclam}\})\text{(NPh)}\text{(O)}]$ (**4**).

An NMR tube-scale reaction of $[\text{U}(\kappa^4\text{-}\{({}^t\text{Bu}_2\text{ArO})_2\text{Me}_2\text{-cyclam}\})\text{(NPh)}\text{(NTol)}]$ (**3**) with CO₂ showed that the bis(imido) complex is quantitatively converted to a mixture of mono-oxo imido $[\text{U}(\kappa^4\text{-}\{({}^t\text{Bu}_2\text{ArO})_2\text{Me}_2\text{-cyclam}\})\text{(NPh)}\text{(O)}]$ (**4**) and $[\text{U}(\kappa^4\text{-}\{({}^t\text{Bu}_2\text{ArO})_2\text{Me}_2\text{-cyclam}\})\text{(NTol)}\text{(O)}]$ (**5**), in approximately 1 : 1 ratio, along with the corresponding aryl isocyanates (Fig. S13 and S14 in ESI[†]). Although most ¹H NMR resonances of the two uranium compounds overlap, it was possible to differentiate resonances that match the phenyl imido ligand of **4** from those of the methyl phenyl imido ligand of **5** (see Fig. S14 in ESI[†]). Two of the ^tBu phenolate resonances of **5** are at 2.08 and 1.86 ppm, with the other



Scheme 2 Synthesis of uranium(vi) oxo-imido complex **4**.

two centered at 1.57 ppm and overlapping with a ^tBu resonance of **4**. ¹³C NMR spectroscopy confirmed formation of the two U(vi) mono-oxo imido complexes, with the appearance of two pairs of carbon resonances for the C–O phenolates of $\{({}^t\text{Bu}_2\text{ArO})_2\text{Me}_2\text{-cyclam}\}^{2-}$ (**4**: 166.75 and 166.56 ppm; **5**: 166.81 and 166.69 ppm).

The reaction of $[\text{U}(\kappa^4\text{-}\{({}^t\text{Bu}_2\text{ArO})_2\text{Me}_2\text{-cyclam}\})\text{(NPh)}_2]$ (**2**) with an excess of CO₂ in toluene resulted in a dark cherry red solid after evaporation of the solvent (Scheme 2). ¹H NMR of the crude solid was consistent with formation of **4** and a minor side product, perhaps from dimerization of phenyl isocyanate.²⁹ Recrystallization of the crude solid from hexane provided single crystals; X-ray diffraction confirmed the molecular structure as oxo imido complex $[\text{U}(\kappa^4\text{-}\{({}^t\text{Bu}_2\text{ArO})_2\text{Me}_2\text{-cyclam}\})\text{(O)}\text{(NPh)}]$ (**4**) (Fig. 1).

Like complexes **2** and **3**, **4** exhibits an octahedral geometry about the uranium center, with the $\{({}^t\text{Bu}_2\text{ArO})_2\text{Me}_2\text{-cyclam}\}^{2-}$ ligand adopting a $\kappa^4\text{-N}_2\text{O}_2$ coordination mode, and the oxo and phenyl imido are *trans*-oriented. The geometric parameters of the $\{\text{RN}=\text{U}=\text{O}\}^{2+}$ core (U1–O3 1.787(3) Å, U1–N5 1.879(3) Å, O3–U1–N5, 176.4(1)°) are comparable to those of previously reported $[\text{U}(\text{O}=\text{NPh}^{\text{IPr}_2})(\text{O})\text{Cl}_2(\text{tppo})_2]$ (U–O 1.778(2), U–N 1.847(3)),³⁰ while the U–O(aryloxide) and U–N(cyclam) bond distances are similar to those of **2** and **3**.

Crystallographically characterized *trans*- $[\text{O}=\text{U}=\text{NR}]^{2+}$ compounds are scarce, mostly prepared by oxygen atom transfer to U(IV) monoimido complexes or reductive cleavage of nitrite by a U(V) imido complex.^{30–32} Treatment of U(vi) *trans*-bis(imido) $[\text{U}(\text{O}=\text{N}^t\text{Bu})_2\text{I}_2(\text{THF})_2]$ with water reagent B(C₆F₅)₃·H₂O also led to formation of a uranyl-like complex, $[\text{U}(\text{O})(\text{O}=\text{N}^t\text{Bu})_2(\text{THF})_2]$.³³

We have shown here the first activation of CO₂ by a U(vi) imido complex, specifically a *trans*-bis(imido), with formation




Fig. 1 Molecular structure of $[\text{U}(\kappa^4\text{-}\{({}^t\text{Bu}_2\text{ArO})_2\text{Me}_2\text{-cyclam}\})\text{(NPh)}\text{(O)}]$ (**4**) with 50% probability thermal ellipsoids.

of a terminal uranium oxo bond. Selective formation of *trans*-[O=U^{VI}=NR]²⁺ complexes **4** and **5** by multiple bond metathesis is presumably driven by a thermodynamic preference for U=O bond formation with release of aryl isocyanate. Complex [U(κ^4 -{(tBu₂ArO)₂Me₂-cyclam})(NPh)(O)] (**4**) did not exhibit reactivity upon CO₂ exposure in benzene-*d*₆, possibly reflecting enhanced stability of complex **4** due to a greater inverse *trans* influence upon U=O bond formation.²⁶

To elucidate mechanistic aspects of the observed reactivity, the ADF program³⁴ was used with the rev-PBE-D3 density functional^{35–37} to compute the CO₂ cycloaddition/isocyanate extrusion reaction pathway (see ESI† for Computational details). The starting reference state A (energy ≡ 0) is the two reactants infinitely separated (Fig. 2). van der Waals interaction between the reactants yields association complex B, which is followed by transition state, TS_B at 103.3 kJ mol⁻¹ above A. The CO₂ insertion into a uranium imido site yields intermediate C at 59.5 kJ mol⁻¹ above A. Cycloaddition *via* TS_B is a donor-acceptor process, as revealed by the frontier orbital in Fig. 3. A second transition state (TS_C) corresponds to cleavage of isocyanate intermediate C and leads to formation of the uranium imido mono-oxo complex **4** and phenyl isocyanate (D). TS_C presents




Fig. 2 Free energy profile for the reaction of [U(κ^4 -{(tBu₂ArO)₂Me₂-cyclam})(NPh)₂] (**2**) with CO₂.




Fig. 3 A clipped detail of the HOMO–5 of TS_B showing the U(5f) → O and N → C donor–acceptor interactions with the incoming CO₂ molecule establishing new bonds.




Fig. 4 Free energy profile for the reaction of [U(κ^4 -{(tBu₂ArO)₂Me₂-cyclam})(NPh)(O)] (**4**) with CO₂.

a substantial barrier of 116 kJ mol⁻¹ above A and 140 kJ mol⁻¹ above B, consistent with the observed slow kinetics at 25 °C. The free energy of 26.2 kJ mol⁻¹ to dissociate complex **4** (steps D–E) is mostly due to the electronic contribution ($\Delta E = +31.2$ kJ mol⁻¹). The interaction in D is presumably dominated by dispersion bonding between the uranium complex and phenyl isocyanate.

Formation of the oxo-imido complex **4** in D is a thermodynamic minimum, which evidently hampers subsequent addition of a second CO₂. The reaction profile for addition of CO₂ to **4** is shown in Fig. 4. Two aspects clearly differ from the profile in Fig. 2: formation of metallacycle intermediate H is more endergonic than for C ($\Delta\Delta G^\circ = +24.9$ kJ mol⁻¹); and both transition states TS_G and TS_H are higher energy compared with TS_B and TS_C. In particular, rate determining transition state TS_H is 8.7 kJ mol⁻¹ higher energy than TS_C. Notably, the exergonicity of step I (–53.4 kJ mol⁻¹) is not as favourable as for step D in Fig. 2 (–81.5 kJ mol⁻¹). Overall, it is apparent that CO₂ addition to complex **4** is both thermodynamically and kinetically hindered *vis-à-vis* CO₂ addition to bis(imido) complex **2**, consistent with observation of only the first process.

¹H NMR spectra were collected during reaction of **2** with CO₂ under pseudo-first-order conditions of a large CO₂ excess at constant concentration. A plot of the ln(molar fraction) of reagent as a function of time yields a pseudo-first-order rate. The rate was obtained for different temperatures such that using the Arrhenius equation and plotting ln *k*_{obs} versus 1/*T* provided a reaction activation energy *E*_a = 102 ± 12 kJ mol⁻¹ (see ESI† for details). This *E*_a is somewhat smaller than the computed value of 140 kJ mol⁻¹ in Fig. 2, which presumably reflects limitations of the computational and/or experimental methodologies (*e.g.* sample homogeneity during NMR acquisition). The effect of the higher pressure used in the experiments should be insignificant.

In conclusion, U(*vi*) *trans*-bis(imido) complexes supported by a bis(phenolate) cyclam ligand react with excess CO₂ to afford the *trans*-[O=U=NR]²⁺ species [U(κ^4 -{(tBu₂ArO)₂Me₂-cyclam})(NAr)(O)]. DFT computational studies suggest that the reaction proceeds *via* endergonic formation of a [2+2] cycloaddition intermediate, with subsequent extrusion of phenyl isocyanate and formation of the U(*vi*) oxo-imido computed to be exergonic.

These reactions are unprecedented examples of activation and cleavage of CO₂ mediated by uranium(vi) imido complexes.

We acknowledge the financial support from the Fundação para a Ciência e a Tecnologia through projects UID/MULTI/00612/2019, UID/MULTI/04046/2019 and UID/MULTI/04349/2019, through the *post doc* fellowships SFRH/BPD/101840/2014 and SFRH/BPD/110419/2015 to LM and NAGB, respectively, and through RNEM – Portuguese Mass Spectrometry Network, ref. LISBOA-01-0145-FEDER-022125, also supported by the Lisboa Regional Operational Programme (Lisboa2020), under the PT2020 Partnership Agreement, through the European Regional Development Fund (ERDF). JKG was supported by U.S. Department of Energy, Basic Energy Sciences, Chemical Sciences, Geosciences, and Biosciences, Heavy Element Chemistry, at LBNL under Contract DE-AC02-05CH1123. The authors thank Dr M. C. Oliveira for the HR-MS analysis.

Conflicts of interest

There are no conflicts to declare.

References

- 1 B. M. Gardner and S. T. Liddle, *Eur. J. Inorg. Chem.*, 2013, 3753–3770.
- 2 H. S. La Pierre and K. Meyer, *Prog. Inorg. Chem.*, 2014, **58**, 303–415.
- 3 P. L. Arnold and Z. R. Turner, *Nat. Rev. Chem.*, 2017, **1**, 2–16.
- 4 I. Castro-Rodriguez, H. Nakai, L. N. Zakharov, A. L. Rheingold and K. Meyer, *Science*, 2004, **305**, 1757–1759.
- 5 I. Castro-Rodriguez and K. Meyer, *J. Am. Chem. Soc.*, 2005, **127**, 11242–11243.
- 6 O. Cooper, C. Camp, J. Pécaut, C. E. Kefalidis, L. Maron, S. Gambarelli and M. Mazzanti, *J. Am. Chem. Soc.*, 2014, **136**, 6716–6723.
- 7 O. T. Summerscales, A. S. P. Frey, F. G. N. Cloke and P. B. Hitchcock, *Chem. Commun.*, 2008, 198–200.
- 8 A.-C. Schmidt, A. V. Nizovtsev, A. Scheurer, F. W. Heinemann and K. Meyer, *Chem. Commun.*, 2012, **48**, 8634–8636.
- 9 N. Tsoureas, L. Castro, A. F. R. Kilpatrick, F. G. N. Cloke and L. Maron, *Chem. Sci.*, 2014, **5**, 3777–3788.
- 10 C. Camp, L. Chatelain, C. E. Kefalidis, J. Pécaut, L. Maron and M. Mazzanti, *Chem. Commun.*, 2015, **51**, 15454–15457.
- 11 K. G. Moloy and T. J. Marks, *Inorg. Chim. Acta*, 1985, **110**, 127–131.
- 12 W. J. Evans, J. R. Walensky and J. W. Ziller, *Organometallics*, 2010, **29**, 945–950.
- 13 S. C. Bart, C. Anthon, F. W. Heinemann, E. Bill, N. M. Edelstein and K. Meyer, *J. Am. Chem. Soc.*, 2008, **130**, 12536–12546.
- 14 C. Lescop, T. Arliguie, M. Lance, M. Nierlich and M. Ephritikhine, *J. Organomet. Chem.*, 1999, **580**, 137–144.
- 15 A.-C. Schmidt, F. W. Heinemann, L. Maron and K. Meyer, *Inorg. Chem.*, 2014, **53**, 13142–13153.
- 16 C. L. Boyd, E. Clot, A. E. Guiducci and P. Mountford, *Organometallics*, 2005, **24**, 2347–2367.
- 17 U. J. Kilgore, F. Basuli, J. C. Huffman and D. J. Mindiola, *Inorg. Chem.*, 2006, **45**, 487–489.
- 18 A. E. Guiducci, C. L. Boyd, E. Clot and P. Mountford, *Dalton Trans.*, 2009, 5960–5979.
- 19 J. C. Anderson and R. B. Moreno, *Org. Biomol. Chem.*, 2012, **10**, 1334–1338.
- 20 P. A. Cleaves, C. E. Kefalidis, B. M. Gardner, F. Tuna, E. J. L. McInnes, W. Lewis, L. Maron and S. T. Liddle, *Chem. – Eur. J.*, 2017, **23**, 2950–2959.
- 21 T. W. Hayton, *Dalton Trans.*, 2010, **39**, 1145–1158.
- 22 T. W. Hayton, *Chem. Commun.*, 2013, **49**, 2956–2973.
- 23 M. A. Boreen and J. Arnold, in *The Heaviest Metals: Science and Technology of the Actinides and Beyond*, ed. W. J. Evans and T. P. Hanusa, Wiley, 2019, pp. 105–122.
- 24 C. J. Tatebe, K. E. Gettys and S. C. Bart, in *Handbook on the Physics and Chemistry of Rare Earths*, ed. J.-C. G. Bünzli and V. K. Pecharsky, Elsevier, 2018, vol. 54, pp. 1–42.
- 25 T. W. Hayton, J. M. Boncella, B. L. Scott, P. D. Palmer, E. R. Batista and P. J. Hay, *Science*, 2005, **310**, 1941–1943.
- 26 R. G. Denning, *J. Phys. Chem. A*, 2007, **111**, 4125–4143.
- 27 L. P. Spencer, P. Yang, B. L. Scott, E. R. Batista and J. M. Boncella, *J. Am. Chem. Soc.*, 2008, **130**, 2930–2931.
- 28 L. Maria, I. C. Santos, V. R. Sousa and J. Marçalo, *Inorg. Chem.*, 2015, **54**, 9115–9126.
- 29 A. R. Katritzky, T.-B. Huang and M. V. Voronkov, *J. Org. Chem.*, 2001, **66**, 1043–1045.
- 30 R. E. Jilek, N. C. Tomson, R. L. Shook, B. L. Scott and J. M. Boncella, *Inorg. Chem.*, 2014, **53**, 9818–9826.
- 31 E. Lu, O. J. Cooper, J. McMaster, F. Tuna, E. J. L. McInnes, W. Lewis, A. J. Blake and S. T. Liddle, *Angew. Chem., Int. Ed.*, 2014, **53**, 6696–6700.
- 32 K. C. Mullane, A. J. Lewis, H. Yin, P. J. Carroll and E. J. Schelter, *Inorg. Chem.*, 2014, **53**, 9129–9139.
- 33 T. W. Hayton, J. M. Boncella, B. L. Scott and E. R. Batista, *J. Am. Chem. Soc.*, 2006, **128**, 12622–12623.
- 34 G. te Velde, F. M. Bickelhaupt, E. J. Baerends, C. Fonseca Guerra, S. J. A. van Gisbergen, J. G. Snijders and T. Ziegler, *J. Comput. Chem.*, 2001, **22**, 931–967.
- 35 Y. Zhang and W. Yang, *Phys. Rev. Lett.*, 1998, **80**, 890.
- 36 J. P. Perdew, K. Burke and M. Ernzerhof, *Phys. Rev. Lett.*, 1996, **77**, 3865–3868.
- 37 L. Goerigk, H. Kruse and S. Grimme, *ChemPhysChem*, 2011, **12**, 3421–3433.
- 38 M. Álvarez-Moreno, C. de Graaf, N. López, F. Maseras, J. M. Poblet and C. Bo, *J. Chem. Inf. Model.*, 2015, **55**, 95–103.