UC Irvine

UC Irvine Previously Published Works

Title

Probing the Interaction of PAF with Human Platelet Membrane Using the Fluorescent Probe Laurdan.

Permalink

<https://escholarship.org/uc/item/7mn6t4m3>

Journal

Platelets, 5(3)

ISSN

0953-7104

Authors

Kantar, A
Giorgi, PL
Gratton, E
et al.

Publication Date

1994

DOI

10.3109/09537109409005527

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

Probing the Interaction of PAF with Human Platelet Membrane Using the Fluorescent Probe Laurdan

A. Kantar, P. L. Giorgi, E. Gratton, R. Fiorini

SUMMARY. Changes in membrane polarity of human platelets during the interaction with PAF were investigated by measuring the steady-state fluorescence emission spectra of 2-dimethylamino(6-lauroyl)naphthalene (Laurdan), which is known to be incorporated at the hydrophobic-hydrophilic interface of the bilayer, displaying spectral sensitivity to the polarity of its surroundings. Laurdan shows a marked steady-state emission red-shift in polar solvents, with respect to non-polar solvents. Our results demonstrate that platelet activation factor (PAF) (10^{-7} M) induces a red-shift of the fluorescence emission spectra of Laurdan. These changes were not observed in the presence of the PAF antagonist, L-659,989. These data suggest that the interaction of PAF with its specific receptor and the signalling pathways involved in platelet activation are accompanied by an increase in polarity at the hydrophobic-hydrophilic interface of human platelet membranes.

Platelet activating factor (PAF, 1-O-alkyl-2-O-acetyl-sn-glycerol-3-phosphocholine) is a potent lipid mediator with a wide spectrum of biological activities in a variety of cells.¹⁻³ PAF exerts its actions through the activation of specific membrane receptors.⁴ Human platelets possess a specific extracellular PAF-receptor.^{5,6} The stimulation of human platelets by PAF is accompanied by a complex cascade of biochemical events that influence the plasma membrane.⁷ These events, initiated by the binding of PAF to its specific receptor, include the stimulation of GTPase,⁸ transient elevation of intracellular Ca^{2+} ,⁹ coupling to both adenylate cyclase and phospholipase C,¹⁰ and protein phosphorylation.¹¹ The response of platelets to PAF is transient and PAF desensitizes its receptor.¹² Fluorescence spectroscopy techniques have been widely employed to study the physico-chemical changes in membrane organization. The

fluorescent membrane probe 2-dimethylamino (6-lauroyl) naphthalene (Laurdan) displays spectral sensitivity to the polarity of its surrounding, showing a red-shift of the emission in polar solvents, with respect to non-polar solvents and a blue-shift in non-polar with respect to polar solvents.¹³

In this study we investigated the effect of PAF on membrane polarity of platelets, by measuring steady-state emission spectra of Laurdan incorporated into the platelets plasma membrane before and after the addition of PAF in the absence or presence of the PAF antagonist L-659,989.¹⁴

Materials and Methods

Preparation of Platelets

10 ml of freshly drawn acid-citrate-dextrose anticoagulated blood was centrifuged at 120 g for 15 min at 37°C to prepare the platelets rich plasma. The platelets were isolated and washed according to Kubina et al.¹⁵ The platelets were finally resuspended in Tyrode's buffer (137 mM NaCl, 2 mM KCl, 12 mM $NaHCO_3$, 0.3 mM NaH_2PO_4 , 1 mM $MgCl_2$,

A. Kantar, P. L. Giorgi, Department of Pediatrics, R. Fiorini, Department of Biochemistry, University of Ancona, Ancona, Italy
E. Gratton, Laboratory of Fluorescence Dynamics, Department of Physics, University of Illinois, Urbana, IL, USA.

Correspondence to: A. Kantar, MD, Department of Pediatrics, via Corridoni 11, I-60123 Ancona, Italy.

2 mM CaCl_2 , 5 mM glucose) with 0.1% bovine serum albumin and apyrase (Sigma Chemical Co., St. Louis, MO, USA). The total platelet count was obtained using a Sysmex E-2500 blood analyser. The platelet suspension was adjusted to a concentration of $4 \times 10^8/\text{ml}$. Phase contrast microscopy (Diaplan, Leica, Milan, Italy) showed that, under the conditions used, control and Laurdan labelled platelets remained discoid and functional, being sensitive to thrombin (1 U/ml).⁷ The platelets were immediately used for fluorescence measurements.

Chemical Agents

6-dodecanoyl-2-dimethylamino-naphthalene, Laurdan was purchased from Molecular Probes Inc. (Eugene, OR, USA) and PAF from Sigma Chemical Co. (St. Louis, MO, USA). L-659,989 was kindly donated by Dr W. H. Parsons, Merck Sharp & Dohme Research Laboratories, NJ, USA.

Fluorescence Measurements

The platelets were labelled with an ethanol solution of Laurdan under N_2 at a final probe concentration of 0.1 μM . All samples were prepared at room temperature and used immediately for fluorescence measurements. Steady-state excitation and emission spectra were measured at 37°C before and during activation of the platelets, with a Perkin Elmer Spectrofluorimeter MPF66, equipped with a Perkin Elmer 7300 Personal Computer for data acquisition and elaboration as previously described.¹⁶ The platelets were activated by injecting PAF (10^{-7} M) into the sample and fluorescence spectra were immediately measured. PAF antagonist, L-659,989 was added at a final concentration of 10^{-6} M before activation with PAF. Fluorescence measurements were performed in ten samples.

Results

The background phospholipid fluorescence of the platelets was checked prior to each measurement and was less than 0.1% of the fluorescence when Laurdan was added. The contribution of the light scattering was negligible in our samples because of the low cell concentration used in this study. The fluorescence intensity of the free probe in buffer solution, in the absence of platelets, was negligible and did not increase upon the addition of PAF at the concentration used in the study; moreover, this concentration (10^{-7} M) is lower than the critical micelle concentration of PAF (2×10^{-5} M) as reported in the literature.¹⁷ After the addition of 0.1 μM Laurdan to the platelets a constant increase in fluorescence was observed, plateauing after 10 min (data not shown). Dilution experiments, which are based on the rapid equilibrium of Laurdan between plasma membrane and buffer solution, have demonstrated

Fig. 1 Normalized emission spectra of Laurdan incorporated in the platelet plasma membrane at 37°C before (—) and after (---●) addition of PAF 10^{-7} M. The spectra were measured at an excitation wavelength of 350 nm.

Fig. 2 Normalized excitation spectra of Laurdan incorporated in the platelet plasma membrane at 37°C before (—) and after (---●) addition of PAF 10^{-7} M. The spectra were measured at an emission wavelength of 450 nm.

that the probe in platelets is located in the plasma membrane during the acquisition of the measurements.

Fluorescence emission and excitation spectra of Laurdan in platelets before and during activation with PAF are reported in Figures 1 and 2, respectively. During activation with PAF (Fig. 1) a 5 nm red-shift of the Laurdan emission spectrum was observed. During basal conditions the maximum of Laurdan emission was at 446 nm, while after activation with PAF it was at 451 nm.

The Laurdan excitation spectra during activation of the platelets with PAF (Fig. 2) showed a decrease in intensity at shorter wavelengths with respect to the excitation spectrum in basal conditions. The spectral changes induced by PAF were observed in all samples. The PAF antagonist L-659,989 had no effect on emission or excitation spectra of Laurdan in unactivated platelets.

In the presence of L-659,989, the addition of PAF did not change fluorescence spectra significantly.

Discussion

The organization of cell membrane components depends on a number of factors, including the

protein-lipid composition of the membrane, receptor occupancies, and activities of the membrane proteins.¹⁸ Cell membrane respond to stimuli by activation of membrane components that lead to signal transduction. These events lead to changes in membrane composition and configuration. Depending on the nature of the stimulus and the state of the membrane, different biochemical events take place at the membrane level during activation of the cell.^{19,20}

The fluorescent probe Laurdan has been recently employed to detect modifications in the composition and physical state of living cells.^{17,21-23} Laurdan has been reported to be incorporated at the hydrophilic-hydrophobic interface of the membrane, with the lauric acid tail anchored in the hydrophobic region of the bilayer.²⁴ It has been demonstrated that Laurdan displays spectral sensitivity to the polarity of its surroundings, showing a red-shift of the emission in polar solvents, with respect to nonpolar solvents.¹³ This behavior is referred to dipolar relaxation phenomena that are related to the physical state and the dynamics of the surrounding phospholipid polar head group.²⁵ In single-phase phospholipid vesicles the dynamics of the surroundings detected by Laurdan is very different in the case of the gel or of the liquid-crystalline phase; the probe shows a marked steady-state emission red-shift in the phospholipid liquid-crystalline phase, with respect to the gel phase. If solvent molecules can move during the fluorescence lifetime, the Laurdan excited-state molecular dipole will orient the neighbouring solvent dipoles, and the steady-state Laurdan emission spectra will be red-shifted.²⁶ On the other hand, Laurdan undergoes a marked blue-shift when incorporated in the gel-crystalline phase, with respect to the liquid phase.²⁶

Our results demonstrate that PAF induced a red-shift of the emission spectra of Laurdan incorporated in platelet membranes, indicating an increase in polarity of the environment surrounding the probe. In basal condition, Laurdan reflects the interactions between the probe and the phospholipid polar head groups. The addition of PAF to platelets increases the motional freedom of the bilayer polar residues, as demonstrated by the presence of an unrelaxed red-shift of the spectrum. In a previous study, using the fluorescent probe 1-(4-trimethylammonium-phenyl)-6-phenyl-1,3,5-hexatriene (TMA-DPH), we have shown that PAF induces a time limited increase in platelet membranes phospholipid packing and a decrease in membrane heterogeneity.⁷ Thus, it is therefore possible that such changes may alter the hydrophobic environment surrounding the Laurdan molecules reflecting an increase in membrane polarity. It is interesting that our previous studies demonstrated that PAF induces a decrease in plasma membrane polarity of human neutrophils.²¹ The differences in the PAF receptor, G proteins and effectors between human platelets and human neutrophils have been reviewed by Hwang.⁵ It is plausible

that the distinct molecular events accompanying the activation of human platelets or neutrophils modulates membrane polarity differently. Our data demonstrated that the anti-PAF L-659,989 blocked the effect of PAF on membrane polarity suggesting that these changes may be attributed to the PAF-receptor interaction and the biochemical events associated with signal transduction.

References

1. Koltai M, Hosford D, Guinot P, Esanu A, Braquet P. Platelet activating factor (PAF): a review of its effects, antagonists and possible future clinical implications (Part I). *Drugs* 1991; 42: 9-29.
2. Koltai M, Hosford D, Guinot P, Esanu A, Braquet P. Platelet activating factor (PAF): a review of its effects, antagonists and possible future clinical implications (Part II). *Drugs* 1991; 42: 174-204.
3. Synder F, Lee T C, Blank M L. Platelet activating factor and related ether lipid mediators. Biological activities, metabolism and regulation. *Ann NY Acad Sci* 1989; 568: 35-43.
4. Shukla S D. Platelet activating factor receptor and signal transduction mechanisms. *FASEB J* 1992; 6: 2296-2301.
5. Hwang S-B. Specific receptors of platelet activating factor, receptor heterogeneity, and signal transduction mechanisms. *J Lipid Mediators* 1990; 2: 123-158.
6. Braquet P, Godfroid J J. Conformational properties of PAF-acether receptor on platelets based on structure-activity studies. In: Synder F, ed. *Platelet activating factor and related lipid mediators*. New York: Plenum Press, 1987; 191-235.
7. Kantar A, Giorgi P L, Curatola G, Fiorini R. Interaction between PAF and human platelet membranes: a fluorescence study. *Mediators of Inflammation* 1992; 1: 127-131.
8. Haslam R J, Williams K A, Davis W, Sherwood J, Van der Meulen J. Roles of GTP-binding proteins and protein kinase C in signal transduction in platelets. *Adv Second Messenger Phosphoprotein Res* 1990; 24: 364-369.
9. Valone F H, Johnson B. Modulation of cytoplasmic calcium in human platelets by phospholipid platelet-activating factor, 1-O-alkyl-2-acetyl-sn-glycero-3-phosphorylcholine. *J Immunol* 1985; 134: 1120-1124.
10. Grouch M F, Lapetina E G. No direct correlation between Ca^{2+} mobilization and dissociation of G_i during platelet phospholipase A2 activation. *Biochem Biophys Res Comm* 1988; 135: 21-30.
11. Lapetina E G, Siegel E L. Shape change induced in human platelets by platelet-activating factor. Correlation with the formation of phosphatidic acid and phosphorylation of a 40 000-dalton protein. *J Biol Chem* 1983; 258: 7241-7245.
12. Henson P M. Activation and Desensitization of platelets by platelet-activating factor (PAF) derived from IgE-sensitized basophils I. characteristics of the sensory response. *J Exp Med* 1976; 143: 937-952.
13. Parasassi T, Conti F, Gratton E. Time-resolved fluorescence emission spectra of Laurdan in phospholipid vesicles by multifrequency phase modulation fluorometry. *Cell Mol Biol* 1986; 32: 103-108.
14. Hwang S-B, Lam M H, Alberts A W, Bugianesi R L, Chabala J C, Ponpipom M M. Biochemical and pharmacological characterization of L-659,989: an extremely potent, selective and competitive receptor antagonist of platelet-activating factor. *J Pharmacol Exp Ther* 1988; 246: 534-541.
15. Kubina M, Lanza F, Cazenave J-P, Laustriat G, Kuhry J-G. Parallel investigation of exocytosis kinetics and membrane fluidity changes in human platelets with the fluorescent probe, trimethylammonio-diphenyl hexatriene. *Biochim Biophys Acta* 1987; 901: 138-146.
16. Fiorini R, Curatola G, Kantar A, Giorgi P L, Gratton E. Use of laurdan fluorescence in studying plasma membrane organization of polymorphonuclear leukocytes during the respiratory burst. *Photochem Photobiol* 1993; 57: 438-441.
17. Varveri F S, Mantaka-Marketou A E, Papadopoulos K, Nikokavouras J. Chemiluminescence in organized molecular assemblies: chemiluminescence of lucigenin in Lyso-PAF (C_{16}). *J Photochem Photobiol* 1992; 66: 113-118.

18. Kinnunen P K J. On the principles of functional ordering in biological membranes. *Chem Phys Lipids* 1991; 57: 375-399.
19. Marcus A J, Safier L B. Thromboregulation: multicellular modulation of platelet reactivity in hemostasis and thrombosis. *FASEB J* 1993; 7: 516-522.
20. Ganong B R. Roles of lipid turnover in transmembrane signal transduction. *Am J Med Sci* 1991; 302: 304-312.
21. Kantar A, Giorgi P L, Fiorini R. Effect of PAF on polymorphonuclear leucocyte plasma membrane polarity: a fluorescence study. *Mediators of Inflammation* 1993; 2: 149-151.
22. Fiorini R, Kantar A. Fluorescence emission spectra of laurdan in hemoglobin free erythrocyte membranes in children with trisomy 21. *Biophys J* 1993; 64: 309 (Abstr).
23. Parasassi T, Di Stefano M, Ravagnan G, Saporà O, Gratton E. Membrane aging during cell growth ascertained by laurdan generalized polarization. *Exp Cell Res* 1992; 202: 432-439.
24. Chong P L G. Effects of hydrostatic pressure on the location of prodan in lipid bilayers and cellular membranes. *Biochemistry* 1988; 27: 399-404.
25. Parasassi T, Conti F, Gratton E. Fluorophores in a polar medium: time dependence of emission spectra detected by multifrequency phase and modulation fluorometry. *Cell Mol Biol* 1986; 32: 99-102.
26. Parasassi T, De Stasio G, D'Ubaldo A, Gratton E. Phase fluctuation in phospholipid membranes revealed by laurdan fluorescence. *Biophys J* 1990; 57: 1179-1186.