

LOCAL WELL-POSEDNESS OF SKEW MEAN CURVATURE FLOW FOR SMALL DATA IN $d \geq 4$ DIMENSIONS

JIAXI HUANG AND DANIEL TATARU

ABSTRACT. The skew mean curvature flow is an evolution equation for d dimensional manifolds embedded in \mathbb{R}^{d+2} (or more generally, in a Riemannian manifold). It can be viewed as a Schrödinger analogue of the mean curvature flow, or alternatively as a quasilinear version of the Schrödinger Map equation. In this article, we prove small data local well-posedness in low-regularity Sobolev spaces for the skew mean curvature flow in dimension $d \geq 4$.

CONTENTS

1. Introduction	1
2. The differentiated equations and the gauge choice	5
3. Function spaces and notations	16
4. Elliptic estimates	21
5. Multilinear and nonlinear estimates	37
6. Local energy decay and the linearized problem	41
7. Well-posedness in the good gauge	48
8. The reconstruction of the flow	58
Acknowledgments	70
References	71

Date: June 5, 2024.

2010 Mathematics Subject Classification. Primary: 35Q55; Secondary: 53E10.

Key words and phrases. Skew mean curvature flow, local well-posedness, low regularity, small data.

J. Huang was partially supported by China Postdoctoral Science Foundation Grant 2021M690223 and the NSFC Grant No. 11771415, and was also sponsored by the China Scholarship Council (No. 201806340044) for one year at University of California, Berkeley.

D. Tataru was supported by the NSF grant DMS-1800294 as well as by a Simons Investigator grant from the Simons Foundation.

1. INTRODUCTION

The skew mean curvature flow (SMCF) is a nonlinear Schrödinger type flow modeling the evolution of a d dimensional oriented manifold embedded into a fixed oriented $d + 2$ dimensional manifold. It can be seen as a Schrödinger analogue of the well studied mean curvature flow. In this article, we consider the small data local well-posedness for the skew mean curvature flow in high dimensions $d \geq 4$, for low regularity initial data.

1.1. The (SMCF) equations. Let Σ^d be a d -dimensional oriented manifold, and $(\mathcal{N}^{d+2}, g_{\mathcal{N}})$ be a $d + 2$ -dimensional oriented Riemannian manifold. Let $I = [0, T]$ be an interval and $F : I \times \Sigma^d \rightarrow \mathcal{N}$ be a one parameter family of immersions. This induces a time dependent Riemannian structure on Σ^d . For each $t \in I$, we denote the submanifold by $\Sigma_t = F(t, \Sigma)$, its tangent bundle by $T\Sigma_t$, and its normal bundle by $N\Sigma_t$ respectively. For an arbitrary vector Z at F we denote by Z^\perp its orthogonal projection onto $N\Sigma_t$. The mean curvature $\mathbf{H}(F)$ of Σ_t can be identified naturally with a section of the normal bundle $N\Sigma_t$.

The normal bundle $N\Sigma_t$ is a rank two vector bundle with a naturally induced complex structure $J(F)$ which simply rotates a vector in the normal space by $\pi/2$ positively. Namely, for any point $y = F(t, x) \in \Sigma_t$ and any normal vector $\nu \in N_y \Sigma_t$, we define $J(F)\nu \in N_y \Sigma_t$ as the unique vector with the same length so that

$$J(F)\nu \perp \nu, \quad \omega(F_*(e_1), F_*(e_2), \dots, F_*(e_d), \nu, J(F)\nu) > 0,$$

where ω is the volume form of \mathcal{N} and $\{e_1, \dots, e_d\}$ is an oriented basis of Σ^d . The skew mean curvature flow (SMCF) is defined by the initial value problem

$$(1.1) \quad \begin{cases} (\partial_t F)^\perp = J(F)\mathbf{H}(F), \\ F(\cdot, 0) = F_0, \end{cases}$$

which evolves a codimension two submanifold along its binormal direction with a speed given by its mean curvature.

The (SMCF) was derived both in physics and mathematics. The one-dimensional (SMCF) in the Euclidean space \mathbb{R}^3 is the well-known vortex filament equation (VFE)

$$\partial_t \gamma = \partial_s \gamma \times \partial_s^2 \gamma,$$

where γ is a time-dependent space curve, s is its arc-length parameter and \times denotes the cross product in \mathbb{R}^3 . The (VFE) was first discovered by Da Rios [6] in 1906 in the study of the free motion of a vortex filament.

The (SMCF) also arises in the study of asymptotic dynamics of vortices in the context of superfluidity and superconductivity. For the Gross-Pitaevskii equation, which models the wave function associated with a Bose-Einstein condensate, physics evidence indicates that the vortices would evolve along the (SMCF). An incomplete verification was attempted by T. Lin [20] for the vortex filaments in three space dimensions. For higher dimensions, Jerrard [14] proved this conjecture when the initial singular set is a codimension two sphere with multiplicity one.

The other motivation is that the (SMCF) naturally arises in the study of the hydrodynamical Euler equation. A singular vortex in a fluid is called a vortex membrane in higher dimensions if it is supported on a codimension two subset. The law of locally induced motion of a vortex membrane can be deduced from the Euler equation by applying the Biot-Savart formula. Shashikanth [24] first investigated the motion of a vortex membrane in \mathbb{R}^4 and

showed that it is governed by the two dimensional (SMCF), while Khesin [18] then generalized this conclusion to any dimensional vortex membranes in Euclidean spaces.

From a mathematical standpoint, the (SMCF) equation is a canonical geometric flow for codimension two submanifolds which can be viewed as the Schrödinger analogue of the well studied mean curvature flow. In fact, the infinite-dimensional space of codimension two immersions of a Riemannian manifold admits a generalized Marsden-Weinstein symplectic structure, and hence the Hamiltonian flow of the volume functional on this space is verified to be the (SMCF). Haller-Vizman [12] noted this fact where they studied the nonlinear Grassmannians. For a detailed mathematical derivation of these equations we refer the reader to the article [28, Section 2.1].

The study of higher dimensional (SMCF) is still at its infancy compared with its one-dimensional case. For the 1-d case, we refer the reader to the survey article of Vega [29]. For the higher dimensional case, Song-Sun [28] proved the local existence of (SMCF) with a smooth, compact oriented surface as the initial data in two dimensions, then Song [27] generalized this result to compact oriented manifolds for all $d \geq 2$ and also proved a corresponding uniqueness result. Recently, Li [19] considered the transversal small perturbations of Euclidean planes under the (SMCF) and proved the global regularity for small initial data. In addition, Song [26] also proved that the Gauss map of a d dimensional (SMCF) in \mathbb{R}^{d+2} satisfies a Schrödinger Map type equation but relative to the varying metric. We remark that in one space dimension this is exactly the classical Schrödinger Map type equation, provided that one chooses suitable coordinates, i.e. the arclength parametrization.

As written above, the (SMCF) equations are independent of the choice of coordinates in $I \times \Sigma$; here we include the time interval I to emphasize that coordinates may be chosen in a time dependent fashion. The manifold Σ^d simply serves to provide a parametrization for the moving manifold Σ_t ; it determines the topology of Σ_t , but nothing else. Thus, the (SMCF) system written in the form (1.1) should be seen as a geometric evolution, with a large gauge group, namely the group of time dependent changes of coordinates in $I \times \Sigma$. In particular, interpreting the equations (1.1) as a nonlinear Schrödinger equation will require a good gauge choice. This is further discussed in Section 2.

In this article we will restrict ourselves to the case when $\Sigma^d = \mathbb{R}^d$, i.e. where Σ_t has a trivial topology. We will further restrict to the case when \mathcal{N}^{d+2} is the Euclidean space \mathbb{R}^{d+2} . Thus, the reader should visualize Σ_t as an asymptotically flat codimension two submanifold of \mathbb{R}^{d+2} .

1.2. Scaling and function spaces. To understand what are the natural thresholds for local well-posedness, it is interesting to consider the scaling properties of the solutions. As one might expect, a clean scaling law is obtained when $\Sigma^d = \mathbb{R}^d$ and $\mathcal{N}^{d+2} = \mathbb{R}^{d+2}$. Then we have the following

Proposition 1.1 (Scale invariance for (SMCF)). *Assume that F is a solution of (1.1) with initial data $F(0) = F_0$. If $\lambda > 0$ then $\tilde{F}(t, x) := \lambda^{-1}F(\lambda^2 t, \lambda x)$ is a solution of (1.1) with initial data $\tilde{F}(0) = \lambda^{-1}F_0(\lambda x)$.*

Proof. Since the induced metric and Christoffel symbols of the immersion \tilde{F} are

$$\tilde{g}_{\alpha\beta}(t, x) = \langle \partial_\alpha \tilde{F}, \partial_\beta \tilde{F} \rangle = g_{\alpha\beta}(\lambda^2 t, \lambda x),$$

and

$$\tilde{\Gamma}_{\alpha\beta}^\gamma(t, x) = \lambda \Gamma_{\alpha\beta}^\gamma(\lambda^2 t, \lambda x).$$

Then by the relation $\mathbf{H}(F) = g^{\alpha\beta}(\partial_{\alpha\beta}^2 F - \Gamma_{\alpha\beta}^\gamma \partial_\gamma F)$, we have

$$\begin{aligned} (\partial_t \tilde{F})^\perp &= \lambda (\partial_t F)^\perp(\lambda^2 t, \lambda x) = \lambda J g^{\alpha\beta}(\lambda^2 t, \lambda x) [(\partial_{\alpha\beta}^2 F - \Gamma_{\alpha\beta}^\gamma \partial_\gamma F)(\lambda^2 t, \lambda x)] \\ &= J \tilde{g}^{\alpha\beta}(\partial_{\alpha\beta}^2 \tilde{F} - \tilde{\Gamma}_{\alpha\beta}^\gamma \partial_\gamma \tilde{F})(t, x). \end{aligned}$$

□

The above scaling would suggest the critical Sobolev space for our moving surfaces Σ_t to be $\dot{H}^{\frac{d}{2}+1}$. However, instead of working directly with the surfaces, it is far more convenient to track the regularity at the level of the curvature $\mathbf{H}(\Sigma_t)$, which scales at the level of $\dot{H}^{\frac{d}{2}-1}$.

1.3. The main result. Our objective in this paper is to establish the local well-posedness of skew mean curvature flow for small data at low regularity. A key observation is that providing a rigorous description of fractional Sobolev spaces for functions (tensors) on a rough manifold is a delicate matter, which a-priori requires both a good choice of coordinates on the manifold and a good frame on the vector bundle (the normal bundle in our case). This is done in the next section, where we fix the gauge and write the equation as a quasilinear Schrödinger evolution in a good gauge. At this point, we content ourselves with a less precise formulation of the main result:

Theorem 1.2 (Small data local well-posedness). *Let $s > \frac{d}{2}$, $d \geq 4$. Then there exists $\epsilon_0 > 0$ sufficiently small such that, for all initial data Σ_0 with metric $\|\partial_x(g_0 - I_d)\|_{H^s} \leq \epsilon_0$ and mean curvature $\|\mathbf{H}_0\|_{H^s(\Sigma_0)} \leq \epsilon_0$, the skew mean curvature flow (1.1) for maps from \mathbb{R}^d to the Euclidean space $(\mathbb{R}^{d+2}, g_{\mathbb{R}^{d+2}})$ is locally well-posed on the time interval $I = [0, 1]$ in a suitable gauge.*

Remark 1.2.1. We remark on the necessity of having a smallness condition on both $g_0 - I_d$ and the mean curvature \mathbf{H}_0 . The combined efforts of E. De Giorgi [7], F. J. Almgren, Jr. [1], and J. Simons [25] led to the following theorem (see Theorem 4.2, [3]):

“If $u : \mathbb{R}^{n-1} \rightarrow \mathbb{R}$ is an entire solution to the minimal surface equation and $n \leq 8$, then u is an affine function.”

However, in 1969 E. Bombieri, De Giorgi, and E. Giusti [2] constructed entire non-affine solutions to the minimal surface equation in \mathbb{R}^9 . Hence the bound $\|\mathbf{H}_0\|_{H^s(\Sigma_0)} \leq \epsilon_0$ on the mean curvature does not necessarily imply that the sub-manifold is almost flat.

Here we only prove the small data local well-posedness, which means that the initial submanifold Σ_0 should be a perturbation of Euclidean plane \mathbb{R}^d . Hence, the bound on metric $\|\partial_x(g_0 - I_d)\|_{H^s} \leq \epsilon_0$ is also necessary in our main result, at least in very high dimension. This condition on metric will insure the existence of global harmonic coordinates (see Proposition 8.2). Later, the mean curvature bound will also yield an estimate $\|\partial_x(g_0 - I_d)\|_{H^{s+1}} \lesssim \epsilon_0$ in harmonic coordinates.

Unlike any of the prior results, which prove only existence and uniqueness for smooth data, here we consider rough data and provide a full, Hadamard style well-posedness result based on a more modern, frequency envelope approach and using a paradifferential form for both the full and the linearized equations. For an overview of these ideas we refer the reader to the expository paper [13]. While, for technical reasons, this result is limited to dimensions

$d \geq 4$, we expect the same strategy to also work in lower dimension; the lower dimensional case will be considered in forthcoming work.

The favourable gauge mentioned in the theorem, defined in the next section, will have two components:

- The harmonic coordinates on the manifolds Σ_t .
- The Coulomb gauge for the orthonormal frame on the normal bundle.

In the next section we reformulate the (SMCF) equations as a quasilinear Schrödinger evolution for a good scalar complex variable ψ , which is exactly the mean curvature but represented in the good gauge. There we provide an alternate formulation of the above result, as a well-posedness result for the ψ equation. In the final section of the paper we close the circle and show that one can reconstruct the full (SMCF) flow starting from the good variable ψ .

One may compare our gauge choices with the prior work in [28] and [27]. There the tangential component of $\partial_t F$ in (1.1) is omitted, and the coordinates on the manifold Σ_t are simply those transported from the initial time. The difficulty with such a choice is that the regularity of the map F is no longer determined by the regularity of the second fundamental form, and instead there is a loss of derivatives which may only be avoided if the initial data is assumed to have extra regularity. This loss is what prevents a complete low regularity theory in that approach.

Once our problem is rephrased as a nonlinear Schrödinger evolution, one may compare its study with earlier results on general quasilinear Schrödinger evolutions. This story begins with the classical work of Kenig-Ponce-Vega [15, 16, 17], where local well-posedness is established for more regular and localized data. Lower regularity results in translation invariant Sobolev spaces were later established by Marzuola-Metcalf-Tataru [21, 22, 23]. The local energy decay properties of the Schrödinger equation, as developed earlier in [4, 5, 8, 9] play a key role in these results. While here we are using some of the ideas in the above papers, the present problem is both more complex and exhibits additional structure. Because of this, new ideas and more work are required in order to close the estimates required for both the full problem and for its linearization.

1.4. An overview of the paper. Our first objective in this article will be to provide a self-contained formulation of the (SMCF) flow, interpreted as a nonlinear Schrödinger equation for a single independent variable. This independent variable, denoted by ψ , represents the trace of the second fundamental form on Σ_t , in complex notation. In addition to the independent variables, we will use several dependent variables, as follows:

- The Riemannian metric g on Σ_t .
- The (complex) second fundamental form λ for Σ_t .
- The magnetic potential A , associated to the natural connection on the normal bundle $N\Sigma_t$, and the corresponding temporal component B .
- The advection vector field V , associated to the time dependence of our choice of coordinates.

These additional variables will be viewed as uniquely determined by our independent variable ψ , provided that a suitable gauge choice was made. The gauge choice involves two steps:

- (i) The choice of coordinates on Σ_t ; here we use harmonic coordinates, with suitable boundary conditions at infinity.

- (ii) The choice of the orthonormal frame on $N\Sigma_t$; here we use the Coulomb gauge, again assuming flatness at infinity.

To begin this analysis, in the next section we describe the gauge choices, so that by the end we obtain

- (a) a nonlinear Schrödinger equation for ψ , see (2.35).
- (b) An elliptic fixed time system (2.36) for the dependent variables $\mathcal{S} = (g, \lambda, V, A, B)$, together with suitable compatibility conditions (constraints).

Setting the stage to solve these equations, in Section 3 we describe the function spaces for both ψ and \mathcal{S} . This is done at two levels, first at fixed time, which is useful in solving the elliptic system (2.36), and then using in the space-time setting, which is needed in order to solve the Schrödinger evolution. The fixed time spaces are classical Sobolev spaces, with matched regularities for all the components. The space-time norms are the so called local energy spaces, as developed in [21, 22, 23].

Using these spaces, in Section 4 we consider the solvability of the elliptic system (2.36). This is first considered and solved without reference to the constraint equations, but then we prove that the constraints are indeed satisfied.

Finally, we turn our attention to the Schrödinger system (2.35), in several stages. In Section 5 we establish several multilinear and nonlinear estimates in our space-time function spaces. These are then used in Section 6 in order to prove local energy decay bounds first for the linear paradifferential Schrödinger flow, and then for a full linear Schrödinger flow associated to the linearization of our main evolution. The analysis is completed in Section 7, where we use the linear Schrödinger bounds in order to (i) construct solutions for the full nonlinear Schrödinger flow, and (ii) to prove the uniqueness and continuous dependence of the solutions. The analysis here broadly follows the ideas introduced in [21, 22, 23], but a number of improvements are needed which allow us to take better advantage of the structure of the (SMCF) equations.

Last but not least, in the last section we prove that the full set of variables (g, λ, V, A, B) suffice in order to uniquely reconstruct the defining function F for the evolving surfaces Σ_t , as H_{loc}^{s+2} manifolds. More precisely, with respect to the parametrization provided by our chosen gauge, F has regularity

$$\partial_t F, \partial_x^2 F \in C[0, 1; H^s].$$

2. THE DIFFERENTIATED EQUATIONS AND THE GAUGE CHOICE

The goal of this section is to introduce our main independent variable ψ , which represents the trace of the second fundamental form in complex notation, as well as the following auxiliary variables: the metric g , the second fundamental form λ , the connection coefficients A, B for the normal bundle as well as the advection vector field V . For ψ we start with (1.1) and derive a nonlinear Schrödinger type system (2.35), with coefficients depending on $\mathcal{S} = (\lambda, h, V, A, B)$, where $h = g - I_d$. Under suitable gauge conditions, the auxiliary variables \mathcal{S} are shown to satisfy an elliptic system (2.36), as well as a natural set of constraints. We conclude the section with a gauge formulation of our main result, see Theorem 2.7.

We remark that H. Gomez ([11, Chapter 4]) introduced the language of gauge fields as an appropriate framework for presenting the structural properties of the surface and the evolution equations of its geometric quantities, and showed that the complex mean curvature

of the evolving surface satisfies a nonlinear Schrödinger-type equation. Here we will further derive the self-contained modified Schrödinger system under harmonic coordinate conditions and Coulomb gauge.

2.1. The Riemannian metric g . Let (Σ^d, g) be a d -dimensional oriented manifold and let $(\mathbb{R}^{d+2}, g_{\mathbb{R}^{d+2}})$ be $(d+2)$ -dimensional Euclidean space. Let $\alpha, \beta, \gamma, \dots \in \{1, 2, \dots, d\}$ and $k \in \{1, 2, \dots, d+2\}$. Considering the immersion $F : \Sigma \rightarrow (\mathbb{R}^{d+2}, g_{\mathbb{R}^{d+2}})$, we obtain the induced metric g in Σ ,

$$(2.1) \quad g_{\alpha\beta} = \partial_{x_\alpha} F \cdot \partial_{x_\beta} F.$$

We denote the inverse of the matrix $g_{\alpha\beta}$ by $g^{\alpha\beta}$, i.e.

$$g^{\alpha\beta} := (g_{\alpha\beta})^{-1}, \quad g_{\alpha\gamma} g^{\gamma\beta} = \delta_\alpha^\beta.$$

Let ∇ be the canonical Levi-Civita connection in Σ associated with the induced metric g . A direct computation shows that on the Riemannian manifold (Σ, g) we have the Christoffel symbols

$$\Gamma_{\alpha\beta}^\gamma = \frac{1}{2} g^{\gamma\sigma} (\partial_\beta g_{\alpha\sigma} + \partial_\alpha g_{\beta\sigma} - \partial_\sigma g_{\alpha\beta}) = g^{\gamma\sigma} \partial_{\alpha\beta}^2 F \cdot \partial_\sigma F.$$

Hence, the Laplace-Beltrami operator Δ_g can be written in the form

$$\begin{aligned} \Delta_g f &= \text{tr } \nabla^2 f = g^{\alpha\beta} (\partial_{\alpha\beta}^2 f - \Gamma_{\alpha\beta}^\gamma \partial_\gamma f) \\ &= g^{\alpha\beta} [\partial_{\alpha\beta}^2 f - g^{\gamma\sigma} (\partial_{\alpha\beta}^2 F \cdot \partial_\sigma F) \partial_\gamma f], \end{aligned}$$

for any twice differentiable function $f : \Sigma \rightarrow \mathbb{R}$. The curvature tensor R on the Riemannian manifold (Σ, g) is given by

$$R(\partial_\alpha, \partial_\beta) \partial_\gamma = (\partial_\alpha \Gamma_{\beta\gamma}^\sigma + \Gamma_{\beta\gamma}^m \Gamma_{\alpha m}^\sigma - \partial_\beta \Gamma_{\alpha\gamma}^\sigma - \Gamma_{\alpha\gamma}^m \Gamma_{\beta m}^\sigma) \partial_\sigma.$$

Hence, we have

$$(2.2) \quad R_{\gamma\alpha\beta}^\sigma = \partial_\alpha \Gamma_{\beta\gamma}^\sigma - \partial_\beta \Gamma_{\alpha\gamma}^\sigma + \Gamma_{\beta\gamma}^m \Gamma_{\alpha m}^\sigma - \Gamma_{\alpha\gamma}^m \Gamma_{\beta m}^\sigma.$$

By $R(X, Y, Z, W) = \langle R(Z, W)Y, X \rangle$ and $R_{\alpha\beta\gamma\sigma} = R(\partial_\alpha, \partial_\beta, \partial_\gamma, \partial_\sigma)$, we get

$$R_{\alpha\beta\gamma\sigma} = \langle R(\partial_\gamma, \partial_\sigma) \partial_\beta, \partial_\alpha \rangle = \langle R_{\beta\gamma\sigma}^\mu \partial_\mu, \partial_\alpha \rangle = g_{\mu\alpha} R_{\beta\gamma\sigma}^\mu,$$

We will also use the Ricci curvature

$$\text{Ric}_{\alpha\beta} = R_{\alpha\sigma\beta}^\sigma = g^{\sigma\gamma} R_{\gamma\alpha\sigma\beta}.$$

2.2. The second fundamental form. Let $\bar{\nabla}$ be the Levi-Civita connection in $(\mathbb{R}^{d+2}, g_{\mathbb{R}^{d+2}})$ and let \mathbf{h} be the second fundamental form for Σ as an embedded manifold. For any vector fields $u, v \in T_*\Sigma$, the Gauss relation is

$$\bar{\nabla}_u F_* v = F_*(\nabla_u v) + \mathbf{h}(u, v).$$

Then we have

$$\begin{aligned} \mathbf{h}_{\alpha\beta} &= \mathbf{h}(\partial_\alpha, \partial_\beta) = \bar{\nabla}_{\partial_\alpha} \partial_\beta F - F_*(\nabla_{\partial_\alpha} \partial_\beta) \\ &= \partial_{\alpha\beta}^2 F + \bar{\Gamma}_{kl} \partial_\alpha F^k \partial_\beta F^l - \Gamma_{\alpha\beta}^\gamma \partial_\gamma F. \end{aligned}$$

By $\bar{\Gamma}_{kl}^j = 0$, this gives the mean curvature \mathbf{H} at $F(x)$,

$$\mathbf{H} = \text{tr}_g \mathbf{h} = g^{\alpha\beta} \mathbf{h}_{\alpha\beta} = g^{\alpha\beta} (\partial_{\alpha\beta}^2 F - \Gamma_{\alpha\beta}^\gamma \partial_\gamma F) = \Delta_g F.$$

Hence, the F -equation in (1.1) is rewritten as

$$(\partial_t F)^\perp = J(F)\Delta_g F = J(F)g^{\alpha\beta}(\partial_{\alpha\beta}^2 F - \Gamma_{\alpha\beta}^\gamma \partial_\gamma F).$$

This equation is still independent of the choice of coordinates in Σ^d , which at this point are allowed to fully depend on t .

2.3. The complex structure equations. Here we introduce a complex structure on the normal bundle $N\Sigma_t$. This is achieved by choosing $\{\nu_1, \nu_2\}$ to be an orthonormal basis of $N\Sigma_t$ such that

$$J\nu_1 = \nu_2, \quad J\nu_2 = -\nu_1.$$

Such a choice is not unique; in making it we introduce a second component to our gauge group, namely the group of sections of an $SU(1)$ bundle over $I \times \mathbb{R}^d$.

The vectors $\{F_1, \dots, F_d, \nu_1, \nu_2\}$ form a frame at each point on the manifold (Σ, g) , where F_α for $\alpha \in \{1, \dots, d\}$ are defined as

$$F_\alpha = \partial_\alpha F.$$

If we differentiate the frame, we obtain a set of structure equations of the following type

$$(2.3) \quad \begin{cases} \partial_\alpha F_\beta = \Gamma_{\alpha\beta}^\gamma F_\gamma + \kappa_{\alpha\beta} \nu_1 + \tau_{\alpha\beta} \nu_2, \\ \partial_\alpha \nu_1 = -\kappa_\alpha^\gamma F_\gamma + A_\alpha \nu_2, \\ \partial_\alpha \nu_2 = -\tau_\alpha^\gamma F_\gamma - A_\alpha \nu_1, \end{cases}$$

where the tensors $\kappa_{\alpha\beta}, \tau_{\alpha\beta}$ and the connection coefficients A_α are defined by

$$\kappa_{\alpha\beta} := \partial_\alpha F_\beta \cdot \nu_1, \quad \tau_{\alpha\beta} := \partial_\alpha F_\beta \cdot \nu_2, \quad A_\alpha = \partial_\alpha \nu_1 \cdot \nu_2.$$

The mean curvature \mathbf{H} can be expressed in term of $\kappa_{\alpha\beta}$ and $\tau_{\alpha\beta}$, i.e.

$$\mathbf{H} = g^{\alpha\beta}(\kappa_{\alpha\beta} \nu_1 + \tau_{\alpha\beta} \nu_2).$$

Next, we complexify the structure equations (2.3) as follows. We define the complex vector m and the complex second fundamental form tensor $\lambda_{\alpha\beta}$ to be

$$m = \nu_1 + i\nu_2, \quad \lambda_{\alpha\beta} = \kappa_{\alpha\beta} + i\tau_{\alpha\beta}.$$

Then we define the *complex scalar mean curvature* ψ as the trace of the second fundamental form,

$$(2.4) \quad \psi := \text{tr } \lambda = g^{\alpha\beta} \lambda_{\alpha\beta}.$$

Our objective for the rest of this section will be to interpret the (SMCF) equation as a nonlinear Schrödinger evolution for ψ , by making suitable gauge choices.

We remark that the action of sections of the $SU(1)$ bundle is given by

$$(2.5) \quad \psi \rightarrow e^{i\theta} \psi, \quad \lambda \rightarrow e^{i\theta} \lambda, \quad m \rightarrow e^{i\theta} m, \quad A_\alpha \rightarrow A_\alpha - \partial_\alpha \theta.$$

for a real valued function θ .

We use the convention for the inner product of two complex vectors, say a and b , as

$$\langle a, b \rangle = \sum_{j=1}^{d+2} a_j \bar{b}_j,$$

where a_j and b_j are the complex components of a and b respectively. Then we get the following relations for the complex vector m ,

$$\langle m, m \rangle = |\nu_1|^2 + |\nu_2|^2 = 2, \quad \langle m, \bar{m} \rangle = \langle \bar{m}, m \rangle = |\nu_1|^2 - |\nu_2|^2 = 0.$$

From these relations we obtain

$$\begin{aligned} \kappa_{\alpha\beta}\nu_1 + \tau_{\alpha\beta}\nu_2 &= \frac{1}{2}(\lambda_{\alpha\beta} + \bar{\lambda}_{\alpha\beta})\frac{1}{2}(m + \bar{m}) + \frac{1}{2i}(\lambda_{\alpha\beta} - \bar{\lambda}_{\alpha\beta})\frac{1}{2i}(m - \bar{m}) \\ &= \frac{1}{2}(\lambda_{\alpha\beta}\bar{m} + \bar{\lambda}_{\alpha\beta}m) = \text{Re}(\lambda_{\alpha\beta}\bar{m}). \end{aligned}$$

Then the structure equations (2.3) are rewritten as

$$(2.6) \quad \begin{cases} \partial_\alpha F_\beta = \Gamma_{\alpha\beta}^\gamma F_\gamma + \text{Re}(\lambda_{\alpha\beta}\bar{m}), \\ \partial_\alpha^A m = -\lambda_\alpha^\gamma F_\gamma, \end{cases}$$

where

$$\partial_\alpha^A = \partial_\alpha + iA_\alpha.$$

2.4. The Gauss and Codazzi relations. The Gauss and Codazzi equations are derived from the equality of second derivatives $\partial_\alpha\partial_\beta F_\gamma = \partial_\beta\partial_\alpha F_\gamma$ for the tangent vectors on the submanifold Σ and for the normal vectors respectively. Here we use the Gauss and Codazzi relations to derive the Riemannian curvature, the first compatibility condition and a symmetry.

By the structure equations (2.6), we get

$$(2.7) \quad \begin{aligned} \partial_{\alpha\beta}^2 F_\gamma &= \partial_\alpha(\Gamma_{\beta\gamma}^\sigma F_\sigma + \text{Re}(\lambda_{\beta\gamma}\bar{m})) \\ &= \partial_\alpha\Gamma_{\beta\gamma}^\sigma F_\sigma + \Gamma_{\beta\gamma}^\sigma(\Gamma_{\alpha\sigma}^\mu F_\mu + \text{Re}(\lambda_{\alpha\sigma}\bar{m})) + \text{Re}(\partial_\alpha\lambda_{\beta\gamma}\bar{m} + \lambda_{\beta\gamma}(iA_\alpha\bar{m} - \bar{\lambda}_\alpha^\mu F_\mu)) \\ &= (\partial_\alpha\Gamma_{\beta\gamma}^\sigma + \Gamma_{\beta\gamma}^\mu\Gamma_{\alpha\mu}^\sigma - \text{Re}(\lambda_{\beta\gamma}\bar{\lambda}_\alpha^\sigma))F_\sigma + \text{Re}[(\partial_\alpha^A\lambda_{\beta\gamma} + \Gamma_{\beta\gamma}^\sigma\lambda_{\alpha\sigma})\bar{m}]. \end{aligned}$$

Then in view of $\partial_\alpha\partial_\beta F_\gamma = \partial_\beta\partial_\alpha F_\gamma$ and equating the coefficients of the tangent vectors, we obtain

$$\partial_\alpha\Gamma_{\beta\gamma}^\sigma + \Gamma_{\beta\gamma}^\mu\Gamma_{\alpha\mu}^\sigma - \partial_\beta\Gamma_{\alpha\gamma}^\sigma - \Gamma_{\alpha\gamma}^\mu\Gamma_{\beta\mu}^\sigma = \text{Re}(\lambda_{\beta\gamma}\bar{\lambda}_\alpha^\sigma - \lambda_{\alpha\gamma}\bar{\lambda}_\beta^\sigma).$$

This gives the Riemannian curvature

$$(2.8) \quad R_{\sigma\gamma\alpha\beta} = \langle R_{\gamma\alpha\beta}^\mu F_\mu, F_\sigma \rangle = \langle R(\partial_\alpha, \partial_\beta)F_\gamma, F_\sigma \rangle = \text{Re}(\lambda_{\beta\gamma}\bar{\lambda}_{\alpha\sigma} - \lambda_{\alpha\gamma}\bar{\lambda}_{\beta\sigma}),$$

which is a complex formulation of the Gauss equation. Correspondingly we obtain the the Ricci curvature

$$(2.9) \quad \text{Ric}_{\gamma\beta} = \text{Re}(\lambda_{\gamma\beta}\bar{\psi} - \lambda_{\gamma\alpha}\bar{\lambda}_\beta^\alpha).$$

After equating the coefficients of the vector m in (2.7), we obtain

$$\partial_\alpha^A\lambda_{\beta\gamma} + \Gamma_{\beta\gamma}^\sigma\lambda_{\alpha\sigma} = \partial_\beta^A\lambda_{\alpha\gamma} + \Gamma_{\alpha\gamma}^\sigma\lambda_{\beta\sigma},$$

By the definition of covariant derivatives, i.e.

$$\nabla_\alpha\lambda_{\beta\gamma} = \partial_\alpha\lambda_{\beta\gamma} - \Gamma_{\alpha\beta}^\sigma\lambda_{\sigma\gamma} - \Gamma_{\alpha\gamma}^\sigma\lambda_{\beta\sigma},$$

we obtain

$$\partial_\alpha^A\lambda_{\beta\gamma} - \Gamma_{\alpha\gamma}^\sigma\lambda_{\beta\sigma} - \Gamma_{\alpha\beta}^\sigma\lambda_{\sigma\gamma} = \partial_\beta^A\lambda_{\alpha\gamma} - \Gamma_{\beta\gamma}^\sigma\lambda_{\alpha\sigma} - \Gamma_{\alpha\beta}^\sigma\lambda_{\sigma\gamma}.$$

This implies the complex formulation of the Codazzi equation, namely

$$(2.10) \quad \nabla_\alpha^A\lambda_{\beta\gamma} = \nabla_\beta^A\lambda_{\alpha\gamma}.$$

As a consequence of this equality, we obtain

Lemma 2.1. *The second fundamental form λ satisfies the Codazzi relations*

$$(2.11) \quad \nabla_\alpha^A \lambda_\beta^\gamma = \nabla_\beta^A \lambda_\alpha^\gamma = \nabla^{A,\gamma} \lambda_{\alpha\beta}.$$

Proof. Here we prove the last equality. By $\nabla_\beta g^{\gamma\sigma} = 0$ and (2.10) we have

$$\nabla_\beta^A \lambda_\alpha^\gamma = g^{\gamma\sigma} \nabla_\beta^A \lambda_{\sigma\alpha} = g^{\gamma\sigma} \nabla_\sigma^A \lambda_{\beta\alpha} = \nabla^{A,\gamma} \lambda_{\alpha\beta}.$$

The first equality can be proved similarly. □

Next, we use the relation $\partial_\alpha \partial_\beta m = \partial_\beta \partial_\alpha m$ in order to derive a compatibility condition between the connection A in the normal bundle and the second fundamental form. Indeed, from $\partial_\alpha \partial_\beta m = \partial_\beta \partial_\alpha m$ we obtain the commutation relation

$$(2.12) \quad [\partial_\alpha^A, \partial_\beta^A]m = i(\partial_\alpha A_\beta - \partial_\beta A_\alpha)m.$$

By (2.6) we have

$$\partial_\alpha^A \partial_\beta^A m = -\partial_\alpha^A (\lambda_\beta^\gamma F_\gamma) = -(\partial_\alpha^A \lambda_\beta^\sigma + \lambda_\beta^\gamma \Gamma_{\alpha\gamma}^\sigma) F_\sigma - \lambda_\beta^\gamma \operatorname{Re}(\lambda_{\alpha\gamma} \bar{m}).$$

Then multiplying (2.12) by m yields

$$\begin{aligned} 2i(\partial_\alpha A_\beta - \partial_\beta A_\alpha) &= \langle [-\lambda_\beta^\gamma \operatorname{Re}(\lambda_{\alpha\gamma} \bar{m}) + \lambda_\alpha^\gamma \operatorname{Re}(\lambda_{\beta\gamma} \bar{m})], m \rangle \\ &= -\lambda_\beta^\gamma \bar{\lambda}_{\alpha\gamma} + \lambda_\alpha^\gamma \bar{\lambda}_{\beta\gamma} = 2i \operatorname{Im}(\lambda_\alpha^\gamma \bar{\lambda}_{\beta\gamma}). \end{aligned}$$

This gives the compatibility condition for the curvature of A ,

$$\partial_\alpha A_\beta - \partial_\beta A_\alpha = \operatorname{Im}(\lambda_\alpha^\gamma \bar{\lambda}_{\beta\gamma}).$$

Using covariant derivative, this can be written as

$$(2.13) \quad \nabla_\alpha A_\beta - \nabla_\beta A_\alpha = \operatorname{Im}(\lambda_\alpha^\gamma \bar{\lambda}_{\beta\gamma}),$$

which can be seen as the complex form of the Ricci equations.

We remark that, by equating the coefficients of the tangent vectors in (2.12), we also obtain

$$\partial_\alpha^A \lambda_\beta^\sigma + \lambda_\beta^\gamma \Gamma_{\alpha\gamma}^\sigma = \partial_\beta^A \lambda_\alpha^\sigma + \lambda_\alpha^\gamma \Gamma_{\beta\gamma}^\sigma,$$

and hence

$$\nabla_\alpha^A \lambda_\beta^\sigma = \nabla_\beta^A \lambda_\alpha^\sigma,$$

which is the same as (2.11).

Next, we state an elliptic system for the second fundamental form $\lambda_{\alpha\beta}$ in terms of ψ , using the Codazzi relations (2.11).

Lemma 2.2 (Div-curl system for λ). *The second fundamental form λ satisfies*

$$(2.14) \quad \begin{cases} \nabla_\alpha^A \lambda_{\beta\gamma} - \nabla_\beta^A \lambda_{\alpha\gamma} = 0, \\ \nabla^{A,\alpha} \lambda_{\alpha\beta} = \nabla_\beta^A \psi. \end{cases}$$

We remark that a-priori solutions λ to the above system are not guaranteed to be symmetric, so we record this as a separate property:

$$(2.15) \quad \lambda_{\alpha\beta} = \lambda_{\beta\alpha}.$$

Finally, we turn our attention to the connection A , for which we have the curvature relations (2.13) together with the gauge group (2.5). In order to both fix the gauge and obtain an elliptic system for A , we impose the Coulomb gauge condition

$$(2.16) \quad \nabla^\alpha A_\alpha = 0.$$

Next, we derive the elliptic A -equations from the Ricci equations (2.13).

Lemma 2.3 (Elliptic equations for A). *Under the Coulomb gauge condition, the connection A solves*

$$(2.17) \quad \nabla^\gamma \nabla_\gamma A_\alpha = \operatorname{Re}(\lambda_\alpha^\sigma \bar{\psi} - \lambda_\alpha^\gamma \bar{\lambda}_\gamma^\sigma) A_\sigma + \nabla^\gamma \operatorname{Im}(\lambda_\gamma^\sigma \bar{\lambda}_{\alpha\sigma}).$$

Proof. Applying ∇^β to (2.13), by curvature and (2.16) we obtain

$$\nabla^\beta \nabla_\beta A_\alpha = \operatorname{Ric}_{\alpha\delta} A^\delta + \nabla^\beta \operatorname{Im}(\lambda_\beta^\sigma \bar{\lambda}_{\alpha\sigma}).$$

Then the equation (2.17) for A is obtained from (2.9). \square

2.5. The elliptic equation for the metric g in harmonic coordinates. Here we take the next step towards fixing the gauge, by choosing to work in harmonic coordinates. Precisely, we will require the coordinate functions $\{x_\alpha, \alpha = 1, \dots, d\}$ to be globally Lipschitz solutions of the elliptic equations

$$(2.18) \quad \Delta_g x_\alpha = 0.$$

This determines the coordinates uniquely modulo time dependent affine transformations. This remaining ambiguity will be removed later on by imposing suitable boundary conditions at infinity. After this, the only remaining degrees of freedom in the choice of coordinates will be given by *time independent* translations and rigid rotations. Thus, once a choice is made at the initial time, the coordinates will be uniquely determined later on (see also Remark 2.5.1).

Here we will interpret the above harmonic coordinate condition at fixed time as an elliptic equation for the metric g (see e.g. [10], [30, P161]). The equations (2.18) may be expressed in terms of the Christoffel symbols Γ , which must satisfy the condition

$$(2.19) \quad g^{\alpha\beta} \Gamma_{\alpha\beta}^\gamma = 0, \quad \text{for } \gamma = 1, \dots, d.$$

This implies

$$(2.20) \quad g^{\alpha\beta} \partial_\alpha g_{\beta\gamma} = \frac{1}{2} g^{\alpha\beta} \partial_\gamma g_{\alpha\beta}, \quad \partial_\alpha g^{\alpha\gamma} = \frac{1}{2} g_{\alpha\beta} g^{\gamma\sigma} \partial_\sigma g^{\alpha\beta}.$$

Let

$$(2.21) \quad \Gamma_{\alpha\beta,\gamma} = \frac{1}{2} (\partial_\alpha g_{\beta\gamma} + \partial_\beta g_{\alpha\gamma} - \partial_\gamma g_{\alpha\beta}) = g_{\gamma\sigma} \Gamma_{\alpha\beta}^\sigma.$$

Then we also have

$$g^{\alpha\beta} \Gamma_{\alpha\beta,\gamma} = g^{\alpha\beta} g_{\gamma\sigma} \Gamma_{\alpha\beta}^\sigma = 0,$$

and

$$R_{\alpha\gamma\beta\sigma} = \partial_\beta \Gamma_{\gamma\sigma,\alpha} - \partial_\sigma \Gamma_{\beta\gamma,\alpha} + \Gamma_{\sigma\alpha,\nu} \Gamma_{\beta\gamma}^\nu - \Gamma_{\beta\alpha,\nu} \Gamma_{\gamma\sigma}^\nu.$$

This leads to an equation for the metric g :

Lemma 2.4 (Elliptic equations of g). *In harmonic coordinates, the metric g satisfies*

$$(2.22) \quad \begin{aligned} g^{\alpha\beta} \partial_{\alpha\beta}^2 g_{\gamma\sigma} &= [-\partial_\gamma g^{\alpha\beta} \partial_\beta g_{\alpha\sigma} - \partial_\sigma g^{\alpha\beta} \partial_\beta g_{\alpha\gamma} + \partial_\gamma g_{\alpha\beta} \partial_\sigma g^{\alpha\beta}] \\ &\quad + 2g^{\alpha\beta} \Gamma_{\sigma\alpha,\nu} \Gamma_{\beta\gamma}^\nu - 2 \operatorname{Re}(\lambda_{\gamma\sigma} \bar{\psi} - \lambda_{\alpha\gamma} \bar{\lambda}_\sigma^\alpha). \end{aligned}$$

Proof. By the definition of Ricci curvature, (2.2) and (2.19), we have

$$\begin{aligned} \operatorname{Ric}_{\gamma\sigma} &= g^{\alpha\beta} R_{\alpha\gamma\beta\sigma} = g^{\alpha\beta} (\partial_\beta \Gamma_{\gamma\sigma,\alpha} - \partial_\sigma \Gamma_{\beta\gamma,\alpha}) + g^{\alpha\beta} \Gamma_{\sigma\alpha,\nu} \Gamma_{\beta\gamma}^\nu - g^{\alpha\beta} \Gamma_{\beta\alpha,\nu} \Gamma_{\gamma\sigma}^\nu \\ &= g^{\alpha\beta} (\partial_\beta \Gamma_{\gamma\sigma,\alpha} - \partial_\sigma \Gamma_{\beta\gamma,\alpha}) + g^{\alpha\beta} \Gamma_{\sigma\alpha,\nu} \Gamma_{\beta\gamma}^\nu \\ &= I + II. \end{aligned}$$

We compute the first term I . By the definition of $\Gamma_{\alpha\beta,\gamma}$ in (2.21), we have

$$\begin{aligned} I &= \frac{1}{2} g^{\alpha\beta} [\partial_\beta (\partial_\gamma g_{\sigma\alpha} + \partial_\sigma g_{\gamma\alpha} - \partial_\alpha g_{\gamma\sigma}) - \partial_\sigma (\partial_\beta g_{\gamma\alpha} + \partial_\gamma g_{\beta\alpha} - \partial_\alpha g_{\beta\gamma})] \\ &= -\frac{1}{2} g^{\alpha\beta} \partial_{\alpha\beta}^2 g_{\gamma\sigma} + \frac{1}{2} g^{\alpha\beta} (\partial_{\beta\gamma}^2 g_{\alpha\sigma} + \partial_{\alpha\sigma}^2 g_{\beta\gamma} - \partial_{\gamma\sigma}^2 g_{\alpha\beta}) \end{aligned}$$

Since, by (2.20) we have

$$g^{\alpha\beta} (\partial_{\gamma\beta}^2 g_{\alpha\sigma} - \frac{1}{2} \partial_{\gamma\sigma}^2 g_{\alpha\beta}) = -\partial_\gamma g^{\alpha\beta} (\partial_\beta g_{\alpha\sigma} - \frac{1}{2} \partial_\sigma g_{\alpha\beta}).$$

Then

$$\begin{aligned} I &= -\frac{1}{2} g^{\alpha\beta} \partial_{\alpha\beta}^2 g_{\gamma\sigma} + \frac{1}{2} [-\partial_\gamma g^{\alpha\beta} (\partial_\beta g_{\alpha\sigma} - \frac{1}{2} \partial_\sigma g_{\alpha\beta}) - \partial_\sigma g^{\alpha\beta} (\partial_\beta g_{\alpha\gamma} - \frac{1}{2} \partial_\gamma g_{\alpha\beta})] \\ &= -\frac{1}{2} g^{\alpha\beta} \partial_{\alpha\beta}^2 g_{\gamma\sigma} + \frac{1}{2} [-\partial_\gamma g^{\alpha\beta} \partial_\beta g_{\alpha\sigma} - \partial_\sigma g^{\alpha\beta} \partial_\beta g_{\alpha\gamma} + \partial_\gamma g_{\alpha\beta} \partial_\sigma g^{\alpha\beta}]. \end{aligned}$$

Hence,

$$\operatorname{Ric}_{\gamma\sigma} = -\frac{1}{2} g^{\alpha\beta} \partial_{\alpha\beta}^2 g_{\gamma\sigma} + \frac{1}{2} [-\partial_\gamma g^{\alpha\beta} \partial_\beta g_{\alpha\sigma} - \partial_\sigma g^{\alpha\beta} \partial_\beta g_{\alpha\gamma} + \partial_\gamma g_{\alpha\beta} \partial_\sigma g^{\alpha\beta}] + g^{\alpha\beta} \Gamma_{\sigma\alpha,\nu} \Gamma_{\beta\gamma}^\nu.$$

By (2.9) this concludes the proof of the Lemma. \square

2.6. The motion of the frame $\{F_1, \dots, F_d, m\}$ under (SMCF). Here we derive the equations of motion for the frame, assuming that the immersion F satisfying (1.1).

We begin by rewriting the SMCF equations in the form

$$(2.23) \quad \partial_t F = J(F) \mathbf{H}(F) + V^\gamma F_\gamma,$$

where V^γ is a vector field on the manifold Σ , which in general depends on the choice of coordinates.

By the definition of m and $\lambda_{\alpha\beta}$, we get

$$J(F) \mathbf{H}(F) = J(F) \operatorname{Re}(\psi \bar{m}) = \operatorname{Re} i(\psi \bar{m}) = -\operatorname{Im}(\psi \bar{m}).$$

Hence, the above F -equation (2.23) is rewritten as

$$(2.24) \quad \partial_t F = -\operatorname{Im}(\psi \bar{m}) + V^\gamma F_\gamma.$$

Then we use this to derive the equations of motion for the frame. Applying ∂_α to (2.24), by the structure equations (2.6) we obtain

$$\begin{aligned}
\partial_t F_\alpha &= \partial_\alpha F_t = \partial_\alpha [-\text{Im}(\psi \bar{m}) + V^\gamma F_\gamma] \\
&= -\text{Im}((\partial_\alpha + iA_\alpha)\psi \bar{m} + \overline{\psi(\partial_\alpha + iA_\alpha)\bar{m}}) + \partial_\alpha V^\gamma F_\gamma + V^\gamma(\Gamma_{\alpha\gamma}^\sigma F_\sigma + \text{Re}(\lambda_{\alpha\gamma} \bar{m})) \\
&= -\text{Im}(\partial_\alpha^A \psi \bar{m} - \psi \bar{\lambda}_\alpha^\gamma F_\gamma) + \partial_\alpha V^\gamma F_\gamma + V^\gamma(\Gamma_{\alpha\gamma}^\sigma F_\sigma + \text{Re}(\lambda_{\alpha\gamma} \bar{m})) \\
&= -\text{Im}(\partial_\alpha^A \psi \bar{m}) + \text{Re}(\lambda_{\alpha\gamma} V^\gamma \bar{m}) + [\text{Im}(\psi \bar{\lambda}_\alpha^\gamma) + \nabla_\alpha V^\gamma] F_\gamma \\
&= -\text{Im}(\partial_\alpha^A \psi \bar{m} - i\lambda_{\alpha\gamma} V^\gamma \bar{m}) + [\text{Im}(\psi \bar{\lambda}_\alpha^\gamma) + \nabla_\alpha V^\gamma] F_\gamma.
\end{aligned}$$

By the orthogonality relation $m \perp F_\alpha = 0$, this implies

$$\begin{aligned}
\langle \partial_t m, F_\alpha \rangle &= \partial_t \langle m, F_\alpha \rangle - \langle m, \partial_t F_\alpha \rangle \\
&= -\langle m, -\text{Im}(\partial_\alpha^A \psi \bar{m} - i\lambda_{\alpha\gamma} V^\gamma \bar{m}) \rangle \\
&= \langle m, \frac{i}{2} \overline{(\partial_\alpha^A \psi - i\lambda_{\alpha\gamma} V^\gamma) m} \rangle \\
&= -i(\partial_\alpha^A \psi - i\lambda_{\alpha\gamma} V^\gamma).
\end{aligned}$$

In order to describe the normal component of the time derivative of m , we also need the temporal component of the connection in the normal bundle. This is defined by

$$B = \langle \partial_t \nu_1, \nu_2 \rangle.$$

We have

$$(\partial_t m)^\perp = (\partial_t(\nu_1 + i\nu_2))^\perp = B\nu_2 - iB\nu_1 = -iB(\nu_1 + i\nu_2) = -iBm.$$

Then we get

$$\partial_t m = -i(\partial^{A,\alpha} \psi - i\lambda_\gamma^\alpha V^\gamma) F_\alpha - iBm,$$

which can be further rewritten as

$$\partial_t^B m = -i(\partial^{A,\alpha} \psi - i\lambda_\gamma^\alpha V^\gamma) F_\alpha.$$

Therefore, we obtain the following equations of motion for the frame

$$(2.25) \quad \begin{cases} \partial_t F_\alpha = -\text{Im}(\partial_\alpha^A \psi \bar{m} - i\lambda_{\alpha\gamma} V^\gamma \bar{m}) + [\text{Im}(\psi \bar{\lambda}_\alpha^\gamma) + \nabla_\alpha V^\gamma] F_\gamma, \\ \partial_t^B m = -i(\partial^{A,\alpha} \psi - i\lambda_\gamma^\alpha V^\gamma) F_\alpha. \end{cases}$$

From this we obtain the evolution equation for the metric g . By the definition of the induced metric g (2.1) and (2.25), we have

$$\begin{aligned}
\partial_t g_{\alpha\beta} &= \partial_t \langle F_\alpha, F_\beta \rangle = \langle \partial_t F_\alpha, F_\beta \rangle + \langle F_\alpha, \partial_t F_\beta \rangle \\
&= \langle -\text{Im}(\partial_\alpha^A \psi \bar{m} - i\lambda_{\alpha\gamma} V^\gamma \bar{m}) + [\text{Im}(\psi \bar{\lambda}_\alpha^\gamma) + \nabla_\alpha V^\gamma] F_\gamma, F_\beta \rangle \\
&\quad + \langle F_\alpha, -\text{Im}(\partial_\beta^A \psi \bar{m} - i\lambda_{\beta\gamma} V^\gamma \bar{m}) + [\text{Im}(\psi \bar{\lambda}_\beta^\gamma) + \nabla_\beta V^\gamma] F_\gamma \rangle \\
&= g_{\gamma\beta} (\text{Im}(\psi \bar{\lambda}_\alpha^\gamma) + \nabla_\alpha V^\gamma) + g_{\alpha\gamma} (\text{Im}(\psi \bar{\lambda}_\beta^\gamma) + \nabla_\beta V^\gamma) \\
&= 2\text{Im}(\psi \bar{\lambda}_{\alpha\beta}) + \nabla_\alpha V_\beta + \nabla_\beta V_\alpha,
\end{aligned}$$

which we record for later reference:

$$(2.26) \quad \partial_t g_{\alpha\beta} = 2\text{Im}(\psi \bar{\lambda}_{\alpha\beta}) + \nabla_\alpha V_\beta + \nabla_\beta V_\alpha.$$

Then we also obtain

$$(2.27) \quad \partial_t g^{\alpha\beta} = -2 \operatorname{Im}(\psi \bar{\lambda}^{\alpha\beta}) - \nabla^\alpha V^\beta - \nabla^\beta V^\alpha,$$

$$(2.28) \quad \partial_t \Gamma_{\alpha\beta}^\gamma = \nabla_\alpha G_\beta^\gamma + \nabla_\beta G_\alpha^\gamma - \nabla^\gamma G_{\alpha\beta},$$

where $G_{\alpha\beta}$ are defined by

$$(2.29) \quad G_{\alpha\beta} = \operatorname{Im}(\psi \bar{\lambda}_{\alpha\beta}) + \frac{1}{2}(\nabla_\alpha V_\beta + \nabla_\beta V_\alpha).$$

So far, the choice of V has been unspecified; it depends on the choice of coordinates on our manifold as the time varies. However, once the latter is fixed via the harmonic coordinate condition (2.19), we can also derive an elliptic equation for the advection field V :

Lemma 2.5 (Elliptic equation for the vector field V). *Under the harmonic coordinate condition (2.19), the advection field V solves*

$$(2.30) \quad \nabla^\alpha \nabla_\alpha V^\gamma = -2 \nabla_\alpha \operatorname{Im}(\psi \bar{\lambda}^{\alpha\gamma}) - \operatorname{Re}(\lambda_\sigma^\gamma \bar{\psi} - \lambda_{\alpha\sigma} \bar{\lambda}^{\alpha\gamma}) V^\sigma + 2(\operatorname{Im}(\psi \bar{\lambda}^{\alpha\beta}) + \nabla^\alpha V^\beta) \Gamma_{\alpha\beta}^\gamma.$$

Proof. Applying ∂_t to $g^{\alpha\beta} \Gamma_{\gamma\beta}^\gamma$, by (2.27) and (2.28) we have

$$\begin{aligned} \partial_t(g^{\alpha\beta} \Gamma_{\alpha\beta}^\gamma) &= -2G^{\alpha\beta} \Gamma_{\alpha\beta}^\gamma + g^{\alpha\beta} (2\nabla_\alpha G_\beta^\gamma - \nabla^\gamma G_{\alpha\beta}) \\ &= -2G^{\alpha\beta} \Gamma_{\alpha\beta}^\gamma + 2\nabla_\alpha \operatorname{Im}(\psi \bar{\lambda}^{\alpha\gamma}) + \Delta_g V^\gamma + [\nabla_\alpha, \nabla^\gamma] V^\alpha. \end{aligned}$$

Since

$$[\nabla_\alpha, \nabla^\gamma] V^\alpha = \operatorname{Ric}_\sigma^\gamma V^\sigma = \operatorname{Re}(\lambda_\sigma^\gamma \bar{\psi} - \lambda_{\alpha\sigma} \bar{\lambda}^{\alpha\gamma}) V^\sigma.$$

By the harmonic coordinate condition (2.19), the above two equalities give the V -equations (2.30). \square

Remark 2.5.1. Consider an arbitrary choice of coordinates (parametrization) $\{x_1, \dots, x_d\}$ for the time evolving manifolds Σ_t for $t \in [0, T]$. This yields a representation of Σ_t as the image of a map

$$F : \mathbb{R}^d \times [0, T] \rightarrow \mathbb{R}^{d+2},$$

restricted to time t . If Σ_t moves along the (SMCF) flow (2.23), then we have the relation

$$\partial_t(g^{\alpha\beta} \Gamma_{\alpha\beta}^\gamma) = (V \text{ equation}).$$

Here we uniquely determine the evolution of the coordinates as the time varies by choosing the advection vector field V , precisely so that it satisfies the V -equation (2.30). For this choice we obtain $\partial_t(g^{\alpha\beta} \Gamma_{\alpha\beta}^\gamma) = 0$. This implies that $g^{\alpha\beta} \Gamma_{\alpha\beta}^\gamma$ is conserved for any $x \in \mathbb{R}^d$, and thus the harmonic gauge condition is propagated in time.

2.7. Derivation of the modified Schrödinger system from SMCF. Here we derive the main Schrödinger equation and the second compatibility condition. We consider the commutation relation

$$[\partial_t^B, \partial_\alpha^A] m = i(\partial_t A_\alpha - \partial_\alpha B) m.$$

In order, for the left-hand side, by (2.6) and (2.25) we have

$$\begin{aligned} \partial_t^B \partial_\alpha^A m &= -\partial_t^B (\lambda_\alpha^\gamma F_\gamma) = -\partial_t^B \lambda_\alpha^\gamma \cdot F_\gamma - \lambda_\alpha^\gamma \cdot \partial_t F_\gamma \\ &= -[\partial_t^B \lambda_\alpha^\sigma + \lambda_\alpha^\sigma (\operatorname{Im}(\psi \bar{\lambda}_\sigma^\gamma) + \nabla_\gamma V^\sigma)] F_\sigma + \lambda_\alpha^\gamma \operatorname{Im}(\partial_\gamma^A \psi \bar{m} - i \lambda_{\gamma\sigma} V^\sigma \bar{m}), \end{aligned}$$

and

$$\begin{aligned}
\partial_\alpha^A \partial_t^B m &= -i \partial_\alpha^A [(\partial^{A,\sigma} \psi - i \lambda_\gamma^\sigma V^\gamma) F_\sigma] \\
&= -i \partial_\alpha^A (\partial^{A,\sigma} \psi - i \lambda_\gamma^\sigma V^\gamma) F_\sigma - i (\partial^{A,\sigma} \psi - i \lambda_\gamma^\sigma V^\gamma) [\Gamma_{\alpha\sigma}^\mu F_\mu + \text{Re}(\lambda_{\alpha\sigma} \bar{m})] \\
&= -i \nabla_\alpha^A (\partial^{A,\sigma} \psi - i \lambda_\gamma^\sigma V^\gamma) F_\sigma - i (\partial^{A,\sigma} \psi - i \lambda_\gamma^\sigma V^\gamma) \text{Re}(\lambda_{\alpha\gamma} \bar{m}).
\end{aligned}$$

Then by the above three equalities, equating the coefficients of the tangent vectors and the normal vector m , we obtain the evolution equation for λ

$$(2.31) \quad \partial_t^B \lambda_\alpha^\sigma + \lambda_\alpha^\gamma (\text{Im}(\psi \bar{\lambda}_\gamma^\sigma) + \nabla_\gamma V^\sigma) = i \nabla_\alpha^A (\partial^{A,\sigma} \psi - i \lambda_\gamma^\sigma V^\gamma),$$

as well as the compatibility condition (curvature relation)

$$\begin{aligned}
\partial_t A_\alpha - \partial_\alpha B &= \frac{1}{2i} \langle \lambda_\alpha^\gamma \text{Im}(\partial_\gamma^A \psi \bar{m} - i \lambda_{\gamma\sigma} V^\sigma \bar{m}) + i (\partial^{A,\sigma} \psi - i \lambda_\gamma^\sigma V^\gamma) \text{Re}(\lambda_{\alpha\sigma} \bar{m}), \bar{m} \rangle \\
&= \frac{1}{2} \lambda_\alpha^\gamma (\bar{\partial}_\gamma^A \bar{\psi} + i \bar{\lambda}_{\gamma\sigma} V^\sigma) + \frac{1}{2} (\partial^{A,\sigma} \psi - i \lambda_\gamma^\sigma V^\gamma) \bar{\lambda}_{\alpha\sigma} \\
&= \frac{1}{2} [\lambda_\alpha^\gamma (\bar{\partial}_\gamma^A \bar{\psi} + i \bar{\lambda}_{\gamma\sigma} V^\sigma) + \bar{\lambda}_\alpha^\gamma (\partial_\gamma^A \psi - i \lambda_{\gamma\sigma} V^\sigma)] \\
&= \text{Re}(\lambda_\alpha^\gamma \bar{\partial}_\gamma^A \bar{\psi}) - \text{Im}(\lambda_\alpha^\gamma \bar{\lambda}_{\gamma\sigma}) V^\sigma,
\end{aligned}$$

which we record for later reference:

$$(2.32) \quad \partial_t A_\alpha - \partial_\alpha B = \text{Re}(\lambda_\alpha^\gamma \bar{\partial}_\gamma^A \bar{\psi}) - \text{Im}(\lambda_\alpha^\gamma \bar{\lambda}_{\gamma\sigma}) V^\sigma.$$

This in turn allows us to use the Coulomb gauge condition (2.16) in order to obtain an elliptic equation for B :

Lemma 2.6 (Elliptic equation of B). *The temporal connection coefficient B solves*

$$(2.33) \quad \nabla^\gamma \nabla_\gamma B = -\nabla^\gamma [\text{Re}(\lambda_\gamma^\sigma \bar{\partial}_\sigma^A \bar{\psi}) - \text{Im}(\lambda_\gamma^\sigma \bar{\lambda}_{\sigma\beta}) V^\beta] + (2 \text{Im}(\psi \bar{\lambda}^{\beta\gamma}) + \nabla^\beta V^\gamma + \nabla^\gamma V^\beta) \partial_\beta A_\gamma.$$

Proof. Applying ∇^α to (2.32) yields

$$\nabla^\gamma \nabla_\gamma B = \nabla^\gamma \partial_t A_\gamma - \nabla^\gamma \text{Re}[\lambda_\gamma^\sigma (\bar{\partial}_\sigma^A \bar{\psi} + i \bar{\lambda}_{\sigma\beta} V^\beta)].$$

By the harmonic coordinates condition (2.19), (2.27) and the Coulomb gauge condition (2.16) the first term in the right hand side is written as

$$\begin{aligned}
\nabla^\gamma \partial_t A_\gamma &= g^{\beta\gamma} \nabla_\beta \partial_t A_\gamma = g^{\beta\gamma} (\partial_\beta \partial_t A_\gamma - \Gamma_{\beta\gamma}^\sigma \partial_t A_\sigma) = g^{\beta\gamma} \partial_\beta \partial_t A_\gamma \\
&= \partial_t (g^{\beta\gamma} \partial_\beta A_\gamma) - \partial_t g^{\beta\gamma} \cdot \partial_\beta A_\gamma \\
&= \partial_t \nabla^\gamma A_\gamma + (2 \text{Im}(\psi \bar{\lambda}^{\beta\gamma}) + \nabla^\beta V^\gamma + \nabla^\gamma V^\beta) \partial_\beta A_\gamma \\
&= (2 \text{Im}(\psi \bar{\lambda}^{\beta\gamma}) + \nabla^\beta V^\gamma + \nabla^\gamma V^\beta) \partial_\beta A_\gamma.
\end{aligned}$$

We then obtain the B -equation. □

Next, we use (2.31) to derive the main equation, i.e. the Schrödinger equation for ψ . By (2.10), the right-hand side of (2.31) is rewritten as

$$\nabla_\alpha^A (\partial^{A,\sigma} \psi - i \lambda_\gamma^\sigma V^\gamma) = \nabla_\alpha^A \partial^{A,\sigma} \psi - i \nabla_\gamma^A \lambda_\alpha^\sigma V^\gamma - i \lambda_\gamma^\sigma \nabla_\alpha V^\gamma.$$

Hence, we have

$$(\partial_t^B - V^\gamma \nabla_\gamma^A) \lambda_\alpha^\sigma + \lambda_\alpha^\gamma \text{Im}(\psi \bar{\lambda}_\gamma^\sigma) + (\lambda_\alpha^\gamma \nabla_\gamma V^\sigma - \lambda_\gamma^\sigma \nabla_\alpha V^\gamma) = i \nabla_\alpha^A \nabla^{A,\sigma} \psi,$$

and then contracting this yields

$$i(\partial_t^B - V^\gamma \nabla_\gamma^A) \psi + \nabla_\alpha^A \nabla^{A,\alpha} \psi = -i\lambda_\sigma^\gamma \text{Im}(\psi \bar{\lambda}_\gamma^\sigma).$$

This can be further written as

$$i(\partial_t + iB - V^\gamma \nabla_\gamma^A) \psi + (\nabla_\alpha + iA_\alpha)(\nabla^\alpha + iA^\alpha) \psi = -i\lambda_\sigma^\gamma \text{Im}(\psi \bar{\lambda}_\gamma^\sigma).$$

Hence, under the harmonic coordinates condition (2.19) and the Coulomb gauge condition (2.16) we obtain the main Schrödinger equation

$$(2.34) \quad \begin{aligned} i\partial_t \psi + g^{\alpha\beta} \partial_\alpha \partial_\beta \psi &= iV^\gamma \nabla_\gamma^A \psi - 2iA_\alpha \nabla^\alpha \psi + (B + A_\alpha A^\alpha - i\nabla_\alpha A^\alpha) \psi - i\lambda_\sigma^\gamma \text{Im}(\psi \bar{\lambda}_\gamma^\sigma) \\ &= iV^\gamma \nabla_\gamma^A \psi - 2iA_\alpha \nabla^\alpha \psi + (B + A_\alpha A^\alpha) \psi - i\lambda_\sigma^\gamma \text{Im}(\psi \bar{\lambda}_\gamma^\sigma). \end{aligned}$$

In conclusion, under the Coulomb gauge condition $\nabla^\alpha A_\alpha = 0$ and the harmonic coordinate condition $g^{\alpha\beta} \Gamma_{\alpha\beta}^\gamma = 0$, by (2.34), (2.14), (2.22), (2.30), (2.17) and (2.33), we obtain the Schrödinger equation for the complex mean curvature ψ

$$(2.35) \quad \begin{cases} i\partial_t \psi + g^{\alpha\beta} \partial_\alpha \partial_\beta \psi = i(V - 2A)_\alpha \nabla^\alpha \psi + (B + A_\alpha A^\alpha - V_\alpha A^\alpha) \psi - i\lambda_\sigma^\gamma \text{Im}(\psi \bar{\lambda}_\gamma^\sigma), \\ \psi(0) = \psi_0, \end{cases}$$

where the metric g , curvature tensor λ , the advection field V , connection coefficients A and B are determined at fixed time in an elliptic fashion via the following equations

$$(2.36) \quad \begin{cases} \nabla_\alpha^A \lambda_{\beta\gamma} - \nabla_\beta^A \lambda_{\alpha\gamma} = 0, & \nabla^{A,\alpha} \lambda_{\alpha\beta} = \nabla_\beta^A \psi, \\ g^{\alpha\beta} \partial_{\alpha\beta}^2 g_{\gamma\sigma} = [-\partial_\gamma g^{\alpha\beta} \partial_\beta g_{\alpha\sigma} - \partial_\sigma g^{\alpha\beta} \partial_\beta g_{\alpha\gamma} + \partial_\gamma g_{\alpha\beta} \partial_\sigma g^{\alpha\beta}] \\ \quad + 2g^{\alpha\beta} \Gamma_{\sigma\alpha,\nu} \Gamma_{\beta\gamma}^\nu - 2 \text{Re}(\lambda_{\gamma\sigma} \bar{\psi} - \lambda_{\alpha\gamma} \bar{\lambda}_\sigma^\alpha), \\ \nabla^\alpha \nabla_\alpha V^\gamma = -2\nabla_\alpha \text{Im}(\psi \bar{\lambda}^{\alpha\gamma}) - \text{Re}(\lambda_\sigma^\gamma \bar{\psi} - \lambda_{\alpha\sigma} \bar{\lambda}^{\alpha\gamma}) V^\sigma \\ \quad + 2(\text{Im}(\psi \bar{\lambda}^{\alpha\beta}) + \nabla^\alpha V^\beta) \Gamma_{\alpha\beta}^\gamma, \\ \nabla^\gamma \nabla_\gamma A_\alpha = \text{Re}(\psi \bar{\lambda}_\alpha^\sigma - \lambda_\alpha^\beta \bar{\lambda}_\beta^\sigma) A_\sigma + \nabla^\gamma \text{Im}(\lambda_\gamma^\sigma \bar{\lambda}_{\alpha\sigma}), \\ \nabla^\gamma \nabla_\gamma B = -\nabla^\gamma [\text{Re}(\lambda_\gamma^\sigma \bar{\partial}_\sigma^A \bar{\psi}) - \text{Im}(\lambda_\gamma^\sigma \bar{\lambda}_{\sigma\beta}) V^\beta] \\ \quad + (2 \text{Im}(\psi \bar{\lambda}^{\beta\gamma}) + \nabla^\beta V^\gamma + \nabla^\gamma V^\beta) \partial_\beta A_\gamma. \end{cases}$$

Fixing the remaining degrees of freedom (i.e. the affine group for the choice of the coordinates as well as the time dependence of the $SU(1)$ connection) we can assume that the following conditions hold at infinity in an averaged sense:

$$\lambda(\infty) = 0, \quad g(\infty) = I_d, \quad V(\infty) = 0, \quad A(\infty) = 0, \quad B(\infty) = 0$$

These are needed to insure the unique solvability of the above elliptic equations in a suitable class of functions. For the metric g it will be useful to use the representation

$$g = I_d + h$$

so that h vanishes at infinity.

Finally, we note that the above system (2.35)-(2.36) is accompanied by a large family of compatibility conditions as follows:

- (i) The trace relation (2.4).
- (ii) The Gauss equations (2.8) connecting the curvature R of g and λ .
- (iii) The symmetry property (2.15).

- (iv) The Ricci equations (2.13) for the curvature of A .
- (v) The Coulomb gauge condition (2.16) for A .
- (vi) The harmonic coordinates condition (2.19) for g .
- (vii) The time evolution (2.26) for the metric g (2.26).
- (viii) The time evolution (2.31) for the second fundamental form λ .
- (ix) The time evolution (2.32) for A .

These conditions will all be shown to be satisfied for small solutions to the nonlinear elliptic system (2.35).

Now we can restate here the small data local well-posedness result for the (SMCF) system in Theorem 1.2 in terms of the above system:

Theorem 2.7 (Small data local well-posedness in the good gauge). *Let $s > \frac{d}{2}$, $d \geq 4$. Then there exists $\epsilon_0 > 0$ sufficiently small such that, for all initial data ψ_0 with*

$$\|\psi_0\|_{H^s} \leq \epsilon_0,$$

the modified Schrödinger system (2.35), with (λ, h, V, A, B) determined via the elliptic system (2.36), is locally well-posed in H^s on the time interval $I = [0, 1]$. Moreover, the mean curvature satisfies the bounds

$$(2.37) \quad \|\psi\|_{l^2 \mathbf{X}^s} + \|(\lambda, h, V, A, B)\|_{\mathcal{E}^s} \lesssim \|\psi_0\|_{H^s}.$$

In addition, the auxiliary functions (λ, h, V, A, B) satisfy the constraints (2.4), (2.8), (2.15), (2.13), (2.16) and (2.19), and the time evolutions (2.26), (2.31) and (2.32).

Here the solution ψ satisfies in particular the expected bounds

$$\|\psi\|_{C[0,1;H^s]} \lesssim \|\psi_0\|_{H^s}.$$

The spaces $l^2 \mathbf{X}^s$ and \mathcal{E}^s , defined in the next section, contain a more complete description of the full set of variables $\psi, \lambda, h, V, A, B$, which includes both Sobolev regularity and local energy bounds.

In the above theorem, by well-posedness we mean a full Hadamard-type well-posedness, including the following properties:

- i) Existence of solutions $\psi \in C[0, 1; H^s]$, with the additional regularity properties (2.37).
- ii) Uniqueness in the same class.
- iii) Continuous dependence of solutions with respect to the initial data in the strong H^s topology.
- iv) Weak Lipschitz dependence of solutions with respect to the initial data in the weaker L^2 topology.
- v) Energy bounds and propagation of higher regularity.

3. FUNCTION SPACES AND NOTATIONS

The goal of this section is to define the function spaces where we aim to solve the (SMCF) system in the good gauge, given by (2.35). Both the spaces and the notation presented in this section are similar to those introduced in [21, 22, 23]. All the function spaces described below will be used with respect to harmonic coordinates determined by our gauge choices described in the previous section. We neither attempt nor need to transfer these spaces to other coordinate frames.

For a function $u(t, x)$ or $u(x)$, let $\hat{u} = \mathcal{F}u$ denote the Fourier transform in the spatial variable x . Fix a smooth radial function $\varphi : \mathbb{R}^d \rightarrow [0, 1]$ supported in $[-2, 2]$ and equal to 1 in $[-1, 1]$, and for any $i \in \mathbb{Z}$, let

$$\varphi_i(x) := \varphi(x/2^i) - \varphi(x/2^{i-1}).$$

We then have the spatial Littlewood-Paley decomposition,

$$\sum_{i=-\infty}^{\infty} P_i(D) = 1, \quad \sum_{i=0}^{\infty} S_i(D) = 1,$$

where P_i localizes to frequency 2^i for $i \in \mathbb{Z}$, i.e.,

$$\mathcal{F}(P_i u) = \varphi_i(\xi) \hat{u}(\xi),$$

and

$$S_0(D) = \sum_{i \leq 0} P_i(D), \quad S_i(D) = P_i(D), \quad \text{for } i > 0.$$

For simplicity of notation, we set

$$u_j = S_j u, \quad u_{\leq j} = \sum_{i=0}^j S_i u, \quad u_{\geq j} = \sum_{i=j}^{\infty} S_i u, \quad \text{for } j \geq 0.$$

For each $j \in \mathbb{N}$, let \mathcal{Q}_j denote a partition of \mathbb{R}^d into cubes of side length 2^j , and let $\{\chi_Q\}$ denote an associated partition of unity. For a translation-invariant Sobolev-type space U , set $l_j^p U$ to be the Banach space with associated norm

$$\|u\|_{l_j^p U}^p = \sum_{Q \in \mathcal{Q}_j} \|\chi_Q u\|_U^p$$

with the obvious modification for $p = \infty$.

Next we define the $l^2 \mathbf{X}^s$ and $l^2 N^s$ spaces, which will be used for the primary variable ψ , respectively for the source term in the Schrödinger equation for ψ . Following [21, 22, 23], we first define the X -norm as

$$\|u\|_X = \sup_{l \in \mathbb{N}} \sup_{Q \in \mathcal{Q}_l} 2^{-\frac{l}{2}} \|u\|_{L^2 L^2([0,1] \times Q)}.$$

Here and throughout, $L^p L^q$ represents $L_t^p L_x^q$. To measure the source term, we use an atomic space N satisfying $X = N^*$. A function a is an atom in N if there is a $j \geq 0$ and a $Q \in \mathcal{Q}_j$ such that a is supported in $[0, 1] \times Q$ and

$$\|a\|_{L^2([0,1] \times Q)} \lesssim 2^{-\frac{j}{2}}.$$

Then we define N as linear combinations of the form

$$f = \sum_k c_k a_k, \quad \sum_k |c_k| < \infty, \quad a_k \text{ atom},$$

with norm

$$\|f\|_N = \inf \left\{ \sum_k |c_k| : f = \sum_k c_k a_k, \quad a_k \text{ atoms} \right\}.$$

For solutions which are localized to frequency 2^j with $j \geq 0$, we will work in the space

$$X_j = 2^{-\frac{j}{2}} X \cap L^\infty L^2,$$

with norm

$$\|u\|_{X_j} = 2^{\frac{j}{2}}\|u\|_X + \|u\|_{L^\infty L^2}.$$

One way to assemble the X_j norms is via the X^s space

$$\|u\|_{X^s}^2 = \sum_{j \geq 0} 2^{2js} \|S_j u\|_{X_j}^2.$$

But we will also add the l^p spatial summation on the 2^j scale to X_j , in order to obtain the space $l_j^p X_j$ with norm

$$\|u\|_{l_j^p X_j} = \left(\sum_{Q \in \mathcal{Q}_j} \|\chi_Q u\|_{X_j}^p \right)^{1/p}.$$

We then define the space $l^p X^s$ by

$$\|u\|_{l^p X^s}^2 = \sum_{j \geq 0} 2^{2js} \|S_j u\|_{l_j^p X_j}^2.$$

For the solutions of Schrödinger equation in (2.35), we will be working primarily in $l^2 \mathbf{X}^s$, which is defined by

$$\|u\|_{l^2 \mathbf{X}^s} = \|u\|_{l^2 X^s} + \|\partial_t u\|_{L^2 H^{s-2}}.$$

We note that the second component, introduced here for the first time, serves the purpose of providing better bounds at low frequencies $j \leq 0$.

We analogously define

$$N_j = 2^{\frac{j}{2}} N + L^1 L^2,$$

which has norm

$$\|f\|_{N_j} = \inf_{f=2^{\frac{j}{2}} f_1 + f_2} (\|f_1\|_N + \|f_2\|_{L^1 L^2}),$$

and

$$\|f\|_{l^p N^s}^2 = \sum_{j \geq 0} 2^{2js} \|S_j f\|_{l_j^p N_j}^2.$$

Here we shall be working primarily with $l^2 N^s$.

We also note that for any $j \in \mathbb{N}$, we have

$$\sup_{Q \in \mathcal{Q}_j} 2^{-\frac{j}{2}} \|u\|_{L^2 L^2([0,1] \times Q)} \leq \|u\|_X,$$

hence

$$\|u\|_N \lesssim 2^{j/2} \|u\|_{l_j^1 L^2 L^2}.$$

This bound will come in handy at several places later on.

For the elliptic system (2.36), at a fixed time we define the \mathcal{H}^s norm,

$$\|(\lambda, h, V, A, B)\|_{\mathcal{H}^s} = \|\lambda\|_{H^s} + \| |D|h \|_{H^{s+1}} + \| |D|V \|_{H^s} + \| |D|A \|_{H^s} + \| |D|B \|_{H^{s-1}}.$$

In addition to the fixed time norms, for the study of the Schrödinger equation for ψ we will also need to bound time dependent norms \mathcal{E}^s and $\boldsymbol{\mathcal{E}}^s$ for the elliptic system (2.36), in terms of similar norms for ψ . For simplicity of notation, we define

$$\|u\|_{Z^{\sigma,s}} = \| |D|^\sigma S_0 u \|_{l_0^2 L^\infty L^2}^2 + \sum_{j>0} 2^{2sj} \|S_j u\|_{l_j^2 L^\infty L^2}^2.$$

Then the $\mathbf{Z}^{\sigma,s}$ spaces are defined by

$$\|u\|_{\mathbf{Z}^{\sigma,s}} = \|u\|_{Z^{\sigma,s}} + \||D|^{\sigma} \partial_t u\|_{L^2 H^{s-\sigma-2}}.$$

For the λ , V , A and B -equations in (2.36), we will be working primarily in $\mathbf{Z}^{0,s}$, $\mathbf{Z}^{1,s+1}$, $\mathbf{Z}^{1,s+1}$ and $\mathbf{Z}^{1,s}$, respectively.

On the other hand, for the metric component $h = g - I_d$ we need to introduce some additional structure which is associated to spatial scales larger than the frequency. Precisely, to measure the portion of h which is localized to frequency 2^j , $j \in \mathbb{Z}$, we decompose $P_j h$ as an atomic summation of components $h_{j,l}$ associated to spatial scales 2^l with $l \geq |j|$, where $h_{j,l}$ still localizes to frequency 2^j , i.e.,

$$P_j h = \sum_{l \geq |j|} h_{j,l}.$$

Then we define the Y_j -norm by

$$\|P_j h\|_{Y_j} = \inf_{P_j h = \sum_{l \geq |j|} h_{j,l}} \sum_{l \geq |j|} 2^{l-|j|} \|h_{j,l}\|_{l^1 L^\infty L^2}.$$

Assembling together the dyadic pieces in an l^2 Besov fashion, we obtain the $Y^{\sigma,s}$ space with norm given by

$$\|h\|_{Y^{\sigma,s}}^2 = \sum_{j \in \mathbb{Z}} 2^{2(\sigma j^- + s j^+)} \|P_j h\|_{Y_j}^2.$$

Then for h -equation in (2.35), we will be working primarily in \mathbf{Y}^{s+2} , whose norm is defined by

$$\|h\|_{\mathbf{Y}^{s+2}} = \|h\|_{Y^{s+2}} + \|\nabla \partial_t h\|_{L^2 H^{s-1}} = \|h\|_{Y^{\frac{d}{2}-1-\delta, s+2}} + \|h\|_{\mathbf{Z}^{1,s+2}},$$

where the space $Y^s = Y^{\frac{d}{2}-1-\delta, s} \cap Z^{1,s}$. Collecting all the components defined above, for the elliptic system (2.36), we define the \mathcal{E}^s norm as

$$\|(\lambda, h, V, A, B)\|_{\mathcal{E}^s} = \|\lambda\|_{Z^{0,s}} + \|h\|_{Y^{s+2}} + \|V\|_{Z^{1,s+1}} + \|A\|_{Z^{1,s+1}} + \|B\|_{Z^{1,s}},$$

and the \mathcal{E}^s norm as

$$\|(\lambda, h, V, A, B)\|_{\mathcal{E}^s} = \|(\lambda, h, V, A, B)\|_{\mathcal{E}^s} + \|\partial_t(\lambda, h, V, A, B)\|_{L^2 \mathcal{H}^{s-2}}.$$

Since we often use Littlewood-Paley decompositions, the next lemma is a convenient tool to see that our function spaces are invariant under the action of some standard classes of multipliers:

Lemma 3.1. *For any Schwartz function $f \in \mathcal{S}$, multiplier $m(D)$ with $\|\mathcal{F}^{-1}(m(\xi))\|_{L^1} < \infty$, and translation-invariant Sobolev-type space U , we have*

$$\|m(D)f\|_U \lesssim \|\mathcal{F}^{-1}(m(\xi))\|_{L^1} \|f\|_U.$$

We will also need the following Bernstein-type inequality:

Lemma 3.2 (Bernstein-type inequality). *For any $j, k \in \mathbb{Z}$ with $j + k \geq 0$, $1 \leq r < \infty$ and $1 \leq q \leq p \leq \infty$, we have*

$$(3.1) \quad \|P_k f\|_{l_j^r L^p} \lesssim 2^{kd(\frac{1}{q} - \frac{1}{p})} \|P_k f\|_{l_j^r L^q},$$

$$(3.2) \quad \|\langle x \rangle^{\alpha-d} * f_{\leq 0}\|_{l_0^p L^\infty L^2} \lesssim \|f_{\leq 0}\|_{l_0^1 L^\infty L^2}, \quad \text{for } p > \frac{d}{d-\alpha}.$$

Proof. We begin with the Bernstein-type inequality (3.1). Using the properties of the Fourier transform, $P_k f$ is rewritten as

$$P_k f = \int_{\mathbb{R}^d} (\mathcal{F}^{-1} \varphi_k)(x-y) P_k f(y) dy = 2^{kd} \int_{\mathbb{R}^d} K(2^k(x-y)) S_k f(y) dy,$$

where $K(x) = \mathcal{F}^{-1} \varphi(x)$. Then

$$\begin{aligned} \|P_k f\|_{l_j^r L^p} &= 2^{kd} \left(\sum_{Q \in \mathcal{Q}_j} \|\chi_Q(x) \int_{\mathbb{R}^d} K(2^k(x-y)) P_k f(y) dy\|_{L^p}^r \right)^{1/r} \\ &\leq 2^{kd} \left(\sum_{Q \in \mathcal{Q}_j} \|\chi_Q(x) \int_{\mathbb{R}^d} K(2^k(x-y)) \mathbf{1}_{<M}(2^k(x-y)) P_k f(y) dy\|_{L^p}^r \right)^{1/r} \\ &\quad + 2^{kd} \|K(2^k x) \mathbf{1}_{>M}(2^k x) * P_k f\|_{l_j^r L^p} \\ &:= I + II, \end{aligned}$$

where $d(Q, \tilde{Q}) = \inf\{|x-y| : x \in Q, y \in \tilde{Q}\}$ and M is a large constant. Since $j+k \geq 0$, for any fixed $Q \in \mathcal{Q}_j$ there are only finite many $\tilde{Q} \in \mathcal{Q}_j$ such that $d(Q, \tilde{Q}) \leq 2^{-k}M$. Then from Young's inequality and $1 + 1/p = 1/q + 1/\tilde{q}$ we can bound I by

$$I \lesssim 2^{kd} \left(\sum_{Q \in \mathcal{Q}_j} \sum_{d(Q, \tilde{Q}) \leq 2^{-k}M, \tilde{Q} \in \mathcal{Q}_j} \|K(2^k x)\|_{L^{\tilde{q}}}^r \|\chi_{\tilde{Q}} P_k f\|_{L^q}^r \right)^{1/r} \lesssim 2^{kd(\frac{1}{q} - \frac{1}{p})} \|P_k f\|_{l_j^r L^q}.$$

On the other hand, since $|K(x)| \lesssim \langle x \rangle^{-N}$ for any large N , for II we have

$$\begin{aligned} II &\lesssim 2^{k(d-N)} \| |2^k x|^{-N} \mathbf{1}_{>M}(2^k x) \|_{L^1} \|S_k f\|_{l_j^r L^p} \\ &\lesssim M^{-N+d} \|P_k f\|_{l_j^r L^q}, \end{aligned}$$

which can be absorbed by the term on the left. These imply the bound (3.1).

Next, we prove the estimate (3.2). The left hand side of (3.2) is decomposed as

$$\begin{aligned} \|\langle x \rangle^{\alpha-d} * f_{\leq 0}\|_{l_0^p L^\infty L^2} &\lesssim \sum_{Q \in \mathcal{Q}_0} \|\chi_Q(x) \int \langle y \rangle^{\alpha-d} f_{\leq 0}(x-y) dy\|_{L^\infty L^\infty}^p \\ &\lesssim \sum_{Q \in \mathcal{Q}_0} \|\chi_Q(x) \int_{|y| \leq 1} \langle y \rangle^{\alpha-d} f_{\leq 0}(x-y) dy\|_{L^\infty L^\infty}^p \\ &\quad + \sum_{Q \in \mathcal{Q}_0} \|\chi_Q(x) \int_{|y| > 1} \langle y \rangle^{\alpha-d} \sum_{\tilde{Q} \in \mathcal{Q}_0} \chi_{\tilde{Q}} f_{\leq 0}(x-y) dy\|_{L^\infty L^\infty}^p \\ &= I_1^p + I_2^p. \end{aligned}$$

Then by (3.1) we bound I_1 by

$$I_1 \lesssim \|f_{\leq 0}\|_{l_0^p L^\infty L^\infty} \lesssim \|f_{\leq 0}\|_{l_0^p L^\infty L^2}.$$

On the other hand, by Hölder's inequality and (3.1), we bound I_2 by

$$\begin{aligned}
I_2 &\lesssim \sum_{\tilde{Q} \in \mathcal{Q}_0} \left(\sum_{Q \in \mathcal{Q}_0} \|\chi_Q(x) \int_{|y|>1} \langle y \rangle^{\alpha-d} \chi_{\tilde{Q}} f_{\leq 0}(x-y) dy\|_{L^\infty L^\infty}^p \right)^{1/p} \\
&\lesssim \sum_{\tilde{Q} \in \mathcal{Q}_0} \left(\sum_{Q \in \mathcal{Q}_0} \int_{|y|>1} \chi_Q \langle y \rangle^{(\alpha-d)p} dy \|\chi_{\tilde{Q}} f_{\leq 0}\|_{L^\infty L^q}^p \right)^{1/p} \\
&\lesssim \|f_{\leq 0}\|_{l_0^1 L^\infty L^q} \left(\int \langle y \rangle^{(\alpha-d)p} dy \right)^{1/p} \\
&\lesssim \|f_{\leq 0}\|_{l_0^1 L^\infty L^2},
\end{aligned}$$

which gives the bound (3.2), and thus completes the proof of the lemma. \square

Finally, we define the frequency envelopes as in [21, 22, 23] which will be used in multilinear estimates. Consider a Sobolev-type space U for which we have

$$\|u\|_U^2 = \sum_{k=0}^{\infty} \|S_k u\|_U^2.$$

A frequency envelope for a function $u \in U$ is a positive l^2 -sequence, $\{a_j\}$, with

$$\|S_j u\|_U \leq a_j.$$

We shall only permit slowly varying frequency envelopes. Thus, we require $a_0 \approx \|u\|_U$ and

$$a_j \leq 2^{\delta|j-k|} a_k, \quad j, k \geq 0, \quad 0 < \delta \ll s - d/2.$$

The constant δ only depends on s and the dimension d . Such frequency envelopes always exist. For example, one may choose

$$(3.3) \quad a_j = 2^{-\delta j} \|u\|_U + \max_k 2^{-\delta|j-k|} \|S_k u\|_U.$$

4. ELLIPTIC ESTIMATES

Here we consider the solvability of the elliptic system (2.36), together with the constraints (2.4), (2.8), (2.15), (2.13), (2.19) and (2.16). We will do this in two steps. First we prove that this system is solvable in Sobolev spaces at fixed time. Then we prove space-time bounds in local energy spaces; the latter will be needed in the study of the Schrödinger evolution (2.35).

For simplicity of notations, we define the set of elliptic variables by

$$\mathcal{S} = (\lambda, h, V, A, B),$$

Later when we compare two solutions for (2.36), we will denote the differences of two solutions or the linearized variable by

$$\delta \mathcal{S} = (\delta \lambda, \delta h, \delta V, \delta A, \delta B).$$

Our fixed time result is as follows:

Theorem 4.1. *a) Assume that ψ is small in H^s for $s > d/2$ and $d \geq 4$. Then the elliptic system (2.36) admits a unique small solution $\mathcal{S} = (\lambda, h, V, A, B)$ in \mathcal{H}^s , with*

$$(4.1) \quad \|\mathcal{S}\|_{\mathcal{H}^s} \lesssim \|\psi\|_{H^s}.$$

In addition this solution has a smooth dependence on ψ in H^s and satisfies the constraints (2.4), (2.8), (2.15), (2.13), (2.19) and (2.16).

b) Let ψ and $(\lambda, h, V, A, B) = \mathcal{S}(\psi)$ be as above. Then for the linearization of the solution map above we also have the bound:

$$(4.2) \quad \|\mathcal{D}\mathcal{S}(\delta\psi)\|_{\mathcal{H}^\sigma} \lesssim \|\delta\psi\|_{H^\sigma}, \quad \sigma \in (d/2 - 3, s].$$

Moreover, assume that \tilde{p}_k and s_k are admissible frequency envelopes for $\psi \in H^\sigma$, $\mathcal{S} \in \mathcal{H}^s$ respectively. Then we have

$$(4.3) \quad \|S_k \mathcal{S}\|_{\mathcal{H}^\sigma} \lesssim \tilde{p}_k + s_k \|\delta\mathcal{S}\|_{\mathcal{H}^\sigma}.$$

c) We also have a similar bound for the Hessian of the solution map,

$$(4.4) \quad \|D^2\mathcal{S}(\delta_1\psi, \delta_2\psi)\|_{\mathcal{H}^\sigma} \lesssim \|\delta_1\psi\|_{H^{\sigma_1}} \|\delta_2\psi\|_{H^{\sigma_2}},$$

with $\sigma, \sigma_1, \sigma_2 \in (d/2 - 3, s], \sigma_1 + \sigma_2 = \sigma + s$.

Remark 4.1.1. Here we solve the elliptic system (2.36) in the function space \mathcal{H}^s for $s > d/2$, which is more suitable for the nonlinear estimates of ψ -equation. Nevertheless, this system can be solved in a similar fashion for the full range of indices s above scaling, namely $s > d/2 - 1$. However, in the additional range $d/2 - 1 < s \leq d/2$ one needs to replace the above solution space \mathcal{H}^s with a slightly larger one,

$$\|\mathcal{S}\|_{\tilde{\mathcal{H}}^s} = \|\lambda\|_{H^s} + \|D|h\|_{H^{\sigma+1}} + \|D|V|\|_{H^\sigma} + \|D|A|\|_{H^\sigma} + \|D|B|\|_{H^{\sigma-1}},$$

where $\sigma = 2s - d/2$. Then the elliptic system (2.36) admits a unique small solution \mathcal{S} in $\tilde{\mathcal{H}}^s$ with $\|\mathcal{S}\|_{\tilde{\mathcal{H}}^s} \lesssim \|\psi\|_{H^s}$.

Proof of Theorem 4.1. a) The proof is based on a perturbative argument. We rewrite the system (2.36) in the form

$$(4.5) \quad \begin{cases} \partial_\alpha \lambda_{\alpha\beta} = \partial_\beta \psi + H_{1\lambda}, \\ \partial_\alpha \lambda_{\beta\gamma} - \partial_\beta \lambda_{\alpha\gamma} = H_{2\lambda}, \\ \Delta g_{\gamma\sigma} = H_g, \\ \Delta V^\gamma = H_V, \\ \Delta A_\alpha = H_A, \\ \Delta B = H_B, \end{cases}$$

where $\Delta = \sum_{\alpha=1}^d \partial_\alpha^2$ and the nonlinear source terms are given by

$$\begin{aligned} H_{1\lambda} &= iA_\beta \psi - h^{\alpha\mu} \partial_\mu \lambda_{\alpha\beta} + \Gamma_{\alpha\beta, \sigma} \lambda^{\alpha\sigma}, \\ H_{2\lambda} &= -iA_\alpha \lambda_{\beta\gamma} + iA_\beta \lambda_{\alpha\gamma} + \Gamma_{\alpha\gamma, \sigma} \lambda_\beta^\sigma - \Gamma_{\beta\gamma, \sigma} \lambda_\alpha^\sigma, \\ H_g &= -h^{\alpha\beta} \partial_{\alpha\beta}^2 g_{\gamma\sigma} - \partial_\gamma g^{\alpha\beta} \partial_\beta g_{\alpha\sigma} - \partial_\sigma g^{\alpha\beta} \partial_\beta g_{\alpha\gamma} + \partial_\gamma g_{\alpha\beta} \partial_\sigma g^{\alpha\beta} \\ &\quad + 2g^{\alpha\beta} \Gamma_{\sigma\alpha, \nu} \Gamma_{\beta\gamma}^\nu - 2\operatorname{Re}(\lambda_{\gamma\sigma} \bar{\psi} - \lambda_{\alpha\gamma} \bar{\lambda}_\sigma^\alpha), \end{aligned}$$

$$\begin{aligned}
H_V &= -\nabla^\alpha \nabla_\alpha V^\gamma + \Delta V^\gamma - 2\nabla_\alpha \operatorname{Im}(\psi \bar{\lambda}^{\alpha\gamma}) - \operatorname{Re}(\lambda_\sigma^\gamma \bar{\psi} - \lambda_{\alpha\sigma} \bar{\lambda}^{\alpha\gamma}) V^\sigma \\
&\quad + 2(\operatorname{Im}(\psi \bar{\lambda}^{\alpha\beta}) + \nabla^\alpha V^\beta) \Gamma_{\alpha\beta}^\gamma, \\
H_A &= -\nabla^\gamma \nabla_\gamma A_\alpha + \Delta A_\alpha + \operatorname{Re}(\psi \bar{\lambda}_\alpha^\sigma - \lambda_\alpha^\beta \bar{\lambda}_\beta^\sigma) A_\sigma + \nabla^\gamma \operatorname{Im}(\lambda_\gamma^\sigma \bar{\lambda}_{\alpha\sigma}), \\
H_B &= -\nabla^\gamma \nabla_\gamma B + \Delta B - \nabla^\gamma \operatorname{Re}[\lambda_\gamma^\sigma (\bar{\partial}_\sigma^A \bar{\psi} + i \bar{\lambda}_{\sigma\beta} V^\beta)] \\
&\quad + (2 \operatorname{Im}(\psi \bar{\lambda}^{\beta\gamma}) + \nabla^\beta V^\gamma + \nabla^\gamma V^\beta) \partial_\beta A_\gamma.
\end{aligned}$$

In order to prove the existence of solutions to (4.5) at a fixed time for small $\psi \in H^s$, we construct solutions to (4.5) iteratively. We define the sets of elliptic variables

$$\mathcal{S}^{(n)} = (\lambda^{(n)}, h^{(n)}, V^{(n)}, A^{(n)}, B^{(n)}),$$

at each step, based on the scheme

$$(4.6) \quad \begin{cases} \partial_\alpha \lambda_{\alpha\beta}^{(n+1)} = \partial_\beta \psi + H_{1\lambda}^{(n)}, \\ \partial_\alpha \lambda_{\beta\gamma}^{(n+1)} - \partial_\beta \lambda_{\alpha\gamma}^{(n+1)} = H_{2\lambda}^{(n)}, \\ \Delta g_{\gamma\sigma}^{(n+1)} = H_g^{(n)}, \\ \Delta V^{(n+1)\gamma} = H_V^{(n)}, \\ \Delta A_\alpha^{(n+1)} = H_A^{(n)}, \\ \Delta B^{(n+1)} = H_B^{(n)}, \end{cases}$$

with the trivial initialization

$$\mathcal{S}^{(0)} = (0, 0, 0, 0, 0), \quad g^{(0)} = h^{(0)} + I_d,$$

where $H_{1\lambda}^{(n)}$, $H_{2\lambda}^{(n)}$, $H_g^{(n)}$, $H_V^{(n)}$, $H_A^{(n)}$ and $H_B^{(n)}$ are defined as $H_{1\lambda}$, $H_{2\lambda}$, H_g , H_V , H_A and H_B with

$$\mathcal{S} = \mathcal{S}^{(n)}.$$

We will inductively show that

$$\|\mathcal{S}^{(n)}\|_{\mathcal{H}^s} \leq C \|\psi\|_{H^s},$$

with a large universal constant C . This trivially holds for our initialization. Then using a standard Littlewood-Paley decomposition, Bernstein's inequality and the smallness of our data $\psi \in H^s$ in order to estimate the source terms $H_{1\lambda}^{(n)}$, $H_{2\lambda}^{(n)}$, $H_g^{(n)}$, $H_V^{(n)}$, $H_A^{(n)}$ and $H_B^{(n)}$, we obtain

$$\|\mathcal{S}^{(n+1)}\|_{\mathcal{H}^s} \lesssim \|\psi\|_{H^s} + \|\mathcal{S}^{(n)}\|_{\mathcal{H}^s}^2 (1 + \|\mathcal{S}^{(n)}\|_{\mathcal{H}^s})^N \lesssim \|\psi\|_{H^s}.$$

From the iterative scheme (4.6) and $\psi \in H^s$ small, we can repeat the same analysis for successive differences in order to obtain a small Lipschitz constant,

$$\|\mathcal{S}^{(n+1)} - \mathcal{S}^{(n)}\|_{\mathcal{H}^s} \ll \|\mathcal{S}^{(n)} - \mathcal{S}^{(n-1)}\|_{\mathcal{H}^s}.$$

Hence the elliptic system (2.36) admits a small solution

$$\mathcal{S} = \lim_{n \rightarrow \infty} \mathcal{S}^{(n)} \in \mathcal{H}^s.$$

The uniqueness and the Lipschitz dependence of the solution on ψ are easily obtained by similar elliptic estimates.

Next, we prove the solution satisfies the constraints (2.4), (2.15), (2.13), (2.16), (2.19) and (2.8). To get started, let us summarize the compatibility conditions we need to verify:

$$\begin{aligned} \psi &= g^{\alpha\beta} \lambda_{\alpha\beta}; & \lambda_{\alpha\beta} &= \lambda_{\beta\alpha}; & \nabla_\alpha A_\beta - \nabla_\beta A_\alpha &= \text{Im}(\lambda_\alpha^\gamma \bar{\lambda}_{\gamma\beta}); & \nabla^\alpha A_\alpha &= 0; \\ g^{\alpha\beta} \Gamma_{\alpha\beta,\delta} &= 0; & \text{Ric}_{\gamma\beta} &= \text{Re}(\lambda_{\gamma\beta} \bar{\psi} - \lambda_{\gamma\alpha} \bar{\lambda}_\beta^\alpha); & R_{\sigma\gamma\alpha\beta} &= \text{Re}(\lambda_{\gamma\beta} \bar{\lambda}_{\sigma\alpha} - \lambda_{\gamma\alpha} \bar{\lambda}_{\sigma\beta}). \end{aligned}$$

We need to show that these constraints are satisfied for solutions to the elliptic system (2.36). We can disregard the B and V equations, which are unneeded here.

To shorten the notations, we define

$$\begin{aligned} C^1 &= \psi - g^{\alpha\beta} \lambda_{\alpha\beta}, & C_{\alpha\beta}^2 &= \lambda_{\alpha\beta} - \lambda_{\beta\alpha}, \\ C_{\alpha\beta}^3 &= \nabla_\alpha A_\beta - \nabla_\beta A_\alpha - \text{Im}(\lambda_{\alpha\gamma} \bar{\lambda}_\beta^\gamma), & C^4 &= \nabla^\alpha A_\alpha, & C_\delta^5 &= g^{\alpha\beta} \Gamma_{\alpha\beta,\delta}, \\ C_{\gamma\beta}^6 &= \text{Ric}_{\gamma\beta} - \text{Re}(\lambda_{\gamma\beta} \bar{\psi} - \lambda_{\gamma\alpha} \bar{\lambda}_\beta^\alpha), & C_{\sigma\gamma\alpha\beta}^7 &= R_{\sigma\gamma\alpha\beta} - \text{Re}(\lambda_{\gamma\beta} \bar{\lambda}_{\sigma\alpha} - \lambda_{\gamma\alpha} \bar{\lambda}_{\sigma\beta}). \end{aligned}$$

Here C^2 and C^3 are antisymmetric, C^6 is symmetric and C^7 inherits all the linear symmetries of the curvature tensor.

Our goal is to show that all these functions vanish. We will prove this by showing that they solve a coupled linear homogeneous elliptic system of the form

$$\begin{aligned} \nabla_\beta^A C^1 &= \nabla^{A,\alpha} C_{\alpha\beta}^2, \\ \Delta_g^A C^2 &= (\lambda + \psi)(C^3 + C^6 + C^7) + (\lambda^2 + \lambda\psi)C^2, \\ \Delta_g C^3 &= RC^3 + \nabla(C^6 A) + \nabla(\lambda\nabla C^2 + \nabla\lambda C^2), \\ \Delta_g C^4 &= \nabla(C^6 A) + RC^3 + \nabla^2(\lambda C^2), \\ \Delta_g C^5 &= RC^5 + \nabla(C^1\psi) + \lambda\nabla C^2 + \nabla\lambda C^2, \\ C_{\gamma\sigma}^6 &= \frac{1}{2}(\nabla_\gamma C_\sigma^5 + \nabla_\sigma C_\gamma^5), \\ \nabla_\delta C_{\sigma\gamma\alpha\beta}^7 + \nabla_\sigma C_{\gamma\delta\alpha\beta}^7 + \nabla_\gamma C_{\delta\sigma\alpha\beta}^7 &= 0, \\ \nabla^\sigma C_{\sigma\gamma\alpha\beta}^7 &= \nabla_\alpha C_{\gamma\beta}^6 - \nabla_\beta C_{\gamma\alpha}^6 + \nabla(\lambda C^1 + \lambda C^2). \end{aligned}$$

Here the covariant Laplace operators Δ_g , respectively Δ_g^A are symmetric and coercive in \dot{H}^1 . We consider these equations as a system in the space

$$(C^1, C^2, C^3, C^4, C^5, C^6, C^7) \in \dot{H}^1 \times \dot{H}^1 \times \dot{H}^1 \times \dot{H}^1 \times \dot{H}^1 \times L^2 \times L^2$$

using \dot{H}^1 bounds for the Laplace operator in the second to fifth equations, and interpreting the last two equations as an elliptic div-curl system in L^2 , with an \dot{H}^{-1} source term. Since the coefficients are all small, the right hand side terms are perturbative and 0 is the unique solution for this system. The details are left for the reader, as they only involve Sobolev embeddings and Hölder's inequality.

To complete the argument, we now successively derive the equations in the above system. In the computations below, it is convenient to introduce several auxiliary notations. The curvature of the connection A acting on complex valued functions is denoted by

$$\mathbf{F}_{\alpha\beta} = \partial_\alpha A_\beta - \partial_\beta A_\alpha$$

so that we have

$$[\nabla_\alpha^A, \nabla_\beta^A]\psi = i\mathbf{F}_{\alpha\beta}\psi.$$

We also set

$$C_{\sigma\gamma\alpha\beta}^7 = R_{\sigma\gamma\alpha\beta} - \tilde{R}_{\sigma\gamma\alpha\beta}, \quad \tilde{R}_{\sigma\gamma\alpha\beta} := \operatorname{Re}(\lambda_{\gamma\beta}\bar{\lambda}_{\sigma\alpha} - \lambda_{\gamma\alpha}\bar{\lambda}_{\sigma\beta}),$$

respectively

$$C_{\gamma\beta}^6 = \operatorname{Ric}_{\gamma\beta} - \widetilde{\operatorname{Ric}}_{\gamma\beta}, \quad \widetilde{\operatorname{Ric}}_{\gamma\beta} := \operatorname{Re}(\lambda_{\gamma\beta}\bar{\psi} - \lambda_{\gamma\alpha}\bar{\lambda}_{\beta}^{\alpha}), \quad \tilde{R} := g^{\gamma\beta}\widetilde{\operatorname{Ric}}_{\gamma\beta},$$

and

$$C_{\alpha\beta}^3 = \mathbf{F}_{\alpha\beta} - \tilde{\mathbf{F}}_{\alpha\beta}, \quad \tilde{\mathbf{F}}_{\alpha\beta} := \operatorname{Im}(\lambda_{\alpha\gamma}\bar{\lambda}_{\beta}^{\gamma}).$$

The equation for C^1 . This equation has the exact form

$$\nabla_{\beta}^A C^1 = \nabla^{A,\alpha} C_{\alpha\beta}^2.$$

This is obtained by (2.14) directly. □

The equation for C^2 . The full system for C^2 has the form

$$(4.7) \quad \Delta_g^A C_{\alpha\beta}^2 = (\lambda + \psi)(C^3 + C^6 + C^7) + (\lambda^2 + \lambda\psi)C^2.$$

By λ -equation (2.14) we have

$$\begin{aligned} \nabla^{A,\gamma}\nabla_{\gamma}^A \lambda_{\alpha\beta} &= [\nabla^{A,\gamma}, \nabla_{\alpha}^A] \lambda_{\gamma\beta} + \nabla_{\alpha}^A \nabla_{\beta}^A \psi \\ &= \operatorname{Ric}_{\alpha\mu} \lambda^{\mu}_{\beta} + R_{\sigma\alpha\beta\mu} \lambda^{\sigma\mu} + iC_{\gamma\alpha}^3 \lambda^{\gamma}_{\beta} + i \operatorname{Im}(\lambda_{\gamma}^{\mu} \bar{\lambda}_{\mu\alpha}) \lambda^{\gamma}_{\beta} + \nabla_{\alpha}^A \nabla_{\beta}^A \psi. \end{aligned}$$

Then we use C^6 , C^7 and C^3 to give

$$\begin{aligned} \Delta_g^A C_{\alpha\beta}^2 &= C_{\alpha\mu}^6 \lambda^{\mu}_{\beta} - C_{\beta\mu}^6 \lambda^{\mu}_{\alpha} + C_{\sigma\alpha\beta\mu}^7 \lambda^{\sigma\mu} - C_{\sigma\beta\alpha\mu}^7 \lambda^{\sigma\mu} \\ &\quad + iC_{\gamma\alpha}^3 \lambda^{\gamma}_{\beta} - iC_{\gamma\beta}^3 \lambda^{\gamma}_{\alpha} + iC_{\alpha\beta}^3 \psi + C^2(\lambda^2 + \lambda\psi). \end{aligned}$$

Hence, the C^2 -equation (4.7) follows. □

The equation for C^3 . This has the form

$$(4.8) \quad \begin{aligned} \Delta_g C_{\alpha\beta}^3 &= \nabla_{\beta}(C_{\alpha\delta}^6 A^{\delta}) - \nabla_{\alpha}(C_{\beta\delta}^6 A^{\delta}) + R_{\beta\alpha\sigma\delta} C^{3,\sigma\delta} + \operatorname{Ric}_{\alpha\delta} C^{3,\delta\beta} - \operatorname{Ric}_{\beta\delta} C^{3,\delta\alpha} \\ &\quad + \nabla^{\gamma} \operatorname{Im}(\lambda_{\gamma}^{\sigma} (\overline{\nabla_{\alpha}^A C_{\sigma\beta}^2} - \overline{\nabla_{\beta}^A C_{\sigma\alpha}^2})) + \nabla_{\gamma}^A \lambda^{\sigma}_{\beta} \bar{C}^2_{\alpha\sigma}. \end{aligned}$$

To prove this, it is convenient to separate the left hand side into two terms,

$$\Delta_g C_{\alpha\beta}^3 = ([\Delta_g, \nabla_{\alpha}] A_{\beta} - [\Delta_g, \nabla_{\beta}] A_{\alpha}) + (\nabla_{\alpha} \Delta_g A_{\beta} - \nabla_{\beta} \Delta_g A_{\alpha} - \Delta_g \tilde{\mathbf{F}}_{\alpha\beta}) := I + II.$$

For the commutator we use the Bianchi identities to compute

$$\begin{aligned} I &= [\nabla^{\sigma} \nabla_{\sigma}, \nabla_{\alpha}] A_{\beta} - [\nabla^{\sigma} \nabla_{\sigma}, \nabla_{\beta}] A_{\alpha} \\ &= \nabla^{\sigma} (R_{\sigma\alpha\beta\delta} A^{\delta} - R_{\sigma\beta\alpha\delta} A^{\delta}) + (R_{\sigma\alpha\beta\delta} - R_{\sigma\beta\alpha\delta}) \nabla^{\sigma} A^{\delta} + R^{\sigma}_{\alpha\sigma\delta} \nabla^{\delta} A_{\beta} - R^{\sigma}_{\beta\sigma\delta} \nabla^{\delta} A_{\alpha} \\ &= \nabla^{\sigma} R_{\beta\alpha\sigma\delta} A^{\delta} + 2R_{\beta\alpha\sigma\delta} \nabla^{\sigma} A^{\delta} + \operatorname{Ric}_{\alpha\delta} \nabla^{\delta} A_{\beta} - \operatorname{Ric}_{\beta\delta} \nabla^{\delta} A_{\alpha} \\ &= (\nabla_{\beta} \operatorname{Ric}_{\alpha\delta} - \nabla_{\alpha} R_{\beta\delta}) A^{\delta} + R_{\beta\alpha\sigma\delta} \mathbf{F}^{\sigma\delta} + \operatorname{Ric}_{\alpha\delta} (\mathbf{F}^{\delta}_{\beta} + \nabla_{\beta} A^{\delta}) - \operatorname{Ric}_{\beta\delta} (\mathbf{F}^{\delta}_{\alpha} + \nabla_{\alpha} A^{\delta}) \\ &= \nabla_{\beta} (\operatorname{Ric}_{\alpha\delta} A^{\delta}) - \nabla_{\alpha} (\operatorname{Ric}_{\beta\delta} A^{\delta}) + R_{\beta\alpha\sigma\delta} \mathbf{F}^{\sigma\delta} + \operatorname{Ric}_{\alpha\delta} \mathbf{F}^{\delta}_{\beta} - \operatorname{Ric}_{\beta\delta} \mathbf{F}^{\delta}_{\alpha}. \end{aligned}$$

On the other hand for the second term we use the A equation in (2.36) to write

$$\begin{aligned} II &= \nabla_\alpha[\widetilde{\text{Ric}}_{\beta\sigma} A^\sigma] - \nabla_\beta[\widetilde{\text{Ric}}_{\alpha\sigma} A^\sigma] \\ &\quad + \nabla_\alpha \nabla^\gamma \text{Im}(\lambda_{\gamma\sigma} \bar{\lambda}^\sigma_\beta) - \nabla_\beta \nabla^\gamma \text{Im}(\lambda_{\gamma\sigma} \bar{\lambda}^\sigma_\alpha) - \nabla_\gamma \nabla^\gamma \text{Im}(\lambda_{\alpha\sigma} \bar{\lambda}^\sigma_\beta) \\ &:= II_1 + II_2. \end{aligned}$$

The first term II_1 combines directly with the first two terms in I . For the second we commute

$$\begin{aligned} II_2 &= R_{\alpha\gamma\gamma\delta} \tilde{\mathbf{F}}_\beta^\delta + R_{\alpha\gamma\beta\delta} \tilde{\mathbf{F}}_\gamma^\delta - R_{\beta\gamma\gamma\delta} \tilde{\mathbf{F}}_\alpha^\delta - R_{\beta\gamma\alpha\delta} \tilde{\mathbf{F}}_\gamma^\delta + \\ &\quad + \nabla^\gamma (\nabla_\alpha \text{Im}(\lambda_{\gamma\sigma} \bar{\lambda}^\sigma_\beta) - \nabla_\beta \text{Im}(\lambda_{\gamma\sigma} \bar{\lambda}^\sigma_\alpha) - \nabla_\gamma \text{Im}(\lambda_{\alpha\sigma} \bar{\lambda}^\sigma_\beta)) \\ &= -\text{Ric}_{\alpha\delta} \tilde{\mathbf{F}}_\beta^\delta + \text{Ric}_{\beta\delta} \tilde{\mathbf{F}}_\alpha^\delta - R_{\beta\alpha\sigma\delta} \tilde{\mathbf{F}}^{\sigma\delta} \\ &\quad + \nabla^\gamma \text{Im}(\lambda_\gamma^\sigma (\overline{\nabla_\alpha^A C_{\sigma\beta}^2} - \overline{\nabla_\beta^A C_{\sigma\alpha}^2}) + \nabla_\gamma^A \lambda_\beta^\sigma \bar{C}^2_{\alpha\sigma}). \end{aligned}$$

Summing up the expressions for I and II we obtain (4.8). \square

The equation for C^4 . This has the form

$$(4.9) \quad \Delta_g C^4 = -\nabla^\sigma (C_{\mu\sigma}^6 A^\mu) - \frac{1}{2} [\nabla^\alpha, \nabla^\gamma] C_{\gamma\alpha}^3 - \frac{1}{2} \nabla^\gamma \nabla^\alpha \text{Im}(C_{\sigma\gamma}^2 \bar{\lambda}_\alpha^\sigma + \lambda_\gamma^\sigma \bar{C}^2_{\alpha\sigma}).$$

To prove it we commute Δ_g with ∇^α

$$\begin{aligned} \Delta_g \nabla^\alpha A_\alpha &= \nabla_\sigma [\nabla^\sigma, \nabla^\alpha] A_\alpha + [\nabla_\sigma, \nabla^\alpha] \nabla^\sigma A_\alpha + \nabla^\alpha \Delta_g A_\alpha \\ &= -\nabla^\sigma (\text{Ric}_{\mu\sigma} A^\mu) + \frac{1}{2} [\nabla^\sigma, \nabla^\alpha] \mathbf{F}_{\sigma\alpha} + \nabla^\alpha (\widetilde{\text{Ric}}_{\alpha\sigma} A^\sigma) + \nabla^\alpha \nabla^\gamma \tilde{\mathbf{F}}_{\alpha\gamma} \end{aligned}$$

In the last term we can symmetrize in α and γ , and the desired equation (4.9) follows. \square

The equation for C^5 . Here we compute

$$(4.10) \quad \Delta_g C_\beta^5 = -[\nabla^\alpha, \nabla_\beta] C_\alpha^5 - \text{Re}(\nabla_\beta (C^1 \bar{\psi}) - 2\bar{\lambda}^{\alpha\sigma} \nabla_\alpha^A C_{\sigma\beta}^2 + \nabla_\beta (\lambda^{\alpha\sigma} C_{\sigma\alpha}^2)).$$

We can rewrite the g equation (2.22) as

$$\text{Ric}_{\alpha\beta} = \widetilde{\text{Ric}}_{\alpha\beta} + \frac{1}{2} (\nabla_\alpha C_\beta^5 + \nabla_\beta C_\alpha^5)$$

which by contraction yields

$$R = \tilde{R} + \nabla^\alpha C_\alpha^5.$$

To get to $\Delta_g C_\beta^5$, by the above two equalities we write

$$\begin{aligned} \frac{1}{2} \Delta_g C_\beta^5 &= \nabla^\alpha (\text{Ric}_{\alpha\beta} - \widetilde{\text{Ric}}_{\alpha\beta}) - \frac{1}{2} [\nabla^\alpha, \nabla_\beta] C_\alpha^5 - \frac{1}{2} \nabla_\beta (R - \tilde{R}) \\ &= (\nabla^\alpha R_{\alpha\beta} - \frac{1}{2} \nabla_\beta R) - \frac{1}{2} [\nabla^\alpha, \nabla_\beta] C_\alpha^5 - (\nabla^\alpha \tilde{R}_{\alpha\beta} - \frac{1}{2} \nabla_\beta \tilde{R}). \end{aligned}$$

The first term drops by twice contracted Bianchi,

$$g^{\mu\nu} g^{\gamma\alpha} (\nabla_\gamma R_{\nu\beta\mu\alpha} + \nabla_\nu R_{\beta\gamma\mu\alpha} + \nabla_\beta R_{\gamma\nu\mu\alpha}) = 0,$$

and the last one is quadratic in λ and yields C^1 and C^2 terms,

$$(\nabla^\alpha \tilde{R}_{\alpha\beta} - \frac{1}{2} \nabla_\beta \tilde{R}) = \text{Re}(\frac{1}{2} \nabla_\beta (C^1 \bar{\psi}) - \bar{\lambda}^{\alpha\sigma} \nabla_\alpha^A C_{\sigma\beta}^2 + \frac{1}{2} \nabla_\beta (\lambda^{\alpha\sigma} C_{\sigma\alpha}^2)).$$

This completes the derivation of (4.10). \square

The equation for C^6 . This has the form

$$(4.11) \quad C_{\gamma\sigma}^6 = \frac{1}{2}(\nabla_\gamma C_\sigma^5 + \nabla_\sigma C_\gamma^5).$$

Indeed, by the g -equation in (2.36) and its proof, we recover the Ricci curvature

$$\operatorname{Re}(\lambda_{\gamma\sigma}\bar{\psi} - \lambda_{\gamma\alpha}\bar{\lambda}_\sigma^\alpha) = \operatorname{Ric}_{\gamma\sigma} - \frac{1}{2}(\partial_\gamma C_\sigma^5 + \partial_\sigma C_\gamma^5) + \Gamma_{\gamma\sigma}^\nu C_\nu^5.$$

This implies the relation (4.11) immediately. \square

The equation for C^7 . By the second Bianchi identities of Riemannian curvature and the following equality

$$\nabla_\delta \operatorname{Re}(\lambda_{\gamma\beta}\bar{\lambda}_{\sigma\alpha} - \lambda_{\gamma\alpha}\bar{\lambda}_{\sigma\beta}) + \nabla_\sigma \operatorname{Re}(\lambda_{\delta\beta}\bar{\lambda}_{\gamma\alpha} - \lambda_{\delta\alpha}\bar{\lambda}_{\gamma\beta}) + \nabla_\gamma \operatorname{Re}(\lambda_{\sigma\beta}\bar{\lambda}_{\delta\alpha} - \lambda_{\sigma\alpha}\bar{\lambda}_{\delta\beta}) = 0,$$

we have the counterpart of the second Bianchi identities

$$\nabla_\delta C_{\sigma\gamma\alpha\beta}^7 + \nabla_\sigma C_{\gamma\delta\alpha\beta}^7 + \nabla_\gamma C_{\delta\sigma\alpha\beta}^7 = 0,$$

which combine with the algebraic symmetries of the same tensor to yield an elliptic system for C^7 . Precisely, using the above relation we have

$$\nabla^\sigma C_{\sigma\gamma\alpha\beta}^7 = \nabla_\alpha C_{\gamma\beta}^6 - \nabla_\beta C_{\gamma\alpha}^6 + \nabla(\lambda C^1 + \lambda C^2),$$

which combined with the previous one yields the desired elliptic system, with C^6 viewed as a source term. \square

b) Assume that \tilde{s}_k and s_k are admissible frequency envelopes for $\delta\mathcal{S} \in \mathcal{H}^\sigma$ and $\mathcal{S} \in \mathcal{H}^s$, respectively. In view of the bound (4.1) and of the smallness of $\|\psi\|_{H^s}$, it suffices to prove the difference or linearized estimate

$$(4.12) \quad \|S_k \delta\mathcal{S}\|_{\mathcal{H}^\sigma} \lesssim \|S_k \delta\psi\|_{H^\sigma} + (\tilde{s}_k \|\psi\|_{H^s} + s_k \|\delta\mathcal{S}\|_{\mathcal{H}^\sigma})(1 + \|\psi\|_{H^s})^N.$$

If this is true, then the bound (4.2) follows. Thus, by the definition of frequency envelope (3.3), (4.2) and the smallness of $\psi \in H^s$, the bound (4.12) with operator $\delta = Id$ and $\sigma = s$ also implies the bound (4.3).

As an intermediate step in the proof of (4.2), we collect in the next Lemma several bilinear estimates. The proof of this Lemma is standard by Littlewood-Paley decompositions and Bernstein inequality.

Lemma 4.2. *Let $d/2 - 3 < \sigma \leq s$, $d \geq 3$, then we have*

$$\begin{aligned} \|\nabla\delta(\tilde{h}h)\|_{H^\sigma} &\lesssim \|\nabla\delta\tilde{h}\|_{H^\sigma}\|\nabla h\|_{H^s} + \|\nabla\tilde{h}\|_{H^s}\|\nabla\delta h\|_{H^\sigma}, \\ \|\delta(\lambda h)\|_{H^\sigma} &\lesssim \|\delta\lambda\|_{H^\sigma}\|\nabla h\|_{H^s} + \|\lambda\|_{H^s}\|\nabla\delta h\|_{H^\sigma}, \\ \|\nabla\delta(Ah)\|_{H^{\sigma-1}} &\lesssim \|\nabla\delta A\|_{H^{\sigma-1}}\|\nabla h\|_{H^s} + \|\nabla A\|_{H^s}\|\nabla\delta h\|_{H^\sigma}. \end{aligned}$$

Now we turn our attention to the proof of (4.12). Here we first prove the estimates for $\delta\lambda$. By λ -equations in (4.5) it suffices to consider the following form

$$\begin{aligned} \partial_\alpha \delta\lambda_{\alpha\beta} &= \partial_\beta \delta\psi + \delta A\psi + A\delta\psi + \delta h\nabla\lambda + h\nabla\delta\lambda + \nabla\delta h\lambda + \nabla h\delta\lambda, \\ \partial_\alpha \delta\lambda_{\beta\gamma} - \partial_\beta \delta\lambda_{\alpha\gamma} &= \delta A\lambda + A\delta\lambda + \nabla\delta h\lambda + \nabla h\delta\lambda. \end{aligned}$$

By the relation

$$(4.13) \quad \widehat{\lambda}(\xi) = |\xi|^{-2}(\widehat{\lambda} \cdot \xi)\xi + |\xi|^{-2}(\widehat{\lambda}\xi^\top - \xi\widehat{\lambda}^\top) \cdot \xi,$$

we obtain

$$\begin{aligned} \|S_k \delta \lambda\|_{H^\sigma} &\lesssim \|S_k \delta \psi\|_{H^\sigma} + \| |D|^{-1} S_k [\delta A(\lambda + \psi) + A(\delta \lambda + \delta \psi) + \delta h \nabla \lambda + h \nabla \delta \lambda \\ &\quad + \nabla \delta h \lambda + \nabla h \delta \lambda] \|_{H^\sigma} \\ &\lesssim \|S_k \delta \psi\|_{H^\sigma} + \tilde{s}_k \|\mathcal{S}\|_{\mathcal{H}^s} + s_k \|\delta \mathcal{S}\|_{\mathcal{H}^\sigma}. \end{aligned}$$

Next we provide the estimate for δA ; the other estimates can be proved similarly. By A -equation in (4.5) and Lemma 4.2, it suffices to consider the following form

$$\begin{aligned} \Delta \delta A &= \delta h \nabla^2 A + h \nabla^2 \delta A + \nabla \delta h \nabla A + \nabla h \nabla \delta A + \nabla \delta h \nabla h A + (\nabla h)^2 \delta A \\ &\quad + \delta \lambda \lambda (A + \nabla h) + \lambda^2 (\delta A + \nabla \delta h) + \nabla \lambda \delta \lambda + \lambda \nabla \delta \lambda. \end{aligned}$$

Using Littlewood-Paley trichotomy and Bernstein inequality, we bound all the nonlinearities except $\nabla \lambda \delta \lambda$ and $\lambda \nabla \delta \lambda$ by

$$\begin{aligned} &\| |D|^{-1} S_k (\delta h \nabla^2 A + h \nabla^2 \delta A + \nabla \delta h \nabla A + \nabla h \nabla \delta A + \nabla \delta h \nabla h A + (\nabla h)^2 \delta A) \|_{H^\sigma} \\ &\quad + \| |D|^{-1} S_k (\delta \lambda \lambda (A + \nabla h) + \lambda^2 (\delta A + \nabla \delta h)) \|_{H^\sigma} \\ &\lesssim (\tilde{s}_k \|\mathcal{S}\|_{\mathcal{H}^s} + s_k \|\delta \mathcal{S}\|_{\mathcal{H}^\sigma}) (1 + \|\mathcal{S}\|_{\mathcal{H}^s}). \end{aligned}$$

For the remainder terms, we can also bound their low-frequency part by

$$\| |D|^{-1} S_0 (\nabla \lambda \delta \lambda + \lambda \nabla \delta \lambda) \|_{L^2} \lesssim \|S_0 (\nabla \lambda \delta \lambda + \lambda \nabla \delta \lambda) \|_{L^1} \lesssim \tilde{s}_0 \|\lambda\|_{H^s},$$

and bound their high-frequency part S_k for $k > 0$ by

$$\| |D|^{-1} S_k (\nabla \lambda \delta \lambda + \lambda \nabla \delta \lambda) \|_{H^\sigma} \lesssim \tilde{s}_k \|\lambda\|_{H^s} + s_k \|\delta \lambda\|_{H^\sigma}.$$

This completes the proof of (4.2).

c) Using the similar argument to b), we have

$$\|D^2 \mathcal{S}(\delta_1 \psi, \delta_2 \psi)\|_{\mathcal{H}^\sigma} \lesssim \|\delta_1 \mathcal{S}\|_{\mathcal{H}^\sigma} \|\delta_2 \mathcal{S}\|_{\mathcal{H}^s} (1 + \|\psi\|_{H^s})^N,$$

and

$$\|D^2 \mathcal{S}(\delta_1 \psi, \delta_2 \psi)\|_{\mathcal{H}^\sigma} \lesssim \|\delta_1 \mathcal{S}\|_{\mathcal{H}^s} \|\delta_2 \mathcal{S}\|_{\mathcal{H}^\sigma} (1 + \|\psi\|_{H^s})^N.$$

Then by the smallness of $\psi \in H^s$, (4.2) and interpolation, the above two bounds imply

$$\|D^2 \mathcal{S}(\delta_1 \psi, \delta_2 \psi)\|_{\mathcal{H}^\sigma} \lesssim \|\delta_1 \psi\|_{H^{\sigma_1}} \|\delta_2 \psi\|_{H^{\sigma_2}}.$$

This completes the proof of (4.4). \square

Next we establish bounds for the above solutions in space-time local energy spaces:

Theorem 4.3. *a) Assume that ψ is small in $l^2 \mathbf{X}^s$ for $s > d/2$, $d \geq 4$. Then the solution (λ, h, A, V, B) for the elliptic system (2.36) given by Theorem 4.1 belongs to \mathcal{E}^s and satisfies the bounds*

$$(4.14) \quad \|\mathcal{S}\|_{\mathcal{E}^s} \lesssim \|\psi\|_{l^2 \mathbf{X}^s},$$

with Lipschitz dependence on the initial data in these topologies. Moreover, assume that p_k is an admissible frequency envelope for $\psi \in l^2 \mathbf{X}^s$, we have the frequency envelope version

$$(4.15) \quad \|\mathcal{S}_k\|_{\mathcal{E}^s} \lesssim p_k.$$

b) In addition, for the linearization of the elliptic system (2.36) we have the bounds

$$(4.16) \quad \|\delta\mathcal{S}\|_{\mathcal{E}^\sigma} \lesssim \|\delta\psi\|_{L^2\mathbf{X}^\sigma},$$

for $\sigma \in (d/2 - 1, s]$.

Proof of Theorem 4.3. For the elliptic system (4.5), we will prove the bound for differences $\delta\mathcal{S}$

$$(4.17) \quad \|\delta\mathcal{S}\|_{\mathcal{E}^\sigma} \lesssim \|\delta\psi\|_{L^2\mathbf{X}^\sigma} + \|\delta\mathcal{S}\|_{\mathcal{E}^\sigma} \|\mathcal{S}\|_{\mathcal{E}^s} (1 + \|\mathcal{S}\|_{\mathcal{E}^s})^N.$$

If this is true, by a continuity argument the bounds (4.14) and (4.16) follow.

Assume that \tilde{s}_k and s_k are admissible frequency envelopes for $\delta\mathcal{S} \in \mathcal{E}^\sigma$ and $\mathcal{S} \in \mathcal{E}^s$, respectively. We can separate the bound (4.17) into two parts, namely

$$\|\partial_t \delta\mathcal{S}\|_{L^2\mathcal{H}^{\sigma-2}} \lesssim \|\delta\psi\|_{L^2\mathbf{X}^\sigma} (1 + \|\partial_t \psi\|_{L^2H^{s-2}})$$

respectively

$$(4.18) \quad \|S_k \delta\mathcal{S}\|_{\mathcal{E}^\sigma} \lesssim p_k + (\tilde{s}_k \|\mathcal{S}\|_{\mathcal{E}^s} + s_k \|\delta\mathcal{S}\|_{\mathcal{E}^\sigma}) (1 + \|\mathcal{S}\|_{\mathcal{E}^s})^N.$$

Here one can think of the first bound as a fixed time bound for the linearization of the elliptic system (2.36), square integrated in time. As such, this is a direct consequence of the bound (4.2) with argument $\partial_t \delta\psi$ and regularity index $\sigma - 2$, and the bound (4.4) with $\delta_1 = \partial_t, \delta_2 = \delta, \sigma_1 = s - 2, \sigma_2 = \sigma$ in Theorem 4.1. So it remains to prove (4.18).

If the bound (4.18) holds, then by the bound (4.3) with $\delta = \partial_t, \sigma = s - 2$ and (4.18) with $\delta = Id, \sigma = s$, the bound (4.15) follows.

As an intermediate step in the proof of (4.18), we collect in the next Lemma several bilinear estimates and equivalent relations.

Lemma 4.4 (Bilinear estimates). *Let $s > d/2$, $0 < \sigma \leq s$, $d \geq 4$, assume that $h \in \mathbf{Y}^s$, then we have*

$$(4.19) \quad \|\tilde{h}h\|_{Y^\sigma} \lesssim \|\tilde{h}\|_{\mathbf{Y}^\sigma} \|h\|_{\mathbf{Y}^s},$$

$$(4.20) \quad \|\lambda h\|_{Z^{0,\sigma}} \lesssim \|\lambda\|_{\mathbf{Z}^{0,\sigma}} \|h\|_{\mathbf{Y}^s},$$

$$(4.21) \quad \|(Ah)\|_{Z^{1,\sigma}} \lesssim \|A\|_{\mathbf{Z}^{1,\sigma}} \|h\|_{\mathbf{Y}^s}.$$

As consequences of these bounds, for $h^{\alpha\beta} = g^{\alpha\beta} - \delta^{\alpha\beta}$, $h_{\alpha\beta} = g_{\alpha\beta} - \delta_{\alpha\beta}$, $\lambda^{\alpha\beta} = g^{\alpha\gamma} \lambda_\gamma^\beta$, $\lambda_\gamma^\beta = g^{\beta\nu} \lambda_{\nu\gamma}$, $V^\alpha = g^{\alpha\beta} V_\beta$ and $A^\alpha = g^{\alpha\beta} A_\beta$, assume that $\|h_{\alpha\beta}\|_{\mathbf{Y}^{\sigma+1}} \ll 1$, we have

$$\begin{aligned} \|h_{\alpha\beta}\|_{\mathbf{Y}^{\sigma+1}} &\approx \|\tilde{h}^{\alpha\beta}\|_{\mathbf{Y}^{\sigma+1}}, \\ \|\lambda^{\alpha\beta}\|_{\mathbf{Z}^{0,\sigma}} &\approx \|\lambda_\alpha^\beta\|_{\mathbf{Z}^{0,\sigma}} \approx \|\lambda_{\alpha\beta}\|_{\mathbf{Z}^{0,\sigma}}, \\ \|V_\alpha\|_{\mathbf{Z}^{1,\sigma}} &\approx \|V^\alpha\|_{\mathbf{Z}^{1,\sigma}}, \\ \|A_\alpha\|_{\mathbf{Z}^{1,\sigma}} &\approx \|A^\alpha\|_{\mathbf{Z}^{1,\sigma}}. \end{aligned}$$

Proof of Lemma 4.4. We do this in several steps:

Proof of the bound (4.19). First, we consider the Y -norm estimates. For the high-low interaction, for any decomposition $P_j \tilde{h} = \sum_{l \geq |j|} \tilde{h}_{j,l}$, we have

$$\left\| \sum_{l \geq |j|} (\tilde{h}_{j,l} h_{\leq j}) \right\|_{Y_j} \lesssim \sum_{l \geq |j|} 2^{l-|j|} \|(\tilde{h}_{j,l} h_{\leq j})\|_{l^1 L^\infty L^2} \lesssim \sum_{l \geq |j|} 2^{l-|j|} \|\tilde{h}_{j,l}\|_{l^1 L^\infty L^2} \|h_{\leq j}\|_{L^\infty L^\infty}.$$

Taking the infimum over the decomposition of \tilde{h}_j yields

$$\left\| \sum_{l \geq |j|} (\tilde{h}_{j,l} h_{\leq j}) \right\|_{Y_j} \lesssim \|P_j \tilde{h}\|_{Y_j} \|h\|_{Z^{1,s}},$$

which is acceptable. Similarly, for the low-high interaction, we have

$$\begin{aligned} \left\| \sum_{l \geq |j|} (P_{\leq j} \tilde{h} h_{j,l}) \right\|_{Y^{\frac{d}{2}-1-\delta,\sigma}} &\lesssim 2^{(\frac{d}{2}-1-\delta)j^- + \sigma j^+} \sum_{k \leq j} 2^{dk/2} \|P_k \tilde{h}\|_{L^\infty L^2} \|P_j h\|_{Y_j} \\ &\lesssim \|\nabla \tilde{h}\|_{L^\infty H^{\sigma-1}} \|P_j h\|_{Y^{d/2-1-\delta,s}}, \end{aligned}$$

which is acceptable.

Next, for the high-high interaction, when $j < 0$ we rewrite it as

$$\sum_{j < j_1 < -j} P_j(P_{j_1} \tilde{h} P_{j_1} h) + \sum_{-j \leq j_1} P_j(P_{j_1} \tilde{h} P_{j_1} h).$$

Then we bound the first term by

$$\begin{aligned} &2^{(d/2-1-\delta)j} \left\| \sum_{j < j_1 < -j} P_j(P_{j_1} \tilde{h} P_{j_1} h) \right\|_{Y_j} \\ &\lesssim 2^{(d-1-\delta)j} \sum_{j < j_1 < -j} \|P_{j_1} \tilde{h} P_{j_1} h\|_{l_{|j_1|}^1 L^\infty L^1} \\ &\lesssim 2^{(d-1-\delta)j} \sum_{j < j_1 < -j} \|P_{j_1} \tilde{h}\|_{l_{|j_1|}^2 L^\infty L^2} \|P_{j_1} h\|_{l_{|j_1|}^2 L^\infty L^2} \\ &\lesssim 2^{(d-3-2\delta)j} \|\nabla \tilde{h}_{\leq 0}\|_{l_0^2 L^\infty L^2} \|\nabla h_{\leq 0}\|_{l_0^2 L^\infty L^2} + 2^{(d-1-\delta)j} \|\tilde{h}\|_{Z^{1,0}} \|h\|_{Z^{1,0}}. \end{aligned}$$

We bound the second term by

$$\begin{aligned} 2^{(d/2-1-\delta)j} \left\| \sum_{-j \leq j_1} P_j(P_{j_1} h P_{j_1} h) \right\|_{Y_j} &\lesssim \sum_{-j \leq j_1} 2^{(d-\delta)j+j_1} \|(P_{j_1} \tilde{h} P_{j_1} h)\|_{l_{j_1}^1 L^\infty L^1} \\ &\lesssim \sum_{-j \leq j_1} 2^{(d-\delta)j+j_1} \|P_{j_1} \tilde{h}\|_{l_{j_1}^2 L^\infty L^2} \|P_{j_1} h\|_{l_{j_1}^2 L^\infty L^2} \\ &\lesssim 2^{(d-\delta)j} \|\tilde{h}\|_{Z^{1,0}} \|h\|_{Z^{1,1}}. \end{aligned}$$

When $j \geq 0$, we have

$$\begin{aligned} &2^{\sigma j} \left\| \sum_{j_1 > j} P_j(P_{j_1} \tilde{h} P_{j_1} h) \right\|_{Y_j} \\ &\lesssim \sum_{j_1 > j} 2^{(\sigma-1+d/2)j+j_1} \|(P_{j_1} \tilde{h} P_{j_1} h)_j\|_{l_{j_1}^1 L^\infty L^1} \\ &\lesssim \sum_{j_1 > j} 2^{(\sigma-1+d/2)(j-j_1)} \|P_{j_1} \tilde{h}\|_{Z^{1,\sigma}} \|P_{j_1} h\|_{Z^{1,s}}, \end{aligned}$$

which is acceptable.

Secondly, we consider the $Z^{1,\sigma+1}$ -norm estimates. For the low-frequency part, we have

$$\begin{aligned} \|\nabla(\tilde{h}h)_{\leq 0}\|_{l_0^2 L^\infty L^2} &\lesssim \|\nabla \tilde{h}_{\leq 0}\|_{l_0^2 L^\infty L^2} \|h_{\leq 0}\|_{L^\infty L^\infty} + \sum_{j>0} \|(\tilde{h}_j h_j)_{\leq 0}\|_{l_0^2 L^\infty L^1} \\ &\lesssim \|\tilde{h}\|_{Z^{1,0}} \|h\|_{Z^{1,s}}. \end{aligned}$$

For the high frequency part, by Littlewood-Paley dichotomy, we have

$$\begin{aligned} &2^{\sigma j} \|(\tilde{h}h)_j\|_{l_j^2 L^\infty L^2} \\ &\lesssim 2^{\sigma j} \|\tilde{h}_j\|_{l_j^2 L^\infty L^2} \|h_{\leq j}\|_{L^\infty L^\infty} + 2^{\sigma j} \|\tilde{h}_{\leq j}\|_{L^\infty L^\infty} \|h_j\|_{l_j^2 L^\infty L^2} + \sum_{l \geq j} 2^{(\sigma+d/2)j} \|(\tilde{h}_l h_l)_j\|_{l_j^2 L^\infty L^1} \\ &\lesssim \|\tilde{h}_j\|_{Z^{1,\sigma}} \|\nabla h\|_{L^\infty H^{s-1}} + \|\tilde{h}\|_{Z^{1,\sigma}} \|h_j\|_{Z^{1,s}} + \sum_{l \geq j} 2^{\sigma(j-l)} 2^{(\sigma+d/2)l} \|\tilde{h}_l\|_{l_l^2 L^\infty L^2} \|h_l\|_{L^\infty L^2}, \end{aligned}$$

which is acceptable. This completes the proof of (4.19).

Proof of the bound (4.20). First we consider the $Z^{\delta,\sigma}$ -norm estimates. For the low-frequency part we have

$$\begin{aligned} \|(h\lambda)_{\leq 0}\|_{l_0^2 L^\infty L^2} &\lesssim \|h_{\leq 0}\|_{L^\infty L^\infty} \|\lambda_{\leq 0}\|_{l_0^2 L^\infty L^2} + \sum_{j>0} 2^{dj/2} \|h_j\|_{L^\infty L^2} \|\lambda_j\|_{l_j^2 L^\infty L^2} \\ &\lesssim \|h\|_{Z^{1,s}} \|\lambda\|_{Z^{0,\sigma}}. \end{aligned}$$

For the high-frequency part, by the Littlewood-Paley dichotomy, we have

$$\begin{aligned} 2^{\sigma j} \|(\lambda h)_j\|_{l_j^2 L^\infty L^2} &\lesssim \sum_{l < j} 2^{\sigma j + dl/2} \|\lambda_l\|_{L^\infty L^2} \|h_j\|_{l_j^2 L^\infty L^2} + 2^{\sigma j} \|\lambda_j\|_{l_j^2 L^\infty L^2} \|h\|_{Z^{1,s}} \\ &\quad + \sum_{l > j} 2^{\sigma(j-l)} 2^{(\sigma+d/2)l} \|h_l\|_{l_l^2 L^\infty L^2} \|\lambda_l\|_{L^\infty L^2}, \end{aligned}$$

which implies

$$\left(\sum_{j>0} 2^{2\sigma j} \|(\lambda h)_j\|_{l_j^2 L^\infty L^2}^2 \right)^{1/2} \lesssim \|h\|_{Z^{1,s}} \|\lambda\|_{Z^{\delta,\sigma}}.$$

This completes the proof of (4.20).

Proof of the bound (4.21). For the low-frequency part, by Bernstein's inequality we have

$$\begin{aligned} \|\nabla(Ah)_{\leq 0}\|_{l_0^2 L^\infty L^2} &\lesssim \|\nabla(A_{\leq 0} h_{\leq 0})\|_{l_0^2 L^\infty L^2} + \sum_{j>0} \|\nabla(A_j h_j)_{\leq 0}\|_{l_0^2 L^\infty L^2} \\ &\lesssim \|\nabla A_{\leq 0}\|_{l_0^2 L^\infty L^2} \|\nabla h_{\leq 0}\|_{L^\infty L^2} + \|\nabla A_{\leq 0}\|_{L^\infty L^2} \|\nabla h_{\leq 0}\|_{l_0^2 L^\infty L^2} \\ &\quad + \sum_{j>0} 2^{dj/2} \|A_j\|_{l_j^2 L^\infty L^2} \|h_j\|_{L^\infty L^2} \\ &\lesssim \|A\|_{Z^{1,0}} \|h\|_{Z^{1,s}}. \end{aligned}$$

For the high-frequency part, by Littlewood-Paley dichotomy we bound the high-low and low-high interactions by

$$\begin{aligned}
& 2^{\sigma k} \|S_k(A_k h_{<k} + A_{<k} h_k)\|_{l_k^2 L^\infty L^2} \\
& \lesssim 2^{\sigma k} (\|A_k\|_{l_k^2 L^\infty L^2} \|h_{<k}\|_{L^\infty L^\infty} + \|A_{<k}\|_{L^\infty L^\infty} \|h_k\|_{l_k^2 L^\infty L^2}) \\
& \lesssim \|A_k\|_{Z^{1,\sigma}} \|h\|_{Z^{1,s}} + \|A\|_{Z^{1,\sigma}} \|h_k\|_{Z^{1,s}},
\end{aligned}$$

which is acceptable. We bound the high-high interaction by

$$\begin{aligned}
& 2^{\sigma k} \sum_{j>k} \|S_k(A_j h_j)\|_{l_k^2 L^\infty L^2} \\
& \lesssim \sum_{j>k} 2^{(\sigma+d/2)k} \|A_j h_j\|_{l_k^2 L^\infty L^1} \\
& \lesssim \sum_{j>k} 2^{\sigma(k-j)} 2^{(\sigma+d/2)j} \|A_j\|_{l_j^2 L^\infty L^2} \|h_j\|_{L^\infty L^2},
\end{aligned}$$

which is also acceptable. Hence, we conclude the proof of the bound (4.21). \square

We now turn our attention to the proof of (4.18).

Step 1. Proof of the elliptic estimates for λ equations. By the λ -equations and Proposition 4.4, it suffices to consider the following simplified form of the equations:

$$\begin{aligned}
\partial_\alpha \delta \lambda_{\alpha\beta} &= \partial_\beta \delta \psi + \delta A \psi + A \delta \psi + \delta h \nabla \lambda + h \nabla \delta \lambda + \nabla \delta h \lambda + \nabla h \delta \lambda, \\
\partial_\alpha \delta \lambda_{\beta\gamma} - \partial_\beta \delta \lambda_{\alpha\gamma} &= \delta A \lambda + A \delta \lambda + \nabla \delta h \lambda + \nabla h \delta \lambda.
\end{aligned}$$

By the relation (4.13) we have for any $k > 0$

$$\begin{aligned}
\|S_k \delta \lambda\|_{Z^{0,\sigma}} &\lesssim \|S_k \mathcal{R} \delta \psi\|_{Z^{0,\sigma}} + \|S_k |D|^{-1} [\delta A(\psi + \lambda) + A(\delta \psi + \delta \lambda) \\
&\quad + \delta h \nabla \lambda + h \nabla \delta \lambda + \nabla \delta h \lambda + \nabla h \delta \lambda]\|_{Z^{0,\sigma}} \\
&\lesssim \tilde{p}_k + \tilde{s}_k \|\mathcal{S}\|_{\mathcal{E}^s} + s_k \|\delta \mathcal{S}\|_{\mathcal{E}^\sigma}.
\end{aligned}$$

In order to bound the low frequency part $k = 0$, we use the relation

$$(4.22) \quad f(t) = f(0) + \int_0^t \partial_s f(s) ds.$$

Then we have

$$\|f\|_{l_0^2 L^\infty L^2} \lesssim \|f(0)\|_{L^2} + \|\partial_t f\|_{L^2 L^2}.$$

Using this idea, by Sobolev embeddings we have

$$\begin{aligned}
\|S_0\delta\lambda\|_{l_0^2L^\infty L^2} &\lesssim \|S_0\mathcal{R}\delta\psi\|_{l_0^2L^\infty L^2} + \|S_0|D|^{-1}[\delta A(\psi + \lambda) + A(\delta\psi + \delta\lambda) \\
&\quad + \delta h\nabla\lambda + h\nabla\delta\lambda + \nabla\delta h\lambda + \nabla h\delta\lambda]\|_{l_0^2L^\infty L^2} \\
&\lesssim \|S_0\delta\psi\|_{\mathbf{Z}^{0,\sigma}} \\
&\quad + \|S_0|D|^{-1}[\delta A(\psi + \lambda) + A(\delta\psi + \delta\lambda) + \delta h\nabla\lambda \\
&\quad + h\nabla\delta\lambda + \nabla\delta h\lambda + \nabla h\delta\lambda](0)\|_{L^2} \\
&\quad + \|S_0|D|^{-1}\partial_t[\delta A(\psi + \lambda) + A(\delta\psi + \delta\lambda) + \delta h\nabla\lambda \\
&\quad + h\nabla\delta\lambda + \nabla\delta h\lambda + \nabla h\delta\lambda]\|_{L^2L^2} \\
&\lesssim \tilde{p}_0 + \tilde{s}_0\|\mathcal{S}\|_{\mathcal{E}^s}.
\end{aligned}$$

The high frequency part is obtained by a standard Littlewood-Paley decomposition and Bernstein inequality. This gives the elliptic estimate for the $\delta\lambda$ -equation.

Step 2. Proof of the elliptic estimates for V , A and B equations. By the V , A , B -equations and Proposition 4.4, it suffices to consider the following form

$$\begin{aligned}
\Delta V &= h\nabla^2V + \nabla h\nabla V + \nabla h\nabla hV + \lambda^2(A + V + \nabla h) + \lambda\nabla\lambda, \\
\Delta A &= h\nabla^2A + \nabla h\nabla A + \nabla h\nabla hA + \lambda^2(A + \nabla h) + \nabla(\lambda^2), \\
\Delta B &= h\nabla^2B + \nabla(\lambda\nabla\lambda + (V + A)\lambda^2) + \lambda^2\nabla A + \nabla h(\lambda\nabla\lambda + (V + A)\lambda^2) \\
&\quad + \nabla V\nabla A + \nabla hV\nabla A.
\end{aligned}$$

The proofs of the three elliptic estimates for the above equations are similar, so we only prove the elliptic estimate for the linearization of A -equation in detail, i.e.

$$\begin{aligned}
\Delta\delta A &= \delta h\nabla^2A + h\nabla^2\delta A + \nabla\delta h\nabla A + \nabla h\nabla\delta A + \nabla\delta h\nabla hA + (\nabla h)^2\delta A \\
&\quad + \delta\lambda\lambda(A + \nabla h) + \lambda^2(\delta A + \nabla\delta h) + \nabla\lambda\delta\lambda + \lambda\nabla\delta\lambda.
\end{aligned}$$

We bound all the nonlinearities except $\nabla\lambda\delta\lambda$ and $\lambda\nabla\delta\lambda$ by

$$\begin{aligned}
&\| |D|^{-2}S_k(\delta h\nabla^2A + h\nabla^2\delta A + \nabla\delta h\nabla A + \nabla h\nabla\delta A + \nabla\delta h\nabla hA + (\nabla h)^2\delta A) \|_{Z^{1,\sigma+1}} \\
&\quad + \| |D|^{-2}S_k(\delta\lambda\lambda(A + \nabla h) + \lambda^2(\delta A + \nabla\delta h)) \|_{Z^{1,\sigma+1}} \\
&\lesssim (\tilde{s}_k\|\mathcal{S}\|_{\mathcal{E}^s} + s_k\|\delta\mathcal{S}\|_{\mathcal{E}^\sigma})(1 + \|\mathcal{S}\|_{\mathcal{E}^s})^N,
\end{aligned}$$

for $\sigma \in (d/2 - 1, s]$. All terms are estimated in a similar fashion, so we only bound the first term $\delta h\nabla^2A$.

For the low-frequency part we use the relation (4.22) to bound the second term $\delta h\nabla^2A$ by

$$\begin{aligned}
&\|\nabla^{-1}(\delta h\nabla^2A)_{\leq 0}\|_{l_0^2L^\infty L^2} \\
&\lesssim \|\nabla^{-1}(\delta h\nabla^2A)_{\leq 0}(0)\|_{L^2} + \|\nabla^{-1}\partial_t(\delta h\nabla^2A)_{\leq 0}\|_{L^2L^2} \\
&\lesssim \|(\delta h\nabla^2A)_{\leq 0}(0)\|_{L^{2d/(d+2)}} + \|\partial_t(\delta h\nabla^2A)_{\leq 0}\|_{L^2L^{2d/(d+2)}} \\
&\lesssim \|\delta h\|_{Z^{1,1}}\|A\|_{Z^{1,1}} + \|\nabla\partial_t\delta h\|_{L^2H^{\sigma-1}}\|A\|_{Z^{1,\sigma+1}} + \|\delta h\|_{Z^{1,\sigma+2}}\|\nabla\partial_tA\|_{L^2H^{\sigma-3}} \\
&\lesssim \|\delta h\|_{\mathbf{Z}^{1,\sigma+2}}\|A\|_{\mathbf{Z}^{1,\sigma+1}}.
\end{aligned}$$

A minor modification of this argument also yields

$$\|\nabla^{-1}(\delta h\nabla^2A)_{\leq 0}\|_{l_0^2L^\infty L^2} \lesssim \tilde{s}_0\|\mathcal{S}\|_{\mathcal{E}^s}.$$

For the high-frequency part, by Littlewood-Paley dichotomy and Bernstein's inequality (3.1), we have

$$\begin{aligned}
& 2^{\sigma j} \||D|^{-1}(\delta h \nabla^2 A)_j\|_{l_j^2 L^\infty L^2} \\
& \lesssim 2^{(\sigma-1)j} (\|\delta h_{<j} \nabla^2 A_j\|_{l_j^2 L^\infty L^2} + \|\delta h_j \nabla^2 A_{<j}\|_{l_j^2 L^\infty L^2} + \sum_{l>j} \|\delta h_l \nabla^2 A_l\|_{l_j^2 L^\infty L^2}) \\
& \lesssim \|\delta h\|_{L^\infty L^\infty} 2^{(\sigma+1)j} \|A_j\|_{l_j^2 L^\infty L^2} + \sum_{l<j} 2^{(l-j)+\sigma j} \|\delta h_j\|_{l_j^2 L^\infty L^2} \|\nabla^{d/2+1} A_l\|_{L^\infty L^2} \\
& \quad + \sum_{l>j} 2^{(\sigma-1)j} 2^{(d/2+2)l} \|\delta h_l\|_{L^\infty L^2} \|A_l\|_{l_l^2 L^\infty L^2} \\
& \lesssim s_j \|\delta \mathcal{S}\|_{\mathcal{E}^\sigma} + \tilde{s}_j \|\mathcal{S}\|_{\mathcal{E}^s}.
\end{aligned}$$

Finally, we bound the last two terms $\nabla \lambda \delta \lambda$ and $\lambda \nabla \delta \lambda$. For low-frequency part, using $d \geq 4$ we have

$$\begin{aligned}
& \||D|^{-1}(\nabla \lambda \delta \lambda)_{\leq 0}\|_{l_0^2 L^\infty L^2} \\
& \lesssim \||D|^{-1}(\nabla \lambda \delta \lambda)_{\leq 0}(0)\|_{L^2} + \||D|^{-1} \partial_t (\nabla \lambda \delta \lambda)_{\leq 0}\|_{L^2 L^2} \\
& \lesssim \|(\nabla \lambda \delta \lambda)_{\leq 0}(0)\|_{L^1} + \|\partial_t (\nabla \lambda \delta \lambda)_{\leq 0}\|_{L^2 L^1} \\
& \lesssim \|\delta \lambda\|_{\mathbf{Z}^{0,\sigma}} \|\lambda\|_{\mathbf{Z}^{0,s}}.
\end{aligned}$$

We also obtain

$$\||D|^{-1}(\nabla \lambda \delta \lambda)_{\leq 0}\|_{l_0^2 L^\infty L^2} \lesssim \tilde{s}_0 \|\mathcal{S}\|_{\mathcal{E}^s}.$$

For the high-frequency part, we have

$$\|\Delta^{-1}(\nabla \lambda \delta \lambda)_j\|_{Z^{1,\sigma+1}} \lesssim \tilde{s}_j \|\mathcal{S}\|_{\mathcal{E}^s} + s_j \|\delta \mathcal{S}\|_{\mathcal{E}^\sigma}.$$

We can also bound the term $\lambda \nabla \delta \lambda$ similarly. This gives the elliptic estimate for δA -equation.

Step 3. Proof of the elliptic estimate for h -equation. By h -equation in (4.5) and Proposition 4.4, it suffices to consider a more general equation of the form

$$\Delta \delta h = \delta h \nabla^2 h + h \nabla^2 \delta h + \nabla \delta h \nabla h + \delta h \nabla h \nabla h + h \nabla h \nabla \delta h + \delta \lambda \lambda.$$

The proof of the $Z^{1,\sigma+2}$ bound is similar to the estimates for V, A, B equations in Step 2, hence we only bound of the $Y^{d/2-1-\delta,\sigma+2}$ -norm. We prove that the following frequency envelope version holds:

$$\|S_j \delta h\|_{Y^{d/2-1-\delta,\sigma+2}} \lesssim (\tilde{s}_j \|\mathcal{S}\|_{\mathcal{E}^s} + s_j \|\delta \mathcal{S}\|_{\mathcal{E}^\sigma}) (1 + \|\mathcal{S}\|_{\mathcal{E}^s})^N.$$

Case 1. The contribution of $\delta \lambda \lambda$. By the Littlewood-Paley dichotomy, it suffices to consider the high-low, low-high and high-high cases for any $j \in \mathbb{Z}$

$$\sum_{l<j+O(1)} P_j(P_j \delta \lambda P_l \lambda), \quad \sum_{l<j+O(1)} P_j(P_l \delta \lambda P_j \lambda), \quad \sum_{l>j+O(1)} P_j(P_l \delta \lambda P_l \lambda).$$

Case 1(a). The contribution of high-low and low-high interaction. The two cases are proved similarly, so we only consider the worst case, namely the low-high interaction. When

$j \leq 0$, by the definition of the Y_j -norm we have

$$\begin{aligned}
2^{(d/2-1-\delta)j} \|\Delta^{-1} \sum_{l < j} P_j(P_l \delta \lambda \cdot P_j \lambda)\|_{Y_j} &\lesssim 2^{(d/2-3-\delta)j} \sum_{l < j} \|(P_l \delta \lambda \cdot P_j \lambda)\|_{l_{|l|}^1 L^\infty L^2} \\
&\lesssim 2^{(d-3-\delta)j} \sum_{l < j} \|P_l \delta \lambda\|_{l_{|l|}^2 L^\infty L^2} \|P_j \lambda\|_{l_{|l|}^2 L^\infty L^2} \\
&\lesssim 2^{(d-3-3\delta)j} \| |D|^\delta \delta \lambda_{\leq 0} \|_{l_0^2 L^\infty L^2} \| |D|^\delta \lambda_{\leq 0} \|_{l_0^2 L^\infty L^2} \\
&\lesssim 2^{(d-3-3\delta)j} \tilde{s}_0 \|\mathcal{S}\|_{\mathcal{E}^s}.
\end{aligned}$$

When $j > 0$, we further divide the low-high interaction into

$$\sum_{l < j} P_j(P_l \delta \lambda \cdot P_j \lambda) = \sum_{-j \leq l < j} P_j(P_l \delta \lambda \cdot P_j \lambda) + \sum_{l < -j} P_j(P_l \delta \lambda \cdot P_j \lambda).$$

For the first term, by Bernstein's inequality we have

$$\begin{aligned}
2^{(\sigma+2)j} \|\Delta^{-1} \sum_{-j \leq l < j} P_j(P_l \delta \lambda \cdot P_j \lambda)\|_{Y_j} &\lesssim 2^{\sigma j} \sum_{-j \leq l \leq j} \|P_l \delta \lambda \cdot P_j \lambda\|_{l_j^1 L^\infty L^2} \\
&\lesssim 2^{\sigma j} \sum_{-j \leq l \leq j} 2^{dl/2} \|P_l \delta \lambda\|_{l_j^2 L^\infty L^2} \|P_j \lambda\|_{l_j^2 L^\infty L^2} \\
&\lesssim s_j \|\delta \mathcal{S}\|_{\mathcal{E}^\sigma}.
\end{aligned}$$

For the second term we have

$$\begin{aligned}
2^{(\sigma+2)j} \|\Delta^{-1} \sum_{l < -j} P_j(P_l \delta \lambda \cdot P_j \lambda)\|_{Y_j} &\lesssim 2^{\sigma j} \sum_{l < -j} 2^{|l|-j} \|P_l \delta \lambda \cdot P_j \lambda\|_{l_{|l|}^1 L^\infty L^2} \\
&\lesssim 2^{(\sigma-1)j} \sum_{l < -j} 2^{(d/2-1)l} \|P_l \delta \lambda\|_{l_{|l|}^2 L^\infty L^2} \|P_j \lambda\|_{l_{|l|}^2 L^\infty L^2} \\
&\lesssim \| |D|^\delta \delta \lambda_{\leq 0} \|_{l_0^2 L^\infty L^2} 2^{(\sigma-1)j} \|P_j \lambda\|_{l_j^2 L^\infty L^2} \\
&\lesssim s_j \|\delta \mathcal{S}\|_{\mathcal{E}^\sigma}.
\end{aligned}$$

Case 1(b). The contribution of high-high interactions. When $j < 0$, we divide this into

$$\sum_{l > j} P_j(P_l \delta \lambda \cdot P_l \lambda) = \sum_{-j \geq l > j} P_j(P_l \delta \lambda \cdot P_l \lambda) + \sum_{l > -j} P_j(P_l \delta \lambda \cdot P_l \lambda).$$

Then we bound the first term by

$$\begin{aligned}
&2^{(d/2-1-\delta)j} \|\Delta^{-1} \sum_{-j \geq l > j} P_j(P_l \delta \lambda \cdot P_l \lambda)\|_{Y_j} \\
&\lesssim 2^{(d-3-\delta)j} \sum_{-j \geq l > j} \|P_l \delta \lambda \cdot P_l \lambda\|_{l_{|l|}^1 L^\infty L^1} \\
&\lesssim 2^{(d-3-2\delta)j} \left(\sum_{0 \geq l > j} 2^{\delta l} \|\delta \lambda_{\leq 0}\|_{l_0^2 L^\infty L^2} \|\lambda_{\leq 0}\|_{l_0^2 L^\infty L^2} + \sum_{-j \geq l > 0} \|\delta \lambda_l\|_{l_l^2 L^\infty L^2} \|\lambda_l\|_{l_l^2 L^\infty L^2} \right) \\
&\lesssim 2^{(d-3-2\delta)j} \tilde{s}_0 \|\mathcal{S}\|_{\mathcal{E}^s}.
\end{aligned}$$

Using the Y_j norm we can also bound the second term by

$$\begin{aligned}
2^{(d/2-1-\delta)j} \|\Delta^{-1} \sum_{l>-j} P_j(P_l \delta \lambda \cdot P_l \lambda)\|_{Y_j} &\lesssim 2^{(d-3-\delta)j} \sum_{l>-j} 2^{l-|j|} \|(P_l \delta \lambda \cdot P_l \lambda)\|_{l^1 L^\infty L^1} \\
&\lesssim 2^{(d-2-\delta)j} \sum_{l>-j} 2^l \|P_l \delta \lambda\|_{l^2 L^\infty L^2} \|P_l \lambda\|_{l^2 L^\infty L^2} \\
&\lesssim 2^{(d-2-4\delta)j} \tilde{s}_0 \|\mathcal{S}\|_{\mathcal{E}^s}.
\end{aligned}$$

Finally, when $j > 0$, using again the Y_j norm we have

$$\begin{aligned}
2^{(\sigma+2)j} \|\Delta^{-1} \sum_{l>j} P_j(P_l \delta \lambda \cdot P_l \lambda)\|_{Y_j} &\lesssim 2^{(\sigma+d/2)j} \sum_{l>j} 2^{l-j} \|(\delta \lambda_l \cdot \lambda_l)_j\|_{l^1 L^\infty L^1} \\
&\lesssim \sum_{j_1>j} 2^{(\sigma+d/2-1)(j-l)} 2^{(\sigma+d/2)l} \|\delta \lambda_l\|_{l^2 L^\infty L^2} \|\lambda_l\|_{l^2 L^\infty L^2} \\
&\lesssim \tilde{s}_j \|\mathcal{S}\|_{\mathcal{E}^s}.
\end{aligned}$$

Case 2. The contribution of $\delta h \nabla^2 h$, $h \nabla^2 \delta h$ and $\nabla \delta h \nabla h$. It suffices to prove that

$$\|\Delta^{-1} S_j(\delta h \nabla^2 h + \nabla^2 \delta h \cdot h + \nabla \delta h \nabla h)\|_{Y^{d/2-1-\delta, \sigma+2}} \lesssim \tilde{s}_j \|\mathcal{S}\|_{\mathcal{E}^s} + s_j \|\delta \mathcal{S}\|_{\mathcal{E}^\sigma}.$$

For the high-low interactions, it suffices to consider the worst case $\nabla^2 P_j \delta h \cdot P_{\leq j} h$. For any decomposition $P_j \delta h = \sum_{l \geq |j|} \delta h_{j,l}$, we have

$$\begin{aligned}
\|\Delta^{-1} \sum_{l \geq |j|} (\nabla^2 \delta h_{j,l} P_{\leq j} h)\|_{Y_j} &\lesssim \sum_{l \geq |j|} 2^{l-|j|-2j} \|(\nabla^2 \delta h_{j,l} P_{\leq j} h)\|_{l^1 L^\infty L^2} \\
&\lesssim \sum_{l \geq |j|} 2^{l-|j|} \|\delta h_{j,l}\|_{l^1 L^\infty L^2} \|P_{\leq j} h\|_{L^\infty L^\infty}
\end{aligned}$$

Taking the infimum over the decomposition of $P_j h$ yields

$$\|\Delta^{-1} (\nabla^2 P_j \delta h P_{\leq j} h)\|_{Y_j} \lesssim \|P_j \delta h\|_{Y_j} \|P_{\leq j} h\|_{L^\infty L^\infty},$$

which is acceptable. The low-high interactions is similar and omitted.

For the high-high interaction, it suffices to estimate $\sum_{l>j} P_j(P_l \nabla \delta h P_l \nabla h)$. By Bernstein's inequality we have

$$\begin{aligned}
&2^{(d/2-1-\delta)j^- + (\sigma+2)j^+} \|\Delta^{-1} \sum_{l>j} P_j(P_l \nabla \delta h P_l \nabla h)\|_{Y_j} \\
&\lesssim 2^{(d-3-\delta)j^- + (\sigma+d/2)j^+} \sum_{l>j} \|P_j(\nabla P_l \delta h P_l \nabla h)\|_{l^1_{|j|} L^\infty L^1} \\
&\lesssim 2^{(d-3-\delta)j^- + (\sigma+d/2)j^+} \sum_{l>j} \|P_l \nabla \delta h\|_{l^2_{|j|} L^\infty L^2} \|P_l \nabla h\|_{l^2_{|j|} L^\infty L^2} \\
&\lesssim 2^{(d-3-2\delta)j^-} (\|\nabla \delta h_{\leq 0}\|_{l^2_0 L^\infty L^2} \|\nabla h_{\leq 0}\|_{l^2_0 L^\infty L^2} + \sum_{l>0} 2^{dl} \|\nabla \delta h_l\|_{l^2 L^\infty L^2} \|\nabla h_l\|_{l^2 L^\infty L^2}) \\
&\quad + \sum_{l>j, j>0} 2^{(\sigma-d/2)(j-l)} 2^{(\sigma+d/2)l} \|\nabla \delta h_l\|_{l^2 L^\infty L^2} \|\nabla h_l\|_{l^2 L^\infty L^2} \\
&\lesssim 2^{(d-3-2\delta)j^-} \tilde{s}_0 \|\mathcal{S}\|_{\mathcal{E}^s} + \tilde{s}_j \|\mathcal{S}\|_{\mathcal{E}^s}.
\end{aligned}$$

Case 3. The contribution of $\delta h \nabla h \nabla h$ and $h \nabla h \nabla \delta h$. It suffices to prove that

$$\|\Delta^{-1} S_j (\delta h \nabla h \nabla h + h \nabla h \nabla \delta h)\|_{Y^{d/2-1-\delta, \sigma+2}} \lesssim \tilde{s}_j \|\mathcal{S}\|_{\mathcal{E}^s}^2.$$

For the low-frequency part, By Bernstein's inequality and $d \geq 4$ we have

$$\begin{aligned} & \|\Delta^{-1} (\delta h \nabla h \nabla h)_{\leq 0}\|_{Y^{d/2-1-\delta, \sigma+2}} \\ & \lesssim \|(\delta h \nabla h \nabla h)_{\leq 0}\|_{l_0^1 L^\infty L^1} \\ & \lesssim \|\delta h_{\leq 0}\|_{L^\infty L^\infty} \|(\nabla h \nabla h)_{\leq 0}\|_{l_0^1 L^\infty L^1} + \sum_{j>0} \|\delta h_j\|_{l_0^2 L^\infty L^2} \|(\nabla h \nabla h)_j\|_{l_0^2 L^\infty L^2} \\ & \lesssim \tilde{s}_0 \|\mathcal{S}\|_{\mathcal{E}^s}^2. \end{aligned}$$

For the high-frequency part, by Bernstein's inequality we also have

$$2^{(\sigma+2)j} \|\Delta^{-1} (\delta h \nabla h \nabla h)_j\|_{Y_j} \lesssim 2^{\sigma j} \|(\delta h \nabla h \nabla h)_j\|_{l_1^1 L^\infty L^2} \lesssim \tilde{s}_j \|\mathcal{S}\|_{\mathcal{E}^s}^2.$$

Thus this completes the proof of $Y^{d/2-1-\delta, \sigma+2}$ bound. \square

5. MULTILINEAR AND NONLINEAR ESTIMATES

This section contains our main multilinear estimates which are needed for the analysis of the Schrödinger equation in (2.35). We begin with the following low-high bilinear estimates of $\nabla h \nabla \psi$.

Lemma 5.1. *Let $s > \frac{d}{2}$, $d \geq 2$ and $k \in \mathbb{N}$. Suppose that $\nabla a(x) \lesssim \langle x \rangle^{-1}$, $h \in \mathbf{Y}^{\sigma+2}$ and $\psi_k \in l^2 X^s$. Then for $-s \leq \sigma \leq s$ we have*

$$(5.1) \quad \|\nabla h_{\leq k} \cdot \nabla \psi_k\|_{l^2 N^\sigma} \lesssim \min\{\|h\|_{\mathbf{Y}^{\sigma+2}} \|\psi_k\|_{l^2 X^s}, \|h\|_{\mathbf{Y}^{s+2}} \|\psi_k\|_{l^2 X^\sigma}\},$$

$$(5.2) \quad \|h_{\leq k} \nabla a \nabla \psi_k\|_{l^2 N^\sigma} \lesssim \min\{\|h\|_{\mathbf{Y}^{\sigma+2}} \|\psi_k\|_{l^2 X^s}, \|h\|_{\mathbf{Y}^{s+2}} \|\psi_k\|_{l^2 X^\sigma}\}.$$

In addition, if $-s \leq \sigma \leq s-1$ then we have

$$(5.3) \quad \|h_{\leq k} \nabla^2 \psi_k\|_{l^2 N^\sigma} \lesssim \|h\|_{\mathbf{Y}^{\sigma+2}} \|\psi_k\|_{l^2 X^s}.$$

Proof. a) The estimates (5.1) and (5.3). The proof of second bound (5.3) is similar to the first, so we only prove the first bound in detail. By duality, it suffices to estimate

$$I_j := \langle \nabla P_j h \nabla \psi_k, z_k \rangle, \quad j \leq k, \quad j \in \mathbb{Z}, \quad k \in \mathbb{N},$$

for any $z_k := S_k z \in l_k^2 X_k$ with $\|z_k\|_{l_k^2 X_k} \leq 1$. For I_j and any decomposition $P_j h = \sum_{l \geq |j|} h_{j,l}$, by duality and Bernstein inequality, we have

$$\begin{aligned} I_j & \lesssim \sum_{l \geq |j|} \sup_{\|z_k\|_{l_k^2 X_k} \leq 1} \langle \nabla h_{j,l} \nabla \psi_k, z_k \rangle \\ & \lesssim \sum_{l \geq |j|} \sup_{\|z_k\|_{l_k^2 X_k} \leq 1} \|\nabla h_{j,l}\|_{l_1^1 L^\infty L^\infty} \|\nabla \psi_k\|_{l_1^\infty L^2 L^2} \|z_k\|_{l_1^\infty L^2 L^2} \\ & \lesssim \sum_{l \geq |j|} 2^l \|\nabla h_{j,l}\|_{l_1^1 L^\infty L^\infty} \|\psi_k\|_{X_k} \\ & \lesssim 2^{(\frac{d}{2}+1)j+|j|} \sum_{l \geq |j|} 2^{l-|j|} \|h_{j,l}\|_{l_1^1 L^\infty L^2} \|\psi_k\|_{X_k}. \end{aligned}$$

Then taking the infimum over the decomposition of $P_j h$ and incorporating the summation over j yield

$$\sum_{j \leq k} 2^{\sigma k} I_j \lesssim \|h\|_{\mathbf{Y}^{d/2+2+\epsilon}} \|\psi_k\|_{X^\sigma},$$

for any $\epsilon > 0$. If $-s \leq \sigma \leq d/2$, we also have

$$\begin{aligned} \sum_{j \leq k} 2^{\sigma k} I_j &\lesssim \sum_{j \leq 0} 2^{dj/2} \|P_j h\|_{Y_j} \|\psi_k\|_{X^\sigma} + \sum_{j > 0} 2^{(d/2+\epsilon-\sigma)(j-k)} 2^{(\sigma+2)j} \|P_j h\|_{Y_j} \|\psi_k\|_{X^s} \\ &\lesssim \|h\|_{\mathbf{Y}^{\sigma+2}} \|\psi_k\|_{X^s}. \end{aligned}$$

Thus the bound (5.1) follows.

Estimate (5.2). By duality, it suffices to bound

$$II_j = \langle P_j h \nabla a \nabla \psi_k, z_k \rangle, \quad j \leq k, \quad j \in \mathbb{Z},$$

for any $z_k \in l_k^2 X_k$ with $\|z_k\|_{l_k^2 X_k} \leq 1$. For any decomposition $P_j h = \sum_{l \geq |j|} h_{j,l}$, by $|\nabla a|(x) \lesssim \langle x \rangle^{-1}$, we consider the two cases $|x| \geq 2^{j/2}$ and $|x| < 2^{j/2}$ respectively and then obtain

$$\begin{aligned} II_j &\lesssim \sum_{l \geq |j|} \sup_{\|z_k\|_{l_k^2 X_k} \leq 1} \langle h_{j,l} \langle x \rangle^{-1} \mathbf{1}_{\leq 2^{l/2}}(x) \nabla \psi_k, z_k \rangle + \sum_{l \geq |j|} \sup_{\|z_k\|_{l_k^2 X_k} \leq 1} \langle h_{j,l} \langle x \rangle^{-1} \mathbf{1}_{> 2^{l/2}}(x) \nabla \psi_k, z_k \rangle \\ &= II_{j1} + II_{j2}. \end{aligned}$$

The first term is bounded by

$$\begin{aligned} II_{j1} &\lesssim \sum_{l \geq |j|} \sup_{\|z_k\|_{l_k^2 X_k} \leq 1} \|h_{j,l}\|_{l_l^1 L^\infty L^\infty} \|\nabla \psi_k\|_{l_{l/2}^\infty L^2 L^2} \|z_k\|_{l_{l/2}^\infty L^2 L^2} \\ &\lesssim \sum_{l \geq |j|} 2^{l/2} \|h_{j,l}\|_{l_l^1 L^\infty L^\infty} \|\psi_k\|_{X_k} \\ &\lesssim 2^{dj/2+|j|/2} \sum_{l \geq |j|} 2^{l-|j|} \|h_{j,l}\|_{l_l^1 L^\infty L^2} \|\psi_k\|_{X_k} \end{aligned}$$

The second term is bounded by

$$\begin{aligned} II_{j2} &\lesssim \sum_{l \geq |j|} 2^{-l/2} \sup_{\|z_k\|_{l_k^2 X_k} \leq 1} \|h_{j,l}\|_{l_l^1 L^\infty L^\infty} \|\nabla \psi_k\|_{l_l^\infty L^2 L^2} \|z_k\|_{l_l^\infty L^2 L^2} \\ &\lesssim \sum_{l \geq |j|} 2^{l/2} \|h_{j,l}\|_{l_l^1 L^\infty L^\infty} \|\psi_k\|_{X_k} \\ &\lesssim 2^{dj/2+|j|/2} \sum_{l \geq |j|} 2^{l-|j|} \|h_{j,l}\|_{l_l^1 L^\infty L^2} \|\psi_k\|_{X_k}. \end{aligned}$$

Then we obtain

$$\begin{aligned} \sum_{j \leq k} 2^{\sigma k} II_j &\lesssim \left(\sum_{j \leq 0} 2^{(d-1)j/2} \|P_j h\|_{Y_j} + \sum_{j > 0} 2^{(d+1)j/2} \|h_j\|_{Y_j} \right) \|\psi_k\|_{X^\sigma} \\ &\lesssim \min\{ \|h\|_{\mathbf{Y}^{\sigma+2}} \|\psi_k\|_{l^2 X^s}, \|h\|_{\mathbf{Y}^{\sigma+2}} \|\psi_k\|_{l^2 X^\sigma} \}. \end{aligned}$$

Thus the bound (5.2) follows. \square

We next prove the remaining bilinear estimates and trilinear estimates.

Proposition 5.2 (Nonlinear estimates). *a) Let $s > \frac{d}{2}$ and $d \geq 3$, assume that p_k and s_k are admissible frequency envelopes for $\psi \in l^2 X^s$, $\mathcal{S} \in \mathcal{E}^s$ respectively. Then we have*

$$(5.4) \quad \|S_k(B\psi)\|_{l^2 N^s} \lesssim s_k \|\psi\|_{l^2 X^s} + p_k \|B\|_{Z^{1,s}},$$

$$(5.5) \quad \|S_k(A^2\psi)\|_{l^2 N^s} \lesssim s_k \|A\|_{Z^{1,s}} \|\psi\|_{l^2 X^s} + p_k \|A\|_{Z^{1,s}}^2,$$

$$(5.6) \quad \|S_k(\lambda^3)\|_{l^2 N^s} \lesssim s_k \|\lambda\|_{Z^{0,s}}^2.$$

b) Assume that \tilde{p}_k and \tilde{s}_k are admissible frequency envelopes for $\psi \in l^2 X^\sigma$, $\mathcal{S} \in \mathcal{E}^\sigma$ respectively. Then for $-s \leq \sigma \leq s$ we have

$$(5.7) \quad \|S_k \nabla(h_{\geq k-4} \nabla \psi)\|_{l^2 N^\sigma} \lesssim \min\{\tilde{s}_k \|\psi\|_{l^2 X^s}, \tilde{p}_k \|h\|_{Z^{1,s+2}}\},$$

$$(5.8) \quad \|S_k(A_{\geq k-4} \nabla \psi)\|_{l^2 N^\sigma} \lesssim \min\{\tilde{s}_k \|\psi\|_{l^2 X^s}, \tilde{p}_k \|A\|_{Z^{1,s+1}}\},$$

and for $-s \leq \sigma \leq s - \delta$ we have

$$(5.9) \quad \|S_k(B\psi)\|_{l^2 N^\sigma} \lesssim \min\{\tilde{s}_k \|\psi\|_{l^2 X^s}, \tilde{p}_k \|B\|_{Z^{1,s}}\},$$

$$(5.10) \quad \|S_k(A^2\psi)\|_{l^2 N^\sigma} \lesssim \min\{\tilde{s}_k \|A\|_{Z^{1,s}} \|\psi\|_{l^2 X^s}, \tilde{p}_k \|A\|_{Z^{1,s}}^2\},$$

$$(5.11) \quad \|S_k(\lambda^3)\|_{l^2 N^\sigma} \lesssim \tilde{s}_k \|\lambda\|_{Z^{0,s}}^2.$$

If $-s \leq \sigma \leq s - 1$, then

$$(5.12) \quad \|S_k(A_{< k-4} \nabla \psi)\|_{l^2 N^\sigma} \lesssim p_k \|A\|_{Z^{1,\sigma+1}}.$$

Proof. We first prove (5.7) and (5.8). These two bounds are proved similarly, here we only prove the first bound in detail. For the high-low case, by (3.1) we have

$$\begin{aligned} \sum_{j_2 \leq k+C} \|S_k \nabla(h_{j_1} \nabla \psi_{j_2})\|_{l^2 N^\sigma} &\lesssim \sum_{j_1 = k+O(1), j_2 \leq k+C} 2^{(\sigma+1)k} \|h_{j_1}\|_{l_k^2 L^2 L^2} \|\nabla \psi_{j_2}\|_{l_k^2 L^\infty L^\infty} \\ &\lesssim \sum_{j_2 \leq k+C} 2^{\sigma k + (d/2+1)j_2} \|\nabla h_k\|_{L^2 L^2} \|\psi_{j_2}\|_{l_{j_2}^2 L^\infty L^2} \\ &\lesssim \min\{\tilde{s}_k \|\psi\|_{l^2 X^s}, \tilde{p}_k \|h\|_{Z^{1,s+2}}\}. \end{aligned}$$

For the high-high case, when $\sigma + d/2 + 1 > \delta$ we have

$$\begin{aligned} &\sum_{j_1 = j_2 + O(1), j_1 > k} \|S_k \nabla(h_{j_1} \nabla \psi_{j_2})\|_{l^2 N^\sigma} \\ &\lesssim \sum_{j_1 = j_2 + O(1), j_1 > k} 2^{(\sigma+1)k + dk/2} \|S_k(h_{j_1} \nabla \psi_{j_2})\|_{L^2 L^1} \\ &\lesssim \sum_{j_1 = j_2 + O(1), j_1 > k} 2^{(\sigma+1+d/2)(k-j_1) + (\sigma+2+d/2)j_1} \|h_{j_1}\|_{L^2 L^2} \|\psi_{j_2}\|_{L^\infty L^2} \\ &\lesssim \min\{\tilde{s}_k \|\psi\|_{l^2 X^s}, \tilde{p}_k \|h\|_{Z^{1,s+2}}\}, \end{aligned}$$

and when $\sigma + d/2 + 1 \leq \delta$ we have

$$\begin{aligned} & \sum_{j_1=j_2+O(1), j_1>k} \|S_k \nabla(h_{j_1} \nabla \psi_{j_2})\|_{l^2 N^\sigma} \\ & \lesssim \sum_{j_1=j_2+O(1), j_1>k} 2^{(\sigma+1+d/2-2\delta)k+(2\delta+1)j_1} \|h_{j_1}\|_{L^2 L^2} \|\psi_{j_2}\|_{L^\infty L^2} \\ & \lesssim \min\{\tilde{s}_k \|\psi\|_{l^2 X^s}, \tilde{p}_k \|h\|_{Z^{1,s+2}}\}, \end{aligned}$$

Next, we prove the bounds (5.4)-(5.6) and (5.9)-(5.11). These bounds can be estimated similarly, we only prove (5.4) and (5.9) in detail. Indeed, by duality we have

$$\|S_k(B\psi)\|_{l^2 N^\sigma} \lesssim 2^{\sigma k} \|S_k(B\psi)\|_{L^2 L^2}.$$

Then using Littlewood-Paley dichotomy to divide this into low-high, high-low and high-high cases. For the low-high case, by Sobolev embedding we have

$$2^{\sigma k} \|S_k(B_{<k} \psi_k)\|_{L^2 L^2} \lesssim \|B_{<k}\|_{L^\infty L^\infty} 2^{\sigma k} \|\psi_k\|_{L^2 L^2} \lesssim \tilde{p}_k \|B\|_{Z^{1,s}}.$$

If $-s \leq \sigma \leq s - \delta$ we also have

$$\begin{aligned} 2^{\sigma k} \|S_k(B_{<k} \psi_k)\|_{L^2 L^2} & \lesssim \mathbf{1}_{[-s, \frac{d}{2})}(\sigma) \sum_{0 \leq l < k} 2^{(d/2+2\delta-\sigma)(l-k)} \|\nabla B_l\|_{L^\infty H^{\sigma-1}} 2^{(d/2+2\delta)k} \|\psi_k\|_{L^2 L^2} \\ & \quad + \mathbf{1}_{[\frac{d}{2}, s-\delta]}(\sigma) \sum_{0 \leq l < k} \|\nabla B_l\|_{L^\infty H^{\sigma-1}} 2^{\sigma k} \|\psi_k\|_{L^2 L^2} \\ & \lesssim \mathbf{1}_{[-s, \frac{d}{2})}(\sigma) \sum_{0 \leq l < k} 2^{(d/2+2\delta-\sigma)(l-k)} 2^{\delta(k-l)} \tilde{s}_k 2^{(d/2+2\delta)k} \|\psi_k\|_{L^2 L^2} \\ & \quad + \mathbf{1}_{[\frac{d}{2}, s-\delta]}(\sigma) \sum_{0 \leq l < k} 2^{\delta(k-l)} \tilde{s}_k 2^{\sigma k} \|\psi_k\|_{L^2 L^2} \\ & \lesssim \tilde{s}_k \|\psi\|_{l^2 X^s}. \end{aligned}$$

The high-low case can be estimated similarly. For the high-high case, by Sobolev embedding when $\sigma + d/2 \geq 0$ we have

$$\begin{aligned} 2^{\sigma k} \|S_k(B_l \psi_l)\|_{L^2 L^2} & \lesssim \sum_{l>k} 2^{(\sigma+d/2+\delta)(k-l)} 2^{(\sigma+d/2+\delta)l} \|B_l\|_{L^\infty L^2} \|\psi_l\|_{L^2 L^2} \\ & \lesssim \min\{\tilde{s}_k \|\psi\|_{l^2 X^s}, \tilde{p}_k \|B\|_{Z^{1,s}}\}, \end{aligned}$$

and when $\sigma + d/2 < 0$ we have

$$\begin{aligned} 2^{\sigma k} \|S_k(B_l \psi_l)\|_{L^2 L^2} & \lesssim \sum_{l>k} 2^{(\sigma+d/2)k} \|B_l\|_{L^\infty L^2} \|\psi_l\|_{L^2 L^2} \\ & \lesssim \min\{\tilde{s}_k \|\psi\|_{l^2 X^s}, \tilde{p}_k \|B\|_{Z^{1,s}}\}, \end{aligned}$$

These imply the bound (5.4) and (5.9).

Finally, we prove the bound (5.12). If $\sigma > d/2 - 1 + \delta$, by duality and Sobolev embedding, we have

$$2^{\sigma k} \|A_{<k} \nabla \psi_k\|_{L^2 L^2} \lesssim \sum_{l \leq k} 2^{(d/2-1)l} \|\nabla A_l\|_{L^\infty L^2} 2^{(\sigma+1)k} \|\psi_k\|_{L^2 L^2} \lesssim p_k \|A\|_{Z^{1,\sigma+1}}.$$

If $\sigma \leq d/2 - 1 + \delta$, we have

$$\begin{aligned} 2^{\sigma k} \|A_{<k} \nabla \psi_k\|_{L^2 L^2} &\lesssim \sum_{0 \leq l < k} 2^{(d/2-1-\sigma+\delta)(l-k)} \|\nabla A_l\|_{L^\infty H^\sigma} 2^{(d/2+\delta)k} \|\psi_k\|_{L^2 L^2} \\ &\lesssim p_k \|A\|_{Z^{1,\sigma+1}}. \end{aligned}$$

Then the bound (5.12) follows. Hence this completes the proof of the lemma. \square

We shall also require the following bounds on commutators.

Proposition 5.3 (Commutator bounds). *Let $s > \frac{d}{2}$, $d \geq 2$. Let $m(D)$ be a multiplier with symbol $m \in S^0$. Assume $h \in \mathbf{Y}^{s+2}$, $A \in Z^{1,s+1}$ and $\psi_k \in l^2 X^s$, frequency localized at frequency 2^k . If $-s \leq \sigma \leq s$ we have*

$$(5.13) \quad \|\nabla[S_{<k-4} h, m(D)] \nabla \psi_k\|_{l^2 N^\sigma} \lesssim \min\{\|h\|_{\mathbf{Y}^{\sigma+2}} \|\psi_k\|_{l^2 X^s}, \|h\|_{\mathbf{Y}^{s+2}} \|\psi_k\|_{l^2 X^\sigma}\},$$

$$(5.14) \quad \|[S_k, A_{<k-4}] \nabla \psi_k\|_{l^2 N^\sigma} \lesssim \min\{\|A\|_{Z^{1,s+1}} \|\psi_k\|_{l^2 X^\sigma}, \|A\|_{Z^{1,\sigma+1}} \|\psi_k\|_{l^2 X^s}\}.$$

Proof. First we estimate (5.13). In [21, Proposition 3.2], it was shown that

$$\nabla[S_{<k-4} g, m(D)] \nabla S_k \psi = L(\nabla S_{<k-4} g, \nabla S_k \psi),$$

where L is a translation invariant operator satisfying

$$L(f, g)(x) = \int f(x+y) g(x+z) \tilde{m}(y+z) dy dz, \quad \tilde{m} \in L^1.$$

Given this representation, as we are working in translation-invariant spaces, by (5.1) the bound (5.13) follows.

Next, for the bound (5.14). Since

$$[S_k, A_{<k}] \nabla \psi = \int_0^1 \int 2^{kd} \tilde{\varphi}(2^k y) 2^k y \nabla A_{<k}(x - sy) 2^{-k} \nabla \psi_{[k-3, k+3]}(x - y) dy ds,$$

By translation-invariance and the similar argument to (5.9), the bound (5.14) follows. This completes the proof of the lemma. \square

6. LOCAL ENERGY DECAY AND THE LINEARIZED PROBLEM

In this section, we consider a linear Schrödinger equation

$$(6.1) \quad \begin{cases} i\partial_t \psi + \partial_\alpha g^{\alpha\beta} \partial_\beta \psi + 2iA^\alpha \partial_\alpha \psi = F, \\ \psi(0) = \psi_0, \end{cases}$$

and, under suitable assumptions on the coefficients, we prove that the solution satisfies suitable energy and local energy bounds.

6.1. The linear paradifferential Schrödinger flow. As an intermediate step, here we prove energy and local energy bounds for a frequency localized linear paradifferential Schrödinger equation

$$(6.2) \quad i\partial_t \psi_k + \partial_\alpha (g_{<k-4}^{\alpha\beta} \partial_\beta \psi_k) + 2iA_{<k-4}^\alpha \partial_\alpha \psi_k = f_k.$$

We begin with the energy estimates, which are fairly standard:

Lemma 6.1 (Energy-type estimate). *Let $d \geq 2$, ψ_k solves the equation (6.2) with initial data $\psi_k(0)$ in the time interval $[0, 1]$. For a fixed $s > \frac{d}{2}$, assume that $A \in Z^{1,s+1}$, $\psi_k \in l_k^2 X_k$, $f_{1k} \in N$ and $f_{2k} \in L^1 L^2$, where $f_k = f_{1k} + f_{2k}$. Then we have*

$$(6.3) \quad \begin{aligned} \|\psi_k\|_{L_t^\infty L_x^2}^2 &\lesssim \|\psi_k(0)\|_{L^2}^2 + \|A\|_{Z^{1,s+1}} \|\psi_k\|_{X_k}^2 + \|\psi_k\|_{X_k} \|f_{1k}\|_{N_k} \\ &\quad + \|\psi_k\|_{L^\infty L^2} \|f_{2k}\|_{L^1 L^2}. \end{aligned}$$

Proof. By (6.2), we have

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} \|\psi_k\|_{L^2}^2 &= \operatorname{Re} \langle \psi_k, \partial_t \psi_k \rangle \\ &= \operatorname{Re} \langle \psi_k, i \partial_\alpha g_{<k-4}^{\alpha\beta} \partial_\beta \psi_k - 2A_{<k-4}^\alpha \partial_\alpha \psi_k - i f_k \rangle \\ &= -\operatorname{Re} \langle \partial_\alpha \psi_k, i g_{<k-4}^{\alpha\beta} \partial_\beta \psi_k \rangle - \operatorname{Re} \int_{\mathbb{R}^d} A_{<k-4}^\alpha \partial_\alpha |\psi_k|^2 dx - \operatorname{Re} \langle \psi_k, i f_k \rangle \\ &= \operatorname{Re} \int_{\mathbb{R}^d} \partial_\alpha A_{<k-4}^\alpha |\psi_k|^2 dx - \operatorname{Re} \langle \psi_k, i f_k \rangle, \end{aligned}$$

and notice that for each $t \in [0, 1]$ we have by duality and Sobolev embedding

$$\begin{aligned} \|\psi_k(t)\|_{L^2}^2 &\lesssim \|\psi_k(0)\|_{L^2}^2 + \int_0^1 \int_{\mathbb{R}^d} |\partial_\alpha A_{<k-4}^\alpha| |\psi_k|^2 dx dt + \|\psi_k\|_{X_k} \|f_{1k}\|_{N_k} \\ &\quad + \|\psi_k\|_{L^\infty L^2} \|f_{2k}\|_{L^1 L^2} \\ &\lesssim \|\psi_k(0)\|_{L^2}^2 + \|A\|_{Z^{1,s+1}} \|\psi_k\|_{X_k}^2 \\ &\quad + \|\psi_k\|_{X_k} \|f_{1k}\|_{N_k} + \|\psi_k\|_{L^\infty L^2} \|f_{2k}\|_{L^1 L^2} \end{aligned}$$

We take the supremum over t on the left hand side and the conclusion follows. \square

Next, we prove the main result of this section, namely the local energy estimates for solutions to (6.2):

Proposition 6.2 (Local energy decay). *Let $d \geq 3$, assume that the coefficients $g^{\alpha\beta} = \delta^{\alpha\beta} + h^{\alpha\beta}$ and A^α in (6.2) satisfy*

$$(6.4) \quad \|h\|_{\mathbf{Y}^{s+2}}, \|A\|_{Z^{1,s+1}} \ll 1$$

for some $s > \frac{d}{2}$. Let ψ_k be a solution to (6.2) which is localized at frequency 2^k . Then the following estimate holds:

$$(6.5) \quad \|\psi_k\|_{l_k^2 X_k} \lesssim \|\psi_{0k}\|_{L^2} + \|f_k\|_{l_k^2 N_k}$$

Proof. The proof is closely related to that given in [21, 22]. However, here we are able to relax the assumptions both on the metric g and on the magnetic potential A . In the latter case, unlike in [21, 22], we treat the magnetic term $2iA_{<k-4}^\alpha \partial_\alpha \psi_k$ as a part of the linear equation, which allows us to avoid bilinear estimates for this term and use only the bound for A in $Z^{1,s+1}$.

As an intermediate step in the proof, we will establish a local energy decay bound in a cube $Q \in \mathcal{Q}_l$ with $0 \leq l \leq k$:

$$(6.6) \quad \begin{aligned} 2^{k-l} \|\psi_k\|_{L^2 L^2([0,1] \times Q)}^2 &\lesssim \|\psi_k\|_{L^\infty L^2}^2 + \|f_k\|_{N_k} \|\psi_k\|_{X_k} \\ &\quad + (2^{-k} + \|A\|_{Z^{1,s+1}} + \|h\|_{\mathbf{Y}^{s+2}}) \|\psi_k\|_{l_k^2 X_k}^2. \end{aligned}$$

The proof of this bound is based on a positive commutator argument using a well chosen multiplier \mathcal{M} . This will be first-order differential operator with smooth coefficients which are localized at frequency $\lesssim 1$. Precisely, we will use a multiplier \mathcal{M} which is a self-adjoint differential operator having the form

$$(6.7) \quad i2^k \mathcal{M} = a^\alpha(x) \partial_\alpha + \partial_\alpha a^\alpha(x)$$

with uniform bounds on a and its derivatives.

Before proving (6.5), we need the following lemma which is used to dismiss the $(g - I_d)$ contribution to the commutator $[\partial_\alpha g^{\alpha\beta} \partial_\beta, \mathcal{M}]$.

Lemma 6.3. *Let $s > \frac{d}{2}$ and $d \geq 3$, assume that $h \in \mathbf{Y}^{s+2}$, $A \in Z^{1,s+1}$ and $\psi \in l_k^2 X_k$, let \mathcal{M} be as (6.7). Then we have*

$$(6.8) \quad \int_0^1 \langle [\partial_\alpha h_{\leq k}^{\alpha\beta} \partial_\beta, \mathcal{M}] \psi_k, \psi_k \rangle ds \lesssim \|h\|_{\mathbf{Y}^{s+2}} \|\psi_k\|_{l_k^2 X_k}^2,$$

$$(6.9) \quad \int_0^1 \operatorname{Re} \langle A_{< k-4}^\alpha \partial_\alpha \psi_k, \mathcal{M} \psi_k \rangle ds \lesssim \|A\|_{Z^{1,s+1}} \|\psi_k\|_{X_k}^2.$$

Proof of Lemma 6.3. By (6.7) and directly computations, we get

$$[\partial_\alpha h^{\alpha\beta} \partial_\beta, \mathcal{M}] \approx 2^{-k} [\nabla(h \nabla a + a \nabla h) \nabla + \nabla h \nabla^2 a + h \nabla^3 a].$$

Then it suffices to estimate

$$2^{-k} \int_0^1 \langle (h_{\leq k} \nabla a + a \nabla h_{\leq k}) \nabla \psi_k, \nabla \psi_k \rangle dt + 2^{-k} \int_0^1 \langle (\nabla h_{\leq k} \nabla^2 a + h_{\leq k} \nabla^3 a) \psi_k, \psi_k \rangle dt$$

The first integral is estimated by (5.1) and (5.2). Using Sobolev embedding, the second integral is bounded by

$$2^{-k} \int_0^1 \langle (\nabla h_{\leq k} + h_{\leq k}) \psi_k, \psi_k \rangle dt \lesssim \| \langle \nabla \rangle h_{\leq k} \|_{L^\infty} 2^{-k} \|\psi_k\|_{L^2 L^2}^2 \lesssim \|\nabla h\|_{L^\infty H^s} \|\psi_k\|_{l_k^2 X_k}^2.$$

Hence, the bound (6.8) follows.

For the second bound (6.9), by (6.7) and integration by parts we rewrite the following term as

$$\begin{aligned} & \operatorname{Re} \langle A^\alpha \partial_\alpha \psi, i \sum_{\beta=1}^d (a_\beta \partial_\beta + \partial_\beta a_\beta) \psi \rangle \\ &= \operatorname{Re} \sum_{\beta=1}^d \int_{\mathbb{R}^d} \left[i \partial_\alpha (\bar{\psi} A^\alpha a_\beta \partial_\beta \psi) - i \bar{\psi} \partial_\alpha A^\alpha a_\beta \partial_\beta \psi - i \bar{\psi} A^\alpha \partial_\alpha a_\beta \partial_\beta \psi - i \bar{\psi} A^\alpha a_\beta \partial_{\alpha\beta}^2 \psi \right. \\ & \quad \left. + i \partial_\beta (A^\alpha \partial_\alpha \bar{\psi} a_\beta \psi) - i \partial_\beta A^\alpha \partial_\alpha \bar{\psi} a_\beta \psi - i A^\alpha \partial_{\alpha\beta}^2 \bar{\psi} a_\beta \psi \right] dx \\ & \approx \int_{\mathbb{R}^d} \langle \nabla \rangle A \psi \nabla \psi dx. \end{aligned}$$

Then we bound the left-hand side of (6.9) by

$$\begin{aligned} \int_0^1 \operatorname{Re} \langle A_{<k-4}^\alpha \partial_\alpha \psi_k, \mathcal{M} \psi_k \rangle ds &\lesssim 2^{-k} \int_0^1 \int_{\mathbb{R}^d} |\langle \nabla \rangle A_{<k} \psi_k \nabla \psi_k| dx ds \\ &\lesssim \|\nabla A\|_{L^\infty H^s} \|\psi_k\|_{L^2 L^2}^2. \end{aligned}$$

This implies the bound (6.9), and hence completes the proof of the lemma. \square

Returning to the proof of (6.6), for the self-adjoint multiplier \mathcal{M} we compute

$$\begin{aligned} \frac{d}{dt} \langle \psi_k, \mathcal{M} \psi_k \rangle &= 2 \operatorname{Re} \langle \partial_t \psi_k, \mathcal{M} \psi_k \rangle \\ &= 2 \operatorname{Re} \langle i \partial_\alpha (g_{<k-4}^{\alpha\beta} \partial_\beta \psi_k) - 2A_{<k-4}^\alpha \partial_\alpha \psi_k - i f_k, \mathcal{M} \psi_k \rangle \\ &= i \langle [-\partial_\alpha g_{<k-4}^{\alpha\beta} \partial_\beta, \mathcal{M}] \psi_k, \psi_k \rangle + 2 \operatorname{Re} \langle -2A_{<k-4}^\alpha \partial_\alpha \psi_k - i f_k, \mathcal{M} \psi_k \rangle \end{aligned}$$

We then use the multiplier \mathcal{M} as in [21, 22] so that the following three properties hold:

(1) Boundedness on frequency 2^k localized functions,

$$\|\mathcal{M}u\|_{L_x^2} \lesssim \|u\|_{L_x^2}.$$

(2) Boundedness in X ,

$$\|\mathcal{M}u\|_X \lesssim \|u\|_X.$$

(3) Positive commutator,

$$i \langle [-\partial_\alpha g_{<k-4}^{\alpha\beta} \partial_\beta, \mathcal{M}]u, u \rangle \gtrsim 2^{k-l} \|u\|_{L_{t,x}^2([0,1] \times Q)}^2 - O(2^{-k} + \|h\|_{\mathbf{Y}^{s+2}}) \|u\|_{L_k^2 X_k}^2.$$

If these three properties hold for $u = \psi_k$, then by (6.9) and (6.4) the bound (6.6) follows.

We first do this when the Fourier transform of the solution ψ_k is restricted to a small angle

$$(6.10) \quad \operatorname{supp} \widehat{\psi}_k \subset \{|\xi| \lesssim \xi_1\}.$$

Without loss of generality due to translation invariance, $Q = \{|x_j| \leq 2^l : j = 1, \dots, d\}$, and we set m to be a smooth, bounded, increasing function such that $m'(s) = \varphi^2(s)$ where φ is a Schwartz function localized at frequencies $\lesssim 1$, and $\varphi \approx 1$ for $|s| \leq 1$. We rescale m and set $m_l(s) = m(2^{-l}s)$. Then, we fix

$$\mathcal{M} = \frac{1}{i2^k} (m_l(x_1) \partial_1 + \partial_1 m_l(x_1)).$$

The properties (1) and (2) are immediate due to the frequency localization of $u = \psi_k$ and m_l as well as the boundedness of m_l . By (6.8) it suffices to consider the property (3) for the operator

$$-\Delta = -\partial_\alpha g_{<k-4}^{\alpha\beta} \partial_\beta + \partial_\alpha h_{<k-4}^{\alpha\beta} \partial_\beta.$$

This yields

$$i2^k [-\Delta, \mathcal{M}] = -2^{-l+2} \partial_1 \varphi^2(2^{-l}x_1) \partial_1 + O(1),$$

and hence

$$i2^k \langle [-\Delta, \mathcal{M}] \psi_k, \psi_k \rangle = 2^{-l+2} \|\varphi(2^{-l}x_1) \partial_1 \psi_k\|_{L^2 L^2}^2 + O(\|\psi_k\|_{L^2 L^2}^2)$$

Utilizing our assumption (6.10), it follows that

$$2^{k-l} \|\varphi(2^{-l}x_1) \psi_k\|_{L^2 L^2}^2 \lesssim i \langle [-\Delta, \mathcal{M}] \psi_k, \psi_k \rangle + 2^{-k} O(\|\psi_k\|_{L^2 L^2}^2)$$

which yields (3) when combined with (6.8).

We proceed to reduce the problem to the case when (6.10) holds. We let $\{\theta_j(\omega)\}_{j=1}^d$ be a partition of unity,

$$\sum_j \theta_j(\omega) = 1, \quad \omega \in \mathbb{S}^{d-1},$$

where $\theta_j(\omega)$ is supported in a small angle about the j -th coordinate axis. Then, we can set $\psi_{k,j} = \Theta_{k,j} \psi_k$ where

$$\mathcal{F}\Theta_{k,j}\psi = \theta_j\left(\frac{\xi}{|\xi|}\right) \sum_{k-1 \leq l \leq k+1} \varphi_l(\xi) \widehat{\psi}(t, \xi).$$

We see that

$$\begin{aligned} & (i\partial_t + \partial_\alpha g_{<k-4}^{\alpha\beta} \partial_\beta) \psi_{k,j} + 2iA_{<k-4}^\alpha \partial_\alpha \psi_{k,j} \\ &= \Theta_{k,j} f_k - \partial_\alpha [\Theta_{k,j}, g_{<k-4}^{\alpha\beta}] \partial_\beta \psi_k - 2i[\Theta_{k,j}, A_{\leq k-4}^\alpha] \partial_\alpha \psi_k. \end{aligned}$$

By applying \mathcal{M} , suitably adapted to the correct coordinate axis, to $\psi_{k,j}$ and summing over j , we obtain

$$\begin{aligned} & 2^{k-l} \|\psi_k\|_{L^2 L^2([0,1] \times Q)}^2 \\ & \lesssim \|\psi_k\|_{L^\infty L^2}^2 + \sum_{j=1}^d \int_0^1 \langle -\Theta_{k,j} f_k, \mathcal{M}\psi_{k,j} \rangle ds \\ & \quad + \sum_{j=1}^d \int \langle [\Theta_{k,j}, \partial_\alpha g_{<k-4}^{\alpha\beta} \partial_\beta] \psi_k + [\Theta_{k,j}, 2iA_{<k-4}^\alpha] \partial_\alpha \psi_k, \mathcal{M}\psi_{k,j} \rangle ds \\ & \quad + (2^{-k} + \|A\|_{Z^{1,s+1}} + \|h\|_{\mathbf{Y}^{s+2}}) \|\psi_k\|_{l_k^2 X_k}^2 \\ & \lesssim \|\psi_k\|_{L^\infty L^2}^2 + \|f_k\|_{N_k} \|\psi_k\|_{X_k} + (2^{-k} + \|A\|_{Z^{1,s+1}} + \|h\|_{\mathbf{Y}^{s+2}}) \|\psi_k\|_{l_k^2 X_k}^2. \end{aligned}$$

The commutator is done via (5.13) and (5.14). Then (6.6) follows.

Next we use the bound (6.6) to complete the proof of Proposition 6.2. Taking the supremum in (6.6) over $Q \in \mathcal{Q}_l$ and over l , we obtain

$$\begin{aligned} 2^k \|\psi_k\|_X^2 & \lesssim \|\psi_k\|_{L^\infty L^2}^2 + \|f_{1k}\|_{N_k} \|\psi_k\|_{X_k} + \|f_{2k}\|_{L^1 L^2} \|\psi_k\|_{L^\infty L^2} \\ & \quad + (2^{-k} + \|A\|_{Z^{1,s+1}} + \|h\|_{\mathbf{Y}^{s+2}}) \|\psi_k\|_{l_k^2 X_k}^2 \\ & \lesssim \|\psi_k\|_{L^\infty L^2}^2 + \|f_{1k}\|_{N_k} \|\psi_k\|_{X_k} + \|f_{2k}\|_{L^1 L^2}^2 \\ & \quad + (2^{-k} + \|A\|_{Z^{1,s+1}} + \|h\|_{\mathbf{Y}^{s+2}}) \|\psi_k\|_{l_k^2 X_k}^2. \end{aligned}$$

Combined with (6.3), we get

$$(6.11) \quad \begin{aligned} \|\psi_k\|_{X_k}^2 & \lesssim \|\psi_k(0)\|_{L^2}^2 + \|f_{1k}\|_{N_k}^2 + \|f_{2k}\|_{L^1 L^2}^2 \\ & \quad + (2^{-k} + \|A\|_{Z^{1,s+1}} + \|h\|_{\mathbf{Y}^{s+2}}) \|\psi_k\|_{l_k^2 X_k}^2. \end{aligned}$$

We now finish the proof by incorporating the summation over cubes. We let $\{\chi_Q\}$ denote a partition via functions which are localized to frequencies $\lesssim 1$ which are associated to cubes

Q of scale $M2^k$. We also assume that $|\nabla^l \chi_Q| \lesssim (2^k M)^{-l}$, $l = 1, 2$. Thus,

$$\begin{aligned} & (i\partial_t + \partial_\alpha g_{<k-4}^{\alpha\beta} \partial_\beta) \chi_Q \psi_k + 2iA_{<k-4}^\alpha \partial_\alpha \chi_Q \psi_k \\ &= \chi_Q f_k + [\partial_\alpha g_{<k-4}^{\alpha\beta} \partial_\beta, \chi_Q] \psi_k + 2iA_{<k-4}^\alpha \partial_\alpha \chi_Q \cdot \psi_k \end{aligned}$$

Applying (6.3) to $\chi_Q \psi_k$, we obtain

$$\begin{aligned} & \sum_Q \|\chi_Q \psi_k\|_{L^\infty L^2}^2 \\ & \lesssim \sum_Q \|\chi_Q \psi_k(0)\|_{L^2}^2 + \|A\|_{Z^{1,s+1}} \sum_Q \|\chi_Q \psi_k\|_{X_k}^2 \\ & \quad + \left(\sum_Q \|\chi_Q f_k\|_{N_k}^2 \right)^{1/2} \left(\sum_Q \|\chi_Q \psi_k\|_{X_k}^2 \right)^{1/2} \\ & \quad + \sum_Q \|[\partial_\alpha g_{<k-4}^{\alpha\beta} \partial_\beta, \chi_Q] \psi_k + 2iA_{<k-4}^\alpha \partial_\alpha \chi_Q \cdot \psi_k\|_{L^1 L^2}^2. \end{aligned}$$

But by (6.4) we have

$$\begin{aligned} (6.12) \quad \sum_Q \|[\nabla g \nabla, \chi_Q] \psi_k\|_{L^1 L^2}^2 & \lesssim \sum_Q \|\nabla g \cdot \nabla \chi_Q \cdot \psi_k + g \nabla(\nabla \chi_Q \cdot \psi_k)\|_{L^1 L^2}^2 \\ & \lesssim (1 + \|h\|_{Z^{1,s+2}}) M^{-2} \sum_Q \|\chi_Q \psi_k\|_{L^\infty L^2}^2, \end{aligned}$$

and also

$$(6.13) \quad \sum_Q \|2iA_{<k-4}^\alpha \partial_\alpha \chi_Q \cdot \psi_k\|_{L^1 L^2}^2 \lesssim (1 + \|A\|_{Z^{1,s}}) M^{-2} \sum_Q \|\chi_Q \psi_k\|_{L^\infty L^2}^2.$$

For M sufficiently large, we can bootstrap the commutator terms, and, after a straightforward transition to cubes of scale 2^k rather than $M2^k$, we observe that

$$(6.14) \quad \|\psi_k\|_{l_k^2 L^\infty L^2}^2 \lesssim \|\psi_k(0)\|_{L^2}^2 + \|A\|_{Z^{1,s+1}} \|\psi_k\|_{l_k^2 X_k}^2 + \|f_k\|_{l_k^2 N_k} \|\psi_k\|_{l_k^2 X_k}.$$

We now apply (6.11) to $\chi_Q \psi_k$, and then by (6.12) and (6.13) we see that

$$\begin{aligned} \sum_Q \|\chi_Q \psi_k\|_{X_k}^2 & \lesssim \|\psi_k(0)\|_{L^2}^2 + \sum_Q \|\chi_Q f_k\|_{N_k}^2 + M^{-2} \sum_Q \|\chi_Q \psi_k\|_{X_k}^2 \\ & \quad + (2^{-k} + \|h\|_{\mathbf{Y}^{s+2}} + \|A\|_{Z^{1,s+1}}) \sum_Q \|\chi_Q \psi_k\|_{l_k^2 X_k}^2. \end{aligned}$$

For $M \gg 1$, we have

$$M^{-d} \|\psi_k\|_{l_k^2 X_k}^2 \lesssim \|\psi_k(0)\|_{L^2}^2 + \|f_k\|_{l_k^2 N_k}^2 + (2^{-k} + \|h\|_{\mathbf{Y}^{s+2}} + \|A\|_{Z^{1,s+1}}) \|\psi_k\|_{l_k^2 X_k}^2.$$

By (6.4), for k sufficiently large (depending on M), we may absorb the last terms in the right-hand side into the left, i.e

$$\|\psi_k\|_{l_k^2 X_k}^2 \lesssim \|\psi_k(0)\|_{L^2}^2 + \|f_k\|_{l_k^2 N_k}^2.$$

On the other hand, for the remaining bounded range of k , we have

$$\|\psi\|_{X_k} \lesssim \|\psi\|_{L^\infty L^2},$$

and then (6.14) and (6.4) gives

$$\begin{aligned}\|\psi_k\|_{l_k^2 X_k}^2 &\lesssim \|\psi_k(0)\|_{L^2}^2 + \|A\|_{Z^{1,s+1}} \|\psi_k\|_{l_k^2 X_k}^2 + \|f_k\|_{l_k^2 N_k} \|\psi_k\|_{l_k^2 X_k} \\ &\lesssim \|\psi_k(0)\|_{L^2}^2 + \|f_k\|_{l_k^2 N_k}^2,\end{aligned}$$

which finishes the proof of (6.5). \square

6.2. The full linear problem. Here we use the bounds for the paradifferential equation in the previous subsection in order to prove similar bounds for the full equation (6.1):

Proposition 6.4 (Well-posedness). *Let $s > \frac{d}{2}$, $d \geq 3$ and $h = g - I_d \in \mathbf{Y}^{s+2}$, assume that the metric g , and the magnetic potential A satisfy*

$$\|h\|_{\mathbf{Y}^{s+2}}, \|A\|_{Z^{1,s+1}} \ll 1.$$

Then the equation (6.1) is well-posed for initial data $\psi_0 \in H^\sigma$ with $-s \leq \sigma \leq s$, and we have the estimate

$$(6.15) \quad \|\psi\|_{l^2 X^\sigma} \lesssim \|\psi_0\|_{H^\sigma} + \|F\|_{l^2 N^\sigma}.$$

Moreover, for $0 \leq \sigma \leq s$ we have the estimate

$$(6.16) \quad \|\psi\|_{l^2 \mathbf{X}^\sigma} \lesssim \|\psi_0\|_{H^\sigma} + \|F\|_{l^2 N^\sigma \cap L^2 H^{\sigma-2}}.$$

Proof. The well-posedness follows in a standard fashion from a similar energy estimate for the adjoint equation. Since the adjoint equation has a similar form, with similar bounds on the coefficients, such an estimate follows directly from (6.15). Thus, we now focus on the proof of the bound (6.15). For ψ solving (6.1), we see that ψ_k solves

$$\begin{cases} i\partial_t \psi_k + \partial_\alpha g_{<k-4}^{\alpha\beta} \partial_\beta \psi_k + 2iA_{<k-4}^\alpha \partial_\alpha \psi_k = F_k + H_k, \\ \psi_k(0) = \psi_{0k}, \end{cases}$$

where

$$\begin{aligned}H_k &= -S_k \partial_\alpha g_{\geq k-4}^{\alpha\beta} \partial_\beta \psi - [S_k, \partial_\alpha g_{<k-4}^{\alpha\beta} \partial_\beta] \psi - 2i[S_k, A_{<k-4}^\alpha] \partial_\alpha \psi \\ &\quad - 2iS_k [A_{\geq k-4}^\alpha \partial_\alpha \psi_k].\end{aligned}$$

If we apply Proposition 6.2 to each of these equations, we see that

$$\|\psi_k\|_{l^2 X^\sigma}^2 \lesssim \|\psi_{0k}\|_{H^\sigma}^2 + \|F_k\|_{l^2 N^\sigma}^2 + \|H_k\|_{l^2 N^\sigma}^2.$$

We claim that

$$\sum_k \|H_k\|_{l^2 N^\sigma}^2 \lesssim (\|h\|_{\mathbf{Y}^{s+2}} + \|A\|_{Z^{1,s+1}})^2 \|\psi\|_{l^2 X^\sigma}^2, \text{ for } -s \leq \sigma \leq s.$$

Indeed, the bound for the terms in H_k follows from (5.7), (5.13), (5.14), (5.8), respectively. Then by the above two bounds, we obtain the estimate (6.15).

Finally, by the ψ -equation (6.1), for time derivative bound it suffices to consider the form

$$\partial_t \psi = \Delta \psi + \nabla(h \nabla \psi) + A \nabla \psi + F.$$

Then by the standard Littlewood-Paley dichotomy and Bernstein's inequality, for $0 \leq \sigma \leq s$ we have the following estimates

$$\|\partial_t \psi\|_{L^2 H^{\sigma-2}} \lesssim \|\psi\|_{L^\infty H^\sigma} + \|F\|_{L^2 H^{\sigma-2}},$$

This, combined with (6.15), yields the bound (6.16), and then completes the proof of the Lemma. \square

6.3. The linearized problem. Here we consider the linearized equation:

$$(6.17) \quad \begin{cases} i\partial_t \Psi + \partial_\alpha g^{\alpha\beta} \partial_\beta \Psi + 2iA^\alpha \partial_\alpha \Psi = F + G, \\ \Psi(0) = \Psi_0, \end{cases}$$

where

$$G = -\nabla(\mathcal{G}\nabla\psi) - 2i\mathcal{A}^\alpha \partial_\alpha \psi,$$

and we prove the following.

Proposition 6.5. *Let $s > \frac{d}{2}$, $0 \leq \sigma \leq s - 1$, $d \geq 3$ and $h = g - I_d \in \mathbf{Y}^{s+2}$, assume that Ψ is a solution of (6.17), the metric g and A satisfy*

$$\|h\|_{\mathbf{Y}^{s+2}}, \|A\|_{Z^{1,s+1}} \ll 1.$$

Then we have the estimate

$$(6.18) \quad \|\Psi\|_{l^2 \mathbf{X}^\sigma} \lesssim \|\Psi_0\|_{H^\sigma} + \|F\|_{l^2 N^\sigma \cap L^2 H^{\sigma-2}} + (\|\mathcal{G}\|_{\mathbf{Y}^{\sigma+2}} + \|\mathcal{A}\|_{Z^{1,\sigma+1}}) \|\psi\|_{l^2 X^s}.$$

Proof. For Ψ solving (6.17), we see that Ψ_k solves

$$\begin{cases} i\partial_t \Psi_k + \partial_\alpha g_{<k-4}^{\alpha\beta} \partial_\beta \Psi_k + 2iA_{<k-4}^\alpha \partial_\alpha \Psi_k = F_k + G_k + H_k, \\ \Psi_k(0) = \Psi_{0k}, \end{cases}$$

where

$$\begin{aligned} G_k &= -S_k(\nabla(\mathcal{G}\nabla\psi) - 2i\mathcal{A}^\alpha \partial_\alpha \psi), \\ H_k &= -S_k \partial_\alpha g_{\geq k-4}^{\alpha\beta} \partial_\beta \Psi - [S_k, \partial_\alpha g_{<k-4}^{\alpha\beta}] \Psi - 2i[S_k, A_{<k-4}^\alpha] \partial_\alpha \Psi \\ &\quad - 2iS_k[A_{\geq k-4}^\alpha \partial_\alpha \Psi_k]. \end{aligned}$$

The proof of (6.18) is similar to that of (6.16). Here it suffices to prove

$$\begin{aligned} \sum_k \|G_k\|_{l^2 N^\sigma}^2 &\lesssim \|\mathcal{G}\|_{\mathbf{Y}^{\sigma+2}}^2 \|\psi\|_{l^2 X^s}^2 + \|\mathcal{A}\|_{Z^{1,\sigma+1}}^2 \|\psi\|_{l^2 X^s}^2, \\ \|G\|_{L^2 H^{\sigma-2}} &\lesssim (\|\mathcal{G}\|_{\mathbf{Y}^{\sigma+2}} + \|\mathcal{A}\|_{Z^{1,\sigma+1}}) \|\psi\|_{l^2 X^s}. \end{aligned}$$

Indeed, the bound for the terms in G_k follows from (5.7), (5.3), (5.8) and (5.12). The second bound follows from a standard Littlewood-Paley decomposition and Bernstein's inequality. This completes the proof of the Lemma. \square

7. WELL-POSEDNESS IN THE GOOD GAUGE

In this section we use the elliptic results in Section 4, the multilinear estimates in Section 5 and the linear local energy decay bounds in Section 6 in order to prove the good gauge formulation of our main result, namely Theorem 2.7.

7.1. The iteration scheme: uniform bounds. Here we seek to construct solutions to (2.35) iteratively, based on the scheme

$$(7.1) \quad \begin{cases} (i\partial_t + \partial_\alpha g^{(n)\alpha\beta} \partial_\beta) \psi^{(n+1)} + 2i(A^{(n)\alpha} - \frac{1}{2}V^{(n)\alpha}) \partial_\alpha \psi^{(n+1)} = F^{(n)}, \\ \psi(0) = \psi_0, \end{cases}$$

with the trivial initialization

$$\psi^{(0)} = 0,$$

where the nonlinearities are

$$(7.2) \quad F^{(n)} = \partial_\alpha g^{(n)\alpha\beta} \cdot \partial_\beta \psi^{(n)} + (B^{(n)} + A_\alpha^{(n)} A^{(n)\alpha} - V^{(n)\alpha} A_\alpha^{(n)}) \psi^{(n)} - i\lambda_\sigma^{(n)\gamma} \text{Im}(\psi^{(n)} \bar{\lambda}_\gamma^{(n)\sigma}),$$

and $\mathcal{S}^{(n)} = (\lambda^{(n)}, h^{(n)}, V^{(n)}, A^{(n)}, B^{(n)})$ are the solutions of elliptic equations (2.36) with $\psi = \psi^{(n)}$.

We assume that ψ_0 is small in H^s . Due to the above trivial initialization, we also inductively assume that

$$\|\psi^{(n)}\|_{l^2 \mathbf{X}^s} \leq C \|\psi_0\|_{H^s},$$

where C is a big constant.

Applying the elliptic estimate (4.14) to (2.36) with $\psi = \psi^{(n)}$ at each step, we obtain

$$\|\mathcal{S}^{(n)}\|_{\mathcal{E}^s} \lesssim \|\psi^{(n)}\|_{l^2 \mathbf{X}^s} \lesssim \|\psi_0\|_{H^s},$$

Applying at each step the local energy bound (6.16) with $\sigma = s$ we obtain the estimate

$$\begin{aligned} \|\psi^{(n+1)}\|_{l^2 \mathbf{X}^s} &\lesssim \|\psi_0\|_{H^s} + \|F^{(n)}\|_{l^2 N^s \cap L^2 H^{s-2}} \\ &\lesssim \|\psi_0\|_{H^s} + \|\mathcal{S}^{(n)}\|_{\mathcal{E}^s} (1 + \|\mathcal{S}^{(n)}\|_{\mathcal{E}^s}) \|\psi^{(n)}\|_{l^2 X^s} \\ &\lesssim \|\psi_0\|_{H^s} + (C \|\psi_0\|_{H^s})^2 (1 + C \|\psi_0\|_{H^s}). \end{aligned}$$

Here the nonlinear terms in $F^{(n)}$ are estimated using (5.1), (5.7), (5.4), (5.5) and (5.6) with $\sigma = s$. Since ψ_0 is small in H^s , the above bound gives

$$(7.3) \quad \|\psi^{(n+1)}\|_{l^2 \mathbf{X}^s} \leq C \|\psi_0\|_{H^s},$$

which closes our induction.

7.2. The iteration scheme: weak convergence. Here we prove that our iteration scheme converges in the weaker H^{s-1} topology. We denote the differences by

$$\begin{aligned} \Psi^{(n+1)} &= \psi^{(n+1)} - \psi^{(n)}, \\ \delta \mathcal{S}^{(n+1)} &= (\Lambda^{(n+1)}, \mathcal{G}^{(n+1)}, \mathcal{V}^{(n+1)}, \mathcal{A}^{(n+1)}, \mathcal{B}^{(n+1)}) = \mathcal{S}^{(n+1)} - \mathcal{S}^{(n)} \end{aligned}$$

Then from (7.1) we obtain the system

$$\begin{cases} i\partial_t \Psi^{(n+1)} + \partial_\alpha (g^{(n)\alpha\beta} \partial_\beta \Psi^{(n+1)}) + 2i(A^{(n)\alpha} - \frac{1}{2}V^{(n)\alpha}) \partial_\alpha \Psi^{(n+1)} = F^{(n)} - F^{(n-1)} + G^{(n)}, \\ \Psi^{(n+1)}(0, x) = 0, \end{cases}$$

where the nonlinearities $G^{(n)}$ have the form

$$G^{(n)} = -\partial_\alpha (\mathcal{G}^{(n)} \partial_\beta \psi^{(n)}) - 2i(\mathcal{A}^{(n)\alpha} - \frac{1}{2}\mathcal{V}^{(n)\alpha}) \partial_\alpha \psi^{(n)},$$

By (4.16) we obtain

$$\|\delta\mathcal{S}^{(n)}\|_{\mathcal{E}^{s-1}} \lesssim \|\Psi^{(n)}\|_{l^2\mathbf{X}^{s-1}}.$$

Applying (6.18) with $\sigma = s - 1$ for the $\Psi^{(n+1)}$ equation we have

$$\|\Psi^{(n+1)}\|_{l^2\mathbf{X}^{s-1}} \lesssim \|F^{(n)} - F^{(n-1)}\|_{l^2N^{s-1}\cap L^2H^{s-3}} + (\|\mathcal{G}^{(n)}\|_{\mathbf{Y}^{s+1}} + \|(\mathcal{V}^{(n)}, \mathcal{A}^{(n)})\|_{\mathbf{Z}^{1,s}})\|\psi^{(n)}\|_{l^2X^s}.$$

Then by (5.1), (5.7), (5.9), (5.10) and (5.11) with $\sigma = s - 1$ we bound the right hand side above by

$$\|\Psi^{(n+1)}\|_{l^2\mathbf{X}^{s-1}} \lesssim C\|\psi_0\|_{H^s}\|(\Psi^{(n)}, \delta\mathcal{S}^{(n)})\|_{l^2\mathbf{X}^{s-1}\times\mathcal{E}^{s-1}} \ll \|\Psi^{(n)}\|_{l^2\mathbf{X}^{s-1}}.$$

This implies that our iterations $\psi^{(n)}$ converge in $l^2\mathbf{X}^{s-1}$ to some function ψ . Furthermore, by the uniform bound (7.3) it follows that

$$(7.4) \quad \|\psi\|_{l^2\mathbf{X}^s} \lesssim \|\psi_0\|_{H^s}.$$

Interpolating, it follows that $\psi^{(n)}$ converges to ψ in $l^2\mathbf{X}^{s-\epsilon}$ for all $\epsilon > 0$. This allows us to conclude that the auxiliary functions $\mathcal{S}^{(n)}$ associated to $\psi^{(n)}$ converge to the functions \mathcal{S} associated to ψ , and also to pass to the limit and conclude that ψ solves the (SMCF) equation (2.35). Thus we have established the existence part of our main theorem.

7.3. Uniqueness via weak Lipschitz dependence. Consider the difference of two solutions

$$(\Psi, \delta\mathcal{S}) = (\psi^{(1)} - \psi^{(2)}, \mathcal{S}^{(1)} - \mathcal{S}^{(2)}).$$

The Ψ solves an equation of this form

$$\begin{cases} i\partial_t\Psi + \partial_\alpha g^{(1)\alpha\beta}\partial_\beta\Psi + 2i(A^{(1)\alpha} - \frac{1}{2}V^{(1)\alpha})\partial_\alpha\Psi = F^{(1)} - F^{(2)} + G, \\ \Psi(0, x) = \psi_0^{(1)}(x) - \psi_0^{(2)}(x), \end{cases}$$

where the nonlinearity G is

$$G = -\partial_\alpha(\mathcal{G}\partial_\beta\psi^{(2)}) - 2i(\mathcal{A}^\alpha - \frac{1}{2}\mathcal{V}^\alpha)\partial_\alpha\psi^{(2)}.$$

By (4.16), we have

$$(7.5) \quad \|\delta\mathcal{S}\|_{\mathcal{E}^{s-1}} \lesssim \|\Psi\|_{l^2\mathbf{X}^{s-1}}.$$

Applying (6.18) with $\sigma = s - 1$ to the Ψ equation, we obtain the estimate

$$\begin{aligned} \|\Psi\|_{l^2\mathbf{X}^{s-1}} &\lesssim \|\Psi_0\|_{H^{s-1}} + \|F^{(1)} - F^{(2)}\|_{l^2N^{s-1}\cap L^2H^{s-3}} + (\|\mathcal{G}\|_{\mathbf{Y}^{s+1}} + \|(\mathcal{V}, \mathcal{A})\|_{\mathbf{Z}^{1,s}})\|\psi^{(2)}\|_{l^2X^s} \\ &\lesssim \|\Psi_0\|_{H^{s-1}} + C\|(\psi_0^{(1)}, \psi_0^{(2)})\|_{H^s}\|(\Psi, \delta\mathcal{S})\|_{l^2\mathbf{X}^{s-1}\times\mathcal{E}^{s-1}}. \end{aligned}$$

Then, by the above bound (7.5), we further have

$$\|\Psi\|_{l^2\mathbf{X}^{s-1}} \lesssim \|\Psi_0\|_{H^{s-1}} + C\|(\psi_0^{(1)}, \psi_0^{(2)})\|_{H^s}\|\Psi\|_{l^2\mathbf{X}^{s-1}}$$

Since the initial data $\psi_0^{(1)}$ and $\psi_0^{(2)}$ are sufficiently small, we obtain

$$(7.6) \quad \|\Psi\|_{l^2\mathbf{X}^{s-1}} \lesssim \|\Psi_0\|_{H^{s-1}}.$$

This gives the weak Lipschitz dependence, as well as the uniqueness of solutions for (2.35).

7.4. Frequency envelope bounds. Here we prove a stronger frequency envelope version of estimate (7.4).

Proposition 7.1. *Let $\psi \in l^2\mathbf{X}^s$ be a small data solution to (2.35), which satisfies (7.4). Let $\{p_{0k}\}$ be an admissible frequency envelope for the initial data $\psi_0 \in H^s$. Then $\{p_{0k}\}$ is also frequency envelope for ψ in $l^2\mathbf{X}^s$.*

Proof. Let p_k and s_k be the admissible frequency envelopes for solution $(\psi, \mathcal{S}) \in l^2\mathbf{X}^s \times \mathcal{E}^s$. Applying S_k to the Schrödinger equation (2.35), we obtain the paradifferential equation

$$\begin{cases} (i\partial_t + \partial_\alpha g_{<k-4}^{\alpha\beta} \partial_\beta) \psi_k + 2i(A - \frac{1}{2}V)_{<k-4}^\alpha \partial_\alpha \psi_k = F_k + J_k, \\ \psi(0, x) = \psi_0(x), \end{cases}$$

where

$$\begin{aligned} J_k = & -S_k \partial_\alpha g_{\geq k-4}^{\alpha\beta} \partial_\beta \psi - [S_k, \partial_\alpha g_{<k-4}^{\alpha\beta} \partial_\beta] \psi - 2i[S_k, (A - \frac{1}{2}V)_{<k-4}^\alpha] \partial_\alpha \psi \\ & - 2iS_k [(A - \frac{1}{2}V)_{\geq k-4}^\alpha \partial_\alpha \psi_k], \end{aligned}$$

and $\mathcal{S} = (\lambda, h, V, A, B)$ is the solution to the elliptic system (2.36). We estimate $\psi_k = S_k \psi$ using Proposition 6.4. By Proposition 5.2, Lemma 5.1 and Lemma 5.3 we obtain

$$\|\psi_k\|_{l^2\mathbf{X}^s} \lesssim p_{0k} + p_k \|\mathcal{S}\|_{\mathcal{E}^s} + s_k \|\psi\|_{l^2\mathbf{X}^s} \lesssim p_{0k} + (p_k + s_k) \|\psi\|_{l^2\mathbf{X}^s}.$$

Then by (4.15), the definition of frequency envelope (3.3) and (7.4), this implies

$$p_k \lesssim p_{0k} + p_k \|\psi\|_{l^2\mathbf{X}^s}.$$

By the smallness of $\psi \in l^2\mathbf{X}^s$, this further gives $p_k \lesssim p_{0k}$, and concludes the proof. \square

7.5. Continuous dependence on the initial data. Here we show that the map $\psi_0 \rightarrow (\psi, \mathcal{S})$ is continuous from H^s into $l^2\mathbf{X}^s \times \mathcal{E}^s$. By (4.16), it suffices to prove $\psi_0 \rightarrow \psi$ is continuous from H^s to $l^2\mathbf{X}^s$.

Suppose that $\psi_0^{(n)} \rightarrow \psi_0$ in H^s . Denote by $p_{0k}^{(n)}$, respectively p_{0k} the frequency envelopes associated to $\psi_0^{(n)}$, respectively ψ_0 , given by (3.3). If $\psi_0^{(n)} \rightarrow \psi_0$ in H^s then $p_{0k}^{(n)} \rightarrow p_{0k}$ in l^2 . Then for each $\epsilon > 0$ we can find some N_ϵ so that

$$\|p_{0, > N_\epsilon}^{(n)}\|_{l^2} \leq \epsilon, \text{ for all } n.$$

By Proposition 7.1 we obtain that

$$(7.7) \quad \|\psi_{> N_\epsilon}^{(n)}\|_{l^2\mathbf{X}^s} \leq \epsilon, \text{ for all } n.$$

To compare $\psi^{(n)}$ with ψ we use (7.6) for low frequencies and (7.7) for the high frequencies,

$$\begin{aligned} \|\psi^{(n)} - \psi\|_{l^2\mathbf{X}^s} & \lesssim \|S_{< N_\epsilon}(\psi^{(n)} - \psi)\|_{l^2\mathbf{X}^s} + \|S_{> N_\epsilon} \psi^{(n)}\|_{l^2\mathbf{X}^s} + \|S_{> N_\epsilon} \psi\|_{l^2\mathbf{X}^s} \\ & \lesssim 2^{N_\epsilon} \|S_{< N_\epsilon}(\psi^{(n)} - \psi)\|_{l^2\mathbf{X}^{s-1}} + 2\epsilon \\ & \lesssim 2^{N_\epsilon} \|S_{< N_\epsilon}(\psi_0^{(n)} - \psi_0)\|_{H^{s-1}} + 2\epsilon. \end{aligned}$$

Letting $n \rightarrow \infty$ we obtain

$$\limsup_{n \rightarrow \infty} \|\psi^{(n)} - \psi\|_{l^2\mathbf{X}^s} \lesssim \epsilon.$$

Letting $\epsilon \rightarrow 0$ we obtain

$$\lim_{n \rightarrow 0} \|\psi^{(n)} - \psi\|_{l^2 \mathbf{X}^s} = 0,$$

which completes the desired result.

7.6. Higher regularity. Here we prove that the solution (ψ, \mathcal{S}) satisfies the bound

$$(7.8) \quad \|(\psi, \mathcal{S})\|_{l^2 \mathbf{X}^\sigma \times \mathcal{E}^\sigma} \lesssim \|\psi_0\|_{H^\sigma}, \quad \sigma \geq s,$$

whenever the right hand side is finite.

Differentiating the original Schrödinger equation (2.35) yields

$$(i\partial_t + \partial_\alpha g^{\alpha\beta} \partial_\beta) \nabla \psi + 2i(A - \frac{V}{2})^\alpha \partial_\alpha \nabla \psi = -\partial_\alpha (\nabla g^{\alpha\beta} \partial_\beta \psi) - 2i\nabla A^\alpha \partial_\alpha \psi + \nabla F,$$

where F is defined as in (7.2) without superscript (n) . Using Proposition 6.5 we obtain

$$\|\nabla \psi\|_{l^2 \mathbf{X}^s} \lesssim \|\nabla \psi_0\|_{H^s} + \|(\nabla \psi, \nabla \mathcal{S})\|_{l^2 \mathbf{X}^s \times \mathcal{E}^s} \|(\psi, \mathcal{S})\|_{l^2 \mathbf{X}^s \times \mathcal{E}^s} (1 + \|(\psi, \mathcal{S})\|_{l^2 \mathbf{X}^s \times \mathcal{E}^s})^N.$$

For elliptic equations, by (4.16) we obtain

$$\|\nabla \mathcal{S}\|_{\mathcal{E}^s} \lesssim \|\nabla \psi\|_{l^2 \mathbf{X}^s}.$$

Hence, by (7.4) and the smallness of ψ_0 in H^s , these imply

$$\|(\nabla \psi, \nabla \mathcal{S})\|_{l^2 \mathbf{X}^s \times \mathcal{E}^s} \lesssim \|\nabla \psi_0\|_{H^s}.$$

Inductively, we can obtain the system for $(\nabla^n \psi, \nabla^n \mathcal{S})$. This leads to

$$\|(\nabla^n \psi, \nabla^n \mathcal{S})\|_{l^2 \mathbf{X}^s \times \mathcal{E}^s} \lesssim \|\psi_0\|_{H^{s+n}} + \|\psi\|_{l^2 \mathbf{X}^{s+n}} \|\psi\|_{l^2 \mathbf{X}^s} (1 + \|\psi\|_{l^2 \mathbf{X}^s})^N,$$

which shows that

$$\|(\psi, \mathcal{S})\|_{l^2 \mathbf{X}^{s+n} \times \mathcal{E}^{s+n}} \lesssim \|\psi_0\|_{H^{s+n}} + \|\psi\|_{l^2 \mathbf{X}^{s+n}} \|\psi\|_{l^2 \mathbf{X}^s} (1 + \|\psi\|_{l^2 \mathbf{X}^s})^N,$$

and hence gives the bound (7.8) by the smallness of ψ in $l^2 \mathbf{X}^s$.

7.7. The time evolution of (λ, g, A) . As part of our derivation of the (SMCF) equations (2.35) for the mean curvature ψ in the good gauge, coupled with the elliptic system (2.36), we have seen that the time evolution of (λ, g, A) is described by the equations (2.31), (2.26) and (2.32). However, our proof of the well-posedness result for the Schrödinger evolution (2.35) does not apriori guarantee that (2.31), (2.26) and (2.32) hold. Here we rectify this omission:

Lemma 7.2. *Assume that $\psi \in C[0, T; H^s]$ solves the SMCF equation (2.35) coupled with the elliptic system (2.36). Then the relations (2.26), (2.31) and (2.32) hold.*

Proof. We recall that, by Theorem 4.1, the solution $\mathcal{S} = (\lambda, h, V, A, B)$ in \mathcal{H}^s for the system (2.36) satisfies the fixed time constraints (2.4), (2.8), (2.15), (2.13), (2.19) and (2.16). On the other hand, in terms of the time evolution, at this point we only have the equation (2.35) for the mean curvature ψ . We will show that this implies (2.26), (2.31) and (2.32).

To shorten the notations, we define the tensors

$$\begin{aligned} T_{\alpha\beta}^1 &= \partial_t g_{\alpha\beta} - 2 \operatorname{Im}(\psi \bar{\lambda}_{\alpha\beta}) - \nabla_\alpha V_\beta - \nabla_\beta V_\alpha, \\ T_\alpha^{2,\sigma} &= (\partial_t^B - V^\gamma \nabla_\gamma^A) \lambda_\alpha^\sigma - i \nabla_\alpha^A \nabla^{A,\sigma} \psi + \lambda_\alpha^\gamma \operatorname{Im}(\psi \bar{\lambda}_\gamma^\sigma) + \lambda_{\alpha\gamma} \nabla^\gamma V^\sigma - \lambda_\gamma^\sigma \nabla_\alpha V^\gamma, \\ T_\alpha^3 &= \partial_t A_\alpha - \partial_\alpha B - \operatorname{Re}(\lambda_\alpha^\gamma \bar{\partial}_\gamma^A \bar{\psi}) + \operatorname{Im}(\lambda_\alpha^\gamma \bar{\lambda}_{\gamma\sigma}) V^\sigma. \end{aligned}$$

We need to show that $T^1 = 0$, $T^2 = 0$, $T^3 = 0$. To do this, we will show that (T^1, T^2, T^3) solve a linear homogeneous coupled elliptic system of the form

$$\begin{cases} \Delta_g T^1 = \nabla(\Gamma T^1) + \lambda^2 T^1 + \lambda T^2, \\ \nabla^{A,\alpha} T_\alpha^{2,\sigma} = \lambda T^3 + \lambda \nabla T^1 + T^1 \nabla \lambda, \\ \nabla_\alpha^A T_\beta^{2,\sigma} - \nabla_\beta^A T_\alpha^{2,\sigma} = \lambda T^3 + \lambda \nabla T^1 + T^1 \nabla \lambda, \\ \nabla^\alpha T_\alpha^3 = T^1 \nabla A, \\ \nabla_\alpha T_\beta^3 - \nabla^\beta T_\alpha^3 = \lambda T^2. \end{cases}$$

Considering this system for $(T^1, T^2, T^3) \in \dot{H}^1 \times L^2 \times L^2$, the smallness condition on the coefficients $(\lambda, h, V, A, B) \in \mathcal{S}$ insures that this system has the unique solution $(T^1, T^2, T^3) = 0$. It remains to derive the system for (T^1, T^2, T^3) .

The equation for T^1 . This has the form

$$(7.9) \quad \begin{aligned} \Delta_g T_{\alpha\beta}^1 &= T_{\delta\beta}^1 \text{Ric}^\delta_\alpha + T_{\delta\alpha}^1 \text{Ric}^\delta_\beta + 2T^{1,\mu\nu} R_{\alpha\mu\beta\nu} - \nabla_\beta(T^{1,\mu\nu} \Gamma_{\mu\nu,\alpha}) - \nabla_\alpha(T^{1,\mu\nu} \Gamma_{\mu\nu,\beta}) \\ &\quad - 2\text{Re}(g_{\sigma\beta} T_\alpha^{2,\sigma} \bar{\psi} + T_{\sigma\beta}^1 \lambda_\alpha^\sigma \psi + \bar{\lambda}_{\alpha\beta} T_\sigma^{2,\sigma} - g_{\sigma\mu} T_\alpha^{2,\sigma} \bar{\lambda}_\beta^\mu - T_{\sigma\mu}^1 \lambda_\alpha^\sigma \bar{\lambda}_\beta^\mu - \bar{\lambda}_{\alpha\sigma} T_\beta^{2,\sigma}). \end{aligned}$$

We start with the first term in T^1 , and compute the expression $\Delta_g \partial_t g_{\alpha\beta}$. We have

$$\begin{aligned} \Delta_g \partial_t g_{\alpha\beta} &= g^{\mu\nu} (\partial_\mu \nabla_\nu \partial_t g_{\alpha\beta} - \Gamma_{\nu\alpha}^\delta \nabla_\nu \partial_t g_{\delta\beta} - \Gamma_{\mu\beta}^\delta \nabla_\nu \partial_t g_{\alpha\delta}) \\ &= [\partial_t (g^{\mu\nu} \partial_\mu \partial_\nu g_{\alpha\beta}) - \partial_t g^{\mu\nu} \partial_\mu \partial_\nu g_{\alpha\beta}] + [-g^{\mu\nu} \Gamma_{\nu\alpha}^\delta \partial_\mu \partial_t g_{\delta\beta} - g^{\mu\nu} \Gamma_{\nu\beta}^\delta \partial_\mu \partial_t g_{\delta\alpha} \\ &\quad - g^{\mu\nu} \partial_\mu \Gamma_{\nu\alpha}^\delta \partial_t g_{\delta\beta} - g^{\mu\nu} \partial_\mu \Gamma_{\nu\beta}^\delta \partial_t g_{\delta\alpha} - g^{\mu\nu} (\Gamma_{\mu\alpha}^\delta \nabla_\nu \partial_t g_{\delta\beta} + \Gamma_{\mu\beta}^\delta \nabla_\nu \partial_t g_{\delta\alpha})] \\ &:= I + II. \end{aligned}$$

We then use covariant derivatives to write II as

$$\begin{aligned} II &= -g^{\mu\nu} \Gamma_{\mu\alpha}^\delta (2\nabla_\nu \partial_t g_{\delta\beta} + \Gamma_{\nu\delta}^\sigma \partial_t g_{\sigma\beta} + \Gamma_{\nu\beta}^\sigma \partial_t g_{\sigma\delta}) \\ &\quad - g^{\mu\nu} \Gamma_{\mu\beta}^\delta (2\nabla_\nu \partial_t g_{\delta\alpha} + \Gamma_{\nu\delta}^\sigma \partial_t g_{\sigma\alpha} + \Gamma_{\nu\alpha}^\sigma \partial_t g_{\sigma\delta}) \\ &\quad - g^{\mu\nu} \partial_\mu \Gamma_{\nu\alpha}^\delta \partial_t g_{\delta\beta} - g^{\mu\nu} \partial_\mu \Gamma_{\nu\beta}^\delta \partial_t g_{\delta\alpha} \\ &= -2g^{\mu\nu} \Gamma_{\mu\alpha}^\delta \nabla_\nu \partial_t g_{\delta\beta} - 2g^{\mu\nu} \Gamma_{\mu\beta}^\delta \nabla_\nu \partial_t g_{\delta\alpha} \\ &\quad - \partial_t g_{\delta\beta} g^{\mu\nu} (\partial_\mu \Gamma_{\nu\alpha}^\delta + \Gamma_{\mu\alpha}^\sigma \Gamma_{\nu\sigma}^\delta) - \partial_t g_{\delta\alpha} g^{\mu\nu} (\partial_\mu \Gamma_{\nu\beta}^\delta + \Gamma_{\mu\beta}^\sigma \Gamma_{\nu\sigma}^\delta) \\ &\quad - 2\partial_t g_{\sigma\delta} g^{\mu\nu} \Gamma_{\mu\alpha}^\delta \Gamma_{\nu\beta}^\sigma. \end{aligned}$$

For I , by the g equation (2.22) we have

$$\begin{aligned} I &= \partial_t [-\partial_\alpha g^{\mu\nu} \partial_\mu g_{\nu\beta} - \partial_\beta g^{\mu\nu} \partial_\mu g_{\nu\alpha} + \partial_\alpha g^{\mu\nu} \partial_\beta g_{\mu\nu}] \\ &\quad + [2\partial_t (g^{\mu\nu} \Gamma_{\mu\alpha,\delta} \Gamma_{\nu\beta}^\delta) - \partial_t g^{\mu\nu} \partial_\mu \partial_\nu g_{\alpha\beta}] - 2\partial_t \widetilde{\text{Ric}}_{\alpha\beta} \\ &:= I_1 + I_2 + I_3. \end{aligned}$$

The expression I_1 is written as

$$\begin{aligned}
I_1 &= -\partial_\alpha \partial_t g^{\mu\nu} \Gamma_{\mu\nu,\beta} - \partial_\beta \partial_t g^{\mu\nu} \Gamma_{\mu\nu,\alpha} - \frac{1}{2} \partial_\beta \partial_t g^{\mu\nu} \partial_\alpha g_{\mu\nu} + \frac{1}{2} \partial_\alpha \partial_t g^{\mu\nu} \partial_\beta g_{\mu\nu} \\
&\quad - \partial_\alpha g^{\mu\nu} \partial_\mu \partial_t g_{\nu\beta} - \partial_\beta g^{\mu\nu} \partial_\mu \partial_t g_{\nu\alpha} + \partial_\alpha g^{\mu\nu} \partial_\beta \partial_t g_{\mu\nu} \\
&= -(\nabla_\alpha \partial_t g^{\mu\nu} - 2\Gamma_{\alpha\delta}^\mu \partial_t g^{\delta\nu}) \Gamma_{\mu\nu,\beta} - (\nabla_\beta \partial_t g^{\mu\nu} - 2\Gamma_{\beta\delta}^\mu \partial_t g^{\delta\nu}) \Gamma_{\mu\nu,\alpha} \\
&\quad + \frac{1}{2} [\nabla_\alpha (\partial_t g^{\mu\nu} \partial_\beta g_{\mu\nu} - \nabla_\beta (\partial_t g^{\mu\nu} \partial_\alpha g_{\mu\nu}))] \\
&\quad - \partial_\alpha g^{\mu\nu} (\nabla_\mu \partial_t g_{\nu\beta} + \Gamma_{\mu\nu}^\delta \partial_t g_{\delta\beta}) - \partial_\beta g^{\mu\nu} (\nabla_\mu \partial_t g_{\nu\alpha} + \Gamma_{\mu\nu}^\delta \partial_t g_{\delta\alpha} + \Gamma_{\mu\alpha}^\delta \partial_t g_{\nu\delta}) \\
&\quad + \partial_\alpha g^{\mu\nu} (\nabla_\beta \partial_t g_{\mu\nu} + \Gamma_{\beta\nu}^\delta \partial_t g_{\mu\delta}) \\
&= \nabla_\alpha \partial_t g^{\mu\nu} (-\Gamma_{\mu\nu,\beta} + \Gamma_{\mu\beta,\nu}) + \nabla_\beta \partial_t g^{\mu\nu} (-\Gamma_{\mu\nu,\alpha} + \Gamma_{\mu\alpha,\nu}) - \nabla_\mu \partial_t g_{\nu\beta} \partial_\alpha g^{\mu\nu} - \nabla_\mu \partial_t g_{\nu\alpha} \partial_\beta g^{\mu\nu} \\
&\quad + 2\partial_t g^{\mu\nu} (\Gamma_{\alpha\mu}^\delta \Gamma_{\delta\nu,\beta} + \Gamma_{\beta\mu}^\delta \Gamma_{\delta\nu,\alpha}) + \partial_t g^{\mu\nu} (-\Gamma_{\alpha\mu}^\delta \partial_\beta g_{\delta\nu} + \Gamma_{\beta\mu}^\delta \partial_\alpha g_{\delta\nu}) \\
&\quad + \partial_t g^{\mu\nu} (-\partial_\beta g_{\mu\sigma} g^{\sigma\delta} \Gamma_{\delta\alpha,\nu} + \partial_\alpha g_{\mu\sigma} g^{\sigma\delta} \Gamma_{\delta\beta,\nu}) \\
&\quad - \partial_t g_{\delta\beta} \partial_\alpha g^{\mu\nu} \Gamma_{\mu\nu}^\delta - \partial_t g_{\delta\alpha} \partial_\beta g^{\mu\nu} \Gamma_{\mu\nu}^\delta
\end{aligned}$$

For I_2 , we first compute

$$\begin{aligned}
2g^{\mu\nu} \partial_t (\Gamma_{\mu\alpha,\delta} \Gamma_{\nu\beta}^\delta) &= g^{\mu\nu} \Gamma_{\nu\beta}^\delta (\nabla_\mu \partial_t g_{\alpha\delta} + \nabla_\alpha \partial_t g_{\mu\delta} - \nabla_\delta \partial_t g_{\mu\alpha}) + 4g^{\mu\nu} \Gamma_{\nu\beta}^\delta \Gamma_{\alpha\mu}^\sigma \partial_t g_{\sigma\delta} \\
&\quad + g^{\mu\nu} \Gamma_{\nu\alpha}^\delta (\nabla_\mu \partial_t g_{\beta\delta} + \nabla_\beta \partial_t g_{\mu\delta} - \nabla_\delta \partial_t g_{\mu\beta}) + 2\partial_t g^{\sigma\delta} g^{\mu\nu} \Gamma_{\mu\alpha,\delta} \Gamma_{\nu\beta,\sigma}
\end{aligned}$$

By the above computations, we collect the $\nabla \partial_t g$ terms from I_1 , I_2 and II

$$\begin{aligned}
&\nabla_\alpha \partial_t g^{\mu\nu} (-\Gamma_{\mu\nu,\beta} + \Gamma_{\mu\beta,\nu}) + \nabla_\beta \partial_t g^{\mu\nu} (-\Gamma_{\mu\nu,\alpha} + \Gamma_{\mu\alpha,\nu}) - \nabla_\mu \partial_t g_{\nu\beta} \partial_\alpha g^{\mu\nu} - \nabla_\mu \partial_t g_{\nu\alpha} \partial_\beta g^{\mu\nu} \\
&\quad + g^{\mu\nu} \Gamma_{\nu\beta}^\delta (\nabla_\mu \partial_t g_{\alpha\delta} + \nabla_\alpha \partial_t g_{\mu\delta} - \nabla_\delta \partial_t g_{\mu\alpha}) + g^{\mu\nu} \Gamma_{\nu\alpha}^\delta (\nabla_\mu \partial_t g_{\beta\delta} + \nabla_\beta \partial_t g_{\mu\delta} - \nabla_\delta \partial_t g_{\mu\beta}) \\
&\quad - 2g^{\mu\nu} \Gamma_{\mu\alpha}^\delta \nabla_\nu \partial_t g_{\delta\beta} - 2g^{\mu\nu} \Gamma_{\mu\beta}^\delta \nabla_\nu \partial_t g_{\delta\alpha},
\end{aligned}$$

where the terms containing $\nabla \partial_t g_{\nu\alpha}$ and $\nabla \partial_t g_{\nu\beta}$ vanish, i.e.

$$\nabla_\mu \partial_t g_{\nu\beta} (-\partial_\alpha g^{\mu\nu} - g^{\nu\delta} \Gamma_{\delta\alpha}^\mu - g^{\delta\mu} \Gamma_{\delta\alpha}^\nu) + \nabla_\mu \partial_t g_{\nu\alpha} (-\partial_\beta g^{\mu\nu} - g^{\nu\delta} \Gamma_{\delta\beta}^\mu - g^{\mu\delta} \Gamma_{\delta\beta}^\nu) = 0,$$

and the terms with $\nabla \partial_t g^{\mu\nu}$ were rewritten as

$$\begin{aligned}
(7.10) \quad &-\nabla_\alpha \partial_t g^{\mu\nu} \Gamma_{\mu\nu,\beta} - \nabla_\beta \partial_t g^{\mu\nu} \Gamma_{\mu\nu,\alpha} \\
&= -\nabla_\alpha (\partial_t g^{\mu\nu} \Gamma_{\mu\nu,\beta}) - \nabla_\beta (\partial_t g^{\mu\nu} \Gamma_{\mu\nu,\alpha}) + \partial_t g^{\mu\nu} (\nabla_\alpha \Gamma_{\mu\nu,\beta} + \nabla_\beta \Gamma_{\mu\nu,\alpha})
\end{aligned}$$

We collect the $\partial_t g$ terms from I and II into

$$\begin{aligned}
&2\partial_t g^{\mu\nu} (\Gamma_{\alpha\mu}^\delta \Gamma_{\delta\nu,\beta} + \Gamma_{\beta\mu}^\delta \Gamma_{\delta\nu,\alpha}) - \partial_t g_{\delta\beta} \partial_\alpha g^{\mu\nu} \Gamma_{\mu\nu}^\delta - \partial_t g_{\delta\alpha} \partial_\beta g^{\mu\nu} \Gamma_{\mu\nu}^\delta \\
&\quad + \partial_t g^{\mu\nu} (2\Gamma_{\mu\alpha,\delta} \Gamma_{\nu\beta}^\delta - \partial_\mu \partial_\nu g_{\alpha\beta}) \\
&\quad - \partial_t g_{\delta\beta} g^{\mu\nu} (\partial_\mu \Gamma_{\nu\alpha}^\delta + \Gamma_{\mu\alpha}^\sigma \Gamma_{\nu\sigma}^\delta) - \partial_t g_{\delta\alpha} g^{\mu\nu} (\partial_\mu \Gamma_{\nu\beta}^\delta + \Gamma_{\mu\beta}^\sigma \Gamma_{\nu\sigma}^\delta).
\end{aligned}$$

Adding the $\partial_t g$ terms together with the third term in (7.10) we obtain

$$\begin{aligned}
&\partial_t g^{\mu\nu} (\nabla_\alpha \Gamma_{\mu\nu,\beta} + \nabla_\beta \Gamma_{\mu\nu,\alpha} + 2\Gamma_{\alpha\mu}^\delta \Gamma_{\delta\nu,\beta} + 2\Gamma_{\beta\mu}^\delta \Gamma_{\delta\nu,\alpha} + 2\Gamma_{\mu\alpha,\delta} \Gamma_{\nu\beta}^\delta - \partial_\mu \partial_\nu g_{\alpha\beta}) \\
&= \partial_t g^{\mu\nu} (\partial_\alpha \Gamma_{\mu\nu,\beta} + \partial_\beta \Gamma_{\mu\nu,\alpha} - \partial_\nu (\Gamma_{\beta\mu,\alpha} + \Gamma_{\alpha\mu,\beta}) + 2\Gamma_{\mu\alpha,\delta} \Gamma_{\nu\beta}^\delta - 2\Gamma_{\alpha\beta}^\delta \Gamma_{\mu\nu,\delta}) \\
&= 2\partial_t g^{\mu\nu} R_{\alpha\mu\beta\nu}.
\end{aligned}$$

Finally, using the harmonic coordinate condition $g^{\mu\nu}\Gamma_{\mu\nu}^\delta = 0$, the terms containing the $\partial_t g_{\delta\alpha}$ expression are written as

$$\begin{aligned} & -\partial_t g_{\delta\beta}\partial_\alpha g^{\mu\nu}\Gamma_{\mu\nu}^\delta - \partial_t g_{\delta\alpha}\partial_\beta g^{\mu\nu}\Gamma_{\mu\nu}^\delta - \partial_t g_{\delta\beta}g^{\mu\nu}(\partial_\mu\Gamma_{\nu\alpha}^\delta + \Gamma_{\mu\alpha}^\sigma\Gamma_{\nu\sigma}^\delta) - \partial_t g_{\delta\alpha}g^{\mu\nu}(\partial_\mu\Gamma_{\nu\beta}^\delta + \Gamma_{\mu\beta}^\sigma\Gamma_{\nu\sigma}^\delta) \\ & = \partial_t g_{\delta\beta}\text{Ric}_\alpha^\delta + \partial_t g_{\delta\alpha}\text{Ric}_\beta^\delta. \end{aligned}$$

Hence, the expression $\Delta_g\partial_t g_{\alpha\beta}$ is written as

$$(7.11) \quad \begin{aligned} \Delta_g\partial_t g_{\alpha\beta} &= -\nabla_\alpha(\partial_t g^{\mu\nu}\Gamma_{\mu\nu,\beta}) - \nabla_\beta(\partial_t g^{\mu\nu}\Gamma_{\mu\nu,\alpha}) + \partial_t g_{\delta\beta}\text{Ric}_\alpha^\delta + \partial_t g_{\delta\alpha}\text{Ric}_\beta^\delta \\ & \quad + 2\partial_t g^{\mu\nu}R_{\alpha\mu\beta\nu} - 2\partial_t \widetilde{\text{Ric}}_{\alpha\beta}. \end{aligned}$$

For the last term $-2\partial_t \widetilde{\text{Ric}}_{\alpha\beta}$, using the expression T^2 we have

$$\begin{aligned} (I_{31}) \quad -2\partial_t \widetilde{\text{Ric}}_{\alpha\beta} &= -2\text{Re}(g_{\sigma\beta}T_\alpha^{2,\sigma}\bar{\psi} + T_{\sigma\beta}^1\lambda_\alpha^\sigma\psi + \bar{\lambda}_{\alpha\beta}T_\sigma^{2,\sigma} - g_{\sigma\mu}T_\alpha^{2,\sigma}\bar{\lambda}_\beta^\mu - T_{\sigma\mu}^1\lambda_\alpha^\sigma\bar{\lambda}_\beta^\mu - \bar{\lambda}_{\alpha\sigma}T_\beta^{2,\sigma}) \\ (I_{32}) \quad & + 2\text{Im}(\nabla_\alpha^A\nabla_\beta^A\psi\bar{\psi} + \nabla_\sigma^A\nabla_\alpha^A\psi\bar{\lambda}_{\alpha\beta} - \nabla_\alpha^A\nabla_\sigma^A\psi\bar{\lambda}_\beta^\sigma - \nabla_\beta^A\nabla_\sigma^A\psi\bar{\lambda}_\alpha^\sigma) \\ (I_{33}) \quad & - 2\widetilde{\text{Ric}}_{\beta\gamma}\nabla_\alpha V^\gamma - 2\widetilde{\text{Ric}}_{\alpha\gamma}\nabla_\beta V^\gamma - 2\nabla_\gamma\widetilde{\text{Ric}}_{\alpha\beta}V^\gamma \\ & - 2\text{Re}(\psi\bar{\lambda}_{\alpha\gamma})\text{Im}(\psi\bar{\lambda}_\beta^\gamma) + 2\text{Re}(\lambda_{\alpha\beta}\bar{\lambda}^{\sigma\gamma})\text{Im}(\psi\bar{\lambda}_{\sigma\gamma}). \end{aligned}$$

Next, we compute

$$\begin{aligned} III &:= -\Delta_g(2\text{Im}(\psi\bar{\lambda}_{\alpha\beta}) + \nabla_\alpha V_\beta + \nabla_\beta V_\alpha) \\ &= -2\nabla^\sigma\nabla_\sigma\text{Im}(\psi\bar{\lambda}_{\alpha\beta}) + [\Delta_g, \nabla_\alpha]V_\beta - [\Delta_g, \nabla_\beta]V_\alpha - \nabla_\alpha\Delta_g V_\beta - \nabla_\beta\Delta_g V_\alpha \\ &= -2\nabla^\sigma\nabla_\sigma\text{Im}(\psi\bar{\lambda}_{\alpha\beta}) - \nabla_\beta\text{Ric}_{\alpha\gamma}V^\gamma - \nabla_\alpha\text{Ric}_{\beta\gamma}V^\gamma - 2\nabla_\gamma\text{Ric}_{\alpha\beta}V^\gamma \\ & \quad - \text{Ric}_{\alpha\gamma}\nabla^\gamma V_\beta - \text{Ric}_{\beta\gamma}\nabla^\gamma V_\alpha + 2R_{\alpha\sigma\beta\delta}(\nabla^\sigma V^\delta + \nabla^\delta V^\sigma) - \nabla_\alpha\Delta_g V_\beta - \nabla_\beta\Delta_g V_\alpha \end{aligned}$$

Using the V -equation (2.30) we write the last two terms as

$$\begin{aligned} & -\nabla_\alpha\Delta_g V_\beta - \nabla_\beta\Delta_g V_\alpha \\ &= 2\nabla_\alpha\nabla_\sigma\text{Im}(\psi\bar{\lambda}_\beta^\sigma) + 2\nabla_\beta\nabla_\sigma\text{Im}(\psi\bar{\lambda}_\alpha^\sigma) + \nabla_\alpha\text{Ric}_{\sigma\beta}V^\sigma + \nabla_\beta\text{Ric}_{\sigma\alpha}V^\sigma \\ & \quad + \text{Ric}_{\sigma\beta}\nabla_\alpha V^\sigma + \text{Ric}_{\sigma\alpha}\nabla_\beta V^\sigma + \nabla_\alpha(\widetilde{\partial_t g^{\mu\nu}}\Gamma_{\mu\nu,\beta}) + \nabla_\beta(\widetilde{\partial_t g^{\mu\nu}}\Gamma_{\mu\nu,\alpha}), \end{aligned}$$

where $\widetilde{\partial_t g^{\mu\nu}}$ denotes the expression

$$\widetilde{\partial_t g^{\mu\nu}} := \partial_t g^{\mu\nu} - T^{1,\mu\nu}.$$

We then add I_{31} together with $\nabla^2\text{Im}(\psi\lambda)$ in III to get

$$\begin{aligned} & I_{31} - 2\nabla^\sigma\nabla_\sigma\text{Im}(\psi\bar{\lambda}_{\alpha\beta}) + 2\nabla_\alpha\nabla_\sigma\text{Im}(\psi\bar{\lambda}_\beta^\sigma) + 2\nabla_\beta\nabla_\sigma\text{Im}(\psi\bar{\lambda}_\alpha^\sigma) \\ &= -2\text{Ric}_{\beta\delta}\text{Im}(\psi\bar{\lambda}_\alpha^\delta) + 2R_{\alpha\sigma\beta\delta}\text{Im}(\psi\bar{\lambda}^{\sigma\delta}) + 2\text{Re}(\lambda_\alpha^\sigma\bar{\psi})\text{Im}(\lambda_{\sigma\mu}\bar{\lambda}_\beta^\mu). \end{aligned}$$

The last term and I_{33} can be further written as

$$\begin{aligned} & 2\text{Re}(\lambda_\alpha^\sigma\bar{\psi})\text{Im}(\lambda_{\sigma\mu}\bar{\lambda}_\beta^\mu) + I_{33} \\ &= 2R_{\alpha\sigma\beta\delta}\text{Im}(\psi\bar{\lambda}^{\sigma\delta}) - 2\text{Ric}_{\alpha\delta}\text{Im}(\psi\bar{\lambda}_\beta^\delta). \end{aligned}$$

Hence, given the expressions of I_3 and III , we obtain

$$\begin{aligned} & I_3 + III \\ &= -2 \operatorname{Re}(g_{\sigma\beta} T_\alpha^{2,\sigma} \bar{\psi} + T_{\sigma\beta}^1 \lambda_\alpha^\sigma \psi + \bar{\lambda}_{\alpha\beta} T_\sigma^{2,\sigma} - g_{\sigma\mu} T_\alpha^{2,\sigma} \bar{\lambda}_\beta^\mu - T_{\sigma\mu}^1 \lambda_\alpha^\sigma \bar{\lambda}_\beta^\mu - \bar{\lambda}_{\alpha\sigma} T_\beta^{2,\sigma}) \\ & \quad - \operatorname{Ric}^\sigma{}_\beta \widetilde{\partial_t g_{\alpha\sigma}} - \operatorname{Ric}^\sigma{}_\alpha \widetilde{\partial_t g_{\beta\sigma}} - 2R_{\alpha\sigma\beta\delta} \widetilde{\partial_t g^{\sigma\delta}} + \nabla_\alpha (\widetilde{\partial_t g^{\mu\nu}} \Gamma_{\mu\nu,\beta}) + \nabla_\beta (\widetilde{\partial_t g^{\mu\nu}} \Gamma_{\mu\nu,\alpha}), \end{aligned}$$

which combined with (7.11) yields the T^1 -equation (7.9). \square

The equation for T^2 . This has the form

$$(7.12) \quad \begin{cases} \nabla^{A,\alpha} T_\alpha^{2,\sigma} = i g^{\sigma\mu} \psi T_\mu^3 - i \lambda^{\beta\sigma} T_\beta^3 + g^{\sigma\delta} \lambda^{\alpha\beta} (-\nabla_\alpha T_{\beta\delta}^1 + \frac{1}{2} \nabla_\delta T_{\alpha\beta}^1) \\ \quad - T^{1,\alpha\beta} (\nabla_\beta^A \lambda_\alpha^\sigma + \Gamma_{\alpha\beta}^\mu \lambda_\mu^\sigma) + T^{1,\sigma\mu} \nabla_\mu^A \psi, \\ \nabla_\alpha^A T_\beta^{2,\sigma} - \nabla_\beta^A T_\alpha^{2,\sigma} = \frac{1}{2} g^{\sigma\gamma} [-\lambda_\beta^\mu (\nabla_\alpha T_{\mu\gamma}^1 + \nabla_\mu T_{\alpha\gamma}^1 - \nabla_\gamma T_{\alpha\mu}^1) \\ \quad + \lambda_\alpha^\mu (\nabla_\beta T_{\mu\gamma}^1 + \nabla_\mu T_{\beta\gamma}^1 - \nabla_\gamma T_{\beta\mu}^1)] - i T_\alpha^3 \lambda_\beta^\sigma + i T_\beta^3 \lambda_\alpha^\sigma. \end{cases}$$

We compute the divergence of T^2 in (7.12) first. Applying $\nabla^{A,\alpha}$ to $T_\alpha^{2,\sigma}$, we have

$$\begin{aligned} \nabla^{A,\alpha} T_\alpha^{2,\sigma} &= [\nabla^{A,\alpha}, \partial_t^B - V^\gamma \nabla_\gamma^A] \lambda_\alpha^\sigma + [\partial_t^B - V^\gamma \nabla_\gamma^A, \nabla^{A,\sigma}] \psi + \nabla^{A,\sigma} (\partial_t^B - V^\gamma \nabla_\gamma^A) \psi \\ & \quad + \nabla^{A,\alpha} (\lambda_\alpha^\gamma \operatorname{Im}(\psi \bar{\lambda}_\gamma^\sigma)) - i \nabla^{A,\alpha} \nabla_\alpha^A \nabla^{A,\sigma} \psi \\ & \quad + \nabla_\gamma^A \psi \nabla^\gamma V^\sigma - \nabla^{A,\sigma} \lambda_{\alpha\gamma} \nabla^\alpha V^\gamma + \lambda^{\alpha\gamma} \nabla_\alpha \nabla_\gamma V^\sigma - \lambda_\gamma^\sigma \Delta_g V^\gamma. \end{aligned}$$

Three of the terms on the right-hand side are written as

$$\begin{aligned} & [\nabla^{A,\alpha}, \partial_t^B - V^\gamma \nabla_\gamma^A] \lambda_\alpha^\sigma - \nabla^{A,\sigma} \lambda_{\alpha\gamma} \nabla^\alpha V^\gamma + \lambda^{\alpha\gamma} \nabla_\alpha \nabla_\gamma V^\sigma \\ &= g^{\alpha\beta} (\nabla_\beta \partial_t \lambda_\alpha^\sigma - \partial_t \nabla_\beta \lambda_\alpha^\sigma) + i g^{\alpha\beta} (\nabla_\beta B - \partial_t A_\beta) \lambda_\alpha^\sigma - \partial_t g^{\alpha\beta} \nabla_\beta^A \lambda_\alpha^\sigma \\ & \quad + \lambda^{\alpha\gamma} \nabla_\alpha \nabla_\gamma V^\sigma - 2 \nabla^{A,\sigma} \lambda_{\alpha\gamma} \nabla^\alpha V^\gamma - V^\gamma [\nabla^\alpha, \nabla_\gamma] \lambda_\alpha^\sigma - i V_\gamma \mathbf{F}^{\alpha\gamma} \lambda_\alpha^\sigma \\ &= -\partial_t g^{\alpha\beta} (\nabla_\alpha^A \lambda_{\beta\sigma} + \Gamma_{\alpha\beta}^\mu \lambda_\mu^\sigma) - \partial_t \Gamma_{\beta\mu}^\sigma \lambda^{\beta\mu} + \lambda^{\alpha\gamma} \nabla_\alpha \nabla_\gamma V^\sigma \\ & \quad - i (\partial_t A_\beta - \nabla_\beta B) \lambda^{\beta\sigma} - i V_\gamma \mathbf{F}^{\alpha\gamma} \lambda_\alpha^\sigma - 2 \nabla^{A,\sigma} \lambda_{\alpha\gamma} \nabla^\alpha V^\gamma - V^\gamma [\nabla^\alpha, \nabla_\gamma] \lambda_\alpha^\sigma \end{aligned}$$

We can further use T^1 to rewrite the last two terms on the first line above as

$$\begin{aligned} & -\partial_t \Gamma_{\alpha\beta}^\sigma \lambda^{\alpha\beta} + \lambda^{\alpha\gamma} \nabla_\alpha \nabla_\gamma V^\sigma \\ &= -\partial_t g^{\sigma\delta} \Gamma_{\alpha\beta,\delta} \lambda^{\alpha\beta} - g^{\sigma\delta} \partial_t (\partial_\alpha g_{\beta\delta} - \frac{1}{2} \partial_\delta g_{\alpha\beta}) \lambda^{\alpha\beta} + \lambda^{\alpha\gamma} \nabla_\alpha \nabla_\gamma V^\sigma \\ &= g^{\sigma\delta} \lambda^{\alpha\beta} (\partial_t g_{\mu\delta} \Gamma_{\alpha\beta}^\mu - \partial_\alpha \partial_t g_{\beta\delta} + \frac{1}{2} \partial_\delta \partial_t g_{\alpha\beta}) + \lambda^{\alpha\gamma} \nabla_\alpha \nabla_\gamma V^\sigma \\ &= g^{\sigma\delta} \lambda^{\alpha\beta} (-\nabla_\alpha \partial_t g_{\beta\delta} + \frac{1}{2} \nabla_\delta \partial_t g_{\alpha\beta}) + \lambda^{\alpha\gamma} \nabla_\alpha \nabla_\gamma V^\sigma \\ &= \lambda_{\mu\nu} (\nabla^\mu T^{1,\nu\sigma} - \frac{1}{2} \nabla^\sigma T^{1,\mu\nu}) \\ & \quad + \lambda^{\alpha\beta} [-2 \nabla_\alpha \operatorname{Im}(\psi \bar{\lambda}_\beta^\sigma) + \nabla^\sigma \operatorname{Im}(\psi \bar{\lambda}_{\alpha\beta}) - [\nabla_\alpha, \nabla^\sigma] V_\beta] \end{aligned}$$

and the following term as

$$-i (\partial_t A_\beta - \nabla_\beta B) \lambda^{\beta\sigma} - i V_\gamma \mathbf{F}^{\alpha\gamma} \lambda_\alpha^\sigma = -i \lambda^{\beta\sigma} T_\beta^3 - i \lambda^{\beta\sigma} \operatorname{Re}(\lambda_\beta^\gamma \overline{\nabla_\gamma^A \psi}).$$

Similarly, we compute the second commutator by

$$\begin{aligned} [\partial_t^B - V^\gamma \nabla_\gamma^A, \nabla^{A,\sigma}] \psi + \nabla_\gamma^A \psi \nabla^\gamma V^\sigma &= \partial_t g^{\sigma\mu} \nabla_\mu^A \psi + i g^{\sigma\mu} \psi T_\mu^3 + i \psi \operatorname{Re}(\lambda^{\sigma\gamma} \overline{\nabla_\gamma^A \psi}) \\ &\quad + \nabla_\gamma^A \psi (\nabla^\gamma V^\sigma + \nabla^\sigma V^\gamma). \end{aligned}$$

Hence, using $T_\alpha^{2,\alpha}$ and the V equation (2.30) we reorganize the expression of $\nabla^{A,\alpha} T_\alpha^{2,\sigma}$ and obtain

$$\begin{aligned} \nabla^{A,\alpha} T_\alpha^{2,\sigma} &= i g^{\sigma\mu} \psi T_\mu^3 - i \lambda^{\beta\sigma} T_\beta^3 + g^{\sigma\delta} \lambda^{\alpha\beta} (-\nabla_\alpha T_{\beta\delta}^1 + \frac{1}{2} \nabla_\delta T_{\alpha\beta}^1) \\ &\quad - \partial_t g^{\alpha\beta} (\nabla_\beta^A \lambda_\alpha^\sigma + \Gamma_{\alpha\beta}^\mu \lambda_\mu^\sigma) + \lambda^{\alpha\beta} [-2 \nabla_\alpha \operatorname{Im}(\psi \bar{\lambda}_\beta^\sigma) + \nabla^\sigma \operatorname{Im}(\psi \bar{\lambda}_{\alpha\beta})] \\ &\quad - i \lambda^{\beta\sigma} \operatorname{Re}(\lambda_\beta^\gamma \overline{\nabla_\gamma^A \psi}) + \partial_t g^{\sigma\mu} \nabla_\mu^A \psi + i \psi \operatorname{Re}(\lambda^{\sigma\gamma} \overline{\nabla_\gamma^A \psi}) \\ &\quad - \nabla^{A,\sigma} (\lambda_\alpha^\gamma \operatorname{Im}(\psi \bar{\lambda}_\gamma^\alpha)) + \nabla^{A,\alpha} (\lambda_\alpha^\gamma \operatorname{Im}(\psi \bar{\lambda}_\gamma^\sigma)) \\ &\quad - i \operatorname{Ric}^\sigma{}_\delta \nabla^{A,\delta} \psi - \nabla_\alpha \mathbf{F}^{\sigma\alpha} \psi - 2 \mathbf{F}^{\sigma\alpha} \nabla_\alpha^A \psi \\ &\quad + \nabla_\gamma^A \psi (\nabla^\gamma V^\sigma + \nabla^\sigma V^\gamma) - 2 \nabla^{A,\sigma} \lambda_{\alpha\gamma} \nabla^\alpha V^\gamma \\ &\quad - 2 \lambda_\gamma^\sigma \nabla_\alpha \operatorname{Im}(\lambda^{\alpha\gamma} \bar{\psi}) + \lambda_\gamma^\sigma \widetilde{\partial_t g^{\alpha\beta}} \Gamma_{\alpha\beta}^\gamma. \end{aligned}$$

Using $T_\alpha^{2,\alpha}$ and the V -equation (2.30), we have

$$\begin{aligned} &\lambda^{\alpha\beta} [-2 \nabla_\alpha \operatorname{Im}(\psi \bar{\lambda}_\beta^\sigma) + \nabla^\sigma \operatorname{Im}(\psi \bar{\lambda}_{\alpha\beta})] - i \lambda^{\beta\sigma} \operatorname{Re}(\lambda_\beta^\gamma \overline{\nabla_\gamma^A \psi}) \\ &\quad + i \psi \operatorname{Re}(\lambda^{\sigma\gamma} \overline{\nabla_\gamma^A \psi}) + \nabla^{A,\alpha} (\lambda_\alpha^\gamma \operatorname{Im}(\psi \bar{\lambda}_\gamma^\sigma)) - i \operatorname{Ric}^\sigma{}_\delta \nabla^{A,\delta} \psi - \nabla_\alpha \mathbf{F}^{\sigma\alpha} \psi - 2 \mathbf{F}^{\sigma\alpha} \nabla_\alpha^A \psi \\ &\quad + 2 \lambda_\gamma^\sigma \nabla^\alpha \operatorname{Im}(\psi \bar{\lambda}_\alpha^\gamma) - \nabla^{A,\sigma} (\lambda_\alpha^\gamma \operatorname{Im}(\psi \bar{\lambda}_\gamma^\alpha)) \\ &= 2 \nabla_\gamma^A \psi \operatorname{Im}(\psi \bar{\lambda}^{\sigma\gamma}) - 2 \nabla^{A,\sigma} \lambda_{\alpha\beta} \operatorname{Im}(\psi \bar{\lambda}^{\alpha\beta}) \end{aligned}$$

Combining these two expressions, we obtain

$$\begin{aligned} \nabla^{A,\alpha} T_\alpha^{2,\sigma} &= i g^{\sigma\mu} \psi T_\mu^3 - i \lambda^{\beta\sigma} T_\beta^3 + g^{\sigma\delta} \lambda^{\alpha\beta} (-\nabla_\alpha T_{\beta\delta}^1 + \frac{1}{2} \nabla_\delta T_{\alpha\beta}^1) \\ &\quad - T^{1,\alpha\beta} (\nabla_\beta^A \lambda_\alpha^\sigma + \Gamma_{\alpha\beta}^\mu \lambda_\mu^\sigma) + T^{1,\sigma\mu} \nabla_\mu^A \psi + \nabla^{A,\sigma} T_\alpha^{2,\alpha}. \end{aligned}$$

Next we compute the *curl* of T^2 in (7.12). By T^2 we have

$$\begin{aligned} &\nabla_\alpha^A T_\beta^{2,\sigma} - \nabla_\beta^A T_\alpha^{2,\sigma} \\ &= [\nabla_\alpha^A, \partial_t^B - V^\gamma \nabla_\gamma^A] \lambda_\beta^\sigma - [\nabla_\beta^A, \partial_t^B - V^\gamma \nabla_\gamma^A] \lambda_\alpha^\sigma + \lambda_\beta^\gamma \nabla_\alpha \operatorname{Im}(\psi \bar{\lambda}_\gamma^\sigma) - \lambda_\alpha^\gamma \nabla_\beta \operatorname{Im}(\psi \bar{\lambda}_\gamma^\sigma) \\ &\quad - i [\nabla_\alpha^A, \nabla_\beta^A] \nabla^{A,\sigma} \psi + \lambda_\beta^\gamma \nabla_\alpha \nabla_\gamma V^\sigma - \lambda_\alpha^\gamma \nabla_\beta \nabla_\gamma V^\sigma \\ &\quad - \lambda_\gamma^\sigma [\nabla_\alpha, \nabla_\beta] V^\gamma - \nabla_\gamma^A \lambda_\alpha^\sigma \nabla_\beta V^\gamma + \nabla_\gamma^A \lambda_\beta^\sigma \nabla_\alpha V^\gamma. \end{aligned}$$

We use T^1 and T^3 to rewrite six of the terms on the right-hand side as

$$\begin{aligned} &[\nabla_\alpha^A, \partial_t^B - V^\gamma \nabla_\gamma^A] \lambda_\beta^\sigma - [\nabla_\beta^A, \partial_t^B - V^\gamma \nabla_\gamma^A] \lambda_\alpha^\sigma + \lambda_\beta^\gamma \nabla_\alpha \nabla_\gamma V^\sigma - \lambda_\alpha^\gamma \nabla_\beta \nabla_\gamma V^\sigma \\ &\quad - \nabla_\gamma^A \lambda_\alpha^\sigma \nabla_\beta V^\gamma + \nabla_\gamma^A \lambda_\beta^\sigma \nabla_\alpha V^\gamma \\ &= \frac{1}{2} g^{\sigma\gamma} [-\lambda_\beta^\mu (\nabla_\alpha T_{\mu\gamma}^1 + \nabla_\mu T_{\alpha\gamma}^1 - \nabla_\gamma T_{\alpha\mu}^1) + \lambda_\alpha^\mu (\nabla_\beta T_{\mu\gamma}^1 + \nabla_\mu T_{\beta\gamma}^1 - \nabla_\gamma T_{\beta\mu}^1)] \\ &\quad - iT_\alpha^3 \lambda_\beta^\sigma + iT_\beta^3 \lambda_\alpha^\sigma \\ &\quad - \lambda_\beta^\mu \nabla_\alpha \operatorname{Im}(\psi \bar{\lambda}_\mu^\sigma) + \lambda_\alpha^\mu \nabla_\beta \operatorname{Im}(\psi \bar{\lambda}_\mu^\sigma) + I_1 + I_2, \end{aligned}$$

where I_1 and I_2 are

$$\begin{aligned} I_1 &:= \lambda_\beta^\mu (-\nabla_\mu \operatorname{Im}(\psi \bar{\lambda}_\alpha^\sigma) + \nabla^\sigma \operatorname{Im}(\psi \bar{\lambda}_{\alpha\mu})) - \lambda_\alpha^\mu (-\nabla_\mu \operatorname{Im}(\psi \bar{\lambda}_\beta^\sigma) + \nabla^\sigma \operatorname{Im}(\psi \bar{\lambda}_{\beta\mu})) \\ &\quad - i \operatorname{Re}(\lambda_\alpha^\gamma \overline{\nabla_\gamma^A \psi}) \lambda_\beta^\sigma + i \operatorname{Re}(\lambda_\beta^\gamma \overline{\nabla_\gamma^A \psi}) \lambda_\alpha^\sigma, \\ I_2 &:= \frac{1}{2} \lambda_\beta^\mu (R_{\alpha\mu\sigma\delta} + R_{\alpha\mu\delta}^\sigma + R_{\mu\alpha\delta}^\sigma) V^\delta - \frac{1}{2} \lambda_\alpha^\mu (R_{\beta\mu\sigma\delta} + R_{\beta\mu\delta}^\sigma + R_{\mu\beta\delta}^\sigma) V^\delta \\ &\quad - V^\gamma R_{\alpha\gamma\sigma\delta} \lambda_\beta^\delta - V^\gamma R_{\alpha\gamma\beta\delta} \lambda_\sigma^\delta + V^\gamma R_{\beta\gamma\sigma\delta} \lambda_\alpha^\delta + V^\gamma R_{\beta\gamma\alpha\delta} \lambda_\sigma^\delta. \end{aligned}$$

Then we use Bianchi identities and compatibility conditions to compute I_1 and I_2 by

$$I_1 = i[\nabla_\alpha^A, \nabla_\beta^A] \nabla^{A,\sigma} \psi$$

and

$$I_2 = V^\gamma R_{\beta\gamma\sigma\delta} \lambda_\alpha^\delta + V^\gamma R_{\beta\gamma\alpha\delta} \lambda_\sigma^\delta = \lambda_\gamma^\sigma [\nabla_\alpha, \nabla_\beta] V^\gamma.$$

Hence, we obtain

$$\begin{aligned} \nabla_\alpha^A T_\beta^{2,\sigma} - \nabla_\beta^A T_\alpha^{2,\sigma} &= \frac{1}{2} g^{\sigma\gamma} [-\lambda_\beta^\mu (\nabla_\alpha T_{\mu\gamma}^1 + \nabla_\mu T_{\alpha\gamma}^1 - \nabla_\gamma T_{\alpha\mu}^1) \\ &\quad + \lambda_\alpha^\mu (\nabla_\beta T_{\mu\gamma}^1 + \nabla_\mu T_{\beta\gamma}^1 - \nabla_\gamma T_{\beta\mu}^1)] - iT_\alpha^3 \lambda_\beta^\sigma + iT_\beta^3 \lambda_\alpha^\sigma. \end{aligned}$$

This completes the derivation of (7.12). \square

The equation for T^3 . This has the form

$$\begin{cases} \nabla^\alpha T_\alpha^3 = -T^{1,\alpha\beta} \partial_\alpha A_\beta, \\ \nabla_\alpha T_\beta^3 - \nabla_\beta T_\alpha^3 = \operatorname{Im}(T_\alpha^{2,\sigma} \bar{\lambda}_{\sigma\beta} + \lambda_\alpha^\sigma \overline{T_{\sigma\beta}^2}). \end{cases}$$

Applying ∇^α to T_α^3 , we then use the Coulomb condition $\nabla^\alpha A_\alpha = 0$ and the B -equation (2.33) to get

$$\begin{aligned} \nabla^\alpha T_\alpha^3 &= \nabla^\alpha \partial_t A_\alpha - \Delta_g B - \nabla^\alpha \operatorname{Re}(\lambda_\alpha^\sigma \overline{\nabla_\sigma^A \psi} + i \lambda_\alpha^\sigma \bar{\lambda}_{\sigma\gamma} V^\gamma) \\ &= g^{\alpha\beta} \partial_\beta \partial_t A_\alpha + \partial_t g^{\beta\gamma} \partial_\beta A_\gamma - T^{1,\beta\gamma} \partial_\beta A_\gamma = -T^{1,\alpha\beta} \partial_\alpha A_\beta. \end{aligned}$$

The *curl* of T^3 is obtained by (2.13) directly. \square

\square

8. THE RECONSTRUCTION OF THE FLOW

In this last section we close the circle of ideas in this paper, and prove that one can start from the good gauge solution given by Theorem 2.7, and reconstruct the flow at the level of d -dimensional embedded submanifolds. For completeness, we provide here another, more complete statement of our main theorem:

Theorem 8.1 (Small data local well-posedness). *Let $s > \frac{d}{2}$, $d \geq 4$. Consider the skew mean curvature flow (1.1) for maps F from \mathbb{R}^d to the Euclidean space $(\mathbb{R}^{d+2}, g_{\mathbb{R}^{d+2}})$ with initial data Σ_0 which, in some coordinates, has a metric g_0 satisfying $\|\partial_x(g_0 - I_d)\|_{H^s} \leq \epsilon_0$ and mean curvature $\|\mathbf{H}_0\|_{H^s(\Sigma_0)} \leq \epsilon_0$.*

If $\epsilon_0 > 0$ is sufficiently small, then there exists a unique solution

$$F : \mathbb{R}^d \times [0, 1] \rightarrow (\mathbb{R}^{d+2}, g_{\mathbb{R}^{d+2}})$$

which, when represented in harmonic coordinates, has regularity

$$\partial_x^2 F, \partial_t F \in C([0, 1]; H^s(\mathbb{R}^d)).$$

and induced metric and mean curvature

$$\partial_x g \in C([0, 1]; H^{s+1}(\mathbb{R}^d)), \quad \mathbf{H} \in C([0, 1]; H^s(\mathbb{R}^d)).$$

In addition the mean curvature satisfies the bounds

$$\|\psi\|_{L^2 \mathbf{X}^s} + \|(\lambda, h, V, A, B)\|_{\mathcal{E}^s} \lesssim \|\psi_0\|_{H^s}.$$

where ψ and λ are expressed using the Coulomb gauge in the normal bundle $N\Sigma_t$.

We complement the theorem with the following remarks:

Remark 8.1.1. Here uniqueness should be interpreted in two steps:

- (i) If $s > \frac{d}{2} + 1$ then we have a direct uniqueness statement for solutions F which in some coordinate system are continuous with values in H^{s+2} .
- (ii) For smaller s , then our solutions can be identified as the unique limits of smooth solutions expressed in harmonic coordinates.

Remark 8.1.2. The only role of the smallness condition on the metric is to exclude large nonflat minimal surfaces; the topology we use there is less essential as long as some critical norm of F is made small. This guarantees that (i) we can find harmonic coordinates on the surface Σ_0 and a Coulomb frame in the normal bundle and (ii) in harmonic coordinates and the Coulomb gauge the surface is uniquely (and smoothly) determined by the mean curvature ψ up to rigid rotations.

We do this in several steps:

8.1. The starting point. Our evolution begins at time $t = 0$, where we need to represent the initial submanifold as parametrized with global harmonic coordinates, represented via the map $F : \mathbb{R}^d \rightarrow \mathbb{R}^{d+2}$, and to construct a Coulomb frame in the normal bundle, leading to the complex mean curvature function ψ . This is the goal of this subsection, which is carried out in Proposition 8.2.

Once this is done, we have the frame F_α in the tangent space and the frame m in the normal bundle. In turn, as described in Section 2, these generate the metric g , the second fundamental form λ with trace ψ and the connection A , all at the initial time $t = 0$.

Moving forward in time, Theorem 2.7 provides us with the time evolution of ψ via the Schrödinger flow (2.35), as well as the functions (λ, g, V, A, B) satisfying the elliptic system (2.36) together with the constraints (2.4), (2.8), (2.15), (2.13), (2.16) and (2.19) and the time evolutions (2.26), (2.31) and (2.32). The objective of the rest of this section is then to use these functions in order to reconstruct the map F which describes the manifold F at later times.

We now return to the question of constructing the harmonic coordinates at the initial time. In order to state the following proposition, we define some notations. Let $F : \mathbb{R}_x^d \rightarrow (\mathbb{R}^{d+2}, g_{\mathbb{R}^{d+2}})$ be an immersion with induced metric $g(x)$. For any change of coordinate $y = x + \phi(x)$, we denote

$$\tilde{F}(y) = F(x(y)),$$

and its induced metric $\tilde{g}_{\alpha\beta}(y) = \langle \partial_{y_\alpha} \tilde{F}, \partial_{y_\beta} \tilde{F} \rangle$. We also denote its Christoffel symbol as $\tilde{\Gamma}$ and $\tilde{h}(y) = \tilde{g}(y) - I_d$.

Proposition 8.2. *Let $d \geq 3$, $s > \frac{d}{2}$, and $F : (\mathbb{R}_x^d, g) \rightarrow (\mathbb{R}^{d+2}, g_{\mathbb{R}^{d+2}})$ be an immersion with induced metric $g = I_d + h$. Assume that $\nabla h(x)$ and mean curvature \mathbf{H} are small in $H^s(dx)$, namely*

$$\|\partial_x h\|_{H^s} \leq \epsilon_0, \quad \|\mathbf{H}\|_{H^s} \leq \epsilon_0.$$

Then there exists a unique change of coordinates $y = x + \phi(x)$ with $\lim_{x \rightarrow \infty} \phi(x) = 0$ and $\nabla \phi$ uniformly small, such that the new coordinates $\{y_1, \dots, y_d\}$ are global harmonic coordinates, namely,

$$\tilde{g}^{\alpha\beta}(y) \tilde{\Gamma}_{\alpha\beta}^\gamma(y) = 0, \quad \text{for any } y \in \mathbb{R}^d.$$

Moreover,

$$(8.1) \quad \|\nabla^2 \phi(x)\|_{H^s(dx)} \lesssim \|\nabla h(x)\|_{H^s(dx)},$$

and, in the new coordinates $\{y_1, \dots, y_d\}$,

$$(8.2) \quad \|\partial_y \tilde{h}\|_{H^s(dy)} \lesssim \|\partial_x h\|_{H^s(dx)}.$$

In addition, for the mean curvature we have equivalent norms,

$$(8.3) \quad \|\mathbf{H}\|_{H^s(dy)} \lesssim \|\mathbf{H}\|_{H^s(dx)},$$

and the bound for complex scalar mean curvature ψ in the Coulomb gauge

$$(8.4) \quad \|\psi\|_{H^s} \lesssim \epsilon_0.$$

Proof. Step 1: Derivation of the ϕ -equations.

We make the following change of coordinates such that the $\{y_1, \dots, y_d\}$ is a global harmonic coordinate

$$\begin{array}{ccccc} \mathbb{R}^d & \longrightarrow & \mathbb{R}^d & \longrightarrow & \mathbb{R}^{d+2} \\ y & \longmapsto & x & \longmapsto & F(x(y)) = \tilde{F}(y) \end{array}$$

where $x + \phi(x) = y$ with $\lim_{x \rightarrow \infty} \phi(x) = 0$ and $\nabla \phi$ small.

To determine the function ϕ , we perform a few computations. For any vector $f = (f_1, \dots, f_d)$, we denote

$$\frac{\partial f}{\partial x} = \begin{pmatrix} \frac{\partial f_1}{\partial x_1} & \cdots & \frac{\partial f_1}{\partial x_d} \\ \vdots & \ddots & \vdots \\ \frac{\partial f_d}{\partial x_1} & \cdots & \frac{\partial f_d}{\partial x_d} \end{pmatrix}.$$

Then we have

$$\frac{\partial x}{\partial y} + \frac{\partial \phi}{\partial x} \frac{\partial x}{\partial y} = I_d.$$

This implies that

$$\frac{\partial x}{\partial y} = I_d - \frac{\partial \phi}{\partial x} + \mathcal{C}(x),$$

where the matrix $\mathcal{C}(x)$ is a higher order term which satisfies

$$\mathcal{C}(x) = \left(\frac{\partial \phi}{\partial x}\right)^2 - \mathcal{C}(x) \frac{\partial \phi}{\partial x},$$

or, equivalently, it is given by

$$\mathcal{C}(x) = \left(\frac{\partial \phi}{\partial x}\right)^2 \left(I - \frac{\partial \phi}{\partial x}\right)^{-1}.$$

We denote

$$\mathcal{P}_\alpha^\mu := -\frac{\partial\phi_\mu}{\partial x_\alpha} + \mathcal{C}_{\mu\alpha}(x).$$

Since $\tilde{F}(y) = F(x(y))$, then we have

$$(8.5) \quad \begin{aligned} \tilde{g}_{\alpha\beta}(y) &= \left\langle \frac{\partial\tilde{F}}{\partial y_\alpha}, \frac{\partial\tilde{F}}{\partial y_\beta} \right\rangle = \left\langle \frac{\partial F}{\partial x_\mu} \frac{\partial x_\mu}{\partial y_\alpha}, \frac{\partial F}{\partial x_\nu} \frac{\partial x_\nu}{\partial y_\beta} \right\rangle \\ &= g_{\mu\nu}(x) (\delta_\alpha^\mu - \partial_\alpha\phi_\mu + \mathcal{C}_{\mu\alpha}) (\delta_\beta^\nu - \partial_\beta\phi_\nu + \mathcal{C}_{\nu\beta}) \end{aligned}$$

and

$$(8.6) \quad \tilde{g}^{\alpha\beta}(y) = g^{\mu\nu} \frac{\partial y_\alpha}{\partial x_\mu} \frac{\partial y_\beta}{\partial x_\nu} = g^{\mu\nu} (\delta_\mu^\alpha + \partial_\mu\phi_\alpha) (\delta_\nu^\beta + \partial_\nu\phi_\beta).$$

We also have

$$(8.7) \quad \frac{\partial\tilde{g}_{\alpha\beta}(y)}{\partial y_\gamma} = \frac{\partial\tilde{g}_{\alpha\beta}(y)}{\partial x_m} \frac{\partial x_m}{\partial y_\gamma} = -g_{\mu\beta} \partial_{\alpha\gamma}^2 \phi_\mu - g_{\alpha\nu} \partial_{\beta\gamma}^2 \phi_\nu + \partial_\gamma g_{\alpha\beta} + \mathcal{K}_{\alpha\beta,\gamma},$$

where the higher order terms $\mathcal{K}_{\alpha\beta,\gamma}$ are defined as

$$\begin{aligned} \mathcal{K}_{\alpha\beta,\gamma} &:= -g_{\mu\nu} \partial_{\alpha\gamma}^2 \phi_\mu \mathcal{P}_\beta^\nu + g_{\mu\nu} \partial_\gamma \mathcal{C}_{\mu\alpha} \frac{\partial x_\nu}{\partial y_\beta} - g_{\mu\nu} \mathcal{P}_\alpha^\mu \partial_{\beta\gamma}^2 \phi_\nu + g_{\mu\nu} \frac{\partial x_\mu}{\partial y_\alpha} \partial_\gamma \mathcal{C}_{\nu\beta} \\ &\quad + \partial_\gamma g_{\alpha\nu} \mathcal{P}_\beta^\nu + \partial_\gamma g_{\mu\nu} \mathcal{P}_\alpha^\mu \frac{\partial x_\nu}{\partial y_\beta} + \partial_{x_m} [g_{\mu\nu}(x) \frac{\partial x_\mu}{\partial y_\alpha} \frac{\partial x_\nu}{\partial y_\beta}] \mathcal{P}_\gamma^m. \end{aligned}$$

The relation $\tilde{g}^{\alpha\beta} \tilde{\Gamma}_{\alpha\beta,\gamma} = 0$ combined with (8.6) and (8.7) implies that

$$\begin{aligned} 0 &= g^{mn} (\delta_m^\alpha + \partial_m\phi_\alpha) (\delta_n^\beta + \partial_n\phi_\beta) \left[-g_{\mu\beta} \partial_{\alpha\gamma}^2 \phi_\mu - g_{\gamma\nu} \partial_{\beta\alpha}^2 \phi_\nu + \partial_\alpha g_{\gamma\beta} + \mathcal{K}_{\gamma\beta,\alpha} \right. \\ &\quad \left. + \frac{1}{2} g_{\mu\beta} \partial_{\alpha\gamma}^2 \phi_\mu + \frac{1}{2} g_{\alpha\nu} \partial_{\beta\gamma}^2 \phi_\nu - \frac{1}{2} \partial_\gamma g_{\alpha\beta} - \frac{1}{2} \mathcal{K}_{\alpha\beta,\gamma} \right]. \end{aligned}$$

This gives the elliptic equations of ϕ ,

$$(8.8) \quad \Delta\phi_\gamma = \mathbf{Non}_\gamma(g, \phi),$$

with the boundary condition $\lim_{x \rightarrow \infty} \phi(x) = 0$, where the nonlinearities $\mathbf{Non}_\gamma(g, \phi)$ are given by

$$\begin{aligned} \mathbf{Non}_\gamma(g, \phi) &:= -h_{\gamma\nu} \Delta\phi_\nu - h^{\alpha\beta} g_{\gamma\nu} \partial_{\alpha\beta}^2 \phi_\nu + g^{\alpha\beta} (\Gamma_{\alpha\beta,\gamma} + \mathcal{K}_{\gamma\beta,\alpha} - \frac{1}{2} \mathcal{K}_{\alpha\beta,\gamma}) \\ &\quad + g^{mn} (\delta_m^\alpha \partial_n\phi_\beta + \partial_m\phi_\alpha \delta_n^\beta + \partial_m\phi_\alpha \partial_n\phi_\beta) \left[-g_{\mu\beta} \partial_{\alpha\gamma}^2 \phi_\mu - g_{\gamma\nu} \partial_{\beta\alpha}^2 \phi_\nu \right. \\ &\quad \left. + \frac{1}{2} g_{\mu\beta} \partial_{\alpha\gamma}^2 \phi_\mu + \frac{1}{2} g_{\alpha\nu} \partial_{\beta\gamma}^2 \phi_\nu + \Gamma_{\alpha\beta,\gamma} + \mathcal{K}_{\gamma\beta,\alpha} - \frac{1}{2} \mathcal{K}_{\alpha\beta,\gamma} \right]. \end{aligned}$$

Step 2: Solve the ϕ -equations (8.8). By the contraction principle, the existence and uniqueness of solution of (8.8) and the bound (8.1) are obtained by the following Lemma.

Lemma 8.3. *Let g be as in Proposition 8.2. Then the map $\phi \rightarrow \mathbf{Non}_\gamma(g, \phi)$ is Lipschitz from*

$$H^{s+2} + \dot{H}^2 \rightarrow H^s$$

with Lipschitz constant ϵ for $\|\nabla^2\phi\|_{H^s} \lesssim \epsilon$.

Proof of Lemma 8.3. In order to prove Lemma 8.3, we consider the following simplified linearization for $\mathbf{Non}_\gamma(g, \phi)$ as a function of ϕ :

$$(8.9) \quad \begin{aligned} \mathcal{T}(g, \phi, \Phi) = & h(1+h)\nabla^2\Phi + g(\nabla h + \delta\mathcal{K}) \\ & + g(\nabla\Phi + \nabla\phi\nabla\Phi)[g\nabla^2\phi + \nabla h + \mathcal{K}] \\ & + g(\nabla\phi + \nabla\phi\nabla\phi)[g\nabla^2\Phi + \delta\mathcal{K}] \end{aligned}$$

where Φ is the linearized variable associated to ϕ , \mathcal{K} has the form

$$\mathcal{K} := g\nabla^2\phi\mathcal{P} + g\nabla\mathcal{C}(1+\mathcal{P}) + \nabla h\mathcal{P}(1+\mathcal{P}) + \nabla[g(1+\mathcal{P})^2]\mathcal{P},$$

and $\delta\mathcal{K}$ is

$$\begin{aligned} \delta\mathcal{K} := & g\nabla^2\Phi\mathcal{P} + g\nabla^2\phi\delta\mathcal{P} + g\nabla\delta\mathcal{C}(1+\mathcal{P}) + g\nabla\mathcal{C}\delta\mathcal{P} + \nabla h\delta\mathcal{P}(1+\mathcal{P}) \\ & + \nabla[g\delta\mathcal{P}(1+\mathcal{P})]\mathcal{P} + \nabla[g(1+\mathcal{P})^2]\delta\mathcal{P}. \end{aligned}$$

Here \mathcal{C} and $\delta\mathcal{C}$ satisfy

$$\mathcal{C} = \nabla\phi\nabla\phi + \mathcal{C}\nabla\phi, \quad \delta\mathcal{C} = \nabla\phi\nabla\Phi + \delta\mathcal{C}\nabla\phi + \mathcal{C}\nabla\Phi,$$

and \mathcal{P} and $\delta\mathcal{P}$ are

$$\mathcal{P} = \nabla\phi + \mathcal{C}, \quad \delta\mathcal{P} = \nabla\Phi + \delta\mathcal{C}.$$

Then for the equation (8.9) we have estimates as follows:

Lemma 8.4 (Elliptic estimates for (8.9)). *Let $d \geq 3$ and $s > d/2$. Assume that $\|\nabla h\|_{H^s} \lesssim \epsilon$ and $\|\nabla^2\phi\|_{H^s} \lesssim \epsilon$, then for the linearized expression (8.9) we have the following estimate*

$$(8.10) \quad \|\mathcal{T}(g, \phi, \Phi)\|_{H^s} \lesssim \|\nabla h\|_{H^s} + \epsilon\|\nabla^2\Phi\|_{H^s}.$$

Proof of Lemma 8.4. First, we bound \mathcal{C} , $\delta\mathcal{C}$, \mathcal{P} and $\delta\mathcal{P}$. By Sobolev embeddings, using also the smallness condition $\|\nabla^2\phi\|_{H^s} \lesssim \epsilon$, we have

$$\|\nabla\mathcal{C}\|_{H^s} \lesssim \|\nabla^2\phi\|_{H^s}^2 + \|\nabla\mathcal{C}\|_{H^s}\|\nabla^2\phi\|_{H^s} \lesssim \epsilon^2 + \epsilon\|\nabla\mathcal{C}\|_{H^s},$$

and

$$\begin{aligned} \|\nabla\delta\mathcal{C}\|_{H^s} & \lesssim \|\nabla^2\phi\|_{H^s}\|\nabla^2\Phi\|_{H^s} + \|\nabla\delta\mathcal{C}\|_{H^s}\|\nabla^2\phi\|_{H^s} + \|\nabla\mathcal{C}\|_{H^s}\|\nabla^2\Phi\|_{H^s} \\ & \lesssim \epsilon\|\nabla^2\Phi\|_{H^s} + \epsilon\|\nabla\delta\mathcal{C}\|_{H^s} + \|\nabla\mathcal{C}\|_{H^s}\|\nabla^2\Phi\|_{H^s}. \end{aligned}$$

These imply

$$(8.11) \quad \|\nabla\mathcal{C}\|_{H^s} \lesssim \epsilon^2, \quad \|\nabla\delta\mathcal{C}\|_{H^s} \lesssim \epsilon\|\nabla^2\Phi\|_{H^s}.$$

Similarly we have

$$(8.12) \quad \|\nabla\mathcal{P}\|_{H^s} \lesssim \epsilon, \quad \|\nabla\delta\mathcal{P}\|_{H^s} \lesssim \|\nabla^2\Phi\|_{H^s}.$$

By Sobolev embedding we bound $\delta\mathcal{K}$ by

$$\begin{aligned} \|\delta\mathcal{K}\|_{H^s} & \lesssim (1 + \|\nabla h\|_{H^s})[\|\nabla^2\Phi\|_{H^s}\|\nabla\mathcal{P}\|_{H^s} + \|\nabla^2\phi\|_{H^s}\|\nabla\delta\mathcal{P}\|_{H^s} \\ & \quad + \|\nabla\delta\mathcal{C}\|_{H^s}(1 + \|\nabla\mathcal{P}\|_{H^s}) + \|\nabla\mathcal{C}\|_{H^s}\|\nabla\delta\mathcal{P}\|_{H^s} \\ & \quad + \|\nabla h\|_{H^s}\|\nabla\delta\mathcal{P}\|_{H^s}(1 + \|\nabla\mathcal{P}\|_{H^s})^2 \\ & \quad + (1 + \|\nabla h\|_{H^s})\|\nabla\delta\mathcal{P}\|_{H^s}\|\nabla\mathcal{P}\|_{H^s}(1 + \|\nabla\mathcal{P}\|_{H^s})]. \end{aligned}$$

This combined with (8.11) and (8.12) implies

$$(8.13) \quad \|\delta\mathcal{K}\|_{H^s} \lesssim \epsilon \|\nabla^2\Phi\|_{H^s}.$$

Similarly, we also have

$$(8.14) \quad \|\mathcal{K}\|_{H^s} \lesssim \epsilon^2.$$

Now by Sobolev embedding we bound $\mathcal{T}(g, \phi, \Phi)$ by

$$\begin{aligned} \|\mathcal{T}\|_{H^s} &\lesssim \|\nabla h\|_{H^s}(1 + \|\nabla h\|_{H^s})(1 + \|\nabla^2\Phi\|_{H^s}) + (1 + \|\nabla h\|_{H^s})\|\delta\mathcal{K}\|_{H^s} \\ &\quad + (1 + \|\nabla h\|_{H^s})\|\nabla^2\Phi\|_{H^s}(1 + \|\nabla^2\phi\|_{H^s}) \\ &\quad \cdot [(1 + \|\nabla h\|_{H^s})\|\nabla^2\phi\|_{H^s} + \|\nabla h\|_{H^s} + \|\mathcal{K}\|_{H^s}] \\ &\quad + (1 + \|\nabla h\|_{H^s})\|\nabla^2\phi\|_{H^s}(1 + \|\nabla^2\phi\|_{H^s})[(1 + \|\nabla h\|_{H^s})\|\nabla^2\Phi\|_{H^s} + \|\delta\mathcal{K}\|_{H^s}]. \end{aligned}$$

By the assumptions, (8.14) and (8.13), this gives

$$\|\mathcal{T}(g, \phi, \Phi)\|_{H^s} \lesssim \|\nabla h\|_{H^s} + \epsilon \|\nabla^2\Phi\|_{H^s}.$$

We conclude the proof of the lemma. \square

We continue to prove Lemma 8.3. With small Lipschitz constant ϵ for $\|\nabla^2\phi\|_{H^s} \lesssim \epsilon$, by (8.10) we have

$$\|\mathbf{Non}_\gamma(g, \phi)\|_{H^s} \lesssim \|\nabla h\|_{H^s} + \epsilon^2,$$

and

$$\|\mathbf{Non}_\gamma(g, \phi) - \mathbf{Non}_\gamma(g, \tilde{\phi})\|_{H^s} \lesssim \epsilon \|\nabla^2(\phi - \tilde{\phi})\|_{H^s}.$$

These give the Lipschitz continuity, completing the proof of Lemma 8.3. \square

Step 3: Prove the bound (8.2). First we prove the following bound

$$(8.15) \quad \|(\partial_y \tilde{h})(y(x))\|_{H^s(dx)} \lesssim \|\partial_x h\|_{H^s(dx)}.$$

By (8.5), it suffices to bound

$$\begin{aligned} \|(1 + \mathcal{P})\partial_x[g(1 + \mathcal{P})^2]\|_{H^s} &\lesssim \|\partial_x[g(1 + \mathcal{P})^2]\|_{H^s}(1 + \|\nabla\mathcal{P}\|_{H^s}) \\ &\lesssim \|\partial_x g\|_{H^s}(1 + \|\nabla\mathcal{P}\|_{H^s})^3 \\ &\quad + \|\partial_x \mathcal{P}\|_{H^s}(1 + \|\partial_x h\|_{H^s})(1 + \|\nabla\mathcal{P}\|_{H^s})^2 \\ &\lesssim (\|\partial_x g\|_{H^s} + \|\partial_x \mathcal{P}\|_{H^s})(1 + \epsilon)^3 \lesssim \|\partial_x h\|_{H^s}. \end{aligned}$$

This gives the bound (8.15).

In order to complete the proof, we also need the following lemma:

Lemma 8.5. *Let the change of coordinates $x + \phi(x) = y$ be as in Proposition 8.2. Define the linear operator T as $T(f)(y) = f(x(y))$ for any function $f \in L^2(dx)$. Then we have*

$$(8.16) \quad \|T(f)(y)\|_{H^\sigma(dy)} \lesssim \|f(x)\|_{H^\sigma(dx)}, \quad \sigma \in [0, [s] + 1].$$

Given this lemma, the bound (8.2) is obtained by (8.15) and (8.16) with $\sigma = s$, and the proof of Proposition 8.2 is concluded. It remains to prove the Lemma.

Proof of Lemma 8.5. Let k be an integer $k \in [0, [s] + 1]$, where $[s]$ is the integer part of s . By the change of coordinates $x + \phi(x) = y$, we have

$$\partial_y^k T(f)(y) = \left[\frac{\partial x}{\partial y} \frac{\partial}{\partial x} \right]^k f(x) \approx [(1 + \mathcal{P})\partial_x]^k f(x).$$

It suffices to consider the following forms

$$\sum_{\substack{1 \leq i \leq k-1, \\ l \geq 1, l_1 \geq \dots \geq l_i \geq 1}} \partial_x^l f \partial_x^{l_1} \mathcal{P} \dots \partial_x^{l_i} \mathcal{P} (1 + \mathcal{P})^{k-i}.$$

By Sobolev embedding, we bound each terms by

$$\begin{aligned} \|\partial_x^l f \partial_x^{l_1} \mathcal{P} \dots \partial_x^{l_i} \mathcal{P} (1 + \mathcal{P})^{k-i}\|_{L^2(dy)} &\lesssim \|\partial_x^l f \partial_x^{l_1} \mathcal{P} \dots \partial_x^{l_i} \mathcal{P} (1 + \mathcal{P})^{k-i} \sqrt{\det(I + \partial_x \phi)}\|_{L^2(dx)} \\ &\lesssim \|\partial_x^l f \partial_x^{l_1} \mathcal{P} \dots \partial_x^{l_i} \mathcal{P}\|_{L^2} (1 + \|\nabla \mathcal{P}\|_{H^s})^{k-i} \|1 + \nabla \phi\|_{L^\infty}^d \\ &\lesssim \|f\|_{H^k} \|\nabla \mathcal{P}\|_{H^s}^i (1 + \|\nabla h\|_{H^s})^{k-i} (1 + \|\nabla h\|_{H^s})^d \\ &\lesssim \epsilon^i \|f\|_{H^k}. \end{aligned}$$

Then we have

$$\|\partial_y^k T(f)(y)\|_{L^2(dy)} \lesssim \sum_{i=0}^{k-1} \epsilon^i \|f(x)\|_{H^k(dx)} \lesssim \|f(x)\|_{H^k(dx)}.$$

This implies

$$\|T(f)(y)\|_{H^k(dy)} \lesssim \|f(x)\|_{H^k(dx)}, \quad \text{for any } k \in [0, [s] + 1].$$

Thus the bound (8.16) is obtained if $\sigma \in [0, [s] + 1]$ is an integer. The similar bound for noninteger σ follows by interpolation. \square

Step 4: Prove the bound (8.3). We first prove that the $\partial_{y_\alpha y_\beta}^2 \tilde{F} \in H^s$ is also small under the above change of coordinates as follows.

Proposition 8.6. *Let $d \geq 3$, $s > \frac{d}{2}$, and $F : (\mathbb{R}_x^d, g) \rightarrow (\mathbb{R}^{d+2}, g_{\mathbb{R}^{d+2}})$ be an immersion as in Theorem 8.1. Under the change of coordinates $y = x + \phi(x)$ as in Proposition 8.2, we also have*

$$(8.17) \quad \|\partial_{y_\alpha y_\beta}^2 \tilde{F}\|_{H^s(dy)} \lesssim \epsilon_0.$$

Once the bound (8.17) holds, by (8.2) and Sobolev embedding we obtain the bound (8.3). Here we turn our attention to the proof of Proposition 8.6 and complete the proof of Proposition 8.2.

Proof of Proposition 8.6. Here we first prove that $\partial^2 F$ is also small in H^s . Precisely, by the smallness of $\partial_x g$ and Sobolev embedding, we have

$$\begin{aligned} \|g^{\alpha\beta} \Gamma_{\alpha\beta}^\gamma \partial_\gamma F\|_{H^s} &\lesssim \|g^{\alpha\beta} \Gamma_{\alpha\beta}^\gamma\|_{H^s} + \|g^{\alpha\beta} \Gamma_{\alpha\beta}^\gamma (\partial_\gamma F - \partial_\gamma F(\infty))\|_{H^s} \\ &\lesssim \|\partial_x h\|_{H^s} (1 + \|\partial_x h\|_{H^s}) (1 + \|\partial^2 F\|_{H^s}) \lesssim \epsilon_0 (1 + \|\partial^2 F\|_{H^s}). \end{aligned}$$

Then we can bound $\partial^2 F$ by

$$\begin{aligned}
\|\partial^2 F\|_{H^s} &= \|\mathcal{R}\Delta F\|_{H^s} \lesssim \|\Delta F\|_{H^s} \lesssim \|g^{\alpha\beta}\partial_{\alpha\beta}^2 F\|_{H^s} + \epsilon_0 \|\partial^2 F\|_{H^s} \\
&\lesssim \|\Delta_g F\|_{H^s} + \|g^{\alpha\beta}\Gamma_{\alpha\beta}^\gamma \partial_\gamma F\|_{H^s} + \epsilon_0 \|\partial^2 F\|_{H^s} \\
&\lesssim \|\mathbf{H}\|_{H^s} + \epsilon_0(1 + \|\partial^2 F\|_{H^s}) \\
&\lesssim \epsilon_0(1 + \|\partial^2 F\|_{H^s}),
\end{aligned}$$

which implies

$$(8.18) \quad \|\partial^2 F\|_{H^s} \lesssim \epsilon_0.$$

Next, we turn to prove the bound (8.17). By the change of coordinates, we have the representation $\partial_{y_\alpha y_\beta}^2 \tilde{F}$ as

$$\partial_{y_\alpha y_\beta}^2 \tilde{F} = \partial_{y_\alpha} (\partial_\gamma F \frac{\partial x_\gamma}{\partial y_\beta}) = \partial_{\sigma\gamma}^2 F \frac{\partial x_\sigma}{\partial y_\alpha} \frac{\partial x_\gamma}{\partial y_\beta} + \partial_\gamma F \frac{\partial}{\partial y_\alpha} \frac{\partial x_\gamma}{\partial y_\beta}.$$

Since $\frac{\partial x_\gamma}{\partial y_\beta}$ is a function depending on x and has the form $\frac{\partial x}{\partial y} = I_d + \mathcal{P}(x)$, we write this as

$$\begin{aligned}
\partial_{y_\alpha y_\beta}^2 \tilde{F} &= \partial_{\sigma\gamma}^2 F (I_d + \mathcal{P})^2 + \partial_\gamma F \partial_x (I_d + \mathcal{P}) \cdot (I_d + \mathcal{P}) \\
&= \partial_{\sigma\gamma}^2 F (I_d + \mathcal{P})^2 + \partial_\gamma F \partial_x \mathcal{P} \cdot (I_d + \mathcal{P}).
\end{aligned}$$

As a vector depends on x , by Sobolev embedding, (8.18) and (8.12) we have

$$\begin{aligned}
\|(\partial_{y_\alpha y_\beta}^2 \tilde{F})(x)\|_{H^s(dx)} &\lesssim \|\partial_{\sigma\gamma}^2 F\|_{H^s} (1 + \|\partial_x \mathcal{P}\|_{H^s}^2) + (1 + \|\partial^2 F\|_{H^s}) \|\partial_x \mathcal{P}\|_{H^s} (1 + \|\partial_x \mathcal{P}\|_{H^s}) \\
&\lesssim \epsilon_0.
\end{aligned}$$

Then by Lemma 8.5, the bound (8.17) follows. \square

Step 5: Prove the bound (8.4). Finally, we construct the initial data ψ_0 in the harmonic coordinates and Coulomb gauge. To obtain the Coulomb gauge, we choose $\tilde{\nu}$ constant uniformly transversal to $T\Sigma_0$; such a ν exists because, by Sobolev embeddings, $\partial_x F$ has a small variation in L^∞ . Projecting $\tilde{\nu}$ on the normal bundle $N\Sigma_0$ and normalizing we obtain some $\tilde{\nu}_1$ with the same regularity as ∂F . Then we choose $\tilde{\nu}_2$ in $N\Sigma_0$ perpendicular to $\tilde{\nu}_1$. We obtain the orthonormal frame $(\tilde{\nu}_1, \tilde{\nu}_2)$ in $N\Sigma_0$, which again has the same regularity and bounds as $\partial_x F$. Then we rotate the frame to get a Coulomb frame (ν_1, ν_2) , i.e. where the Coulomb gauge condition (2.16) is satisfied. Projecting the mean curvature \mathbf{H} on the Coulomb frame as in Section 2.3 we obtain the complex mean curvature $\psi \in H^s$.

In order to get the bound for ψ , we recall that the second fundamental form λ satisfies

$$\lambda_{\alpha\beta} = (\partial_{\alpha\beta}^2 F)^\perp \cdot \nu_1 + i(\partial_{\alpha\beta}^2 F)^\perp \cdot \nu_2.$$

We easily have

$$\|\lambda\|_{L^2} \lesssim \|\partial^2 F\|_{L^2} \lesssim \epsilon_0.$$

Then it suffices to bound the \dot{H}^s norm of λ . If $s \in \mathbb{N}$, we have

$$\begin{aligned}
\|\lambda\|_{\dot{H}^s} &\lesssim \sum_{\nu \in \{\nu_1, \nu_2\}; n_1+n_2=s} \|\partial^{n_1}(\partial_{\alpha\beta}^2 F - \Gamma_{\alpha\beta}^\gamma F_\gamma) \cdot \partial^{n_2} \nu\|_{L^2} \\
&\lesssim \sum_{\nu \in \{\nu_1, \nu_2\}} \|(\nabla_x + A)^s \nabla_\alpha \partial_\beta F \cdot \nu\|_{L^2} \\
&\lesssim \|(\nabla_x + A)^s \nabla_\alpha \partial_\beta F\|_{L^2} \\
&\lesssim \|\partial^2 F\|_{H^s} (1 + \|\nabla A\|_{H^s}^s).
\end{aligned}$$

If $s \notin \mathbb{N}$, let $\frac{1}{p} + \frac{1}{q} = \frac{1}{2}$ we also have

$$\begin{aligned}
\|\lambda\|_{\dot{H}^s} &\lesssim \sum_{\nu \in \{\nu_1, \nu_2\}; n_1+n_2=[s]-1} \| |D|^{1+s-[s]} (\partial^{n_1}(\partial_{\alpha\beta}^2 F - \Gamma_{\alpha\beta}^\gamma F_\gamma) \cdot \partial^{n_2} \nu) \|_{L^2} \\
&\lesssim \sum_{\nu \in \{\nu_1, \nu_2\}} \| |D|^{1+s-[s]} ((\nabla_x + A)^{[s]-1} \nabla_\alpha \partial_\beta F \cdot \nu) \|_{L^2} \\
&\lesssim \sum_{\nu \in \{\nu_1, \nu_2\}} (\| |D|^{1+s-[s]} (\nabla_x + A)^{[s]-1} \nabla_\alpha \partial_\beta F \cdot \nu \|_{L^2} \\
&\quad + \| (\nabla_x + A)^{[s]-1} \nabla_\alpha \partial_\beta F \cdot |D|^{1+s-[s]} \nu \|_{L^2}) \\
&\lesssim \| |D|^{1+s-[s]} (\nabla_x + A)^{[s]-1} \nabla_\alpha \partial_\beta F \|_{L^2} \\
&\quad + \| (\nabla_x + A)^{[s]-1} \nabla_\alpha \partial_\beta F \|_{L^p} \sum_{\nu \in \{\nu_1, \nu_2\}} \| |D|^{1+s-[s]} \nu \|_{L^q} \\
&= I_1 + I_2.
\end{aligned}$$

We bound the first term by

$$I_1 \lesssim \|\partial^2 F\|_{H^s} (1 + \|\nabla A\|_{H^s}^{[s]}) \lesssim \epsilon_0 (1 + \|\nabla A\|_{H^s}^{[s]}).$$

For the second term, we choose integer $k = \lceil \frac{d+1}{2} \rceil$ and $\frac{1}{p} = \frac{k-1-(s-[s])}{d}$, then we have

$$\begin{aligned}
I_2 &\lesssim \|(\nabla_x + A)^{[s]-1} \nabla_\alpha \partial_\beta F\|_{H^{1+s-[s]}} \sum_{\nu \in \{\nu_1, \nu_2\}} \|\partial^k \nu\|_{L^2} \\
&\lesssim \|\partial^2 F\|_{H^s} (1 + \|\nabla A\|_{H^s}^{[s]}) (1 + \|A\|_{H^s} + \|\lambda\|_{H^s})^k.
\end{aligned}$$

Therefore, by the elliptic estimates of the div-curl system (2.13)-(2.16) for A we obtain

$$\|\lambda\|_{H^s} \lesssim \epsilon_0 (1 + \|\nabla A\|_{H^s}^{[s]}) (1 + \|A\|_{H^s} + \|\lambda\|_{H^s})^k \lesssim \epsilon_0 (1 + \|\lambda\|_{H^s})^{4[s]+1} \lesssim \epsilon_0 + \epsilon_0 \|\lambda\|_{H^s}^{4[s]+1}.$$

By continuity method, this implies the bound

$$\|\lambda\|_{H^s} \lesssim \epsilon_0.$$

which combined with the smallness of $\partial_x g \in H^s$ also gives the bound (8.4) for ψ . \square

8.2. The moving frame. Once we have the initial data ψ_0 which is small in H^s , Theorem 2.7 yields the good gauge local solution ψ , along with the associated derived variables (λ, h, V, A, B) . But this does not yet give us the actual maps F .

Here we undertake the task of reconstructing the frame (F_α, m) . For this we use the system consisting of (2.6) and (2.25), viewed as a linear ode. We recall these equations here:

$$(8.19) \quad \begin{cases} \partial_\alpha F_\beta = \Gamma_{\alpha\beta}^\gamma F_\gamma + \text{Re}(\lambda_{\alpha\beta} \bar{m}), \\ \partial_\alpha^A m = -\lambda_\alpha^\gamma F_\gamma, \end{cases}$$

respectively

$$(8.20) \quad \begin{cases} \partial_t F_\alpha = -\text{Im}(\partial_\alpha^A \psi \bar{m} - i\lambda_{\alpha\gamma} V^\gamma \bar{m}) + [\text{Im}(\psi \bar{\lambda}_\alpha^\gamma) + \nabla_\alpha V^\gamma] F_\gamma, \\ \partial_t^B m = -i(\partial^{A,\alpha} \psi - i\lambda_\gamma^\alpha V^\gamma) F_\alpha, \end{cases}$$

where $(\psi, \lambda, g, V, A, B)$ is the unique solution of (2.35)-(2.36) with initial data ψ_0 small.

We start with the frame at time $t = 0$, which already is known to solve (8.19), and has the following properties:

- (i) *Orthogonality*, $F_\alpha \perp m$, $\langle m, m \rangle = 2$, $\langle m, \bar{m} \rangle = 0$ and consistency with the metric $g_{\alpha\beta} = \langle F_\alpha, F_\beta \rangle$.
- (ii) *Integrability*, $\partial_\beta F_\alpha = \partial_\alpha F_\beta$.
- (iii) *Consistency* with the second fundamental form and the connection A :

$$\partial_\alpha F_\beta \cdot m = \lambda_{\alpha\beta}, \quad \langle \partial_\alpha m, m \rangle = -2iA_\alpha.$$

Next we extend this frame to times $t > 0$ by simultaneously solving the pair of equations (8.19) and (8.20). To avoid some technical difficulties, we first do this for regular solutions, i.e. $s > d/2 + 2$, and then pass to the limit to obtain the frame for rough solutions.

8.2.1. The frame associated to smooth solutions. The system consisting of (8.19) and (8.20) is overdetermined, and the necessary and sufficient condition for existence of solutions is provided by Frobenius' theorem. We now verify these compatibility conditions in two steps:

a) Compatibility conditions for the system (8.19) at fixed time. Here, by $C_{\alpha\beta}^2 = 0$, $C_{\alpha\beta}^3 = 0$ and $C_{\alpha\beta\mu\nu}^7 = 0$ we have

$$\partial_\alpha(\Gamma_{\beta\gamma}^\sigma F_\sigma + \text{Re}(\lambda_{\beta\gamma} \bar{m})) - \partial_\beta(\Gamma_{\alpha\gamma}^\sigma F_\sigma + \text{Re}(\lambda_{\alpha\gamma} \bar{m})) = C_{\sigma\gamma\alpha\beta}^7 F^\sigma = 0,$$

and

$$\partial_\alpha(iA_\beta m + \lambda_\beta^\sigma F_\sigma) - \partial_\beta(iA_\alpha m + \lambda_\alpha^\sigma F_\sigma) = iC_{\alpha\beta}^3 m = 0,$$

as needed.

b) Between the system (8.19) and (8.20). By (8.19) and (8.20) we have

$$\partial_t(iA_\alpha m + \lambda_\alpha^\sigma F_\sigma) - \partial_\alpha(iBm + i(\partial^{A,\sigma} \psi - i\lambda_\gamma^\sigma V^\gamma) F_\sigma) = iT_\alpha^3 m + T_\alpha^{2\sigma} F_\sigma$$

and

$$(8.21) \quad \begin{aligned} & \partial_\beta[-\text{Im}(\partial_\alpha^A \psi \bar{m} - i\lambda_{\alpha\gamma} V^\gamma \bar{m}) + [\text{Im}(\psi \bar{\lambda}_\alpha^\gamma) + \nabla_\alpha V^\gamma] F_\gamma] - \partial_t[\Gamma_{\beta\alpha}^\gamma F_\gamma + \text{Re}(\lambda_{\beta\alpha} \bar{m})] \\ &= -\text{Re}[(g_{\sigma\alpha} T_\beta^{2\sigma} + \lambda_\beta^\sigma T_{\sigma\alpha}^1) \bar{m}] - T^{1\gamma\sigma} \Gamma_{\beta\alpha,\sigma} F_\gamma - \frac{1}{2}(\partial_\beta T_{\alpha\sigma}^1 + \partial_\alpha T_{\beta\sigma}^1 - \partial_\sigma T_{\beta\alpha}^1) F^\sigma. \end{aligned}$$

The first equality is obtained directly. For the second equality (8.21), by (8.19) and (8.20) we compute this by

$$\begin{aligned} \text{LHS}(8.21) &= -\text{Re}[(g_{\sigma\alpha}T_{\beta}^{2\sigma} + \lambda_{\beta}^{\sigma}T_{\sigma\alpha}^1)\bar{m}] + \nabla_{\beta}(\text{Im}(\psi\bar{\lambda}_{\sigma\alpha}) + \nabla_{\alpha}V_{\sigma})F^{\sigma} \\ &\quad + \text{Im}(\nabla_{\alpha}^A\psi\bar{\lambda}_{\sigma\beta} - \nabla_{\sigma}^A\psi\bar{\lambda}_{\alpha\beta})F^{\sigma} - \tilde{R}_{\sigma\alpha\beta\gamma}V^{\gamma}F^{\sigma} - \partial_t\Gamma_{\beta\alpha}^{\gamma}F_{\gamma}. \end{aligned}$$

By T^1 and the notation $G_{\alpha\beta}$ (2.29) we compute the last term by

$$\begin{aligned} -\partial_t\Gamma_{\beta\alpha}^{\gamma}F_{\gamma} &= -(T^{1\gamma\sigma} - 2G^{\gamma\sigma})\Gamma_{\beta\alpha,\sigma}F_{\gamma} - \frac{1}{2}[\partial_{\beta}(T_{\alpha\sigma}^1 + 2G_{\alpha\sigma})]F^{\sigma} \\ &\quad - \frac{1}{2}[\partial_{\alpha}(T_{\beta\sigma}^1 + 2G_{\beta\sigma})]F^{\sigma} + \frac{1}{2}[\partial_{\sigma}(T_{\beta\alpha}^1 + 2G_{\beta\alpha})]F^{\sigma} \\ &= -T^{1\gamma\sigma}\Gamma_{\beta\alpha,\sigma}F_{\gamma} - \frac{1}{2}(\partial_{\beta}T_{\alpha\sigma}^1 + \partial_{\alpha}T_{\beta\sigma}^1 - \partial_{\sigma}T_{\beta\alpha}^1)F^{\sigma} \\ &\quad + [-\nabla_{\beta}\text{Im}(\psi\bar{\lambda}_{\alpha\sigma}) - \text{Im}(\nabla_{\alpha}^A\psi\bar{\lambda}_{\beta\sigma}) + \text{Im}(\nabla_{\sigma}^A\psi\bar{\lambda}_{\beta\alpha})] \\ &\quad - \frac{1}{2}(\nabla_{\alpha}\nabla_{\beta} + \nabla_{\beta}\nabla_{\alpha})V_{\sigma} - \frac{1}{2}[\nabla_{\beta}, \nabla_{\sigma}]V_{\alpha} - \frac{1}{2}[\nabla_{\alpha}, \nabla_{\sigma}]V_{\beta}]F^{\sigma}. \end{aligned}$$

Then by Bianchi identities and (2.8), we collect the terms above containing V and have

$$\begin{aligned} &\frac{1}{2}([\nabla_{\beta}, \nabla_{\alpha}]V_{\sigma} - [\nabla_{\beta}, \nabla_{\sigma}]V_{\alpha} - [\nabla_{\alpha}, \nabla_{\sigma}]V_{\beta}) - \tilde{R}_{\sigma\alpha\beta\gamma}V^{\gamma} \\ &= \frac{1}{2}(R_{\beta\alpha\sigma\gamma} - R_{\beta\sigma\alpha\gamma} - R_{\alpha\sigma\beta\gamma} - 2R_{\sigma\alpha\beta\gamma})V^{\gamma} = 0. \end{aligned}$$

From the above expressions the equality (8.21) follows.

Once the compatibility conditions in Frobenius' theorem are verified, we obtain the frame (F_{α}, m) for $t \in [0, 1]$. For this we can easily obtain the regularity

$$\partial_x(F_{\alpha}, m) \in L^{\infty}H^{s+2}, \quad \partial_t(F_{\alpha}, m) \in L^{\infty}H^{s+1}.$$

Finally, we show that the properties (i)-(iii) above also extend to all $t \in [0, 1]$. The properties (ii) and (iii) follow directly from the equations (8.19) and (8.20) once the orthogonality conditions in (i) are verified. For (i) we denote

$$\tilde{g}_{00} = \langle m, m \rangle, \quad \tilde{g}_{\alpha 0} = \langle F_{\alpha}, m \rangle, \quad \tilde{g}_{\alpha\beta} = \langle F_{\alpha}, F_{\beta} \rangle.$$

Then by (8.20) and $T_{\alpha\beta}^1 = 0$, we have

$$\begin{aligned} \partial_t\tilde{g}_{\alpha 0} &= -\frac{i}{2}(\overline{\partial_{\alpha}^A\psi} + i\bar{\lambda}_{\alpha\gamma}V^{\gamma})(\tilde{g}_{00} - 2) - i(\overline{\partial^{A,\sigma}\psi} + i\bar{\lambda}_{\gamma}^{\sigma}V^{\gamma})(g_{\alpha\sigma} - \tilde{g}_{\alpha\sigma}) \\ &\quad + \frac{i}{2}(\partial_{\alpha}^A\psi + i\lambda_{\alpha\gamma}V^{\gamma})\langle \bar{m}, m \rangle + (\text{Im}(\psi\bar{\lambda}_{\alpha}^{\gamma}) + \nabla_{\alpha}V^{\gamma})\tilde{g}_{\gamma 0} + iB\tilde{g}_{\alpha 0}, \\ \partial_t(\tilde{g}_{00} - 2) &= 2\text{Im}(\partial^{A,\alpha}\psi - i\lambda_{\gamma}^{\alpha}V^{\gamma})\tilde{g}_{\alpha 0}, \\ \partial_t\langle m, \bar{m} \rangle &= -iB\langle m, \bar{m} \rangle - i(\partial^{A,\alpha}\psi - i\lambda_{\gamma}^{\alpha}V^{\gamma})\tilde{g}_{\alpha 0}, \\ \partial_t(g_{\alpha\beta} - \tilde{g}_{\alpha\beta}) &= (\text{Im}(\psi\bar{\lambda}_{\alpha}^{\gamma}) + \nabla_{\alpha}V^{\gamma})(g_{\beta\gamma} - \tilde{g}_{\beta\gamma}) + (\text{Im}(\psi\bar{\lambda}_{\beta}^{\gamma}) + \nabla_{\beta}V^{\gamma})(g_{\alpha\gamma} - \tilde{g}_{\alpha\gamma}) \\ &\quad + \text{Im}(\partial_{\alpha}^A\psi\tilde{g}_{\beta 0} - i\lambda_{\alpha\gamma}V^{\gamma}\tilde{g}_{\beta 0}) + \text{Im}(\partial_{\beta}^A\psi\tilde{g}_{\alpha 0} - i\lambda_{\beta\gamma}V^{\gamma}\tilde{g}_{\alpha 0}). \end{aligned}$$

Viewed as a linear system of ode's in time, these equations allow us to propagate (i) in time.

8.2.2. *The frame associated to rough solutions.* Here we use our approximation of rough solutions with smooth solutions for the ψ equation in order to construct the frame in the rough case. Precisely, given a small initial data $\psi_0 \in H^s$, there exists a sequence $\{\psi_{0n}\} \in H^{s+2}$ such that $\|\psi_{0n} - \psi_0\|_{H^s} \rightarrow 0$. By Theorem 2.7, the Schrödinger system (2.35) coupled with (2.36) admits solutions ψ_n with $\psi_n(0) = \psi_{0n}$ and

$$\|\psi_n\|_{H^{s+2}} \lesssim \|\psi_{0n}\|_{H^{s+2}}, \quad \|\psi_n - \psi\|_{H^s} \lesssim \|\psi_{0n} - \psi_0\|_{H^s} \rightarrow 0.$$

A-priori, we do not know whether the initial data ψ_{0n} is associated to a frame at the initial time. Hence we first use (8.19) to construct the frame $(F_\alpha^{(n)}, m^{(n)})$ associated with ψ_{0n} at $t = 0$. At some point x_0 , we choose $F_\alpha^{(n)}(x_0)$ and $m^{(n)}(x_0)$ so that they are orthogonal, and $\langle m^{(n)}, m^{(n)} \rangle = 2$, $\langle m^{(n)}, \bar{m}^{(n)} \rangle = 0$ and $\langle F_\alpha^{(n)}, F_\beta^{(n)} \rangle = g_{\alpha\beta}^{(n)}$ hold. With this initial data, we view (8.19) as a linear ode with continuous coefficients. As above, the necessary and sufficient condition for solvability, as provided by Frobenius' theorem, is a consequence of the relations $C^2 = 0$, $C^3 = 0$ and $C^7 = 0$, which are in turn a consequence of Theorem 4.1.

The above construction determines the frame $(F_\alpha^{(n)}, m^{(n)})$ up to symmetries (rigid rotations and translations). Hence, the frame $(F_\alpha^{(n)}, m^{(n)})$ at $t = 0$ is uniquely determined by the condition

$$\lim_{x \rightarrow \infty} (F_\alpha^{(n)}, m^{(n)})(x, 0) = \lim_{x \rightarrow \infty} (F_\alpha, m)(x, 0).$$

In this construction, the properties (i)-(iii) above also extend to all x . The properties (ii) and (iii) follow directly from equation (8.19) once the orthogonality conditions in (i) are verified. For (i) we use (8.19) to compute

$$\begin{aligned} \partial_\alpha \tilde{g}_{\beta 0} &= \Gamma_{\alpha\beta}^\gamma \tilde{g}_{\gamma 0} + \frac{1}{2} \lambda_{\alpha\beta} \langle \bar{m}, m \rangle + \frac{1}{2} \bar{\lambda}_{\alpha\beta} (\tilde{g}_{00} - 2) + \bar{\lambda}_\alpha^\gamma (g_{\beta\gamma} - \tilde{g}_{\beta\gamma}) + i A_\alpha \tilde{g}_{\beta 0}, \\ \partial_\alpha (\tilde{g}_{00} - 2) &= -2 \operatorname{Re}(\lambda_\alpha^\gamma \tilde{g}_{\gamma 0}), \\ \partial_\alpha \langle m, \bar{m} \rangle &= -2i A_\alpha \langle m, \bar{m} \rangle - 2 \operatorname{Re} \lambda_\alpha^\gamma \tilde{g}_{\gamma 0}, \\ \partial_\alpha (g_{\beta\gamma} - \tilde{g}_{\beta\gamma}) &= \Gamma_{\alpha\beta}^\sigma (g_{\sigma\gamma} - \tilde{g}_{\sigma\gamma}) + \Gamma_{\alpha\gamma}^\sigma (g_{\sigma\beta} - \tilde{g}_{\sigma\beta}) + \operatorname{Re}(\bar{\lambda}_{\beta\alpha} \tilde{g}_{\gamma 0} + \bar{\lambda}_{\gamma\alpha} \tilde{g}_{\beta 0}). \end{aligned}$$

By ode uniqueness and the choice of the initial data, the desired properties for the frame are propagated spatially.

Once we have the frames $(F_\alpha^{(n)}, m^{(n)})$ at $t = 0$, we can invoke the smooth case analysis above, using (8.20) and $\psi_n \in H^{s+2}$ to extend the frame $(F_\alpha^{(n)}, m^{(n)})$ to $t > 0$ with initial data $(F_\alpha^{(n)}, m^{(n)})(x, 0)$.

In order to obtain a limiting frame (F_α, m) we study the properties of the regular frames $(F_\alpha^{(n)}, m^{(n)})$ in three steps:

a) *Uniform bounds.* By (8.19), (2.37) and Sobolev embeddings we have

$$\begin{aligned} \|\partial_x F_\alpha^{(n)}\|_{H^s} &\lesssim \|\Gamma^{(n)} F_\gamma^{(n)} + \lambda^{(n)} m^{(n)}\|_{H^s} \\ &\lesssim \|\psi_n\|_{H^s} (|F_\alpha(\infty)| + |m(\infty)| + \|\partial_x (F_\alpha^{(n)}, m^{(n)})\|_{H^s}) \end{aligned}$$

and

$$\begin{aligned} \|\partial_x m^{(n)}\|_{H^s} &\lesssim \|A^{(n)} m^{(n)} + \lambda^{(n)} F_\alpha^{(n)}\|_{H^s} \\ &\lesssim \|\psi_n\|_{H^s} (|F_\alpha(\infty)| + |m(\infty)| + \|\partial_x (F_\alpha^{(n)}, m^{(n)})\|_{H^s}) \end{aligned}$$

Then, by the smallness of $\psi_n \in H^s$, we obtain

$$\|\partial_x(F_\alpha^{(n)}, m^{(n)})\|_{H^s} \lesssim \|\psi_n\|_{H^s}.$$

b) Sobolev and uniform convergence at $t = 0$. Using an argument similar to that in *a)*, by (8.19) and Theorem 4.1 *b)* we have

$$\|\partial_x(F_\alpha^{(n)} - F_\alpha, m^{(n)} - m)\|_{H^s} \lesssim \|\psi_{0n} - \psi_0\|_{H^s} + \|\psi_0\|_{H^s} \|\partial_x(F_\alpha^{(n)} - F_\alpha, m^{(n)} - m)\|_{H^s}.$$

By the smallness of ψ_0 , this implies the H^s convergence. The uniform convergence at $t = 0$ also follows by Sobolev embeddings.

c) a.e. convergence for $t > 0$. Here we use (8.20) as an ode in time. The coefficients converge in L_t^2 for a.e. x , so the frames $(F_\alpha^{(n)}, m^{(n)})$ will also converge uniformly in time for a.e. x . This can be rectified to uniform convergence in view of the uniform Sobolev bounds in (i). This yields the desired limiting frames (F_α, m) .

By (8.19) we also have

$$\|\partial_x(F_\alpha^{(k)} - F_\alpha^{(l)}, m^{(k)} - m^{(l)})\|_{L_t^\infty H^s} \lesssim \|\psi_k - \psi_l\|_{L_t^\infty H^s} \lesssim \|\psi_{0k} - \psi_{0l}\|_{H^s}.$$

This shows that the limiting frame satisfies both equations (8.20) and (8.19), as well the uniform bounds in (a).

8.3. The moving manifold Σ_t . Here we propagate the full map F by simply integrating (2.24), i.e.

$$F(t) = F(0) + \int_0^t -\text{Im}(\psi \bar{m}) + V^\gamma F_\gamma ds.$$

Then by (8.19), we have

$$\partial_\alpha F(t) = \partial_\alpha F(0) + \int_0^t -\text{Im}(\partial_\alpha^A \psi \bar{m} - i \lambda_{\alpha\gamma} V^\gamma \bar{m}) + [\text{Im}(\psi \bar{\lambda}_\alpha^\gamma) + \nabla_\alpha V^\gamma] F_\gamma ds,$$

which is consistent with above definition of F_α .

8.4. The (SMCF) equation for F . Here we establish that F solves (1.1). Using the relation $\lambda_{\alpha\beta} = \partial_{\alpha\beta}^2 F \cdot m$ we have

$$\begin{aligned} -\text{Im}(\psi \bar{m}) &= -\text{Im}(g^{\alpha\beta} \partial_{\alpha\beta}^2 F \cdot (\nu_1 + i\nu_2) (\nu_1 - i\nu_2)) \\ &= (\Delta_g F \cdot \nu_1) \nu_2 - (\Delta_g F \cdot \nu_2) \nu_1 \\ &= J(\Delta_g F)^\perp = \mathbf{JH}(F). \end{aligned}$$

This implies that the F solves (1.1).

ACKNOWLEDGMENTS

J. Huang would like to thank Prof. Lifeng Zhao for many inspirations and discussions, and Dr. Ze Li for carefully reading the manuscript, helpful discussions and comments.

REFERENCES

- [1] F. J. Almgren, Jr. Some interior regularity theorems for minimal surfaces and an extension of Bernstein's theorem. *Ann. of Math. (2)* **84** (1966) 277–292.
- [2] E. Bombieri, E. De Giorgi and E. Giusti, Minimal cones and the Bernstein problem. *Invent. Math.* **7** (1969) 243–268.
- [3] T. H. Colding and W. P. Minicozzi, Minimal submanifolds. *Bull. London Math. Soc.* **38** (2006), no. 3, 353–395.
- [4] P. Constantin and J.-C. Saut, Local smoothing properties of dispersive equations. *J. Amer. Math. Soc.* **1** (1989), 413–446.
- [5] W. Craig, T. Kappeler, and W. Strauss, Microlocal dispersive smoothing for the Schrödinger equation. *Comm. Pure Appl. Math.* **48**, No. 8 (1995), 769–860.
- [6] L. Da Rios, On the motion of an unbounded fluid with a vortex filament of any shape. *Rend. Circ. Mat. Palermo* **22** (1906) 117–135.
- [7] E. De Giorgi, Frontiere orientate di misura minima. *Sem. Mat. Scuola Norm. Sup. Pisa*, (1961) 1–56.
- [8] S. Doi, Remarks on the Cauchy problem for Schrödinger-type equations. *Comm. Partial Differential Equations* **21** (1996), 163–178.
- [9] S. Doi, Smoothing effects for Schrödinger evolution equation and global behavior of geodesic flow. *Math. Ann.* **318** (2000), 355–389.
- [10] V. Fock, The theory of space, time and gravitation. New York: The Macmillan Co., 1964
- [11] H. H. Gomez, Binormal motion of curves and surfaces in a manifold. ProQuest LLC, Ann Arbor, MI, 2004. Thesis (Ph.D.)—University of Maryland, College Park.
- [12] S. Haller and C. Vizman, Non-linear Grassmannians as coadjoint orbits. *Math. Ann.* **329**(4) (2004) 771–785.
- [13] M. Ifrim and D. Tataru, Local well-posedness for quasilinear problems: a primer. Preprint, arXiv:2008.05684
- [14] R. Jerrard, Vortex filament dynamics for Gross–Pitaevsky type equations. *Ann. Sc. Norm. Super. Pisa Cl. Sci. (5)* **1**(4) (2002) 733–768.
- [15] C. E. Kenig, G. Ponce, and L. Vega, Small solutions to nonlinear Schrödinger equations. *Ann. Inst. H. Poincaré Anal. Non Linéaire* **10** (1993), 255–288.
- [16] C. E. Kenig, G. Ponce, and L. Vega, Smoothing effects and local existence theory for the generalized nonlinear Schrödinger equations. *Invent. Math.* **134** (1998), 489–545.
- [17] C. E. Kenig, G. Ponce, and L. Vega, The Cauchy problem for quasi-linear Schrödinger equations. *Invent. Math.* **158** (2004), 343–388.
- [18] B. Khesin, Symplectic structures and dynamics on vortex membranes. *Mosc. Math. J.* **12**(2) (2012) 413–434, 461–462.
- [19] Z. Li, Global transversal stability of Euclidean planes under skew mean curvature flow evolutions. *Calc. Var. Partial Differential Equations* **60** (2021), no. 1, Paper No. 57, 19 pp.
- [20] T. Lin, Rigorous and generalized derivation of vortex line dynamics in superfluids and superconductors. *SIAM J. Appl. Math.* **60**(3) (2000) 1099–1110.
- [21] J. Marzuola, J. Metcalfe, and D. Tataru, Quasilinear Schrödinger equations I: Small data and quadratic interactions. *Adv. Math.* **231** (2012), no. 2, 1151–1172.
- [22] J. Marzuola, J. Metcalfe, and D. Tataru, Quasilinear Schrödinger equations, II: Small data and cubic nonlinearities. *Kyoto J. Math.* **54** (2014), no. 3, 529–546.
- [23] J. Marzuola, J. Metcalfe, and D. Tataru, Quasilinear Schrödinger equations, III: large data and short time. *Arch. Ration. Mech. Anal.* **242** (2021), no. 2, 1119–1175.
- [24] B. N. Shashikanth, Vortex dynamics in \mathbb{R}^4 . *J. Math. Phys.* **53** (2012) 013103.
- [25] J. Simons, Minimal varieties in Riemannian manifolds. *Ann. of Math. (2)* **88** (1968) 62–105.
- [26] C. Song, Gauss map of the skew mean curvature flow. *Proc. Amer. Math. Soc.* **145** (2017), no. 11, 4963–4970.
- [27] C. Song, Local existence and uniqueness of skew mean curvature flow. *J. Reine Angew. Math.* **776** (2021), 1–26.
- [28] C. Song, J. Sun, Skew mean curvature flow, *Commun. Contemp. Math.* **21**(1):1750090, 29, (2019).

- [29] L. Vega, The dynamics of vortex filaments with corners. *Commun. Pure Appl. Anal.* **14** (2015), no. 4, 1581–1601.
- [30] S. Weinberg, *Gravitation and Cosmology: Principles and Applications of the General Theory of Relativity*. New York: Wiley, 1972.

BEIJING INTERNATIONAL CENTER FOR MATHEMATICAL RESEARCH, PEKING UNIVERSITY, BEIJING 100871, P. R. CHINA

Email address: `huangjiaxi@bicmr.pku.edu.cn`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, BERKELEY, BERKELEY, CA 94720, USA

Email address: `tataru@math.berkeley.edu`