

UCLA

UCLA Previously Published Works

Title

Correction to: Differentiation of RPE cells from integration-free iPS cells and their cell biological characterization

Permalink

<https://escholarship.org/uc/item/7nz786pz>

Journal

Stem Cell Research & Therapy, 10(1)

ISSN

1757-6512

Authors

Hazim, Roni A

Karumbayaram, Saravanan

Jiang, Mei

et al.

Publication Date

2019-12-01

DOI

10.1186/s13287-019-1147-7

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

CORRECTION

Open Access



Correction to: Differentiation of RPE cells from integration-free iPS cells and their cell biological characterization

Roni A. Hazim^{1†}, Saravanan Karumbayaram^{2,3,4†}, Mei Jiang¹, Anupama Dimashkie³, Vanda S. Lopes¹, Douran Li^{1,5}, Barry L. Burgess¹, Preethi Vijayaraj^{4,6}, Jackelyn A. Alva-Ornelas⁷, Jerome A. Zack^{2,3,4,8}, Donald B. Kohn^{2,3,4,6}, Brigitte N. Gomperts^{3,4,6}, April D. Pyle^{2,3,4}, William E. Lowry^{3,4,5} and David S. Williams^{1,9,10,11*}

Correction to: *Stem Cell Res Ther* (2017) 8:217
<https://doi.org/10.1186/s13287-017-0652-9>

The original article [1] contains an error in the legend of Fig. 5 whereby the descriptions for panels 5d and 5e are incorrect; as such, the corrected legend can be viewed below with its respective figure images.

Author details

¹Stein Eye Institute and Department of Ophthalmology, David Geffen School of Medicine at UCLA, 100 Stein Plaza, Los Angeles, CA 90095, USA.

²Department of Microbiology Immunology and Molecular Genetics, Los Angeles, CA, USA. ³Eli and Edythe Broad Center of Regenerative Medicine and Stem Cell Research at UCLA, Los Angeles, CA, USA. ⁴Jonsson Comprehensive Cancer Center, Los Angeles, CA, USA. ⁵Department of Molecular Cell and Developmental Biology, Los Angeles, CA, USA.

⁶Department of Pediatrics, David Geffen School of Medicine, Los Angeles, CA, USA. ⁷Department of Population Sciences, City of Hope National Medical Center, Duarte, CA, USA. ⁸Department of Medicine, David Geffen School of Medicine, Los Angeles, CA, USA. ⁹Department of Neurobiology, David Geffen School of Medicine, Los Angeles, CA, USA. ¹⁰Molecular Biology Institute, Los Angeles, CA, USA. ¹¹Brain Research Institute, University of California, Los Angeles, CA, USA.

Received: 24 January 2019 Revised: 24 January 2019
Accepted: 24 January 2019 Published online: 12 February 2019

Reference

1. Hazim RA, et al. Differentiation of RPE cells from integration-free iPS cells and their cell biological characterization. *Stem Cell Res Ther*. 2017;8:217. <https://doi.org/10.1186/s13287-017-0652-9>.

* Correspondence: dswilliams@ucla.edu

[†]Roni A. Hazim and Saravanan Karumbayaram contributed equally to this work.

¹Stein Eye Institute and Department of Ophthalmology, David Geffen School of Medicine at UCLA, 100 Stein Plaza, Los Angeles, CA 90095, USA

⁹Department of Neurobiology, David Geffen School of Medicine, Los Angeles, CA, USA

Full list of author information is available at the end of the article






Fig. 5 Microtubule organization and trafficking of endolysosomes in iPSC-RPE cells. **a** Microtubule organization in the iPSC-RPE cell bodies illustrated by immunostaining of alpha tubulin. Single-plane confocal microscopy images (2- μm apart) represent the apical and basal regions of the cell bodies, plus one plane in between (middle). The apical region of the cells is dominated by horizontal microtubules while the basal region is dominated by vertical microtubules. A z projection of the three panels is shown in the fourth panel. Below are images in two z planes at the yellow lines in the z-projection image, showing primary cilia (indicated by white arrowheads) emanating from the apical surface of the RPE cells. **b** Still image from a movie of iPSC-RPE cells that were incubated with red LysoTracker to label endolysosomes (see Additional file 4 for a similar movie). **c** Trajectory and movement analysis of a population of endolysosomes, using a spots and tracks analysis (Imaris), following movie acquisition over a 25-s interval. The tracks represent the trajectories of the organelles, while their colors are indicative of how far (in terms of time) they are with respect to the 25-s movie, with cool colors being closer to the beginning of the movie, and hot colors being closer to the end of the movie. **d** The average speed of endolysosomes was determined by analyzing the tracks of the organelles, and was found to be similar among the RPE cells derived from three independent iPSC lines. **e** Time-lapse images from a movie showing vertical movement of a labeled organelle (yellow arrowhead). Each panel represents the same z plane at different times. The organelle moves out of the plane after 2 s, indicating that it is traversing different z planes. Scale bars: **a, b**, 20 μm ; **c**, 25 s (time), 5 μm (distance); **e**, 5 μm . iPSC induced pluripotent stem cell, RPE retinal pigment epithelium