UC Irvine

UC Irvine Previously Published Works

Title

Seasonal dynamics and microgeographical spatial heterogeneity of malaria along the China-Myanmar border.

Permalink

<https://escholarship.org/uc/item/7p75z0kh>

Authors

Hu, Yue
Zhou, Guofa
Ruan, Yonghua
et al.

Publication Date

2016-05-01

DOI

10.1016/j.actatropica.2016.01.022

Peer reviewed

Seasonal dynamics and microgeographical spatial heterogeneity of malaria along the China–Myanmar border

Yue Hu^{a,b,1}, Guofa Zhou^{c,1}, Yonghua Ruan^{b,1}, Ming-chieh Lee^c, Xin Xu^a, Shuang Deng^b, Yao Bai^a, Jie Zhang^a, James Morris^d, Huaie Liu^a, Ying Wang^e, Qi Fan^f, Peipei Li^f, Yanrui Wu^a, Zhaoqing Yang^{a,*}, Guiyun Yan^{c,*}, Liwang Cui^{g,*}

^a Department of Pathogen Biology and Immunology, Kunming Medical University, Kunming, China

^b Department of Pathology, Kunming Medical University, Kunming, China

^c Program in Public Health, University of California at Irvine, Irvine, CA, USA

^d Eukaryotic Pathogens Innovation Center, Department of Genetics and Biochemistry, Clemson University, SC, USA

^e Institute of Tropical Medicine, Third Military Medical University, Chongqing, China

^f Dalian Institute of Biotechnology, Dalian, Liaoning, China

^g Department of Entomology, The Pennsylvania State University, University Park, PA 16802, USA

ARTICLE INFO

Article history:

Received 29 July 2015

Received in revised form

29 December 2015

Accepted 21 January 2016

Available online 23 January 2016

Keywords:

Clinical malaria

Seasonality

Hotspot

China–Myanmar border

ABSTRACT

Malaria transmission is heterogeneous in the Greater Mekong Subregion with most of the cases occurring along international borders. Knowledge of transmission hotspots is essential for targeted malaria control and elimination in this region. This study aimed to determine the dynamics of malaria transmission and possible existence of transmission hotspots on a microgeographical scale along the China–Myanmar border. Microscopically confirmed clinical malaria cases were recorded in five border villages through a recently established surveillance system between January 2011 and December 2014. A total of 424 clinical cases with confirmed spatial and temporal information were analyzed, of which 330 (77.8%) were *Plasmodium vivax* and 88 (20.8%) were *Plasmodium falciparum*, respectively. The *P. vivax* and *P. falciparum* case ratio increased dramatically from 2.2 in 2011 to 4.7 in 2014, demonstrating that *P. vivax* malaria has become the predominant parasite species. Clinical infections showed a strong bimodal seasonality. There were significant differences in monthly average incidence rates among the study villages with rates in a village in China being 3–8 folds lower than those in nearby villages in Myanmar. Spatial analysis revealed the presence of clinical malaria hotspots in four villages. This information on malaria seasonal dynamics and transmission hotspots should be harnessed for planning targeted control.

© 2016 Elsevier B.V. All rights reserved.

1. Introduction

Malaria is a serious vector-borne disease and a major public health problem in many tropical countries (WHO, 2014). Intensive intervention during the past two decades has led to a dramatic decrease in global malaria incidence. Countries like China and Thailand have entered the malaria elimination phase (Cui et al., 2012a,b; Zhou et al., 2014a). As malaria transmission further declines, control measures will increasingly rely on accurate

knowledge of risk factors as well as the ability to define high-risk areas and populations for targeted interventions (Bousema et al., 2010; Bousema et al., 2012; Marsh, 2010; Moonen et al., 2010; Mosha et al., 2014).

Intensified malaria control has also resulted in a major change in malaria epidemiology with an increasing proportion of *Plasmodium vivax* malaria. Of the four human malaria species, *P. vivax* is the most common and has the widest geographic range (Battle et al., 2012; Gething et al., 2012; Guerra et al., 2010). It is mostly found outside of Africa and is especially prevalent in Southeast Asia. This parasite is less responsive to control interventions and much more difficult to eliminate when compared to *Plasmodium falciparum*, partly due to the noted ability of *P. vivax* to form dormant liver stages that cause relapses (Chaves et al., 2008). The increased proportions of *P.*

* Corresponding authors. Fax: +1 8148653048.

E-mail addresses: zhaoqingy92@hotmail.com (Z. Yang), guiyuny@uci.edu (G. Yan), luc2@psu.edu (L. Cui).

¹ These authors contributed equally to the work.

vivax malaria in Southeast Asia will be a formidable challenge for the malaria elimination course in these countries.

In the Greater Mekong Subregion (GMS), Myanmar has the highest regional malaria burden and is a key source of malaria exportation (Cui et al., 2012a,b; Delacollette et al., 2009). As a consequence of shared borders, low malaria transmission countries such as China (Zhang et al., 2014b) and Thailand will continually face the challenges of managing cross-border malaria introduction (Parker et al., 2015; Wangdi et al., 2015; Zhou et al., 2005, 2014a). China is entering the malaria elimination phase with a very low malaria incidence rate (<0.3 cases per 1,000,000 person year in 2013). However, the China–Myanmar border area has the highest malaria incidence rate (1.2 and ~10.3 cases per 1,000,000 person year in Yunnan Province and in four border counties, respectively). Therefore, border malaria control is crucial to malaria elimination in China (Zhang et al., 2014a). Southwestern Yunnan, which shares ~4000 km of border with Myanmar, Laos and Vietnam, is the most malaria prevalent province in China (Feng et al., 2015; Zhang et al., 2014a). Border malaria—high malaria transmission along the international borders is a common phenomenon in all the GMS nations, which requires in-depth studies and coordinated international efforts. Since border regions represent the biggest reservoirs for malaria, and frequent malaria introductions by migratory human populations are extremely difficult to monitor, border malaria constitutes one of the biggest obstacles for malaria elimination in these countries. Some border regions in Myanmar were not reached by the national malaria control program, and internal political and military conflicts further exacerbate malaria situations in these border regions. Influx of refugees from areas of higher malaria endemicity as the result of military conflicts may inevitably accompany massive malaria introduction. Therefore, timely, detailed mapping of malaria epidemiology in the border regions is essential for identifying malaria transmission hotspots and guiding targeted control efforts. However, malaria epidemiology studies conducted so far in this region mostly focused on malaria in China on large scales (Bi et al., 2013a; Clements et al., 2009; Hui et al., 2009; Li et al., 2013; Lin et al., 2009; Xu et al., 2015; Zhou et al., 2014a). Though these studies depicted the overall trends of malaria situation, better understanding of malaria transmission on a microgeographic level (e.g., villages) would provide unprecedented opportunities for local health department officials to execute targeted control efforts. In this study, as our continued efforts to monitor malaria transmission in the China–Myanmar border area, we performed longitudinal malaria case detections in villages along the border in order to map malaria transmission on a much finer scale. It will offer more accurate information for local disease control organizations to carry out malaria control operations including mosquito control, drug distribution, and health education.

2. Materials and methods

2.1. Study area and population

The total population (permanent residents) in Lai Zar and Na Bang (NB) is about 20,000 and 1500 (from the 2012 census), respectively. Malaria surveillance was carried out between January 2011 and December 2014 in five closely located villages along the China–Myanmar border: JHK, MSY, SSL, SY in Kachin State, Myanmar, and NB in Yingjiang County, Yunnan Province, China (Fig. 1). JHK, MSY and SY are the rural Myanmar villages in the study area, and SSL is a randomly selected suburban area of Lai Zar town. NB consists of three natural villages, where malaria cases were analyzed in this study. The catchment population of each of the four Myanmar villages ranged from ~120 to ~480. The selected villages all had higher malaria incidence in the past. This region has a sub-

tropical climate with January through March as the dry and cold months, and June–August as the rainy season. Annual rainfall is about 1500 mm, and average annual temperature is 22.7 °C. Average monthly temperature is the lowest (~15 °C) in January, which gradually rises to 24 °C in May, and remains at that level through September before gradually declining in October–December.

2.2. Malaria surveillance

Demographic data of the study villages in Myanmar were surveyed and updated monthly through seasonal census surveys and weekly/biweekly household visits. Information gathered includes age, sex, onset date of a febrile illness, parasite species, final diagnosis, asexual stage parasitemia and gametocytemia. The latitude and longitude of each household was taken using a handset GPS device. Malaria infection data included clinical malaria infections both from passive case surveillance (PCS) and active case surveillance (ACS). PCS was conducted at five clinics, and only patients from the study villages were included in the final data. ACS was conducted through weekly (April–September) and biweekly (October–March) home visits to assess clinical malaria. During the survey, individuals who were diagnosed with malaria by clinicians (for PCS) or who showed malaria symptoms (for ACS), and who agreed to sign a consent and/or assent form (for minors under the age of 18) were included in the study. We collected all cases occurred during the survey period, and cases potentially resulted from relapses were treated as new cases. Trained nurses collected blood samples by a standard finger-prick method, and thick and thin smears were prepared on labeled slides for parasite species identification. A clinical malaria case was defined as an individual with malaria-related symptoms (fever, i.e., axillary temperature ≥ 37.5 °C, chills, severe malaise, headache or vomiting) at the time of examination or 1–2 days prior to the examination with a *Plasmodium*-positive blood smear (Afrane et al., 2013; Zhou et al., 2015). Parasites were identified microscopically by two experienced technologists. For quality control purpose, a third microscopist confirmed parasite identification by random analysis of ~5% of the slides (Afrane et al., 2013; Zhou et al., 2015).

2.3. Statistical and spatial analysis

Clinical malaria incidence rates were calculated as cases per 1000 populations per year (month) based on the populations from the August 2012 demographic survey. Differences in monthly malaria incidence rates among villages were compared using nonparametric Kruskal–Wallis ANOVA and Median test, while Duncan's range test was used to test the pair-wise difference between villages. Friedman's nonparametric ANOVA was used to test inter-year differences in malaria incidence in MSY. The odds ratio with a 95% CI was used. Analysis of spatial clustering in clinical malaria was performed by using hot spot analysis Getis–Ord's $G_i^*(d)$ statistics (Cartabia et al., 2012; Dhimal et al., 2014; Haque et al., 2014, 2012; Kelly–Hope et al., 2009; Ruktanonchai et al., 2014; Saxena et al., 2012). Statistical analysis and spatial clustering were conducted using STATISTICA 10.0 (StatSoft Inc., Tulsa, OK, USA) and ArcGIS 10.3 for Desktop (ESRI, Redlands, CA, USA), respectively.

2.4. Ethical approval

Ethical clearance was granted by the institutional review boards of Kunming Medical University, China, University of California, Irvine, and Pennsylvania State University, USA, and the local Bureau of Health, Kachin State, Myanmar. Informed consent was obtained during the survey both orally and in writing from adult individuals, while for children it was obtained from their parents or guardians.

Fig. 1. Locations of study villages SSL (A), SY (B), MSY (C), NB (D), and JHK (E) along the China–Myanmar border.

3. Results

3.1. Malaria incidence and parasite prevalence

A newly established malaria surveillance system was in place at the study villages in order to monitor malaria transmission on the international border. A total of 424 clinical malaria cases were detected by microscopy during the study period. Among them, 271 (64%) and 153 (36%) were from male and female patients, respectively, and the median age was 19. Of the 424 cases, 330 (77.8%) were *P. vivax*, 88 (20.8%) *P. falciparum*, and six (1.4%) mixed *P. vivax* and *P. falciparum* infections. More than 90% of malaria cases were diagnosed at the clinicians (for PCS), of which 340 cases had household locations. Since this was a retrospective study, household locations could not be obtained for 84 cases (19.8%, 84/424).

Clinical malaria incidence rates varied drastically among villages (Table 1) (Kruskal–Wallis ANOVA with the Median test $\chi^2 = 26.57$, d.f. = 4, $P < 0.0001$). JHK (96.0 cases per 1000 person year) had a 7-fold higher malaria incidence rate than NB (18.1 cases per 1000 person year). NB had the lowest malaria incidence rate among the study villages. Overall, villages in Myanmar had significantly higher incidence rates than NB in China (Duncan's range test, $P < 0.05$).

Table 1

Clinical cases from 2011 to 2014 by village and parasite species.

Site	Population ^a	Blood slides examined	Clinical cases				Case rate ^b
			Total	Pv	Pf	Pv + Pf	
JHK	354	743	136	37	96	3	96.0
MSY	384	964	104	23	79	2	67.7
SSL	476	1713	84	8	76	0	44.1
SY ^c	123	135	18	5	12	1	48.8
NB ^d	1500	170	82	15	67	0	13.7

Pv, *P. vivax*; Pf, *P. falciparum*, Pv + Pf, mixed infections of *P. vivax* and *P. falciparum*.

^a Population from a 2012 demographic survey.

^b Case rate: cases per 1000 population per year.

^c 2014 cases were not available.

^d Population included only permanent residents, while cases included all suspected malaria infections.

3.2. Temporal trend in clinical malaria

The overall clinical malaria incidence showed distinct seasonality with the peak season from April to October (Fig. 2). Despite that these villages are closely located from each other, the peak malaria month and year varied among villages. Specifically, the

Fig. 2. Temporal trends of malaria incidence rate (cases per 1000 persons per year) in different villages from 2011 to 2014.

highest peak occurred in SY in 2012, JHK in 2013, and SSL in 2014, respectively (Fig. 2). MSY was the only village where incidence rates increased continuously from 2011 to 2014, with average incidence rates being 23.4, 41.7, 93.7, and 112.0 cases per 1000 person years, respectively (Friedman ANOVA $\chi^2 = 8.07$, d.f. = 3, $P < 0.05$).

Changes in *P. vivax*/*P. falciparum* (Pv/Pf) ratio were significant over the four-year period (Table 2). In Myanmar, the Pv/Pf ratio increased from 1.6 in 2011 to 2.7, 3.6, and 4.91 in 2012, 2013 and 2014, respectively. The odds ratio also increased from 2.4 (95% CI [0.7, 7.9], $P = 0.14$) in 2011 to 23.3 (95% CI [12.2, 44.2], $P < 0.0001$) in 2014. In China, however, the Pv/Pf ratio decreased from 2011 to 2013 (Table 2).

3.3. Hotspots of clinical malaria

Spatial distribution of malaria cases varied significantly both among villages and within a particular village (Fig. 3). The locations of *P. falciparum* and *P. vivax* cases are mapped in Fig. 3. The median number of cases in each village was JHK 1.5, MSY 0, SY 0, SSL 0, and NB 0, respectively. In NB, SY, and SSL, there was at most one case per household, while most households had no malaria. In contrast, in both JHK and MSY, malaria cases at the household level ranged from zero to more than 10 cases during the study period.

Table 2
Changes in *P. vivax* and *P. falciparum* (Pv/Pf) incidence ratio over time.

Site	Year	Pv/Pf	Odds ratio [95% CI]	P value
Overall	2011	2.20	4.84 [1.68, 13.93]	0.0027
	2012	3.21	8.75 [3.90, 19.61]	<0.0001
	2013	3.47	11.64 [7.02, 19.28]	<0.0001
	2014	4.66	21.16 [12.38, 36.19]	<0.0001
	Total	3.70	14.35 [10.47, 19.69]	<0.0001
Myanmar	2011	1.56	2.42 [0.74, 7.91]	0.1407
	2012	2.73	6.18 [2.46, 15.51]	<0.0001
	2013	3.63	12.60 [7.17, 22.15]	<0.0001
	2014	4.91	23.26 [12.24, 44.21]	<0.0001
	Total	3.63	12.30 [8.56, 17.66]	<0.0001
China	2011	8.00	64.00 [3.38, 1210.61]	0.0034
	2012	5.00	25.00 [4.33, 144.31]	<0.0001
	2013	2.88	8.27 [2.65, 25.79]	0.0001
	2014	4.10	16.81 [6.32, 44.68]	<0.0001
	Total	3.95	15.64 [8.07, 30.30]	<0.0001

Fig. 3. Spatial distribution of malaria cases in each village at the household level. Malaria cases were pooled from 2011 to 2014. Symbol sizes correspond to the numbers of malaria cases. *P. falciparum* and *P. vivax* cases are shown as blue triangles and red circles, respectively. (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article.)

Furthermore, most households in JHK had experienced at least one episode of malaria during the study period.

Getis-Ord's analysis showed the presence of evident malaria hotspots ($z > 2.58$, $P < 0.01$) in four villages, JHK, MSY, SSL, and NB, whereas clustering of malaria cases was marginal ($1.65 < z < 1.96$, $P < 0.10$) in SY (Fig. 4). Interestingly, malaria cases clustered in the village center of JHK, whereas the clusters were located at the gateways of the remaining four villages.

4. Discussion

Motivated by recent achievements in malaria control, several nations in the GMS are pursuing malaria elimination with China taking the lead and aiming to achieve this goal by 2020. Malaria morbidity has continued to decline over the past two decades in China, with the majority of malaria concentrated in western Yunnan (Bi et al., 2013a). International border areas have become key control points for malaria elimination in China (Cui et al., 2012a,b; Zhang et al., 2014a; Zhou et al., 2014a, 2008). Investigation of malaria incidence in border counties of Yunnan province, China revealed that clinical *P. falciparum* malaria was closely associated with travel to Myanmar, whereas local *P. vivax* transmission remained (Xu et al., 2015; Zhou et al., 2014a). This highlights the need for both strengthened local control efforts as well as tightened monitoring near international borders to prevent malaria introduction (Mosha et al., 2014; Wangdi et al., 2015).

At the last stage of malaria elimination, control programs must shift focus from high coverage of interventions to seeking out infections and interrupting transmission. Targeted intervention is probably the most cost-effective way to rid of the source of parasites (Bousema et al., 2010, 2012; Cotter et al., 2013; Mosha et al., 2014). However, targeted intervention requires accurate and timely information for identifying the target areas (Bousema et al., 2010, 2012; Cotter et al., 2013). In China, a malaria elimination surveillance system was set up with a "1-3-7" approach to meet the challenges in communicating and monitoring surveillance and response activities (Cao et al., 2014). The disease control programs will have massive data gathering and dissemination responsibilities, and timely analysis and communications are essential (Cao et al., 2013; Zhou et al., 2014b). In this aspect, spatial and temporal maps to track populations at risk of malaria at local levels would be of great value.

Our analysis of malaria epidemiology on a micro-geographical scale revealed significant malaria transmission heterogeneity in both time and space. First, malaria in this area has clear seasonality with the peak malaria season coincident with the rainy season, which is likely a reflection of the climatic pattern. Yet, it is surprising that peak malaria incidence also had considerable variations among different villages and in different years and months. This may be attributed to an insufficient malaria management system that focuses on epidemics rather than systematic interventions. In several villages, annual malaria incidence in 2012–2014 was much higher than 2011, suggesting of insufficient control

Fig. 4. Hot and cold spots of malaria incidents detected by Getis-Ord G_i^* analysis at the household level. Dot colors indicate different significance levels. Reddish dots indicate hot spots, bluish dots indicate cold spots, and yellow dots represent “not significant”. (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article.)

efforts, probably as a consequence of military conflicts taken place in recent years between the local and central governments. Second, our spatial analysis clearly showed the presence of malaria transmission hotspots in most of the villages. The strongly aggregated distribution of malaria in JHK and MSY may be attributed to the existence of mosquito breeding habitats, which were associated with high *Anopheles* mosquito abundance as detected in our mosquito surveys in 2011–2013. In this area, twenty species of *Anopheles* mosquitoes were identified, with *An. minimus* s.l. accounting for 85% of the total collections (54–91% of total captures in different villages). Mosquito densities varied from 0.05 to 3.00 females per trap per night, with strong seasonality in all sites and a density peak from June to August. Species richness peaked from April to August according to our previously reports (Bi et al., 2013b; Wang et al., 2015; Yu et al., 2013).

Interestingly, most of the hotspots were located in the cross-road or junctions areas. For example, the hotspot in JHK coincides with the major junction of roads to China, the camps for internally displaced people, and inland of Myanmar. It is possible that such hotspots may facilitate local and distant spread of malaria. Finally, the increase of malaria incidence in these villages in recent years accompanied the increase of Pv/Pf ratio, indicating increased transmission of vivax malaria in these villages. The emergence of drug-resistant parasites in this region (Yuan et al., 2014) and less stringent uses of primaquine for radical cure because the presence of a significant portion of the local ethnic population with glucose-6-phosphate dehydrogenase deficiency (Li et al., 2015) may be

partially responsible for the increased vivax cases in this region, which underlines the urgency for implementation of more effective *P. vivax* control measures (Waltmann et al., 2015; Zhou et al., 2014a). Furthermore, current treatment of *P. falciparum* was effective, which may contribute to the decrease in falciparum malaria, similar to what has been found in Thailand (Phimpraphi et al., 2008; Wang et al., 2015; Zhou et al., 2005). The migration flow between two countries is huge, estimated to be at least 20-fold higher than Lai Zar’s residents, and 300-fold higher than NB’s residents based on local records from the border control (unpublished). Our study is limited in that we did not collect data on cross-border movement of the study participants, and therefore could not differentiate indigenous from imported cases. Such information is important for a better understanding of malaria transmission at the border region.

In conclusion, this study investigated temporal and spatial patterns of malaria transmission in villages along the China–Myanmar border. While malaria here has an overall seasonal trend, there were considerable variations in the peak transmission month between the villages. Spatial analysis identified clear hotspots of malaria clinical cases within most studied villages. Malaria incidence rates were significantly higher in the Myanmar villages than in the neighboring village in China. These findings further depicted a scenario of border malaria in the GMS. Difference in malaria transmission levels between the two sides of the border emphasizes the threat of cross-border malaria introduction. These findings are vital for health workers on both sides of the border to imple-

ment targeted malaria intervention and to develop a strategy for controlling and eliminating “border malaria” in the GMS countries.

Acknowledgements

We thank the local communities for willingness to participate in this research. This project was funded by grants from the National Institutes of Health (U19 AI089672), the National Natural Science Foundation of China (U1202226 and 31260508 to ZY). Yanrui Wu was supported by grants (2014YNPHXT05 and 2014FB005) from the Yunnan province.

References

- Afrane, Y.A., Zhou, G., Githeko, A.K., Yan, G., 2013. Utility of health facility-based malaria data for malaria surveillance. *PLoS One* 8, e54305.
- Battle, K.E., Gething, P.W., Elyazar, I.R., Moyes, C.L., Sinka, M.E., Howes, R.E., Guerra, C.A., Price, R.N., Baird, J.K., Hay, S.I., 2012. The global public health significance of *Plasmodium vivax*. *Adv. Parasitol.* 80, 1–111.
- Bi, Y., Hu, W., Yang, H., Zhou, X.N., Yu, W., Guo, Y., Tong, S., 2013a. Spatial patterns of malaria reported deaths in Yunnan Province, China. *Am. J. Trop. Med. Hyg.* 88, 526–535.
- Bi, Y., Yu, W., Hu, W., Lin, H., Guo, Y., Zhou, X.N., Tong, S., 2013b. Impact of climate variability on *Plasmodium vivax* and *Plasmodium falciparum* malaria in Yunnan Province, China. *Parasites Vectors* 6, 357.
- Bousema, T., Drakeley, C., Gesase, S., Hashim, R., Magesa, S., Mosha, F., Otieno, S., Carneiro, I., Cox, J., Msuya, E., Kleinschmidt, I., Maxwell, C., Greenwood, B., Riley, E., Sauerwein, R., Chandramohan, D., Gosling, R., 2010. Identification of hot spots of malaria transmission for targeted malaria control. *J. Infect. Dis.* 201, 1764–1774.
- Bousema, T., Griffin, J.T., Sauerwein, R.W., Smith, D.L., Churcher, T.S., Takken, W., Ghani, A., Drakeley, C., Gosling, R., 2012. Hitting hotspots: spatial targeting of malaria for control and elimination. *PLoS Med.* 9, e1001165.
- Cao, J., Zhou, S.S., Zhou, H.Y., Yu, Y.B., Tang, L.H., Gao, Q., 2013. Malaria from control to elimination in China: transition of goal, strategy and interventions. *Chin. J. Schistosomiasis Control* 25, 439–443.
- Cao, J., Sturrock, H.J., Cotter, C., Zhou, S., Zhou, H., Liu, Y., Tang, L., Gosling, R.D., Feachem, R.G., Gao, Q., 2014. Communicating and monitoring surveillance and response activities for malaria elimination: China's 1–3–7 strategy. *PLoS Med.* 11, e1001642.
- Cartabia, M., Campi, R., Clavenna, A., Bortolotti, A., Fortino, I., Merlino, L., Bonati, M., 2012. Geographical epidemiology of antibacterials in the preschool age. *Int. J. Health Geogr.* 11, 52.
- Chaves, L.F., Kaneko, A., Taleo, G., Pascual, M., Wilson, M.L., 2008. Malaria transmission pattern resilience to climatic variability is mediated by insecticide-treated nets. *Malar. J.* 7, 100.
- Clements, A.C., Barnett, A.G., Cheng, Z.W., Snow, R.W., Zhou, H.N., 2009. Space-time variation of malaria incidence in Yunnan province, China. *Malar. J.* 8, 180.
- Cotter, C., Sturrock, H.J., Hsiang, M.S., Liu, J., Phillips, A.A., Hwang, J., Gueye, C.S., Fullman, N., Gosling, R.D., Feachem, R.G., 2013. The changing epidemiology of malaria elimination: new strategies for new challenges. *Lancet* 382, 900–911.
- Cui, L., Yan, G., Sattabongkot, J., Cao, Y., Chen, B., Chen, X., Fan, Q., Fang, Q., Jongwutiwes, S., Parker, D., Sirichaisinthop, J., Kyaw, M.P., Su, X.Z., Yang, H., Yang, Z., Wang, B., Xu, J., Zheng, B., Zhong, D., Zhou, G., 2012a. Malaria in the Greater Mekong Subregion: heterogeneity and complexity. *Acta Trop.* 121, 227–239.
- Cui, L., Yan, G., Sattabongkot, J., Chen, B., Cao, Y., Fan, Q., Parker, D., Sirichaisinthop, J., Su, X.Z., Yang, H., Yang, Z., Wang, B., Zhou, G., 2012b. Challenges and prospects for malaria elimination in the Greater Mekong Subregion. *Acta Trop.* 121, 240–245.
- Delacollette, C., D'Souza, C., Christophel, E., Thimasarn, K., Abdur, R., Bell, D., Dai, T.C., Gopinath, D., Lu, S., Mendoza, R., Ortega, L., Rastogi, R., Tantinimitkul, C., Ehrenberg, J., 2009. Malaria trends and challenges in the Greater Mekong Subregion. *Southeast Asian J. Trop. Med. Public Health* 40, 674–691.
- Dhimall, M., O'Hara, R.B., Karki, R., Thakur, G.D., Kuch, U., Ahrens, B., 2014. Spatio-temporal distribution of malaria and its association with climatic factors and vector-control interventions in two high-risk districts of Nepal. *Malar. J.* 13, 457.
- Feng, J., Xiao, H., Xia, Z., Zhang, L., Xiao, N., 2015. Analysis of malaria epidemiological characteristics in the People's Republic of China, 2004–2013. *Am. J. Trop. Med. Hyg.* 93, 293–299.
- Gething, P.W., Elyazar, I.R., Moyes, C.L., Smith, D.L., Battle, K.E., Guerra, C.A., Patil, A.P., Tatem, A.J., Howes, R.E., Myers, M.F., George, D.B., Horby, P., Wertheim, H.F., Price, R.N., Mueller, I., Baird, J.K., Hay, S.I., 2012. A long neglected world malaria map: *Plasmodium vivax* endemicity in 2010. *PLoS Negl. Trop. Dis.* 6, e1814.
- Guerra, C.A., Howes, R.E., Patil, A.P., Gething, P.W., Van Boeckel, T.P., Temperley, W.H., Kabaria, C.W., Tatem, A.J., Manh, B.H., Elyazar, I.R., Baird, J.K., Snow, R.W., Hay, S.I., 2010. The international limits and population at risk of *Plasmodium vivax* transmission in 2009. *PLoS Negl. Trop. Dis.* 4, e774.
- Haque, U., Scott, L.M., Hashizume, M., Fisher, E., Haque, R., Yamamoto, T., Glass, G.E., 2012. Modelling malaria treatment practices in Bangladesh using spatial statistics. *Malar. J.* 11, 63.
- Haque, U., Overgaard, H.J., Clements, A.C., Norris, D.E., Islam, N., Karim, J., Roy, S., Haque, W., Kabir, M., Smith, D.L., Glass, G.E., 2014. Malaria burden and control in Bangladesh and prospects for elimination: an epidemiological and economic assessment. *Lancet Glob. Health* 2, e98–e105.
- Hui, F.M., Xu, B., Chen, Z.W., Cheng, X., Liang, L., Huang, H.B., Fang, L.Q., Yang, H., Zhou, H.N., Yang, H.L., Zhou, X.N., Cao, W.C., Gong, P., 2009. Spatio-temporal distribution of malaria in Yunnan Province, China. *Am. J. Trop. Med. Hyg.* 81, 503–509.
- Kelly-Hope, L.A., Hemingway, J., McKenzie, F.E., 2009. Environmental factors associated with the malaria vectors *Anopheles gambiae* and *Anopheles funestus* in Kenya. *Malar. J.* 8, 268.
- Li, N., Parker, D.M., Yang, Z., Fan, Q., Zhou, G., Ai, G., Duan, J., Lee, M.C., Yan, G., Matthews, S.A., Cui, L., Wang, Y., 2013. Risk factors associated with slide positivity among febrile patients in a conflict zone of north-eastern Myanmar along the China–Myanmar border. *Malar. J.* 12, 361.
- Li, Q., Yang, F., Liu, R., Luo, L., Yang, Y., Zhang, L., Liu, H., Zhang, W., Fan, Z., Yang, Z., Cui, L., He, Y., 2015. Prevalence and molecular characterization of glucose-6-phosphate dehydrogenase deficiency at the China–Myanmar border. *PLoS One* 10, e0134593.
- Lin, H., Lu, L., Tian, L., Zhou, S., Wu, H., Bi, Y., Ho, S.C., Liu, Q., 2009. Spatial and temporal distribution of falciparum malaria in China. *Malar. J.* 8, 130.
- Marsh, K., 2010. Research priorities for malaria elimination. *Lancet* 376, 1626–1627.
- Moonen, B., Cohen, J.M., Snow, R.W., Slutsker, L., Drakeley, C., Smith, D.L., Abeyasinghe, R.R., Rodriguez, M.H., Maharaj, R., Tanner, M., Targett, G., 2010. Operational strategies to achieve and maintain malaria elimination. *Lancet* 376, 1592–1603.
- Mosha, J.F., Sturrock, H.J., Brown, J.M., Hashim, R., Kibiki, G., Chandramohan, D., Gosling, R.D., 2014. The independent effect of living in malaria hotspots on future malaria infection: an observational study from Misungwi, Tanzania. *Malar. J.* 13, 445.
- Parker, D.M., Matthews, S.A., Yan, G., Zhou, G., Lee, M.C., Sirichaisinthop, J., Kiattibutr, K., Fan, Q., Li, P., Sattabongkot, J., Cui, L., 2015. Microgeography and molecular epidemiology of malaria at the Thailand–Myanmar border in the malaria pre-elimination phase. *Malar. J.* 14, 198.
- Phimpraphi, W., Paul, R.E., Yimsamran, S., Puangsart-art, S., Thanyavanich, N., Maneboonyang, W., Prommongkol, S., Sornklom, S., Chaimungkun, W., Chavez, I.F., Blanc, H., Looareesuwan, S., Sakuntabhai, A., Singhasivanon, P., 2008. Longitudinal study of *Plasmodium falciparum* and *Plasmodium vivax* in a Karen population in Thailand. *Malar. J.* 7, 99.
- Ruktanonchai, C.W., Pindolia, D.K., Striley, C.W., Odolina, F.T., Cottler, L.B., 2014. Utilizing spatial statistics to identify cancer hot spots: a surveillance strategy to inform community-engaged outreach efforts. *Int. J. Health Geogr.* 13, 39.
- Saxena, R., Nagpal, B.N., Das, M.K., Srivastava, A., Gupta, S.K., Kumar, A., Jeyaseelan, A.T., Baraik, V.K., 2012. A spatial statistical approach to analyze malaria situation at micro level for priority control in Ranchi district, Jharkhand. *Indian J. Med. Res.* 136, 776–782.
- Waltmann, A., Darcy, A.W., Harris, I., Koepfli, C., Lodo, J., Vahi, V., Piziki, D., Shanks, G.D., Barry, A.E., Whittaker, M., Kazura, J.W., Mueller, I., 2015. High rates of asymptomatic, sub-microscopic *Plasmodium vivax* infection and disappearing *Plasmodium falciparum* malaria in an area of low transmission in Solomon Islands. *PLoS Negl. Trop. Dis.* 9, e0003758.
- Wang, Y., Zhong, D., Cui, L., Lee, M.C., Yang, Z., Yan, G., Zhou, G., 2015. Population dynamics and community structure of *Anopheles* mosquitoes along the China–Myanmar border. *Parasites Vectors* 8, 445.
- Wangdi, K., Gatton, M.L., Kelly, G.C., Clements, A.C., 2015. Cross-border malaria: a major obstacle for malaria elimination. *Adv. Parasitol.* 89, 79–107.
- WHO, 2014. World Malaria Report 2014.
- Xu, J.W., Liu, H., Zhang, Y., Guo, X.R., Wang, J.Z., 2015. Risk factors for border malaria in a malaria elimination setting: a retrospective case-control study in Yunnan, China. *Am. J. Trop. Med. Hyg.* 92, 546–551.
- Yu, G., Yan, G., Zhang, N., Zhong, D., Wang, Y., He, Z., Su, F., Fu, W., Yang, F., Chen, B., 2013. The *Anopheles* community and the role of *Anopheles minimus* on malaria transmission on the China–Myanmar border. *Parasites Vectors* 6, 264.
- Yuan, L., Wang, Y., Parker, D.M., Gupta, B., Yang, Z., Liu, H., Fan, Q., Cao, Y., Xiao, Y., Lee, M.C., Zhou, G., Yan, G., Baird, J.K., Cui, L., 2014. Therapeutic responses of *Plasmodium vivax* malaria to chloroquine and primaquine treatment in Northeastern Myanmar. *Antimicrob. Agents Chemother.* 59, 1230–1235.
- Zhang, L., Feng, J., Xia, Z.G., 2014a. Malaria situation in the People's Republic of China in 2013. *Chin. J. Parasitol. Parasitic Dis.* 32, 407–413.
- Zhang, Q., Lai, S., Zheng, C., Zhang, H., Zhou, S., Hu, W., Clements, A.C., Zhou, X.N., Yang, W., Hay, S.I., Yu, H., Li, Z., 2014b. The epidemiology of *Plasmodium vivax* and *Plasmodium falciparum* malaria in China, 2004–2012: from intensified control to elimination. *Malar. J.* 13, 419.
- Zhou, G., Sirichaisinthop, J., Sattabongkot, J., Jones, J., Bjornstad, O.N., Yan, G., Cui, L., 2005. Spatio-temporal distribution of *Plasmodium falciparum* and *P. Vivax* malaria in Thailand. *Am. J. Trop. Med. Hyg.* 72, 256–262.
- Zhou, S.S., Wang, Y., Fang, W., Tang, L.H., 2008. Malaria situation in the People's Republic of China in 2007. *Chin. J. Parasitol. Parasitic Dis.* 26, 401–403.
- Zhou, G., Sun, L., Xia, R., Duan, Y., Xu, J., Yang, H., Wang, Y., Lee, M.C., Xiang, Z., Yan, G., Cui, L., Yang, Z., 2014a. Clinical malaria along the China–Myanmar border

- Yunnan Province, China, January 2011–August 2012. *Emerg. Infect. Dis.* 20, 675–678.
- Zhou, W.G., Qu, Y., Wang, W.G., Tang, S.Y., 2014b. Application of health education of house-to-house visit in malaria prevention and control. *Chin. J. Schistosomiasis Control* 26, 517–521.
- Zhou, G., Afrane, Y.A., Malla, S., Githeko, A.K., Yan, G., 2015. Active case surveillance, passive case surveillance and asymptomatic malaria parasite screening illustrate different age distribution, spatial clustering and seasonality in western Kenya. *Malar. J.* 14, 41.