

# Lawrence Berkeley National Laboratory

## Lawrence Berkeley National Laboratory

### Title

THE ENERGY DEPENDENCE OF  $^{209}\text{Bi}$  FRAGMENTATION IN RELATIVISTIC NUCLEAR COLLISIONS

### Permalink

<https://escholarship.org/uc/item/7pk4c5xt>

### Author

Aleklett, K.

### Publication Date

1980-07-01



# Lawrence Berkeley Laboratory

UNIVERSITY OF CALIFORNIA

Submitted to Physical Review C


THE ENERGY DEPENDENCE OF  $^{209}\text{Bi}$  FRAGMENTATION  
IN RELATIVISTIC NUCLEAR COLLISIONS

K. Aleklett, D.J. Morrissey, W. Loveland,  
P.L. McGaughey, and G.T. Seaborg

July 1980

**TWO-WEEK LOAN COPY**

This is a Library Circulating Copy  
which may be borrowed for two weeks.  
For a personal retention copy, call  
Tech. Info. Division, Ext. 6782



LBL-11273  
c.2

The Energy Dependence of  $^{209}\text{Bi}$  Fragmentation  
in Relativistic Nuclear Collisions

K. Aleklett<sup>†</sup>, D.J. Morrissey, W. Loveland<sup>††</sup>, P.L. McGaughey,  
and G.T. Seaborg

Nuclear Science Division, Lawrence Berkeley Laboratory  
Berkeley, California 94720

NUCLEAR REACTIONS  $^{209}\text{Bi}$  ( $^{12}\text{C},\text{X}$ ) Au,  $E = 4.8, 25.2$  GeV;  $^{209}\text{Bi}$  ( $^{20}\text{Ne},\text{X}$ ) Au,  $E = 8.0$  GeV; measured Au isotopic distributions, relativistic heavy ions, target fragmentation, Ge(Li) spectroscopy.

ABSTRACT

The results of cross section measurements for the reactions  $^{209}\text{Bi}$  ( $^{12}\text{C},\text{X}$ ) Au,  $E = 4.8$  and  $25.2$  GeV and  $^{209}\text{Bi}$  ( $^{20}\text{Ne},\text{X}$ ) Au,  $E = 8.0$  GeV are reported. The observed yields of the gold isotopes show a similar dependence on mass number for each reaction, differing slightly in the position of the centroid of the distribution. As the projectile energy increases, the inferred excitation energy of the primary residues remains the same or decreases slightly. This observation is in agreement with the predictions of the intranuclear cascade model of relativistic heavy ion collisions.

gamma ray emitting nuclei were identified using standard techniques<sup>4</sup> and the production cross sections were calculated for those activities observed in the unseparated foils. The activities observed in the chemical samples were normalized to those seen in the unseparated foils. However, this normalization was not possible for the 25.2 GeV  $^{12}\text{C}$  data because only one foil was irradiated and chemically processed and therefore, these production cross sections are expressed in arbitrary units. The contribution of secondary induced reactions to the Au product yields was studied by comparison of the activities of the two unseparated foils from the 4.8 GeV  $^{12}\text{C} + ^{209}\text{Bi}$  bombardment which differed by a factor of  $\sim 2$  in thickness. This comparison showed that  $\sim 30\%$  of the  $^{198}\text{Au}$  yield in the thickest target ( $\sim .25\%/ \text{mg/cm}$ ) was due to such effects, while the lighter nuclidic activities had little or no secondary contributions. This

secondary reaction correction was applied to the  $A > 198$  product yields for the two reactions with  $\sim 135 \text{ mg/cm}^2$  targets, while all other secondary reaction contributions were assumed to be negligible.

The measured cumulative yield cross sections (which include contributions from nuclei produced by radioactive decay) were corrected for this decay feeding by an iterative fitting of Gaussian distributions to the decay-corrected (and independent yield) cross sections.<sup>4</sup> The resulting calculated and measured independent yield cross sections  $\sigma(Z,A)$  are shown in Fig. 1 and included in Table I. The uncertainties given for  $\sigma(Z,A)$  reflect the contribution from statistical uncertainties from the measurements and an adhoc factor of 20% of the  $\beta$ -decay feeding correction where applied.<sup>5</sup> The yield of  $^{194}\text{Au}$  is depressed relative to the other nuclidic yields in all the reactions for reasons that are not clear.

The number of Au isotopes that were observed in each reaction was limited by two factors; the low activity levels of the neutron rich species and the short half lives (relative to the chemical separation time and length of the bombardment) of the neutron deficient species. Cursory inspection of Fig. 1 shows that the distributions of Au fragments are very similar for the three reactions studied in this work. A more detailed examination of the data shows that there is a slight ( $\lesssim 1$  amu) shift of the distribution centroids toward larger A values as the projectile energy increases. Thus, although the incident projectile energy changes by over 20 GeV, the excitation energy imparted to target fragments changes by less than 10 MeV. The ratio of the magnitude of the integrated Au isotopic yields from the 0.40 GeV/A  $^{20}\text{Ne}$  bombardment to that from the 0.40 GeV/A  $^{12}\text{C}$  bombardment is in rough agreement with the factorization prediction<sup>6</sup> of  $\sigma_R(^{20}\text{Ne}+^{209}\text{Bi})/\sigma_R(^{12}\text{C}+^{209}\text{Bi}) = 3644 \text{ mb}/3030 \text{ mb} = 1.20$ . Both of these results are consistent with the observations of Cumming et al.<sup>7</sup> who found that the target fragment isobaric and isotopic distributions were identical for many reactions with copper targets.

We are left with the question of whether this apparent saturation in energy transfer is predicted by current models of fragmentation reactions. One highly developed model of such collisions which makes predictions about the fragmentation products is the intranuclear cascade model of Yariv and Fraenkel.<sup>8</sup> In this model the collision process is described by the superposition of a series of nucleon-nucleon collisions with proper accounting of the nucleon-nucleon cross section, angular distributions, Fermi energy distributions and particle-hole excitations

in the fragments before de-excitation. The calculated positions of the centroids of the gold isotopic distributions using this model are contained in Table I along with the experimental results. The calculations are in approximate agreement with the results of our study but predict values of the centroids of the distributions which are systematically more neutron deficient ( $\sim 1-2$  amu) than our data. (The excitation energies of the primary fragments were  $\sim 125-150$  MeV.) Thus, the experimentally observed limiting fragmentation behavior of high mass nuclei is predicted by this model of relativistic nucleus-nucleus collisions.

What universal feature of RHI reactions is responsible for this behavior? In this case involving very peripheral collisions (impact parameter  $b \sim 9$  fm  $> R_{Bi} \sim 7.0$  fm according to firestreak model<sup>10</sup> calculations), it is simply a matter that the predicted changes in the experimental distributions are very small compared to the sensitivity of the experimental measurement. Unfortunately, selection of events involving more central collisions (i.e., fragments where  $A_{\text{target}} - A_{\text{fragment}} \geq 20$ ) where the changes in the distributions would be larger suffers from the fact that in such events the primary reaction processes are observed by the subsequent de-excitation of the primary fragments.<sup>9</sup> Thus we conclude that although we have picked a relatively favorable case to study, our observation of limiting fragmentation like that of others could mean either a saturation in the transferred energy with increasing projectile energy or that changes in the transferred energy that are too small to detect with current techniques of fragment yield measurement.

We would like to thank Mr. K. J. Moody for his assistance in evaluating the secondary corrections, and Mr. M. Varney, Drs. K. E. Thomas and R. J. Otto for their aid in performing the chemical separations.


Two of us (WL, KA) wish to thank their home institutions for financial support during this work which was supported in part by U.S. Department of Energy under Contract W-7405-ENG-48.

## References

- <sup>†</sup> Permanent Address: Studsvik Scientific Research Laboratory, Nyköping, Sweden.  
<sup>††</sup> Permanent Address: Dept. of Chemistry, Oregon State University, Corvallis, Oregon 97331.
1. Y.P. Vijoyi, T.J.M. Symons, P. Doll, D.E. Greiner, H.H. Heckman, D.L. Hendrie, P.J. Lindstrom, J. Mahoney, D.K. Scott, K. Van Bibber, G.D. Westfall, H. Wieman, H.J. Crawford, C. McParland, and C.K. Gelbke, Phys. Rev. Lett., 42, 33 (1979).
  2. N.T. Porile, G.D. Cole and C.R. Rudy, Phys. Rev. C19, 2288 (1979).
  3. The bismuth metal foils were dissolved in conc. HCl-conc. HNO<sub>3</sub> (1:2) with carriers. After dissolution, excess NO<sub>2</sub> was driven off, the solution was cooled and diluted with water. The Au was extracted twice into ether, washed twice with 1.5 M HCl and plated on Pt. This procedure required approximately 25 minutes.
  4. D.J. Morrissey, D. Lee, R.J. Otto and G.T. Seaborg, Nucl. Inst. Methods, 158, 499 (1979).
  5. A similar procedure was adopted by T.H. Ku and P.J. Karol, Phys. Rev. C16, 1984 (1977).
  6. Hard sphere reaction cross sections from H.H. Heckman, D.E. Greiner, P.J. Lindstrom and H. Shwe, Phys. Rev. C17, 1735 (1978).
  7. J.B. Cumming, R.W. Stoenner, and P.E. Haustein, Phys. Rev. C14, 1554 (1976); J.B. Cumming, P.E. Haustein, R.W. Stoenner, L. Mausner, and R.A. Naumann, Phys. Rev. C10, 739 (1974); J.B. Cumming, P.E. Haustein, T.J. Ruth and G.J. Virtes, Phys. Rev. C17, 1632 (1978).
  8. Y. Yariv and Z. Fraenkel, Phys. Rev. C20, 2227 (1979).
  9. D.J. Morrissey, L.F. Oliveira, J.O. Rasmussen, G.T. Seaborg, Y. Yariv, and Z. Fraenkel, Phys. Rev. Lett. 43, 1179 (1979).
  10. W.D. Myers, Nucl. Phys. A296, 117 (1978); J. Gossett, J.I. Kapusta, and G.D. Westfall, Phys. Rev. C18, 844 (1978).

Table I  
Experimental Conditions and Results

Projectile	$^{12}\text{C}$	$^{20}\text{Ne}$	$^{12}\text{C}$
Energy (GeV)	4.8	8.0	25.2
<i>Target Thickness (mg/cm<sup>2</sup>)</i>			
(1) unseparated	112.6, 227.7	116	---
(2) chemical	132.9	137	~50
<i>Bombardment Statistics (total particles [hours])</i>			
(1) unseparated	$6.19 \times 10^{13}$ [13.7]	$3.64 \times 10^{13}$ [13.5]	---
(2) chemical	$6.19 \times 10^{13}$ [13.7]	$3.75 \times 10^{13}$ [13.5]	$5.3 \times 10^{12}$ [2.7]
<i>Independent Yield Cross Sections</i>			
	(mb)	(mb)	(arbitrary units)
$^{190}\text{Au}$	$8.2 \pm 1.4$	$13.6 \pm 2.5$	---
$^{191}\text{Au}$	$8.1 \pm 1.4$	$10.4 \pm 1.9$	$14.2 \pm 1.4$
$^{192}\text{Au}$	$7.7 \pm 1.3$	$9.4 \pm 1.9$	$15.5 \pm 1.5$
$^{193}\text{Au}$	$7.7 \pm 1.3$	$8.2 \pm 1.6$	$14.6 \pm 1.5$
$^{194}\text{Au}$	$1.7 \pm 0.2$	$1.8 \pm 0.2$	$4.6 \pm 0.5$
$^{195}\text{Au}$	$3.4 \pm 0.3$	$3.7 \pm 0.4$	---
$^{196\text{m}+g}\text{Au}$	$1.21 \pm 0.1$	$1.24 \pm 0.94$	$4.2 \pm 0.5$
$^{198\text{m}+g}\text{Au}$	$0.65 \pm .052$	$0.51 \pm .039$	$2.3 \pm 0.2$
$^{199}\text{Au}$	$0.54 \pm .058$	$0.24 \pm .026$	$1.4 \pm 0.1$
<i>Isotopic Distribution Centroids &lt;A&gt; for Z = 79</i>			
Experimental	$191.4 \pm 0.3$	$191.6 \pm 0.5$	$192.1 \pm 0.3$
Cascade Model <sup>8</sup>	$189.4 \pm 0.3$	$189.5 \pm 0.5$	$190.6 \pm 0.5$



XBL 807-10700

Fig. 1. The isotopic production cross sections for gold products from the reaction of 4.8 and 25.2 GeV  $^{12}\text{C}$  and 8.0 GeV,  $^{20}\text{Ne}$  with  $^{209}\text{Bi}$  are shown. The lines represent the best Gaussian fit to each distribution with the arrows showing the distribution centroids.

This report was done with support from the United States Energy Research and Development Administration. Any conclusions or opinions expressed in this report represent solely those of the author(s) and not necessarily those of The Regents of the University of California, the Lawrence Berkeley Laboratory or the United States Energy Research and Development Administration.