Lawrence Berkeley National Laboratory

Recent Work

Title

MOMENTUM DEPENDENCE OF PHASE SHIFTS

Permalink

<https://escholarship.org/uc/item/7pw485hv>

Authors

Goebel, Charles J.
Karplus, Robert
Ruderman, Malvin A.

Publication Date

1955-06-10

UNIVERSITY OF
CALIFORNIA

*Radiation
Laboratory*

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 5545*

BERKELEY, CALIFORNIA

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

UNIVERSITY OF CALIFORNIA

Radiation Laboratory
Berkeley, California

Contract No. W-7405-eng-48

MOMENTUM DEPENDENCE OF PHASE SHIFTS

Charles J. Goebel, Robert Karplus, and Malvin A. Ruderman

June 10, 1955

MOMENTUM DEPENDENCE OF PHASE SHIFTS

Charles J. Goebel, Robert Karplus, and Malvin A. Ruderman*

Radiation Laboratory and Physics Department
University of California, Berkeley, California

June 10, 1955

ABSTRACT

A lower limit is found for the momentum derivative of the scattering phase shifts of a relativistic neutral two-particle system when the interaction is of finite range.

* This work was supported in part by the United States Atomic Energy Commission.

MOMENTUM DEPENDENCE OF PHASE SHIFTS

Charles J. Goebel, Robert Karplus, and Malvin A. Ruderman

Some restrictions on the momentum dependence of the phase shifts for a nonrelativistic scattering problem have been discussed recently by Wigner.¹ In this note we derive a somewhat stronger restriction for the phase shifts for the relativistic scattering of neutral particles confined to a single channel. The latter condition essentially means that we are dealing with a first quantized field.

The proof is based upon the unitarity of the partial S-matrix referring to a single phase shift and to the completeness, outside some radius a_0 , of the asymptotic wave functions describing the scatterer and scattered particle in the center-of-mass system in a given state of angular momentum. We shall restrict our discussion to S-waves; analogous theorems are valid for other angular momenta.

For nonrelativistic S-wave scattering, unitarity and completeness have been shown^{2,3} to lead to the conclusion that $S(k)e^{2iak}$ $[a > a_0]$, if it has an analytic continuation,⁴ is a regular function of k for $\text{Im } k > 0$ as long as no bound states are present. If there are no branch points, Ning Hu³ has shown that one can therefore write

$$S(k) = e^{2i\delta(k)} = e^{-2iak} \prod_s \frac{k - k_s^*}{k - k_s} \frac{k + k_s}{k + k_s^*}, \quad (1)$$

$$a > a_0, \quad \text{Im } k_s < 0.$$

The k_s for different s are not assumed to be necessarily distinct.

The form of Eq. (1) is dictated by the absence of poles in the upper half plane and by the condition

-4-

$$S(k) = e^{2i\delta(k)} = e^{-2i\delta(-k)} = S^*(-k). \quad (2)$$

For a relativistic particle the energy is a double-valued function, $E = \pm(k^2 + \mu^2)^{\frac{1}{2}}$; in general the scattering amplitude depends both on k and E so that cuts must be introduced in the k -plane along the imaginary k axis from $i\mu$ to $i\infty$ and $-i\mu$ to $-i\infty$. Following Hu,³ we can write

$$S(k, E) = e^{-2iak} \prod_s \frac{(k_s + k_s^*)(E - E_s^*) - (E_s^* - E_s)(k - k_s^*)}{(k_s + k_s^*)(E - E_s) - (E_s - E_s^*)(k - k_s)} \quad (3)$$

$S(k)$ has a pole at $k = k_s$, $E = E_s = \sqrt{k_s^2 + \mu^2}$ but need not have a pole at $k = k_s$, $E = -E_s$; we have $S^*(k, e) = S(-k, E)$ in accord with unitarity but there is no simple relation between $S(k, E)$ and $S(k, -E)$. But for a real field (neutral particles only) $S(k, E) = S(k, -E)$, so that for every factor in Eq. (3) contributing a pole at k_s, E_s there is one contributing a pole at $k_s, -E_s$. When the pair of factors with E_s are multiplied, Eq. (3) reduces to the form of Eq. (1).

We may solve Eq. (1) for the phase shift and obtain

$$\delta(k) = -ak + \sum_s \tan^{-1} \alpha_s(k) = -ak + \sum_s \theta_s(k), \quad (4)$$

where we have introduced the angle

$$\theta_s(k) = \tan^{-1} \alpha_s(k), \quad (5)$$

and the function $\alpha_s(k)$,

$$\alpha_s(k) = 2k \operatorname{Im} k_s / (k^2 - |k_s|^2) . \quad (5')$$

The properties of the derivative of $S(k)$ with respect to k , denoted by a dot, are related to those of the derivatives of θ and α ,

$$\dot{\delta} = -a + \sum_s \dot{\theta}_s = -a + \sum_s \frac{\dot{\alpha}_s}{1 + \alpha_s^2} . \quad (6)$$

Because the poles of S are all in the lower half plane, however, $\dot{\alpha}$ is always positive,

$$\dot{\alpha}_s = (-2 \operatorname{Im} k_s) \frac{k^2 + |k_s|^2}{(k^2 - |k_s|^2)^2} > 0 , \quad (7)$$

whence

$$\dot{\delta} > -a . \quad (8)$$

Actually, the properties of α_s permit a still stronger statement on the lower limit of $\dot{\delta}$. By comparing Eqs. (5') and (7), we may conclude that

$$\dot{\alpha}_s > |\alpha_s|/k . \quad (9)$$

But this implies that θ_s has the lower limit

$$\dot{\theta}_s > \frac{1}{k} \frac{|\alpha_s|}{1 + \alpha_s^2} = \frac{1}{2k} |\sin 2\theta_s| \quad (10)$$

and that, in turn, $\dot{\delta}$ has to be greater than $-a$ by a positive definite amount

-6-

$$\dot{\delta} > -a + \frac{1}{2k} \sum_s |\sin 2\theta_s| \geq -a + \frac{1}{2k} |\sin 2(\delta + ka)|. \quad (11)$$

The last part of the inequality can be easily deduced by induction, starting with a sum over two terms,

$$\begin{aligned} |\sin(\theta_1 + \theta_2)| &= |\sin\theta_1 \cos\theta_2 + \cos\theta_1 \sin\theta_2| \\ &\leq |\sin\theta_1| |\cos\theta_2| + |\sin\theta_2| |\cos\theta_1| \leq |\sin\theta_1| + |\sin\theta_2|. \end{aligned} \quad (12)$$

Equation (11) above corresponds to Eq. (5a) of Reference 1, but differs in that the oscillating term always makes a positive contribution.

When bound states are present $S(k)$ of Eq. (1) has simple poles in the upper half plane at $k = i\kappa_\lambda$ such that the energy of the bound state is $\sqrt{\mu^2 - \kappa_\lambda^2}$. Then the rhs of Eq. (1) will have a factor

$$\prod_\lambda \frac{k + i\kappa_\lambda}{k - i\kappa_\lambda} \quad (13)$$

and Eq. (12) becomes

$$\dot{\delta} > -a + \frac{1}{2k} |\sin(2\delta + 2ka)| - \sum_\lambda \frac{\kappa_\lambda}{k^2 + \kappa_\lambda^2}. \quad (14)$$

Another inequality which can be stronger than Eq. (11) and is valid for real fields even if inelastic processes enter at higher k , follows from an application of Cauchy's theorem to $\exp(2i\delta(k) + 2ika)$. Since the integrand has no poles in the finite half plane if there are no bound states,

Cauchy's theorem yields

$$e^{2i[\delta(k) + ka]} = \frac{P}{\pi i} \int_{-\infty}^{\infty} \frac{dt}{t - k} e^{2i[\delta(t) + ta]} \quad (15)$$

By differentiating Eq. (15) logarithmically and using $\delta(k) = -\delta(-k)$, we obtain

$$\dot{\delta}(k) = -a + \frac{1}{\pi} \int_{-\infty}^{\infty} \frac{dt \sin^2(\delta(t) - \delta(k) + at - ak)}{(t - k)^2} \quad (16)$$

$> -a$.

The integrand of the rhs is always positive. Therefore a knowledge of $\delta(k)$, even over a limited region of k , contributes a lower limit to the entire integral in Eq. (16).⁵

For the partial S-matrix of Eq. (3), which describes the scattering of a charged relativistic particle, even the inequality (8) will not hold in general; it does not seem possible, therefore, to apply the inequalities (8), (12), and (16) to the scattering of charged mesons. Furthermore the possibility of charge exchange makes even Eq. (3) inadequate for the S-wave scattering of π^0 -mesons by nucleons. Under the transformation $E \rightarrow -E$ the partial S-matrix $e^{2i\delta_3}$ for $\pi^+ + p \rightarrow \pi^+ + p$ becomes that for $\pi^- + p \rightarrow \pi^- + p$.⁶ In a charge-independent theory, for instance, $\pi^- + p \rightarrow \pi^0 + n$ can take place, $e^{2i\delta_3(-E)}$ can not be unitary, and $\delta_3(-E)$ cannot be real.

Nevertheless some information about the range of interaction can be inferred for the meson-nucleon system. Completeness and unitarity give the relation

-8-

$$\int_{-\infty}^{\infty} dk e^{ikr} S(k, +\sqrt{k^2 + \mu^2}) + \int_{-\infty}^{\infty} dk e^{ikr} S(k, -\sqrt{k^2 + \mu^2}) = 0$$

where $S(k, +\sqrt{k^2 + \mu^2})$ may be the partial S-matrix for $\pi^{\pm} + p \rightarrow \pi^{\pm} + p$ or $\pi^{\pm} + p \rightarrow \pi^0 + n$. Charge independence relates the charge exchange amplitude to the two elastic amplitudes, and the π^+ amplitude becomes the complex conjugate of that for π^- when the sign of the energy is changed. Completeness, unitarity, and charge independence lead to the single relation

$$2e^{2i\phi_3(k)} + e^{2i\phi_1(k)} = \frac{P}{\pi i} \int_{-\infty}^{\infty} \frac{2e^{2i\phi_3(t)} + e^{2i\phi_1(t)}}{t - k} dt \quad (17)$$

where

$$\phi_3(k) = \delta_3(k) + ka \quad \text{and} \quad \phi_1(k) = \delta_1(k) + ka.$$

Differentiating both sides of Eq. (17) with respect to k and then setting $k = 0$, we obtain

$$2\delta_3(0) + \delta_1(0) + 3a = \pi \int_{-\infty}^{\infty} \frac{2\sin^2 \phi_3(t) + \sin^2 \phi_1(t)}{t^2} dt. \quad (18)$$

With Orear's extrapolated phase shifts,

$$\delta_3 = - .11 \frac{k}{\mu c} \quad \text{and} \quad \delta_1 = + .16 \frac{k}{\mu c}, \quad (19)$$

the complete neglect of the positive definite integral yields $a \geq .02 \frac{\pi}{\mu c}$.

A stronger limit on a is easily obtained if we use a linear approximation to the S-wave phase shifts.⁷ The integration in Eq. (18) can be carried out, with the result

-9-

$$2 \dot{\phi}_3(0) + \dot{\phi}_1(0) = 2 \left| \dot{\phi}_3(0) \right| + \left| \dot{\phi}_1(0) \right| . \quad (20)$$

From this we can infer

$$a \geq 0.11 \frac{\pi}{\mu c} . \quad (21)$$

REFERENCES

1. E. P. Wigner, Phys. Rev. 98, 145 (1955).
2. W. Heisenberg, Z. Naturforschung 1, 608 (1946).
3. Ning Hu, Phys. Rev. 74, 131 (1948).
4. N. G. van Kampen, Phys. Rev. 89, 1072 (1953) and Phys. Rev. 91, 1267 (1953).
5. If multiple processes occur at momenta greater than k , Eq. (16) still holds, with the $\sin^2 ()$ of the integrand replaced by $e^{-2\delta_2} \sin^2 () + e^{-\delta_2} \sinh (\delta_2)$, where δ_2 is the imaginary part of δ .
6. M. Goldberger, Phys. Rev. (in press). We are indebted to Professor Goldberger for furnishing us a copy of his paper before publication.
7. J. Orear, Phys. Rev. 96, 176 (1954).