

UC Davis

UC Davis Previously Published Works

Title

Multicenter Study of Isavuconazole MIC Distributions and Epidemiological Cutoff Values for the *Cryptococcus neoformans*-*Cryptococcus gattii* Species Complex Using the CLSI M27-A3 Broth Microdilution Method

Permalink

<https://escholarship.org/uc/item/7q44s7s7>

Journal

Antimicrobial Agents and Chemotherapy, 59(1)

ISSN

0066-4804

Authors

Espinel-Ingroff, A
Chowdhary, A
Gonzalez, GM
et al.

Publication Date

2015

DOI

10.1128/aac.04055-14

Peer reviewed

Multicenter Study of Isavuconazole MIC Distributions and Epidemiological Cutoff Values for the *Cryptococcus neoformans*-*Cryptococcus gattii* Species Complex Using the CLSI M27-A3 Broth Microdilution Method

A. Espinel-Ingróff,^a A. Chowdhary,^b G. M. Gonzalez,^c J. Guinea,^d F. Hagen,^e J. F. Meis,^{e,f} G. R. Thompson III,^g J. Turnidge^h

VCU Medical Center, Richmond, Virginia, USA^a; Vallabh Patel Chest Institute, University of Delhi, Delhi, India^b; Universidad Autónoma de Nuevo León, Monterrey, Nuevo León, Mexico^c; Hospital General Universitario Gregorio Marañón, School of Medicine-Universidad Complutense, Madrid, Spain^d; Department of Medical Microbiology & Infectious Diseases, Canisius Wilhelmina Hospital, Nijmegen, The Netherlands^e; Department of Medical Microbiology, Radboud University Medical Center, Nijmegen, The Netherlands^f; University of California Davis Medical Center, Davis, California, USA^g; and University of Adelaide, Adelaide, Australia^h

Epidemiological cutoff values (ECVs) of isavuconazole are not available for *Cryptococcus* spp. The isavuconazole ECVs based on wild-type (WT) MIC distributions for 438 *Cryptococcus neoformans* nongenotyped isolates, 870 isolates of genotype VNI, and 406 *Cryptococcus gattii* isolates from six laboratories and different geographical areas were 0.06, 0.12, and 0.25 µg/ml, respectively. These ECVs may aid in detecting non-WT isolates with reduced susceptibilities to isavuconazole.

Infections caused by members of the *Cryptococcus neoformans*-*Cryptococcus gattii* species complex are considered the second most common severe mycoses among immunocompromised and nonimmunocompromised hosts in certain regions. Although infections caused by *C. neoformans* var. *grubii* (serotype A) and, to a lesser degree, by *C. neoformans* var. *neoformans* (serotype D) are seen worldwide, those by *C. gattii* (serotypes B and C) are more geographically restricted. Irrespective of the species, cryptococcal disease is associated with high mortality rates ($\geq 12.7\%$) (1–4). *C. neoformans* var. *grubii* comprises genotypes VNI, VNII/VNB, VNII (AFLP1, AFLP1A, AFLP1B), *C. neoformans* var. *neoformans* comprises genotypes VNIV (AFLP2) and VNIII (AFLP3), and *C. gattii* comprises genotypes VGI (AFLP4), VGII (AFLP6), VGIV (AFLP10), VGIII (AFLP5), and VGIV (AFLP7) (1, 5–7). Amphotericin B, its lipid formulations, and fluconazole are recommended as the primary alternative induction treatments and/or as salvage consolidation therapies for infections caused by *C. neoformans* and *C. gattii* (3, 8). Isavuconazole (BAL4815, codeveloped by Basilea Pharmaceutica International Ltd. [Basel, Switzerland] and Astellas Pharma [Tokyo, Japan]) is a new water-soluble triazole with favorable pharmacodynamic and pharmacokinetic parameters and is under clinical evaluation (phase III) for the treatment of invasive aspergillosis and other mycoses. Isavuconazole's mode of action is the inhibition of ergosterol biosynthesis (enzymes 14- α -sterol demethylases A and B, which are encoded by the *cyp51* [*ERG11*] gene), similar to that of the other azoles. The *in vitro* activity of isavuconazole is also similar to that of most licensed triazoles against *Aspergillus* spp., *Candida* spp., and *Cryptococcus* spp.; it also has activity comparable to that of posaconazole and, to a certain extent, itraconazole against some *Mucorales* species (9).

The Clinical and Laboratory Standards Institute (CLSI) has established standard conditions for testing the susceptibilities of *Candida* spp. and *Aspergillus* spp. to isavuconazole (10). In the absence of clinical breakpoints (CBPs) (the ultimate means of differentiating treatable and nontreatable isolates), epidemiological cutoff values (ECVs) have been defined for the more prevalent *Candida* spp. and the *C. neoformans*-*C. gattii* species complex ver-

sus those of the four licensed triazoles (11, 12). The ECV (also known as the ECOFF or CO_{WT}) is the highest wild-type (WT)-susceptible value; it is based on MIC distributions where there are two distinct populations, (i) the WT population of isolates/MICs with no detectable acquired or mutational resistance to the drug being evaluated and (ii) the non-WT population or isolates that harbor one or more resistance markers (13). The ECV based on pooled data from multiple laboratories serves as an early indication of emerging changes in the patterns of organism susceptibilities to the agent being evaluated or separates WT from non-WT isolates. Although the *in vitro* activity of isavuconazole against members of the *C. neoformans*-*C. gattii* species complex has been evaluated by CLSI M27-A3 methodology (14), ECVs based on data from multiple laboratories, as well as different geographical areas, have not been established for this complex. While data for the present study were submitted from a total of seven laboratories, the distribution data from one of the laboratories were abnormal (mode at the lower concentration tested) and were not included in the analysis. In addition, MICs were not submitted by each laboratory for each of the cryptococcal groups evaluated. Therefore, the purpose of this study was to define isavuconazole ECVs for three groups within the *C. neoformans*-*C. gattii* species complex for which MICs originated from three or four of the six collaborating laboratories.

Each isolate originated from a unique clinical specimen. The

Received 7 August 2014 Returned for modification 1 October 2014
Accepted 7 October 2014

Accepted manuscript posted online 13 October 2014

Citation Espinel-Ingróff A, Chowdhary A, Gonzalez GM, Guinea J, Hagen F, Meis JF, Thompson GR, III, Turnidge J. 2015. Multicenter study of isavuconazole MIC distributions and epidemiological cutoff values for the *Cryptococcus neoformans*-*Cryptococcus gattii* species complex using the CLSI M27-A3 broth microdilution method. *Antimicrob Agents Chemother* 59:666–668. doi:10.1128/AAC.04055-14.

Address correspondence to A. Espinel-Ingróff, avingrof@vcu.edu.

Copyright © 2015, American Society for Microbiology. All Rights Reserved.

doi:10.1128/AAC.04055-14

TABLE 1 Pooled MIC distributions and ECVs for the *Cryptococcus neoformans*-*Cryptococcus gattii* species complex and isavuconazole using the CLSI M27-A3-RPMI microdilution method^a

Species	No. of isolates	No. of isolates with an MIC ^b (μg/ml) of:								MIC range (μg/ml)	Mode (μg/ml) ^d	ECV (μg/ml) ^c	
		0.008	0.016	0.03	0.06	0.12	0.25	0.5	95%			97.5%	
<i>C. neoformans</i> VNI ^e (AFLP1)	870	68	270	274	213	37	7	1	0.008–0.5	0.03	0.12	0.12	
<i>C. neoformans</i> (nongenotyped)	438	10	135	170	96	19	6	2	0.008–0.5	0.03	0.06	0.12	
<i>C. gattii</i> ^e	406	23	78	106	87	87	25	7	0.008–0.5	0.03	0.25	0.25	

^a MICs were pooled from three or four of the six collaborating laboratories that submitted qualifying data for each cryptococcal group.

^b MICs as determined by the CLSI broth microdilution method using standard RPMI 1640 broth.

^c ECVs comprising ≥95% and ≥97.5% of the statistically modeled MIC population.

^d Most frequent MIC.

^e *C. gattii* comprises the VGI through VGIV (also known as AFLP4 through AFLP7 and AFLP10) genotypes.

CLSI broth microdilution isavuconazole MICs used for ECV definition were obtained from the VCU Medical Center (Richmond, VA), the Vallabhbhai Patel Chest Institute, University of Delhi (Delhi, India), Universidad Autónoma de Nuevo León (Monterrey, Nuevo León, Mexico), the Department of Medical Microbiology and Infectious Diseases, Canisius-Wilhelmina Hospital (Nijmegen, The Netherlands), Department of Medical Microbiology, Radboud University Medical Center (Nijmegen, The Netherlands), and the University of California Davis Medical Center (Davis, CA). Species, serotype, and molecular type identifications were performed at each medical center using standard methodologies (15–19). We have aggregated CLSI MICs for 870 *C. neoformans* (VNI molecular genotype), 438 *C. neoformans* nontyped, and 406 *C. gattii* isolates (comprising the 5 molecular genotypes). We also received a set of isavuconazole MICs for *C. gattii* environmental isolates that were not incorporated in the analysis. The MIC ranges for the control isolates (*Candida parapsilosis* ATCC 22019 and *Candida krusei* ATCC 6258) were within 3 to 4 dilutions.

Our MICs were determined using CLSI testing parameters (with standard RPMI 1640 broth [0.2% dextrose], final inoculum concentrations that ranged from 0.4×10^3 to 5×10^3 CFU/ml, and 72 h of incubation); the MICs were the lowest drug concentrations that produced ≥50% growth inhibition compared to that of the growth control (14). MIC distributions obtained in at least three of the seven coded laboratories for each group were listed in Excel spread sheets and aggregated for the ECV statistical calculation by a previously described method (13); the MICs that captured at least 95% and 97.5% of the modeled WT population (instead of the observed values) were obtained. ECVs are not estimated when the distribution is grossly skewed, which precludes statistical fitting, and a laboratory's data are excluded when the distribution from a single laboratory is truncated (mode at the lower or upper end of the distribution). At least 100 MICs from three independent laboratories are required for each group to establish an ECV. Because of that, we pooled the MICs for five molecular genotypes of *C. gattii* and used only the distributions for the *C. neoformans* VNI genotype; data for the other genotypes ranged from 13 to 172 MICs for VGI to VGIV genotypes and from 13 to 46 for VNII to VNIV genotypes, mostly inadequate sample sizes for ECV calculation. In addition, 1 of the 4 *C. neoformans* nontyped distributions received was truncated and was not pooled for the analysis.

The ECV is the highest MIC of the WT population, while the CBP is the MIC that predicts the clinical outcome of therapy (13,

20, 21). Therefore, the ECV detects the non-WT strains with reduced susceptibility (due to mutations) to the agent being evaluated. Aggregated isavuconazole MIC distributions for *C. neoformans* (genotype VNI), *C. neoformans* nontyped, and *C. gattii* (pooled genotypes) isolates are shown in Table 1. Each of the three pooled distributions had the same MIC range (0.008 to 0.5 μg/ml) and mode (0.03 μg/ml), and these values reflect an earlier report of isavuconazole data for *C. neoformans* (22). Comparable results have been observed with a larger number of isolates and the licensed triazoles for the three groups evaluated in the present study: similar posaconazole and voriconazole ranges and modes 1 or 2 dilutions higher (0.06 to 0.12 μg/ml) (12). Our isavuconazole ECVs comprising ≥95% of the modeled population were 0.06 μg/ml (*C. neoformans* nontyped), 0.12 μg/ml (genotype VNI), and 0.25 μg/ml (*C. gattii*); the ECVs that comprised ≥97.5% of the population were either the same or 1 dilution higher (Table 1). Recently established itraconazole, posaconazole, and voriconazole ECVs for the same groups were 1 dilution higher (0.25 to 1 μg/ml), with the highest ECVs for *C. gattii* (12). It is noteworthy that the isavuconazole mode for the environmental *C. gattii* isolates was 0.12 μg/ml. Preliminary clinical data regarding isavuconazole treatment of cryptococcal infections was recently reported (F. Queiroz-Telles, O. A. Cornely, J. Perfect, L. Kovanda, B. Zeiher, and J. Vazquez, presented at the ICCAC Congress, Washington, DC, 5 to 9 September, 2014). Nine patients with either central nervous system or disseminated cryptococcal disease were treated with isavuconazole (200 mg three times daily for 2 days followed by 200 mg every day [intravenous or oral]; VITAL phase III, open-label, multicenter trial for the efficacy and safety of isavuconazole). Of the eight patients evaluated on the 84th day, six were treatment successes (including all *C. gattii* infections), and two were treatment failures. CLSI isavuconazole MICs for the seven available infecting *C. neoformans* and *C. gattii* isolates were ≤0.12 μg/ml, which may be considered WT according to our proposed ECVs (Table 1). These results underline the potential values of ECV data in the clinical setting.

In conclusion, data originating in three laboratories for each of the three groups enabled us to propose species-specific isavuconazole ECVs of 0.06 to 0.25 μg/ml for the *C. neoformans*-*C. gattii* species complex evaluated. The availability of standard CLSI parameters for testing isavuconazole against these species and the ECVs calculated in the present study will aid in monitoring isavuconazole resistance in cryptococcal species and distinguishing non-WT from WT isolates.

REFERENCES

1. Bovers M, Hagen F, Boekhout T. 2008. Diversity of the *Cryptococcus neoformans*-*Cryptococcus gattii* species-complex. *Rev Iberoam Micol* 25: S4–S12. [http://dx.doi.org/10.1016/S1130-1406\(08\)70019-6](http://dx.doi.org/10.1016/S1130-1406(08)70019-6).
2. Harris JR, Lockhart SR, Debess E, Marsden-Haug N, Goldoft M, Wohrle R, Lee S, Smelser C, Park B, Chiller T. 2011. *Cryptococcus gattii* in the United States: clinical aspects of infection with an emerging pathogen. *Clin Infect Dis* 53:1188–1195. <http://dx.doi.org/10.1093/cid/cir723>.
3. Perfect JR, Dismukes WE, Dromer F, Goldman DL, Graybill JR, Hamill RJ, Harrison TS, Larsen RA, Lortholary O. 2010. Clinical practice guidelines for the management of cryptococcal disease: 2010 update by the Infectious Diseases Society of America. *Clin Infect Dis* 50:291–322. <http://dx.doi.org/10.1086/649858>.
4. Desnos-Ollivier M, Patel S, Spaulding AR, Charlier C, Garcia-Hermoso D, Nielsen K, Dromer F. 2010. Mixed infections and *in vivo* evolution in the human fungal pathogen *Cryptococcus neoformans*. *mBio* 1:e00091-10. <http://dx.doi.org/10.1128/mBio.00091-10>.
5. Byrnes EJ, Li W, Lewit Y, Ma H, Voelz K, Ren P, Carter DA, Chatuvedi D, Bildfell RJ. 2010. Emergence and pathogenicity of highly virulent *Cryptococcus gattii* genotypes in the northwest United States. *PLoS Pathog* 6:e1000850. <http://dx.doi.org/10.1371/journal.ppat.1000850>.
6. Byrnes EJ, Wenjun L, Ren P, Lewit Y, Voelz K, Fraser J, Dietrich FS, May RC, Chatuvedi S, Chatuvedi D, Heitman J. 2011. A diverse population of *Cryptococcus gattii* molecular type VGIII in southern Californian HIV/AIDS patients. *PLoS Pathog* 7:e1002205. <http://dx.doi.org/10.1371/journal.ppat.1002205>.
7. Hagen F, Colom MF, Swinne D, Tintelnot K, Iatta R, Montagna MT, Torres-Rodriguez JM, Cogliati M, Velegraiki A, Burggraaf A, Kamer-mans A, Sweere JM, Meis JF, Klaassen CH, Boekhout T. 2012. Autochthonous and dormant *Cryptococcus gattii* infections in Europe. *Emerg Infect Dis* 18:1618–1624. <http://dx.doi.org/10.3201/eid1810.120068>.
8. Ostrosky-Zeichner L, Marr KA, Rex JH, Cohen SH. 2003. Amphotericin B: time for a new “gold standard.” *Clin Infect Dis* 37:415–425. <http://dx.doi.org/10.1086/376634>.
9. Thompson GR, III, Wiederhold NP. 2010. Isavuconazole: a comprehensive review of spectrum of activity of a new triazole. *Mycopathologia* 170: 291–313. <http://dx.doi.org/10.1007/s11046-010-9324-3>.
10. Clinical and Laboratory Standards Institute. 2014. Minutes and agenda: annual meeting of the Subcommittee for Antifungal Susceptibility Testing. San Antonio, TX, 11 January 2014.
11. Espinel-Ingroff A, Pfaller MA, Bustamante B, Canton E, Fothergill A, Fuller J, Gonzalez GM, Lass-Flörl C, Lockhart SR, Martin-Mazuelos E, Meis JF, Melhem MSC, Ostrosky-Zeichner L, Pelaez T, Szesz MW, St-Germain G, Bonfietti LX, Guarro J, Turnidge J. 2014. Multilaboratory study of epidemiological cutoff values for detection of resistance in eight *Candida* species to fluconazole, posaconazole, and voriconazole. *Antimicrob Agents Chemother* 58:2006–2012. <http://dx.doi.org/10.1128/AAC.02615-13>.
12. Espinel-Ingroff A, Aller I, Canton E, Castanon-Olivares LR, Chowdhary A, Cordoba S, Cuenca-Estrella M, Fothergill A, Fuller J, Govender N, Hagen F, Illnait-Zaragoza MT, Johnson E, Kidd S, Lass-Flörl C, Lockhart SR, Martins MA, Meis JF, Melhem MSC, Ostrosky-Zeichner L, Pelaez T, Pfaller MA, Schell WA, St-Germain G, Trilles L, Turnidge J. 2012. *Cryptococcus neoformans*-*Cryptococcus gattii* species complex: an international study of wild-type susceptibility endpoint distributions and epidemiological cutoff values fluconazole, itraconazole, posaconazole, and voriconazole. *Antimicrob Agents Chemother* 56:5898–5906. <http://dx.doi.org/10.1128/AAC.01115-12>.
13. Turnidge J, Kahmeter G, Kronvall G. 2006. Statistical characterization of bacterial wild-type MIC value distributions and the determination of epidemiological cut-off values. *Clin Microbiol Infect* 12:418–425. <http://dx.doi.org/10.1111/j.1469-0691.2006.01377.x>.
14. Clinical and Laboratory Standards Institute. 2008. Reference method for broth dilution antifungal susceptibility testing of yeasts, 3rd ed. CLSI document M27-A3. Clinical and Laboratory Standards Institute, Wayne, PA.
15. Chowdhary A, Randhawa HS, Sundar G, Kathuria S, Prakash A, Khan Z, Sun S, Xu J. 2011. *In vitro* antifungal susceptibilities and genotypes of 308 clinical and environmental isolates of *Cryptococcus neoformans* var. *grubii* and *C. gattii* serotype B from North West India. *J Med Microbiol* 60:961–967. <http://dx.doi.org/10.1099/jmm.0.029025-0>.
16. Guinea J, Hagen F, Pelaez T, Boekhout T, Tahoune H, Torres-Narbona M, Bouza E. 2010. Antifungal susceptibility, serotyping, and genotyping of clinical *Cryptococcus neoformans* isolates collected during 18 years in a single institution in Madrid, Spain. *Med Mycol* 48:942–948. <http://dx.doi.org/10.3109/13693781003690067>.
17. Hagen F, Illnait-Zaragozi MT, Meis JF, Chew WHM, Curfs-Breuker I, Mouton JW, Hoepelman AI, Spanjaard L, Verweij PE, Kampinga GA, Kuijper EJ, Boekhout T, Klaassen CHW. 2012. Extensive genetic diversity within the Dutch clinical *Cryptococcus neoformans* population. *J Clin Microbiol* 50:1918–1926. <http://dx.doi.org/10.1128/JCM.06750-11>.
18. Hagen F, Illnait-Zaragozi MT, Bartlett KH, Swinne D, Geertsens E, Klaassen CH, Boekhout T, Meis JF. 2010. *In vitro* antifungal susceptibilities and amplified fragment length polymorphism genotyping of a worldwide collection of 350 clinical, veterinary, and environmental *Cryptococcus gattii* isolates. *Antimicrob Agents Chemother* 54:5139–5149. <http://dx.doi.org/10.1128/AAC.00746-10>.
19. Thompson GR, III, Wiederhold NP, Fothergill AW, Vallor AC, Wickes BL, Patterson TF. 2009. Antifungal susceptibilities among different serotypes of *Cryptococcus gattii* and *Cryptococcus neoformans*. *Antimicrob Agents Chemother* 53:309–311. <http://dx.doi.org/10.1128/AAC.01216-08>.
20. Turnidge J, Paterson DL. 2007. Setting and revising antibacterial susceptibility breakpoints. *Clin Microbiol Rev* 20:391–408. <http://dx.doi.org/10.1128/CMR.00047-06>.
21. Dalhoff A, Ambrose PG, Mouton JW. 2009. A long journey from minimum inhibitory concentration testing to clinically predictive breakpoints: deterministic and probabilistic approaches in deriving breakpoints. *Infection* 37: 296–305. <http://dx.doi.org/10.1007/s15010-009-7108-9>.
22. Pfaller MA, Messer SA, Rhomberg PR, Jones RN, Castanheira M. 2013. *In vitro* activities of isavuconazole and comparator antifungal agents tested against a global collection of opportunistic yeasts and molds. *J Clin Microbiol* 51:2608–2616. <http://dx.doi.org/10.1128/JCM.00863-13>.