

UC Irvine

UC Irvine Previously Published Works

Title

Gregarious settlement by the larvae of *Hydroides dianthus* (Polychaeta; Serpulidae)

Permalink

<https://escholarship.org/uc/item/7s459447>

Authors

Loudon, C
Scheltema, RS
Williams, IP
[et al.](#)

Publication Date

1981

Peer reviewed

Gregarious Settlement by the Larvae of *Hydroides dianthus* (Polychaeta: Serpulidae)*

R. S. Scheltema, I. P. Williams, M. A. Shaw and C. Loudon

Woods Hole Oceanographic Institution, Woods Hole, Massachusetts 02543, USA

ABSTRACT: Larval development of the serpulid polychaete worm, *Hydroides dianthus* Verrill 1893 parallels that of other closely related species. The larvae prefer to settle upon surfaces already inhabited by other members of their own species rather than to colonize surfaces without such conspecific residents. However, very wide variations in the intensity of settlement are observed. These differences in the amount of settlement cannot presently be attributed to any single variable. Gregarious settlement appears to be typical for sessile species which do not have the ability to reproduce asexually after settlement.

INTRODUCTION

Gregariousness may be defined as the interactions between planktonic larvae and attached resident members of their own species that result in settlement and metamorphosis of the larvae. Intra-specific settlement responses are now well known for a number of invertebrate taxa. For example, it has been shown that cyprids of the barnacles *Elminius modestus* and *Balanus balanoides* settle gregariously in response to members of their own species (Knight-Jones and Stephenson, 1950; Knight-Jones, 1953), and veligers of oysters belonging to *Ostrea edulis* and *Crassostrea virginica* also settle in response to attached members of their own kind (Knight-Jones, 1951; Crisp, 1967; Bayne, 1969; Hidu, 1969). At least one serpulid polychaete (*sensu stricto*), namely *Pomatoleios kraussi* from the Arabian Gulf, probably also settles gregariously (Crisp, 1977). Biological literature on gregarious settlement of other marine species has been summarized by Crisp (1974) and Scheltema (1974).

The present account describes a gregarious settlement response heretofore unknown in the sessile serpulid tube-worm *Hydroides dianthus*. This species secretes a calcareous tube upon submerged marine surfaces and is a key species in certain faunal assemblages (Haines and Maurer, 1980). The experiments

that show gregariousness in *Hydroides dianthus* were conducted by using larvae grown in the laboratory.

CULTURE METHODS

Sexually mature individuals attached to small cobbles were dredged during mid-June from a single population off Nobska Point, Vineyard Sound, Massachusetts. Eggs obtained from ripe females were artificially fertilized in finger bowls. The young trochophores were transferred 12-15 h after fertilization to larger 3-l jars of filtered sea water, gently aerated, and fed unicellular algae from cultures of *Dunaliella tertiolecta* and *Thalassiosira pseudonana* at a ratio of 1:1. The initial total concentration was 10^5 cells ml^{-1} . Inoculum for the unialgal cultures was kindly provided by Dr. R. R. Guillard. In some instances *Isochrysis galbana* was also used in a mixture of 2:1:1 *T. pseudonana*, *D. tertiolecta* and *I. galbana* respectively. Every second day the larvae were transferred to fresh filtered sea water by passing the entire culture through a fine-meshed 35- μm sieve constructed from Nytex cloth attached to one end of a 3-cm length of polyvinyl chloride pipe 7 cm in diameter. The surface of the screen was kept submerged during the transfer to prevent damage to the trochophore larvae. After the water was changed, sufficient new algal culture was added to bring the concentration to 10^5 cells ml^{-1} . The time required for completion develop-

* Contribution No. 4785, Woods Hole Oceanographic Institution



Fig. 2 *Hydroides dianthus*. Settled larva. Metamorphosis has begun and a rudimentary branchial crown has formed

EXPERIMENTAL METHODS

The experiments to demonstrate gregarious settlement were conducted in large clean finger bowls, 19-cm in diameter, containing approximately 0.8 l sea

water. The larvae were presented with a control surface – a 1.5×1.5 cm slate, left in a running seawater table for 2 days or more until it had acquired a microbial film – and an experimental surface – a slate of similar size to which were attached resident individuals of *Hydroides dianthus*. The slates were then placed on the bottom of the finger bowl, and larvae competent to settle were introduced. The concentration was such that there always remained an excess of larvae for the duration of the experiment. The finger bowls were placed in an evenly lighted culture cabinet, and pairs of control and experimental slates were placed in such a way that the results would not be affected by a bias related to their position in the bowls. No consistent differences were in fact observed between the pairs of slates from opposite sides of the bowls. No settlement occurred on the glass surfaces of the finger bowls during the course of the experiments because they were cleaned at the beginning of the experiment and had not acquired a suitable microbial film. In a few experiments slates of 3.0×1.5 cm or glass slides of either 3.8×2.5 cm or 2.0×2.5 cm were used. The results of the experiments were recorded by drawing the distributions both of settled larvae (Fig. 3) and of resident species on squared paper by means of a camera lucida on a binocular stereomicroscope (Wild M-5) at 25X. This method provided a permanent record for later analysis.




Fig. 3 *Hydroides dianthus*. Settlement of larvae on slates 3.0×1.5 cm; each dot indicates a newly settled individual. Larvae were given a choice of two surfaces. Right: Filmed surface without resident individuals of *H. dianthus*; settlement is sparse and occurred only after 12 h. Left: Surface with 72 resident individuals as shown by outlines of tubes; settlement is heavy, more than 20 times that on the right and most settlement occurred within the first 8 h. Filled circles: Larvae settled within the first 8 h; open circles: larvae settled between the 8th and 13th h of experiment; half filled circles: larvae settled between 13 and 18 h after beginning of experiment. Distribution was recorded on squared paper using a camera lucida on a stereo-microscope at 25X.

RESULTS

The outcome of the experiments were quickly apparent and in most instances quite striking; an example is shown in Figure 3. Here the right-hand rectangle represents a 3.0×1.5 cm control slate on which 73 *Hydroides dianthus* larvae had settled after exposure for 18 h. Each newly settled individual is represented by a small circle. On the left is an experimental slate of the same dimensions initially populated, as indicated by the tubes, by 72 residents of *H. dianthus*; here 2001 colonizing larvae settled during the same time interval. Clearly, when given a choice most colonizers selected the slate on the left. Of 2074 colonizers, 96.4 % preferred to settle near resident members of their own species; only 3.6 % settled on the control slate. This result is further supported by 49 additional experiments summarized in Figure 4. Here the horizontal axis shows intervals indicating the percentage of total settlement (for each experiment) that occurred on the slates having resident individuals; the height of




Fig. 4. *Hydroides dianthus*. Distribution of results from 49 experiments showing percentages of the total number in each experiment that settled on slates or glass slides having resident members of their own species. Included are 34 experiments using 1.5×1.5 cm slates, 6 experiments using 3.8×2.5 cm glass slides. Horizontal axis shows percent of total larvae settling on resident slide; vertical axis: number of experiments (see also Table 1)

the bars shows the number of experiments in which the indicated percentage of settlement occurred. In approximately $\frac{2}{3}$ of all experiments, more than 90 % of the colonizing larvae settled on slates or slides with resident individuals of their own species rather than on the controls, and in only two instances did less than half of the colonizers settle on slates with resident *H. dianthus*.

The non-parametric Mann-Whitney U test utilizing data from 34 experiments and values for the number (rather than percentage) of larvae that settled shows a highly significant difference ($P < 0.001$) between experimental and control slates. Only data from slates of similar size, namely 1.5×1.5 cm, were used and are tabulated (Table 1).

There are large differences – up to 40-fold – in the intensity of larval settlement between experiments concealed by treating the data as in Figure 4. These differences (indicated in Table 1) may be attributed to a number of possible variables which include: (1) the number of attached resident *Hydroides dianthus* on the experimental slates; (2) the number of *H. dianthus* larvae available for settlement; (3) the age of the resident *H. dianthus* on experimental slates; (4) differences in the state of development of *H. dianthus* larvae, or – more precisely – the time elapsed since larvae




Fig. 5. *Hydroides dianthus*. Numbers of larvae settling in relation to the number of residents on 1.5×1.5 cm slates. Data are from Table 1. A single point has been omitted – viz. settlement of 204 larvae on slate with 133 residents of *Hydroides dianthus*

Table 1. *Hydroides dianthus*. Numbers of larvae settling gregariously on 1.5 × 1.5 cm slates with and without resident members of their own species

A Slates with resident individuals	B Control slates	C Total (A + B)	D Percent of total on resident slates (A/C X100)	E No. of resident individuals
818	4	822	99.5	4
647	20	667	97.0	52
633	156	789	80.2	25
510	125	635	80.3	39
466	39	505	92.3	30
368	25	393	93.6	12
343	1	344	99.7	127
340	15	355	95.8	30
306	25	331	92.4	15
300	4	304	98.6	30
274	125	399	68.7	27
221	1	222	99.5	50
206	0	206	100.0	29
204	0	204	100.0	133
196	1	197	99.4	59
186	156	342	54.4	20
160	11	171	93.6	15
155	16	171	90.6	37
135	2	137	98.5	28
135	0	135	100.0	52
133	0	133	100.0	31
133	5	138	96.4	39
115	53	168	68.5	17
102	37	139	73.4	13
76	4	80	95.0	9
68	0	68	100.0	61
63	0	63	100.0	116
38	5	43	88.3	12
30	0	30	100.0	45
29	0	29	100.0	74
29	4	33	87.9	11
27	0	27	100.0	65
25	0	25	100.0	74
19	0	19	100.0	55

became physiologically competent to settle; and (5) qualitative differences in slime film on the panels.

Surprisingly, an examination of the data shows no striking relationship between the number of resident individuals and settlement of larvae (Fig. 5), at least not within the range of the resident population size used in our experiments. Inspection reveals no significant correlation between number of larvae on the vertical axis and residents in the horizontal axis ($r = -0.17$, not significant by analysis of variance). Because *Hydroides dianthus* larvae were always in excess in all experiments, it seems unlikely that their numbers affected the settlement intensity, although there is no direct evidence to support this contention. A test for an analysis of variance shows no relationship between settlement intensity and age of residents. Although there were differences in the mean settlement between experiments using larvae of 4, 5, 6, and 7 d (measured from time of fertilization) there

appeared to be no trend in the data, and older larvae did not consistently settle more readily than younger ones as might be expected. The variable most difficult to assess is qualitative differences in the microbial slime film affecting larval settlement. We know of no way to control readily the composition of such films.

DISCUSSION

Two adaptations allow sessile marine forms to colonize successfully and rapidly the surfaces of hard substrata. On the one hand there are those species which, after settlement, rapidly reproduce asexually to form colonies that can completely pre-empt all available surface; examples are ascidians such as *Botryllus schlosseri* and the cheilostomatous bryozoan *Schizoporella errata*. On the other hand there are those forms that do not reproduce asexually after attachment; instead such species settle together in very large

numbers, forming dense populations that arise from a gregarious response of the larvae (*vide* Jackson, 1977, p. 750; his Table 3). Known examples are barnacles such as *Balanus balanoides* and oysters such as *Crassostrea virginica*. Colonial species tend to have lecithotrophic larvae with a short non-feeding planktonic existence; if the species can readily overgrow other forms, only a few larvae need to settle successfully in order for such species to become established. Solitary species that have a gregarious settlement-response must produce large numbers of larvae in order to be numerous enough to settle together to form aggregations. Such species tend to have planktotrophic larvae that have a planktonic life 1 to 3 weeks; their establishment on a surface depends on massive colonization by large numbers of larvae. To this latter group of gregariously settling species may now be added the serpulid *Hydroides dianthus*.

Acknowledgements. This research was supported by the US Office of Naval Research, Contract No. N00014-79-C-0071 NR083-004 and by a grant from the US National Science Foundation EAR-7926381 to the first author. M. A. Shaw and C. Loudon wish to acknowledge support by Woods Hole Oceanographic Institution summer fellowships. We thank A. H. Scheltema for critically reading the manuscript.

LITERATURE CITED

- Bayne, B. L. (1969). The gregarious behaviour of the larvae of *Ostrea edulis* L. at settlement. *J. mar. biol. Ass. U. K.* 49: 327-356
- Crisp, D. J. (1961). Territorial behaviour in barnacle settlement. *J. exp. Biol.* 39: 429-446
- Crisp, D. J. (1967). Chemical factors inducing settlement in *Crassostrea virginica* (Gmelin). *J. animal Ecol.* 36: 329-336
- Crisp, D. J. (1974). Factors influencing the settlement of marine invertebrate larvae. In: Grant, P. T., Mackie, A. N. (eds) *Chemoreception in marine organisms*. Academic Press, New York, pp. 177-265
- Crisp, M. (1977). The development of the serpulid *Pomatoleios kraussii* (Annelida, Polychaeta). *J. Zool., Lond.* 183: 147-160
- Haines, J. L., Maurer, D. (1980) Quantitative faunal associates of the serpulid polychaetes *Hydroides dianthus*. *Mar. Biol.* 56: 43-47
- Hidu, H. (1969). Gregarious setting in the American oyster *Crassostrea virginica* Gmelin. *Chesapeake Sci.* 10: 85-92
- Jackson, J. B. C. (1977). Competition on marine hard substrata: The adaptive significances of solitary and colonial strategies. *Am. Nat.* 111: 743-767
- Knight-Jones, E. W., Stephenson, J. P. (1950). Gregariousness in the barnacle *Elminius modestus* Darwin. *J. mar. biol. Ass. U. K.* 29: 281-297
- Knight-Jones, E. W. (1951). Aspects of the setting behaviour of larvae of *Ostrea edulis* on the Essex oyster beds. *Rapp. P.-v. Réun. Cons. int. Explor. Mer* 128: 30-34
- Knight-Jones, E. W. (1953). Laboratory experiments on gregariousness during setting in *Balanus balanoides* and other barnacles. *J. exp. Biol.* 30: 584-598
- Scheltema, R. S. (1974). Biological interactions determining larval settlement of marine invertebrates. *Thalassia jugosl.* 10: 263-296
- Segrove, F. (1941). The development of the serpulid *Pomatoceros triqueter* L. *Q. Jl microsc. Sci.* 82: 468-540
- Wisely, B. (1958). The development and settling of a serpulid worm, *Hydroides norvegica* Gunnerus (Polychaeta). *Aust. J. mar. Freshw. Res.* 9: 351-361

This paper was presented by Dr. R. Scheltema; it was accepted for printing on January 20, 1981