

# Lawrence Berkeley National Laboratory

## Recent Work

### Title

POTENTIOMETRIC TITRATION OF PHOTOSYSTEM II FLUORESCENCE DECAY KINETICS IN SPINACH CHLOROPLASTS

### Permalink

<https://escholarship.org/uc/item/7s52h1bp>

### Authors

Karukstis, K.K.  
Sauer, K.

### Publication Date

1982-09-01



# Lawrence Berkeley Laboratory

UNIVERSITY OF CALIFORNIA

## CHEMICAL BIODYNAMICS DIVISION

RECEIVED  
LAWRENCE  
BERKELEY LABORATORY

FEB 18 1983

LIBRARY AND  
DOCUMENTS SECTION

Submitted to *Biochimica et Biophysica Acta*


POTENTIOMETRIC TITRATION OF PHOTOSYSTEM II  
FLUORESCENCE DECAY KINETICS IN SPINACH CHLOROPLASTS

Kerry K. Karukstis and Kenneth Sauer

September 1982

### TWO-WEEK LOAN COPY

*This is a Library Circulating Copy  
which may be borrowed for two weeks.  
For a personal retention copy, call  
Tech. Info. Division, Ext. 6782.*



LBL-15492  
c. 2

## DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

POTENTIOMETRIC TITRATION OF PHOTOSYSTEM II FLUORESCENCE DECAY  
KINETICS IN SPINACH CHLOROPLASTS

Kerry K. Karukstis and Kenneth Sauer

Department of Chemistry and Laboratory of Chemical Biodynamics  
Lawrence Berkeley Laboratory, University of California,  
Berkeley, California 94720

KEY WORDS: REDUCTION POTENTIAL; FLUORESCENCE INDUCTION;  
FLUORESCENCE LIFETIMES; PHOTOSYSTEM II; ELECTRON  
ACCEPTOR; (SPINACH CHLOROPLASTS)

ABBREVIATIONS: DCMU: 3-(3',4'-DICHLOROPHENYL)-1,1-DIMETHYL UREA;  
HEPES: N-2-HYDROXYETHYLPIPERAZINE-N'-2-ETHANESULFONIC  
ACID  
 $E_m$ : REDUCTION POTENTIAL MIDPOINT

This manuscript printed from originals provided by the authors.

## SUMMARY

The fluorescence yield of chloroplasts reflects the redox state of the electron acceptor of the Photosystem II reaction center, with increasing yield as the acceptor is reduced. Chemical reductive titrations of fluorescence yield in chloroplasts at room temperature indicate two distinct midpoint potentials, suggesting the possibility of Photosystem II electron acceptor heterogeneity. We have carried out a potentiometric titration of the fluorescence decay kinetics in spinach chloroplasts using a continuous mode-locked dye laser with low-intensity excitation pulses and a picosecond resolution single-photon timing system. At all potentials the fluorescence decay is best described by three exponential components. As the potential is lowered, the slow phase changes 30-fold in yield with two distinct midpoint potentials, accompanied by a modest (3-fold) increase in the lifetime. The titration curve for the slow component of the fluorescence decay of spinach chloroplasts is best characterized by two single-electron redox reactions with midpoint potentials at pH 8.0 of +119 mV and -350 mV, with corresponding relative contributions to the fluorescence yield of 49% and 51%, respectively. There is little change in the fast and middle components of the fluorescence decay. We found that the oxidized form of the redox mediator 2-hydroxy-1,4-naphthoquinone preferentially quenches the fluorescence, causing an anomalous decrease in the apparent midpoint of the high potential transition. This effect accounts for a significant difference between the midpoint potentials that we observe and some of those previously reported. The selective effect of reduction potentials on particular fluorescence decay components provides useful information about the organization and distribution of the Photosystem II electron acceptor.

## INTRODUCTION

The fluorescence yield of chloroplasts at room temperature reflects the redox state of the secondary electron acceptor, Q, of the Photosystem II reaction center, with increasing yield as the acceptor is reduced [1]. In general, chemical potentiometric titrations of fluorescence yield in chloroplasts at room temperature have indicated two distinct redox transitions, establishing the existence of two quenchers of Photosystem II fluorescence [2-9]. The source of the Photosystem II electron acceptor heterogeneity remains to be established. Various models of the Photosystem II photosynthetic unit suggest that the quenchers represent either alternative acceptors in the same Photosystem II reaction center [4,5,10] or identical acceptors of physically distinct Photosystem II reaction centers [4,10,11].

Conflicting measurements of the potentiometric titration curve of Photosystem II chlorophyll fluorescence render the interpretation of Q heterogeneity difficult. All reports agree in the biphasic character of the curve. However, there are considerable differences in the fluorescence parameters used to determine midpoint potentials and in the actual midpoint potentials measured. The titration curve obtained by Horton and Croze [4] for pea chloroplasts in the presence of redox mediators used the ratio of variable to maximum fluorescence,  $F_{var}/F_{max}$ , to characterize the titration of Q. The curve consisted of two transitions with midpoint potentials at pH 7.8,  $E_{m,7.8}$ , of -45 mV for a high-potential component accounting for 70% of the total variable fluorescence and -247 mV for a low-potential component attributed to the remaining 30% of variable fluorescence. A titration of the initial level of fluorescence upon illumination,  $F_i$ , of pea chloroplasts with comparable redox mediators was reported by Malkin and Barber [5] and provided evidence for two components contributing equally to the variable

fluorescence, with  $E_{m,7.6}$  values of +25 mV and -270 mV. Golbeck and Kok [6] also used  $F_i$  to monitor Q reduction in chloroplasts from *Scenedesmus* mutant No.8 in the presence of redox mediators. They observed a high-potential component with  $E_{m,7.2} = +68$  mV, accounting for 67% of Q, and a low-potential component with an  $E_{m,7.2}$  near -300 mV, accounting for the remaining 33% of Q. In a potentiometric titration of the ratio  $F_{max}/F_i$  in tobacco chloroplasts in the absence of redox mediators, Thielen and van Gorkom [9] assigned an  $E_{m,8.3}$  value of +115 mV to the high-potential transition and an  $E_{m,8.3}$  below -300 mV to the low-potential transition.

The discrepancies in these measurements of the redox titration of the Photosystem II electron acceptor Q are further confused by the heterogeneity observed in the kinetics of the fluorescence induction curve in chloroplasts inhibited with DCMU [1,12-18]. The shape of the induction curve is not characteristic of a first-order single electron transfer, but instead consists of two phases, a slow exponential phase and a fast sigmoidal phase. The slow exponential phase, attributed to Photosystem II reaction centers of low photochemical efficiency ( $\beta$ -centers) [15], titrates as a one-electron component with an  $E_{m,8.3}$  of approximately +115 mV [9] and an  $E_{m,7}$  of +120 mV [7]. One-electron reduction of the component responsible for the fast sigmoidal phase of fluorescence induction, a component ascribed to Photosystem II reaction centers of high photochemical efficiency ( $\alpha$ -centers) occurs at about -300 mV [9]. The functional heterogeneity of Q in  $\alpha$ - and  $\beta$ -centers has been ascribed [15-19] to organizational differences.  $\alpha$ -centers are characterized as Photosystem II reaction centers randomly embedded in an array of chlorophyll a/b light-harvesting antenna and thus capable of communication via excitation transfer, while the  $\beta$ -centers are isolated Photosystem II reaction centers each with its own chlorophyll a/b antenna

complex and thus constituting a separate unit in terms of excitation transfer.

A correlation between the heterogeneity observed in the fluorescence induction kinetics and that measured in potentiometric titrations of fluorescence seems a logical means of attempting to explain the heterogeneity of Q [4,7-9,19]. The results of Thielen and van Gorkom [9] are compatible with the assignment of  $Q_H$  and  $Q_L$  as the electron acceptors in  $\beta$ - and  $\alpha$ -centers, respectively; however, earlier investigations by others [7,8] were not.

In this paper we attempt to characterize more fully the nature of the Photosystem II electron acceptor heterogeneity by extending the analysis to a potentiometric titration of the Photosystem II fluorescence decay kinetics in spinach chloroplasts using low-intensity excitation pulses from a continuous mode-locked dye laser and a picosecond-resolution single-photon counting fluorescence lifetime system. The fluorescence decay kinetics of spinach chloroplasts at room temperature is best described by three exponential phases [20,21]. A fast phase of approximately 50-100 ps and a middle phase of approximately 400-800 ps arise from excitation lost prior to reaching the Photosystem II reaction center. The fast phase is attributed to excitation lost from the chlorophyll a antenna closely associated with Photosystem II reaction centers and the middle phase reflects excitation lost from the chlorophyll a/b light-harvesting antenna. There may also be a contribution from the antenna of Photosystem I. A slow phase of 1-2 ns is attributed to fluorescence from the radical pair recombination of the oxidized primary electron donor in Photosystem II,  $P680^+$ , and the reduced pheophytin primary electron acceptor, denoted  $I^-$ , a recombination which arises as a consequence of a reduced state of the quinone secondary electron acceptor, Q. The changes in the fluorescence yields and lifetimes of the decay components as a function of redox potential reveal valuable information about Q organization and



distribution. Furthermore, comparisons to steady-state potentiometric titrations of fluorescence yield will be made in an effort to explain the contradictions appearing in the literature.

#### MATERIALS AND METHODS

Broken chloroplasts were isolated from market spinach by grinding depetioled leaves in a blender for 10 s in a medium of 0.4 M sucrose, 50 mM HEPES-NaOH, pH 7.5, and 10 mM NaCl, followed by centrifugation at 6000 x g for 7 min. A wash with fresh grinding medium was followed by centrifugation under the same conditions. The pellet was resuspended in a medium of 0.1 M sucrose, 10 mM HEPES-NaOH, pH 7.5, and 10 mM NaCl, and then centrifuged at 6000 x g for 7 min. The isolated chloroplasts were resuspended in a medium of 0.1 M sucrose, 50 mM HEPES-NaOH, pH 7.5 or 8.0, 5 mM NaCl, and 5 mM MgCl<sub>2</sub> to give approximately 1 mg chlorophyll ml<sup>-1</sup>. For fluorescence measurements the chloroplast suspension was diluted with this final buffer, deaerated with argon gas, to a concentration of 17 µg chlorophyll ml<sup>-1</sup>.

For the potentiometric titrations the potential of the medium was measured by means of a platinum electrode with a Ag/AgCl electrode (saturated KCl solution) as reference, calibrated against a quinhydrone electrode. All reduction potentials were measured with a PAR model 173 potentiostat and are reported with respect to the standard hydrogen electrode (pH 0). The reduction potential of the suspension was adjusted under anaerobic conditions by additions of solid dithionite or small aliquots of 250 mM potassium ferricyanide (equilibration time - 15 min). Redox mediators, where present, included indophenol ( $E_{m,7} = +228$  mV); 1,4-naphthoquinone (+60 mV); duroquinone (0 mV); indigotetrasulfonate (-46 mV); 2,5-dihydroxybenzoquinone (-60 mV); indigotrisulfonate (-81 mV); indigodisulfonate (-125 mV); 2-hydroxy-1,4-naphthoquinone (-137 mV); anthraquinone-2,6-disulfonate (-184 mV); anthraquinone-2-sulfonate

(-225 mV); and methyl viologen (-430 mV). Titrations performed in the absence of redox mediators required longer equilibration times (e.g., 20-30 min) to insure adjustment of the reduction potential by  $\text{Na}_2\text{S}_2\text{O}_4$  and  $\text{K}_3\text{Fe}(\text{CN})_6$  alone.

A Spectra Physics synchronously-pumped mode-locked dye laser (SP 171 argon ion laser, SP 362 mode locker, and modified SP 375 dye laser) was used for the fluorescence decay kinetics measurements. Samples were excited with pulses at 620 nm and with a full width at half maximum of 12 ps. The intensity of excitation was kept low to monitor the initial level of fluorescence and prevent appreciable steady-state reduction of Q. Fluorescence was detected at right angles at 680 nm. The single-photon timing system and numerical analysis methods have been described previously [20,21]. All fluorescence decay data were resolved into a sum of exponential decays with a lifetime resolution limit of 25 ps.

Fluorescence induction curves were recorded with an instrument similar in design to that previously constructed by Shimony et al [22]. The chloroplast samples were excited in the blue region (Corning 3-67 and 4-96 filters in series) with low-intensity light ( $<0.05 \text{ J m}^{-2} \text{ s}^{-1}$ ), and fluorescence was detected at wavelengths longer than 620 nm (Corning 2-60 and 2-61 filters in series) using a Hamamatsu R928 photomultiplier tube. Induction curves were stored with a Nicolet Explorer IIIA digital oscilloscope with 500  $\mu\text{s}$ /address sweep timing. Chloroplasts were incubated in darkness at different redox potentials, and then 10  $\mu\text{M}$  DCMU was added 2 min prior to recording the induction curve. The initial level of fluorescence upon illumination is denoted  $F_i$ ; the minimum level of  $F_i$  is observed when all Photosystem II reaction centers are open (Q oxidized) and is denoted  $F_0$ . The maximum level of fluorescence upon continuous illumination is labelled  $F_{\text{max}}$ , and the quantity  $F_{\text{max}} - F_i$  is termed the variable fluorescence,  $F_{\text{var}}$ .

## RESULTS

The fluorescence induction curve in chloroplasts inhibited by DCMU was determined at different reduction potentials. Figure 1 shows the redox titration at pH 7.5 of the ratio  $F_{var}/F_{max}$ , in the absence of redox mediators, indicating the two distinct one-electron transitions previously attributed to the high- and low-potential components of  $Q$ ,  $Q_H$  and  $Q_L$ . The ratio  $F_{var}/F_{max}$  has a maximal value (about 0.6) when  $Q_H$  and  $Q_L$  are both completely oxidized and a minimum (0) when both acceptors are fully reduced. The  $E_{m,7.5}$  values are +98 mV and -381 mV for  $Q_H$  and  $Q_L$ , respectively, with  $Q_H$  accounting for 45% of the variable fluorescence and  $Q_L$  for 55%.

Analysis of the fluorescence lifetime decay kinetics of spinach chloroplasts as a function of reduction potential resulted in a characterization of the decay as a sum of three exponential phases. Figure 2 presents the potentiometric titration of the total fluorescence yield and that for each decay component at pH 8.0 in the presence of the specified mediators. Data were obtained for both oxidative and reductive titrations; the process appears to be completely reversible. The yields of the fast and middle lifetime components of the fluorescence decay are essentially insensitive to the state of the Photosystem II reaction center. The slow phase, however, significantly increases in yield as the potential is lowered, with two distinct redox transitions. Thus the two-step decrease in the total fluorescence yield reflects the two reductive titrations of the components responsible for the slow phase. The curve through the experimental total yield data represents a composite of two Nernst equations for one-electron redox reactions consistent with midpoint reduction potentials  $E_{m,8.0} = +116$  mV and  $-343$  mV and with relative contributions to the variable fluorescence yield of 50% each. A similar attempt to fit the experimental data for the slow phase to two

one-electron Nernst equations results in midpoint potentials  $E_{m,8.0} = +119$  mV and  $-350$  mV for  $Q_H$  and  $Q_L$ , respectively, with corresponding relative contributions to the fluorescence yield of 49% and 51%.

The lifetimes of each component of the fluorescence decay as a function of reduction potential are plotted in Fig. 3. The slow phase shows an increase in lifetime during both  $Q_H$  and  $Q_L$  reduction. A modest increase in the lifetime of the middle component occurs only during  $Q_H$  reduction, while the lifetime of the fast phase is apparently insensitive to the state of the reaction center.

Figure 4 presents an analogous potentiometric titration of the total fluorescence yield in the absence of mediators, but buffered at pH 7.5. The titration curve is well-characterized by two single-electron redox reactions making equal contributions to the total variable fluorescence, with midpoint potentials  $E_{m,7.5} = +106$  mV and  $-384$  mV. The small differences in the corresponding  $E_m$  values for the high- and low-potential transitions of the total fluorescence yield in Figs. 2 and 4 most likely result from pH differences, slight mediator artifacts, and the estimated error in the midpoint potential. The lifetime behavior of each component as a function of reduction potential is similar to that found in the presence of mediators (Fig. 3) and, therefore, is not shown.

The similarities of Figs. 2 and 4 suggest the absence of any significant fluorescence quenching by the specified redox mediators at the low concentration level of  $2.5 \mu\text{M}$ . However, sizable fluorescence quenching was detected in the presence of the oxidized form of 2-hydroxy-1,4-naphthoquinone, even at low concentrations. Figure 5 illustrates the effect of  $2.5 \mu\text{M}$  2-hydroxy-1,4-naphthoquinone in producing a significant lowering of the midpoint potential for  $Q_H$ , from  $+96$  mV to  $-68$  mV. This is attributed to strong quenching of the  $Q_H$  fluorescence by the oxidized form of the mediator. Malkin

and Barber [5] observed quenching of chlorophyll fluorescence by this mediator in similar steady-state redox titrations.

Figure 6 presents the potentiometric titration of the fluorescence decay components in the presence of DCMJ. The curve for the total fluorescence yield is described by two single-electron transitions with  $E_{m,7.5}$  values of +58 mV and -386 mV, contributing 31% and 69%, respectively, to the total variable fluorescence. Only the slow phase yield is influenced by reduction potential, exhibiting two transitions at  $E_{m,7.5} = +64$  mV and -392 mV with relative fluorescence yield contributions of 26% and 74%, respectively. A small effect of reduction potential on the slow phase lifetime is depicted in Fig. 7, while the lifetimes of the middle and fast phases are not altered by reduction potential.

Table 1 summarizes the results obtained for the potentiometric titrations of fluorescence induction and decay kinetics.

#### DISCUSSION

The major features of the potentiometric titration of the fluorescence decay kinetics in chloroplasts at room temperature in the absence of DCMJ are 1) the absence of any significant effect of reduction potential on the fluorescence yield of the fast and middle phases but the presence of two distinct redox transitions in the slow phase yield (and consequently in the total yield), with midpoint potentials  $E_{m,7.5} = +106$  mV and -384 mV; and 2) a moderate increase in the lifetime of the slow phase during both  $Q_H$  and  $Q_L$  reduction and in the lifetime of the middle component during  $Q_H$  reduction, but no effect of reduction potential on the lifetime of the fast phase.

The choice of redox mediators and their concentration is critical in order to accurately monitor changes in chlorophyll fluorescence induced by changes in the redox state of Q. The ability of a number of quinone

mediators to quench chlorophyll fluorescence directly when in their oxidized forms is well-documented [5,23-25]. Distortion of the titration curve is a possible consequence of such quenching. In this work the mediator 2-hydroxy-1,4-naphthoquinone was found to have a high quenching ability, significantly lowering the midpoint potential of the high-potential component in measurements of total fluorescence yield and the slow phase yield of fluorescence decay. Titrations in the absence of 2-hydroxy-1,4-naphthoquinone but either in the presence or absence of other redox mediators compare favorably in midpoint potentials and relative amplitudes of the two transitions.

Changes in the lifetime of the slow phase can be used to determine the extent of excitation energy transfer between reaction centers and thus discriminate between connected or isolated reaction centers. Energy transfer between photosynthetic units implies chlorophyll a/b light harvesting antennae serving more than one reaction center. The lifetime of an excited state in the antenna of a closed reaction center is a function of the ability of that reaction center to communicate with other reaction centers. Communication via the chlorophyll a/b light-harvesting antenna increases the lifetime of such an excited state; the extent of such an increase is dependent on the fraction of open reaction centers. Thus, we expect the lifetime of the slow phase of fluorescence decay of an  $\alpha$ -center, a Photosystem II reaction center capable of communication via excitation transfer, to be dependent on the state of the reaction center as monitored by the effect of reduction potential.  $\beta$ -centers, Photosystem II reaction centers that are isolated in terms of excitation transfer, should exhibit a slow phase lifetime that is independent of reduction potential. On the basis of the lifetime of the slow phase of fluorescence decay in Fig. 3, both  $Q_H$  and  $Q_L$  act as electron acceptors in  $\alpha$ -centers. This result

is in agreement with conclusions made by Horton [8] in fluorescence induction studies of pea chloroplasts. No distinction can be made here as to whether each  $\alpha$ -center contains both  $Q_H$  and  $Q_L$  or whether the  $\alpha$ -centers of a domain of communicating reaction centers contain either  $Q_H$  or  $Q_L$  as an electron acceptor. The electron acceptor of  $\beta$ -centers also remains undetermined.

Such an assignment of electron acceptors to  $\alpha$ - and  $\beta$ -centers has been attempted [7,8,27] by correlating the sigmoidicity of the fluorescence induction curve with the relative proportions of  $Q_H$  and  $Q_L$  observed in redox titrations of chloroplast membranes under a variety of conditions. The sigmoidicity of the fluorescence induction curve can be explained either by energy transfer between the antenna of a domain of reaction centers or by a requirement for two successive photoreactions to close the reaction center [4,7,8,19,27]. Recent experiments on Photosystem II particles from the blue-green alga Phormidium laminosum [28] favor the former explanation. Studies of the fluorescence decay kinetics of a variety of chloroplast membrane systems provide a means to distinguish between these two explanations for sigmoidal induction. Our ongoing work involves monitoring the fluorescence decay kinetics during potentiometric titrations of  $Q$  in chloroplast systems under conditions where the probability for energy transfer between reaction centers can be varied and in systems of variable relative proportions of  $Q_H$  and  $Q_L$ .

A number of features of this study are comparable to recent investigations in this laboratory on the state of the Photosystem II reaction center as determined by illumination conditions and the presence of various inhibitors [20,21]. That the yield of only the slow phase of the fluorescence decay is affected by reduction potential is consistent with the interpretation of the decay components [20,21]; the fast and middle components of the fluorescence decay arise from excitation lost prior to reaching the Photosystem II reaction

center and, therefore, should be largely independent of the redox state of the reaction center. The increase in the lifetime of the middle phase has previously been tentatively attributed to changes in energy distribution regulation mechanisms as a consequence of changes in the redox state of Q [20,21]. However, some differences are apparent in the effect of reduction potential and of illumination conditions on the state of the Photosystem II reaction center. For example, we find that chemical reduction of Q induces as much as an 8-fold increase in total fluorescence yield in chloroplasts at pH 8.0 (Fig. 2), whereas closing the Photosystem II reaction center by high-intensity excitation effected only a 4-fold increase in total fluorescence yield [20,21]. Furthermore, we measure the lifetime of the slow component of fluorescence decay to be significantly greater when the closure of the Photosystem II reaction centers is achieved by low reduction potential than by illumination [20,21]. Our present understanding of the Photosystem II photosynthetic unit and the associated fluorescence do not adequately account for these differences. Future experiments will focus attention on these details.

Despite the similarities in the results of the potentiometric titrations of fluorescence induction and of the fluorescence decay kinetics, a conclusive interpretation of the observed Photosystem II electron acceptor heterogeneity is not possible at this point. Potentiometric titrations using chloroplasts with compositional and structural differences, such as intermittent light-grown chloroplasts which do not contain the light-harvesting chlorophyll a/b protein, may provide insights into the question of Photosystem II electron acceptor heterogeneity.

#### CONCLUSION

Analysis of the potentiometric titration of the Photosystem II fluorescence decay kinetics in spinach chloroplasts at room temperature in the presence and



absence of DCMU indicates two distinct redox transitions in the total fluorescence yield. The yield of the slow phase of the fluorescence decay is also sensitive to the state of the Photosystem II reaction center, but the fast and middle lifetime components of the fluorescence decay are not appreciably affected by the reduction potential of Q. The midpoint reduction potentials for the two fluorescence quenchers,  $Q_H$  and  $Q_L$ , correlate well with those measured by Thielen and van Gorkom [9] for  $Q_B$  and  $Q_A$ , respectively; however, changes in the lifetime of the slow component of fluorescence decay suggest the possible assignment of  $Q_H$  and  $Q_L$  as the electron acceptors in  $\alpha$ -centers. A significant difference in the  $E_m$  value measured for  $Q_H$  from other previous reports is attributable to quenching of fluorescence in the presence of the oxidized form of the mediator 2-hydroxy-1,4-naphthoquinone.

#### ACKNOWLEDGEMENTS

This work was supported by the Director, Office of Energy Research, Office of Basic Energy Sciences, Division of Biological Energy Conversion and Conservation, of the Department of Energy under Contract No. DE-AC03-76SF00098, by National Science Foundation Grant No. PCM 79-11251, and by National Institutes of Health National Research Service Award No. 1 F32 GM08617-01.

TABLE 1  
 CHARACTERISTICS OF THE POTENTIOMETRIC TITRATION  
 OF THE ELECTRON ACCEPTOR Q IN PHOTOSYSTEM II

Chloroplasts, isolated as described in Materials and Methods, were suspended in a medium of 0.1 M sucrose, 50 mM HEPES-NaOH (buffered at the stated pH), 5 mM NaCl, and 5 mM MgCl<sub>2</sub> to give a chlorophyll concentration of 17 µg ml<sup>-1</sup>. Fluorescence samples were dark-adapted under anaerobic conditions for 15 min at the appropriate reduction potential before addition of DCMU or illumination.

Fluorescence Induction Experiments,  $F_{var}/F_{max}$

pH	[DCMU]	$E_m^a$ (mV)	Contribution to $F_{var}$ (%)
7.5	10 µM	+98 <sup>b</sup>	45
		-381	55

Fluorescence Decay Kinetics,  $F_i$  level

pH	[DCMU]	$E_m^a$ (mV)	Contribution to $F_{var}$ (%)
8.0	0	+116 <sup>c</sup>	50
		-343	50
7.5	0	+106 <sup>b</sup>	50
		-384	50
7.5	0	+96 <sup>c,e</sup>	-
7.5	0	-68 <sup>d,e</sup>	-
7.5	10 µM	+58 <sup>b</sup>	31
		-386	69

<sup>a</sup>  $E_m$  values relative to standard hydrogen electrode.

<sup>b</sup> No redox mediators present.

<sup>c</sup> Redox mediators present (2.5 µM); 2-OH-1,4-naphthoquinone not included as a redox mediator.

<sup>d</sup> Redox mediators present (2.5 µM); 2-OH-1,4-naphthoquinone included as a redox mediator.

<sup>e</sup>  $E_m$  value determined for Q<sub>H</sub> only.

## FIGURE CAPTIONS

- Fig. 1 Redox titration of the ratio of variable to maximum fluorescence,  $F_{var}/F_{max}$ , of chloroplasts at pH 7.5 in the presence of 10  $\mu$ M DCMU and in the absence of redox mediators.
- Fig. 2 Potentiometric titration of the total fluorescence yield and the yield of the components of the fluorescence decay in spinach chloroplasts at pH 8.0 in the absence of DCMU. Redox mediators present at 2.5  $\mu$ M included indophenol; 1,4-naphthoquinone; duroquinone; indigotetrasulfonate; 2,5-dihydroxybenzoquinone; indigotrisulfonate; indigodisulfonate; anthraquinone-2,6-disulfonate; and methyl viologen. The symbols are defined as follows: (X) total fluorescence yield; ( $\square$ ) yield of the slow phase; ( $\diamond$ ) yield of the middle phase; and (o) yield of the fast phase.
- Fig. 3 Lifetimes of the components of the fluorescence decay in spinach chloroplasts at pH 8.0 in the absence of DCMU as a function of the reduction potential of the medium. Redox mediators as in Fig. 2. The symbols are defined as follows: (X) lifetime of the slow phase; ( $\square$ ) lifetime of the middle phase; and ( $\diamond$ ) lifetime of the fast phase.
- Fig. 4 Potentiometric titration of the total fluorescence yield in spinach chloroplasts at pH 7.5 in the absence of redox mediators and DCMU.

Fig. 5 Potentiometric titration of the total fluorescence yield in spinach chloroplasts at pH 7.5 in the presence ( $\square$ ) and absence (X) of the redox mediator 2-hydroxy-1,4-naphthoquinone.


Fig. 6 Potentiometric titration of the total fluorescence yield and the yield of the components of the fluorescence decay in spinach chloroplasts at pH 7.5 in the absence of redox mediators and in the presence of 10  $\mu$ M DCMU. The symbols are defined as follows: (X) total fluorescence yield; ( $\square$ ) yield of the slow phase; ( $\diamond$ ) yield of the middle phase; and (o) yield of the fast phase.

Fig. 7 Lifetimes of the components of the fluorescence decay in spinach chloroplasts at pH 7.5 in the presence of 10  $\mu$ M DCMU as a function of reduction potential of the medium. The symbols are defined as follows: (X) lifetime of the slow phase; ( $\square$ ) lifetime of the middle phase; and ( $\diamond$ ) lifetime of the fast phase.


## REFERENCES

1. Dvysens, L. N. M., and Sweers, H. E. (1963) in Studies on Microalgae and Photosynthetic Bacteria (Japanese Society of Plant Physiologists, ed.), pp. 353-372, University of Tokyo Press, Tokyo.
2. Cramer, W. A. and Butler, W. L. (1969) *Biochim. Biophys. Acta* 172, 503-510.
3. Ke, B., Hawkrigde, F. M., and Sahu, S. (1976) *Proc. Natl. Acad. Sci. USA* 73, 2211-2215.
4. Horton, P. and Croze, E. (1979) *Biochim. Biophys. Acta* 545, 188-201.
5. Malkin, R. and Barber, J. (1979) *Arch. Biochem. Biophys.* 193, 169-178.
6. Golbeck, J. H., and Kok, B. (1979) *Biochim. Biophys. Acta* 547, 347-360.
7. Horton, P. (1981) *Biochim. Biophys. Acta* 635, 105-110.
8. Horton, P. (1981) *Biochim. Biophys. Acta* 637, 152-158.
9. Thielen, A. P. G. M. and van Gorkom, H. J. (1981) *FEBS Letters* 129, 205-209.
10. Joliot, P. and Joliot, A. (1979) *Biochim. Biophys. Acta* 546, 93-105.
11. Diner, B. A. (1978) in Proceedings of the 4th International Congress on Photosynthesis (Hall, D. O., Coombs, J., and Goodwin, T. W., eds), pp. 359-372, Biochemical Society, London.
12. Morin, P. (1964) *J. Chim. Phys.* 61, 674-680.
13. Joliot, A. and Joliot, P. (1964) *C. R. Acad. Sci. Paris, Ser. D* 258, 4622-4625.
14. Döschek, W. W. and Kok, B. (1972) *Biophys. J.* 12, 832-838.
15. Melis, A. and Homann, P. H. (1975) *Photochem. Photobiol.* 21, 431-437.
16. Melis, A. and Homann, P. H. (1978) *Arch. Biochem. Biophys.* 190, 523-530.
17. Melis, A. and Dvysens, L. N. M. (1979) *Photochem. Photobiol.* 29, 373-382.

18. Melis, A. and Schreiber, U. (1979) *Biochim. Biophys. Acta* 547, 47-57.
19. Melis, A. (1978) *FEBS Letters* 95, 202-206.
20. Haehnel, W., Nairn, J. A., Reisberg, P., and Sauer, K. (1982) *Biochim. Biophys. Acta* 680, 161-173.
21. Nairn, J. A., Haehnel, W., Reisberg, P., and Sauer, K. (1982) *Biochim. Biophys. Acta* (in press).
22. Shimony, C., Spencer, J., and Govindjee (1967) *Photosynthetica* 1, 113-125.
23. Thomas, J. B., Voskuil, W., Olsman, H., and De Boois, H. M. (1962) *Biochim. Biophys. Acta* 59, 224-226.
24. Arnon, D. I., Tsujimoto, H. Y., and McSwain, B. D. (1965) *Proc. Natl. Acad. Sci. U.S.* 54, 927-934.
25. Amesz, J. and Fork, D. C. (1967) *Biochim. Biophys. Acta* 143, 97-107.
26. Vernotte, C., Etienne, A.-L. and Briantais, J.-M. (1979) *Biochim. Biophys. Acta* 545, 519-527.
27. Horton, P. and Naylor, B. (1979) *Photochem. Photobiophys.* 1, 17-83.
28. Bowes, J.M. and Horton, P. (1982) *Biochim. Biophys. Acta* 680, 127-133.




XBL 828-11049




XBL 828-11050







XBL 828-11051




XBL 828-11052



XBL 828-11053



XBL 828-11054



XBL 828-11055

This report was done with support from the Department of Energy. Any conclusions or opinions expressed in this report represent solely those of the author(s) and not necessarily those of The Regents of the University of California, the Lawrence Berkeley Laboratory or the Department of Energy.

Reference to a company or product name does not imply approval or recommendation of the product by the University of California or the U.S. Department of Energy to the exclusion of others that may be suitable.

TECHNICAL INFORMATION DEPARTMENT  
LAWRENCE BERKELEY LABORATORY  
UNIVERSITY OF CALIFORNIA  
BERKELEY, CALIFORNIA 94720