

UC Davis

UC Davis Previously Published Works

Title

MnSOD in Oxidative Stress Response-Potential Regulation via Mitochondrial Protein Influx

Permalink

<https://escholarship.org/uc/item/7sb78376>

Journal

Antioxidants & Redox Signaling, 20(10)

ISSN

1523-0864

Authors

Candas, Demet

Li, Jian Jian

Publication Date

2014-04-01

DOI

10.1089/ars.2013.5305

Peer reviewed



FORUM REVIEW ARTICLE

MnSOD in Oxidative Stress Response-Potential Regulation *via* Mitochondrial Protein Influx

Demet Candas¹ and Jian Jian Li^{1,2}

Abstract

Significance: The mitochondrial antioxidant manganese superoxide dismutase (MnSOD) is encoded by genomic DNA and its dismutase function is fully activated in the mitochondria to detoxify free radical $O_2^{\bullet-}$ generated by mitochondrial respiration. Accumulating evidence shows an extensive communication between the mitochondria and cytoplasm under oxidative stress. Not only is the MnSOD gene upregulated by oxidative stress, but MnSOD activity can be enhanced *via* the mitochondrial protein influx (MPI). **Recent Advances:** A cluster of MPI containing cytoplasmic/nuclear proteins, such as cyclins, cyclin-dependent kinases, and p53 interact with and alter MnSOD activity. These proteins modulate MnSOD superoxide scavenging activity *via* post-translational modifications in the mitochondria. In addition to well-established pathways in gene expression, recent findings suggest that MnSOD enzymatic activity can also be enhanced by phosphorylation of specific motifs in mitochondria. This review attempts to discuss the pre- and post-translational regulation of MnSOD, and how these modifications alter MnSOD activity, which induces a cell adaptive response to oxidative stress. **Critical Issues:** MnSOD is biologically significant to aerobic cells. Its role in protecting the cells against the deleterious effects of reactive oxygen species is evident. However, the exact network of MnSOD-associated cellular adaptive reaction to oxidative stress and its post-translational modifications, especially its enzymatic enhancement *via* phosphorylation, is not yet fully understood. **Future Directions:** The broad discussion of the multiple aspects of MnSOD regulation, including gene expression, protein modifications, and enzymatic activity, will shed light onto the unknown mechanisms that govern the prosurvival networks involved in cellular and mitochondrial adaptive response to genotoxic environment. *Antioxid. Redox Signal.* 20, 1599–1617.

Introduction

IN MAMMALS, AN OXIDATIVE STRESS condition is created in cells as a result of an imbalance between the generation and the detoxification of reactive oxygen species (ROS). Superoxide dismutase (SOD) scavenger enzymes are the major ROS detoxifying enzymes protecting the cells from potential damage caused by excessive amounts of ROS (56). In mammalian cells, manganese superoxide dismutase (MnSOD) is the essential mitochondrial antioxidant enzyme that detoxifies the free radical superoxide, the major by-product of mitochondrial respiration. Studies from the early 1970s using *Escherichia coli* and yeast as model organisms provide the evidence indicating the importance of MnSOD for the survival in

aerobic environments (61). Studies using mice (182) and fruit flies (40) demonstrated that MnSOD gene knockout is lethal; while MnSOD heterozygous mice with haplo-insufficiency can survive but are more susceptible to oxidative injury (169, 171). The crucial role of MnSOD in protecting cells against oxidative stress has been extensively studied and thoroughly reviewed (75, 89, 123, 130, 150, 161), pointing to the critical roles of MnSOD in maintaining cellular physiology in response to genotoxic conditions, such as oxidative stress. In addition to MnSOD gene transcription, accumulating evidence indicates that MnSOD enzymatic activity is enhanced *via* post-translational modifications. This review will focus on the unique pathways that MnSOD activity is regulated by the mitochondrial protein influx (MPI), which, together with

¹Department of Radiation Oncology and ²NCI-Designated Comprehensive Cancer Center, University of California Davis, Sacramento, California.

transcriptional/post-transcriptional regulations of MnSOD, may present a MnSOD-centered signaling network to protect cells from the injury of oxidative environments.

Oxidative Stress and MPI

An imbalance between the generation of ROS and the detoxification of the reactive intermediates is defined as oxidative stress. Under aerobic conditions, ROS are generated as by-products of the oxygen metabolism during oxidative phosphorylation in the mitochondria (55, 74, 89). Apart from their deleterious effects, ROS functions as signaling regulators. ROS can modulate mitochondrial functions *via* regulating electron transfer chain enzymes and mitochondrial membrane potential (121, 196). ROS are crucial for various cellular processes, including cell growth (18), apoptosis (2, 82, 184), cell adhesion and immune responses (38). Some ROS also act as second messengers in redox signaling (52). When the normal redox state of a cell is impaired, highly reactive peroxides (hydrogen peroxide [H_2O_2]), and free radicals ($\text{O}_2^{\bullet-}$) with the ability to damage any component of the cell, including proteins, lipids, and DNA, accumulates (68). Disruption of the ROS production/detoxification cycle, contributes to the development of human pathologies, including age-related diseases, such as cancer (58, 68, 80, 178), cardiovascular (38, 139), neurodegenerative (178), and genetic disorders (33, 88). ROS can also be generated by exogenous sources like ionizing radiation (IR) (75, 110), through direct interactions with either cellular targets or with water, resulting in DNA damage and gene mutations (45, 69) or cell death (110). However, under oxidative stress, ROS induces the intrinsic antioxidant enzymes to protect cells from ROS-induced toxicity (64).

Adaptive response to oxidative stress

Adaptive response has been observed in both malignant (116) and nonmalignant (117, 118) cells that were exposed to thiol-containing drugs, oxidative stress, such as hyperoxia (80%–95% O_2) or H_2O_2 (160) as well as reactive nitrogen species (RNS or RNOS), namely, nitric oxide (NO) (143). However, this review will focus on the adaptive response to IR; radioadaptive response. Chronic exposure of cells to IR induces an adaptive response that results in enhanced tolerance to the subsequent IR cytotoxicity (2, 47, 64). Exposure to low dose IR (LDIR) generates beneficial effects for mammalian cells with respect to the maintenance of genomic integrity and the ability to repair damaged DNA (49). Radioadaptive resistance is a feature of many other species, including *E. coli*, algae, higher plant cells, and insect cells (94). In mice, whole-body preirradiation with LDIR reduced the incidence of thymic lymphoma from 46% to 16% with a challenging high dose of radiation, indicating the radioadaptive response (13). Similarly, in rabbits, Liu *et al.* reported LDIR-induced radio-protection (98). Human lymphocytes exposed to LDIR showed fewer chromatid breaks when later exposed to larger doses (156). Pretreatment of normal (3, 47) or cancer cells (64) with LDIR (1–10cGy) induced the radioadaptive response by elevating the expression and activity of antioxidant enzymes, primarily MnSOD (116). The above data and many other reports suggest that radioadaptive response is conserved in all species (2); and is also a feature of tumor cells, in which therapeutic radiation induces adaptive response (64, 75, 79),

contributing to the overall ineffectiveness of cancer radiotherapy. However, the exact mechanism of MnSOD induction governing the radioadaptive response of tumor cells is still largely unknown.

Mitochondria

Mitochondria generate cellular fuel for aerobic cells to maintain normal functions (150, 153). The mitochondrial bioenergetics and the number of mitochondria per cell are related to the overall energy demands of the cells. In mammalian cells, about 90% of the ATP is generated by the mitochondria *via* oxidative phosphorylation. As a result of their role in metabolism, mitochondria are the primary source of ROS in aerobic cells. Apart from this critical function in cell respiration, mitochondria are also involved in other fundamental cellular functions, such as apoptotic response (2, 58, 80), cardioprotection (178), and neurodegenerative disorders (38, 178). To perform and regulate such a wide range of cellular roles, mitochondria are engaged in an extensive communication with their cytoplasmic environment *via* a constant flux of inorganic ions, metabolites, and proteins that traffic between the cytosol and mitochondria.

Mitochondria contain their own genome, a single circular DNA, that encodes 2 ribosomal RNAs, 22 tRNAs and 13 polypeptides, which all integrate into the mitochondrial respiratory chain complexes (153). However, the majority of mitochondrial proteins are encoded by nuclear DNA, synthesized as precursors on cytosolic ribosomes, and then transported into the mitochondria. Such peptides are imported into the mitochondria by a complex multistep mechanism involving the recognition, translocation and membrane insertion of these precursors, which then become predominant regulators of mitochondrial functions (152). The number of polypeptides in a single mitochondrion had been estimated to be over 1000 (151) until recent proteomic analyses identified 500–800 proteins (153). Among these are the proteins involved in the biogenesis of mitochondria and in various other biochemical activities of mitochondria. It is important to note that the composition of the mitochondria and relative abundance of proteins in them is highly cell type/tissue/organism specific. The exact functions of many of these mitochondrial proteins are unknown.

Mitochondrial Protein Import/Influx

Mitochondria are the site of oxidative phosphorylation and carry the electron transport chain complexes in their inner membrane, but they also accommodate the Krebs cycle in the matrix, and are involved in fatty-acid metabolism, urea cycle, and programmed cell death. These diverse functions require the transport of metabolites and proteins to and from the cytosol. Mitochondrial influx of proteins is implicated in a wide spectrum of mitochondrial functions, including apoptosis and ATP production (78, 131); and the influxed proteins are targeted to all mitochondrial compartments (9, 72), including the cytoplasm-facing outer surface of the outer membrane (66). This review will focus on a specific cluster of MPI that translocate into the mitochondrial matrix under oxidative stress conditions to regulate MnSOD function.

Mitochondria are membrane-enclosed organelles with two membranes: an outer and an inner membrane. Other compartments include the intermembrane space, cristae space

formed by the convolution and folding of the inner membrane, and the matrix (Fig. 1 insert). Proteins destined to be imported into mitochondria face the challenge of being routed to their correct sub-mitochondrial compartments. Moreover, the proteins targeted to the mitochondrial matrix, the innermost compartment, must be transported across two membranes. These challenges are overcome by two specific machineries on the outer and inner mitochondrial membranes. The transfer of mitochondrial proteins from their site of synthesis into the mitochondria occurs post-translationally through interactions with the mitochondrial import machinery located at the outer membrane (Fig. 1) (135). However, some precursor proteins enter the import channels co-translationally (57). Mitochondria-targeted proteins can be divided into two main classes; proteins with presequences and proteins with internal targeting sequences (180). Presequences are N-terminal cleavable sequences, which form positively charged extensions that interact with the mitochondrial import receptors; whereas internal sequences are not cleavable, do not necessarily contain charged amino-acid residues, and are incorporated into the mature protein. Apart from positively charged amino-acids like arginine and lysine, presequences also contain hydrophobic residues and hydroxylated amino-acids, such as serine and threonine (146). The presequences are usually devoid of acidic residues (146); and they show a high tendency to form an amphipathic α -helix that presents one positively charged surface and one

hydrophobic surface (145), both of which are recognized by distinct receptor subunits of the translocase of the outer membrane (TOM) complex. On the other hand, the internal sequences usually exist in clusters and can be charged or uncharged. The individual signals can function independently but the import is more efficient when they cooperate (135). Mitochondrial localization of proteins can be predicted based on the presence of N-terminal targeting sequences using bioinformatics approaches; however, the consensus sequence for N-terminal presequences is not completely satisfying and the internal sequences are yet to be characterized.

Nuclear-encoded proteins cross the outer membrane of the mitochondria at the TOM complex, from where they are directed to one of two functionally different translocase of the inner membrane (TIM) complexes to transport across the inner membrane. TIM 23 guides the import of proteins with N-terminal presequences and TIM 22 is involved in the import of proteins with internal targeting sequences (34). The TOM complex consists of seven subunits; three receptors Tom20, Tom22, Tom70; the channel forming protein Tom40; and three small proteins Tom5, Tom6, Tom7 (180). Proteins with presequences are recognized by the receptor proteins within the TOM complex, whereas the proteins with internal targeting sequences usually interact with cytosolic chaperons, Hsp70 and Hsp90, which dock onto the receptors of TOM complex (193). Three other cytosolic factors that act on




FIG. 1. Mitochondrial import machinery. Many nuclear-encoded proteins are imported into the mitochondria through TOM and TIM complexes. Proteins with N-terminal presequences are directly recognized by the receptors at the TOM complex and upon entry into the intermembrane space, these proteins are sorted to their final locations by TIM23 complex. Proteins with internal targeting sequences, on the other hand, interact with cytoplasmic chaperons, such as Hsp 70, Hsp 90, and 14-3-3, which then dock on the receptors of TOM complex (R) to initiate the import of these proteins. They are then directed to the TIM 22 complex and sent to the matrix, IMS or IM as their final destinations. TOM, translocase of the outer membrane; TIM, translocase of the inner membrane; IMS, intermembrane space; IM, inner membrane; OM, outer membrane; R, receptor.

mitochondrial preproteins have been described: mitochondrial fusion protein targeting factor from yeast (24), presequence binding factor from rabbit (114) and mitochondrial import stimulating factor from rat (67). Among these, mitochondrial import stimulating factor is a member of the 14-3-3 protein family, a ubiquitous eukaryotic protein family with a wide range of chaperone properties (5). 14-3-3 proteins may form complexes with Hsp70 to facilitate the mitochondrial localization of cytosolic proteins (109). The transfer of proteins across the inner membrane also requires the mitochondrial membrane potential ($\Delta\psi$) (106). The mitochondrial processing peptidase (MPP) cleaves off the presequences from the proteins in the matrix before the protein folds into its active form with the help of chaperons (71). Once the proteins pass through TOM and TIM complexes and the presequences are cleaved off, they are sorted according to the proteins' sorting signals into inner membrane, intermembrane space, outer membrane, or the matrix.

Although the mechanism and the machinery responsible for mitochondrial protein import have been extensively studied, little is known about the factors that affect the influx of proteins under physiological or pathological conditions. It has long been believed that MPI is an unregulated event, until recent evidence from plant studies showed that mitochondrial internalization of proteins is influenced by light-dark cycle and developmental cues (39). The next section will address the regulation of MPI in oxidative stress conditions.

Regulation of MPI under oxidative stress

Studies in mammalian systems have shown that oxidizing conditions decrease the mitochondrial protein import and cause the accumulation of precursor proteins outside mitochondria. The precursor proteins that have failed to be imported end up being degraded by proteasomes (187). Interestingly, Tom 20, a core component of the mitochondrial outer membrane protein import receptor, was also among the proteins that are degraded by proteasomes under oxidative stress (187). These results suggested that oxidizing conditions lead to a mitochondrial influx lesion resulting in the decreased number of mitochondrial proteins. This decrease may represent the mitochondrial response to the oxidative stress. A relatively recent study showed that IR affects the mitochondrial protein import based on the radiation dose. At lower doses (10cGy), IR enhanced the mitochondrial import; while at higher dose (4 Gy), mitochondrial import was decreased (134). These results suggested that altered mitochondrial protein import by LDIR may be linked to the adaptive response of the cells, where increased protein import is aimed to protect the cells. Translocation of proteins across the inner membrane requires potential difference ($\Delta\psi$) across the membrane. IR may disrupt the membrane potential and affect the MPI. Though, changes in membrane potential alone were not enough to explain the mitochondrial import deficiencies observed in this recent study (134). Further studies showed that some components of the mitochondrial import machinery may be damaged by IR, resulting in deficient protein import (10). Defective protein import, in turn, is shown to amplify the effects of oxidative stress and cause several health defects, including degenerative diseases and metabolic disorders.

Superoxide Dismutases

Under normal circumstances, the deleterious effects of ROS are avoided by several antioxidant enzyme systems that detoxify ROS produced in the cells (92). Among the antioxidants, the SOD scavenger enzymes convert superoxide radicals into H_2O_2 and molecular oxygen ($2\text{O}_2^{\bullet-} + 2\text{H}^+ \rightarrow \text{H}_2\text{O}_2 + \text{O}_2$) (56). The resultant H_2O_2 is largely removed by peroxiredoxins (Prx) (125), glutathione peroxidases (104) and catalase (56, 69). There are three SOD enzymes expressed in mammalian cells. Copper/zinc-containing superoxide dismutase (Cu/ZnSOD, SOD1) is primarily localized to cytosol (159), although it has been found in the intermembrane space of mitochondria (128, 179). MnSOD (SOD2) is the mitochondrial antioxidant that exists in homotetramers (19, 140, 183) and localizes in the mitochondrial matrix (128, 179). The third SOD is the extracellular superoxide dismutase (ECsOD, SOD3) is localized to the extracellular space (51). The presence of SOD within the mitochondria suggests that superoxides generated in the mitochondria do not readily cross membranes; and due to their deleterious effects on biomolecules, they must be dismutated immediately by a SOD, preferably at the site of their production.

Manganese Superoxide Dismutase

Human MnSOD is a tetrameric enzyme with four identical subunits each harboring a Mn^{+3} atom (19, 183). Human MnSOD gene is located on the 6th chromosome, 6q25.3 region (174) encoding a ~223 amino acid/26 kD precursor monomer containing a mitochondria targeting sequence of 26 amino acids (183) that is required for mitochondrial localization. Human MnSOD resides in the mitochondrial matrix (19, 183), where the mitochondrial oxidative phosphorylation complexes that generate superoxide are accessible. MnSOD is a highly conserved protein with over 40% sequence homology among human, yeast and *E. coli* (11). Each MnSOD monomer has two distinct domains: an N-terminal helical hairpin domain and a C-terminal α/β domain, containing five α helices and three-stranded antiparallel β sheets (19). As MnSOD is taken up by the mitochondria, the mitochondria targeting sequence is clipped off, leaving a 22 kD monomer, which later incorporates an Mn^{+3} ion and assembles into an 88 kD homotetramer in the mitochondria (183). Residues D159, H163, H26, H74, and a water molecule from each subunit contribute to the manganese metal-binding site, namely, the active site (19).

Mitochondrial import/processing of MnSOD

MnSOD is encoded by nuclear genome, synthesized in the cytosol, imported into the mitochondrial matrix post-translationally *via* TOM and TIM23 complexes and assembled into active enzyme with the incorporation of a manganese ion in the mitochondrial matrix (183). Human MnSOD is translated as a precursor peptide with a leader sequence that is proteolytically removed by MPP in the mitochondrial matrix during the processing of the mature enzyme. Similar to other mitochondrial leading sequences, the presequence of MnSOD has been shown to form amphipathic α -helices with positive charged and hydrophobic residues on opposite faces of the helix (146, 183). This amphipathic helix mediates its interaction with the mitochondrial import machinery and facilitates

the import process. Furthermore, the mitochondrial transport of MnSOD precursor has been long known to be energy dependent (183). Energy derived from ATP is required for uptake and processing of not only MnSOD, but also most precursor mitochondrial proteins (146). It is believed that mitochondrial proteins integrate cofactors as a terminal step of their assembly because cofactors recognize the tertiary structure of the proteins (144). MnSOD is no exception to that; it is imported into the mitochondria as an inactive apoprotein, which later incorporates its cofactor Mn^{2+} ion in the matrix.

A human genetic polymorphism encodes for either alanine (Ala) or valine (Val) in the mitochondrial targeting sequence of MnSOD protein and is shown to modulate MnSOD activity and its mitochondrial influx (165, 166). This polymorphism is at codon 16 of the human precursor protein, which is also located at position -9 of the mature protein; therefore, it is often referred to as either the Ala9Val or Ala16Val MnSOD dimorphism (166). Initial studies on the effect of this dimorphism in MnSOD mitochondrial influx were based on the computer models that can predict the configuration of polypeptides from their amino acid sequences. A partial α -helix configuration was predicted for Ala-MnSOD, while a single β -sheet structure was predicted for the Val-MnSOD. The expanded β -sheet structure was estimated to physically slow down the import of MnSOD through the TIM23 channel and cause partial stalling of the Val-MnSOD in the inner membrane (158). Further studies showed that mitochondrial import and activity of MnSOD is affected by the Ala9Val dimorphism in its MTS (107, 165, 166). The Ala-MnSOD variant with an α -helical MTS is easily imported into the matrix and has high mitochondrial activity, whereas the Val-MnSOD variant with a β -sheet structure in its MTS is arrested within the inner membrane import complex TIM23 and partially degraded in the proteasome. Therefore, the Val-MnSOD variant showed lower enzymatic activity (165, 166). However, conflicting data have been reported from Bastaki *et al.* and Martin *et al.*, showing that Val-MnSOD variant has greater MnSOD activity than the Ala-MnSOD variant (107). The discrepancy in these reports may stem from different *in vitro* and *in vivo* models used in these studies, suggesting a cell type/content dependent effect for this dimorphism. Nevertheless, more studies are needed to elucidate the reasons for the differential results, and to further understand the effect of the dimorphism in the MnSOD activity. Most importantly, the clinical relevance of this dimorphism and whether it could modulate susceptibility to various human diseases requires further investigation.

Oxidative stress-mediated regulation of mitochondrial processing of MnSOD

Mitochondrial processing of human precursor MnSOD was shown to be compromised by superoxide generating compound paraquat (186), which is a redox-cycling agent that transfers electrons to molecular oxygen to form superoxide (54). It has been demonstrated that the mitochondrial processing of human MnSOD from precursor to the mature form was inhibited by paraquat treatment. This, in turn, resulted in lower MnSOD activity. The mechanism of paraquat-mediated inhibition is not known; however, it is suggested that paraquat induces a specific lesion in the MnSOD processing machinery (186). It is believed that paraquat treatment leads to

oxidation of critical sulfhydryl groups (cysteine [Cys] residues) of the proteins of the mitochondrial import machinery (185). Hyperoxia is another well-known condition that increases superoxide generation in the mitochondria (53). Similarly, hyperoxic conditions were shown to inhibit MnSOD processing by interfering with the import machinery. These studies suggest the mitochondrial import machinery as a susceptible cellular target of superoxide radicals and support a role for ROS in decreasing mitochondrial import and activation of MnSOD enzyme under oxidizing conditions.

Regulation of MnSOD enzymatic activity within mitochondria under oxidative stress

MnSOD is essential and biologically significant to aerobic cells. Studies from the early 1970s using *E. coli* and yeast as model organisms provide the early evidence for the importance of MnSOD expression for surviving in aerobic environments (61). Later studies demonstrated clearly that total MnSOD gene knockout is lethal in mice (182) and fruit flies (40); and heterozygous mice with lower MnSOD activity are more susceptible to oxidative injury (169, 171). These, along with numerous other studies, indicated the crucial role of MnSOD in protecting against oxidative stress (85, 123, 195), including radiation insult (47, 64, 79, 130, 195). Although the exact mechanisms associated with the overexpression and/or activation of MnSOD are not fully understood, there are a great deal of reports exploring the role of post-translational as well as transcriptional and post-transcriptional regulation of MnSOD in governing its superoxide dismutase activity in the adaptive response against oxidative stress.

MnSOD activation via MPI-mediated phosphorylation

Post-translational modifications of proteins control many biological processes through a variety of mechanisms that include the changes in protein activity, interactions and subcellular localizations. Post-translational modifications of mitochondrial proteins by reversible phosphorylation/dephosphorylation events play essential roles in a number of cell signaling pathways regulating the mitochondrial respiration as well as the apoptosis (77, 132). The significance of reversible phosphorylation in cellular function can be estimated by the amplitude of the genome dedicated to kinases and phosphatases (6). The involvement of mitochondrial proteins in phosphorylation events has been studied for years and more than 60 mitochondrial proteins have been identified as phosphoproteins (9, 72). The recent advances of mass spectrometry along with the development of new reagents, including phospho-specific antibodies and dyes enabled the characterization of the post-translational modifications of MnSOD and many other mitochondrial phosphoproteins (77).

The first evidence suggesting MnSOD is phosphorylated came from the studies with bacteria (7, 173). Mass spectrometric analysis of phosphoproteins in *Campylobacter jejuni* identified SOD as one of the cytoplasmic phosphoproteins (173). Archambaud *et al.* reported that MnSOD from the cytoplasmic *Listeria monocytogenes* was phosphorylated on serine and threonine residues and the phosphorylated MnSOD was less active when the bacteria were in stationary phase of growth (7). This study was not only the pioneering work showing that MnSOD can be phosphorylated, but also the first to indicate that phosphorylation of MnSOD controls its

activity, although the relevant biological effects of this activity remained to be investigated. Likewise, later studies predicted mitochondrial MnSOD as a target of phosphorylation. Phosphorylation of MnSOD was first reported in potato mitochondria but no activity measurements were conducted (21). Hopper *et al.* observed phosphorylated MnSOD in porcine heart mitochondria using phosphoproteome technique (77). Rat MnSOD is also subjected to phosphorylation shown by *in vitro* kinase assays using mitochondrial extracts as the kinase source (25). Taken together, these initial studies suggested that MnSOD is regulated by phosphorylation in a large scale of species, from bacteria to mammals.

Parallel studies demonstrate that the phosphorylation of antioxidant proteins is linked with the growth in mammalian cells (27). Chang *et al.* showed that phosphorylation of the human antioxidant peroxiredoxin I (Prx 1) by cyclin-dependent kinase 1 (Cdk1) resulted in the inactivation of this antioxidant and affected cell cycle progression (27). Prx are the enzymes involved in the elimination of H_2O_2 and are present in organisms from all kingdoms (142). At least six Prx enzymes are present in mammals (142, 155) and these are distributed differentially within cells: Prx I and II are localized to the cytosol (125), where Prx III and V to the mitochondria (155). Prx I contains a consensus site for Cdk1 phosphorylation and the phosphorylation of Prx I at Thr90 reduces its peroxidase activity (27). In addition to Cdk1; Cdk2, Cdk4, and Cdk6 can also phosphorylate Prx I *in vitro*. The regulation of Prx I with Cdk-mediated phosphorylation may shed light onto the regulation of MnSOD antioxidant enzyme through similar mechanisms.

Human MnSOD contains a minimal Cdk1 phosphorylation consensus sequence (Serine/Threonine-proline [Ser/Thr-Pro]) (170) at Ser106. The fact that Ser106 is very accessible in the final homotetrameric conformation of MnSOD encouraged the further analysis of the possibility of phosphorylation at this residue (Fig. 2). We recently found that substantial amounts of CyclinB1/Cdk1 (22, 120) localized into the mitochondria following LDIR. The presence of any canonical MTS within these proteins was explored using the web-based software MitoProt II (<http://ihg.gsf.de/ihg/mitoprot.html>). The results showed that a potential MTS is indeed present in a 42-residue amino-terminal (N-terminal) region of CyclinB1 but not in Cdk1. Further investigations are needed to determine whether the predicted MTS is involved in mitochondrial influx of CyclinB1 and whether Cdk1 contains any internal targeting signals. Various Cdks form complexes with different cyclins to perform specific functions during each phase of the cell cycle progression. Cdk1 and CyclinB1 form a complex at late growth two (G2) phase to regulate the transition from G2 to mitosis (M) (112). Cdks are serine-threonine kinases that phosphorylate their substrates on serine and threonine residues (113). The enhanced Cdk1/MnSOD complexes and phosphorylated MnSOD were detected in mitochondrial fractions isolated from the irradiated human normal and cancer cells as well as mouse tissues (22). *In vitro* kinase assay using wild-type and mutant MnSOD (Serine106 was replaced with Alanine) verified that Cdk1-phosphorylates MnSOD at Ser106. The phosphorylation of MnSOD resulted in enhanced enzymatic activity (about 2-fold) and protein stability such that the phosphorylated MnSOD was found predominantly in tetrameric conformation, unlike the nonphosphorylated MnSOD that prefer to stay as a monomer and, therefore, was




FIG. 2. Serine 106 of MnSOD is surface accessible. Analysis of the crystal structure of MnSOD (PDB entry 1N0J) reveals that Serine 106 is located at the surface of the homotetrameric MnSOD, renders it solvent accessible in the final fold. (red: Ser106 residue; purple, blue, cyan, grey: MnSOD monomers; pink: manganese ion). MnSOD, manganese superoxide dismutase.

more prone to degradation (half-life of 4 h for phosphorylated MnSOD *versus* 3 h for unphosphorylated MnSOD). The increased activity of MnSOD improved mitochondrial functions and cellular resistance to radiation-induced apoptosis (22). These results demonstrated a unique prosurvival mechanism by which cells enhance survival *via* CyclinB1/Cdk1-mediated MnSOD activation under genotoxic stress conditions.

Additional results suggest that cells may utilize Cdk1-mediated phosphorylation of MnSOD in normal progression of cell cycle: the transient inhibition of Prx 1 activity *via* Cdk1-dependent phosphorylation is thought to cause H_2O_2 accumulation, which then stimulate the transition from G2 to M phase (27). Thus, it is logical to assume that the enhanced activity of Cdk1-phosphorylated MnSOD will further contribute to the accumulation of H_2O_2 , assisting the progression through G2 phase (Fig. 3). Nevertheless, further studies are needed to fully understand the role of Cdk1-mediated phosphorylation of MnSOD in cell cycle progression and other critical cellular outcomes.

Recent studies indicate that several cytosolic kinases translocate into the mitochondria (76, 78). A phosphorylation site identification program, NetPhosK, (www.expasy.org/www.cbs.dtu.dk/services/NetPhosK), which looks for kinase specific phosphorylation sites in eukaryotic proteins, predicted a number of kinases that can phosphorylate human MnSOD. Table 1 shows a list of kinases and phosphatases that are reported to localize to mitochondria and/or are predicted with the NetPhosK program. These proteins, along with many possible, yet to be identified ones, are likely upstream kinases/phosphatases of MnSOD. Among these, extracellular regulated kinase (ERK) has been shown to be induced by ROS,




FIG. 3. Coregulation of MnSOD and Prx1 antioxidants by Cdk1 during G2/M transition. Cell cycle kinase, Cdk1, phosphorylates both MnSOD and Prx1 antioxidants, which results in the inhibition of Prx 1 activity and increase in the MnSOD activity, both of which contribute to the accumulation of H_2O_2 , which then stimulates the transition from G2 to M phase. Cdk1, cyclin-dependent kinase 1; G2, growth two; M, mitosis; H_2O_2 , hydrogen peroxide; Prx, peroxiredoxin; ROS, reactive oxygen species.

namely H_2O_2 (177). ERK is a proline-directed protein kinase, which phosphorylates serine or threonine residues that are neighbors to prolines. Pro-Leu-Ser/Thr-Pro is the most stringent consensus sequence for ERK phosphorylation, however, the sequence Ser/Thr-Pro can also be recognized (59). This broad nature of its substrate recognition, the fact that human MnSOD contains the minimal recognition sequence for ERK, and that ERK is induced by ROS, supports the concept that ERK is an alternative upstream kinase of MnSOD. Furthermore, a recent proteomic study identified kinases interacting with MnSOD in the mitochondria following LDIR (42). Among these, hexokinase 2 (HK2), Cdk11A and pyruvate dehydrogenase kinase 3 (PDK3) are previously reported to localize into the mitochondria, suggesting them as plausible upstream kinases of MnSOD. Hexokinase 2 was previously shown to localize to the outer surface of the outer membrane (20), and it has a hydrophobic N-terminal sequence capable of mitochondrial targeting (164); however, whether it also localizes into the matrix under low dose radiation or their interaction with MnSOD is an experimental artifact requires further investigation. Cdk11A was found to translocate into the mitochondria upon death signal stimulation (50) and PDK3 is part of the mitochondrial multienzyme complex, pyruvate dehydrogenase (63); however, the details of the nature of their interaction with MnSOD remains elusive. Although the exact kinases/phosphatases involved and the sites of phosphorylation of MnSOD remain to be determined, activation/inactivation of MnSOD may provide tools to protect against diseases associated with superoxide, such as cancer, cardiovascular and neurodegenerative diseases.

MnSOD regulation via other post-translational modifications in mitochondria

As recent developments in mass spectrometry enabled detailed structural analysis of covalent modifications of proteins, the identification of post-translational modifications are largely facilitated as well. MnSOD is one of the many proteins whose covalent modifications are widely studied, especially due to its vital role as the mitochondrial antioxidant and the urgent need to elucidate the regulatory mechanisms of its dismutase activity. Besides phosphorylation, the post-translational modifications of MnSOD discovered so far include nitration, acetylation, glutathionylation, methylation

TABLE 1. PROTEIN KINASES AND PHOSPHATASES IN THE MITOCHONDRIA^a ALONG WITH MnSOD UPSTREAM KINASES PREDICTED BY NETPHOS KINASE DATABASE

	Mitochondrial location	References
Protein kinases		
Src	Inner membrane/IMS	(148), NetPhos-predicted
Abl	Unknown	(93)
EGFR	Inner membrane	(17), NetPhos-predicted
RSK	Unknown	NetPhos-predicted
RAF1	Outer membrane	(176)
MEK	Outer membrane	(78)
GSK3 β	Unknown	(14), NetPhos-predicted
PKA	Unknown	(154), NetPhos-predicted
ERK	Outer membrane/IMS/Matrix	(87)
JNK	Outer membrane	(44)
Cdk11A	Matrix	(50)
Cdk1	Matrix	(120), NetPhos-predicted
Cdk5	Unknown	NetPhos-predicted
Akt	Inner membrane/Matrix	(15)
Mtor	Outer membrane	(35)
PDK3	Matrix	(76)
HK2	Matrix	(42)
p38	Unknown	(44), NetPhos-predicted
Protein phosphatases		
MKP1	Outer membrane	(147)
PTP1D	Outer membrane	(23)
PTPMT1	Inner membrane	(132)
SHP2	Inner membrane/IMS	(149)
PP2A	Outer membrane	(78)
PDPs	Matrix	(76)
BCKDP	Matrix	(188)

^aAdapted from (131).

IMS, intermembrane space; MnSOD, manganese superoxide dismutase.

and metal incorporation. The amino acid residues of MnSOD that are involved in post-translational modifications, along with the executing molecules, are summarized in Table 2.

Nitration. The nitrogen monoxide molecule ($\bullet NO$) is a ubiquitous cellular messenger, whose overproduction is associated with several diseases. The pathological effects of NO are linked to the generation of RNS, such as nitrogen oxides (NO_x) and peroxynitrite ($ONOO^-$), as a result of its reaction with oxygen and superoxide, respectively (41). Of these, peroxynitrite stands out as the biologically relevant nitrating agent (62). The reaction of proteins with nitrating agents results in the covalent modification of their tyrosine (Tyr) residues (41, 62). Tyr nitration has been shown to severely diminish the function of proteins when the nitrated Tyr residues are in the active site of the target peptide (41, 62). However, protein nitration is a highly selective process since a relatively limited number of proteins are preferential targets of nitration and within these proteins; only one or a few specific Tyr residues have been identified to be nitrated *in vivo* (8).

In situations of increased oxidative stress, NO is found to react with $O_2^{\bullet -}$ and forms the highly reactive peroxynitrite

TABLE 2. POST-TRANSLATIONAL MODIFICATIONS OF MnSOD

Residue	Post-translational modification	Mediators	References
Serine 106	Phosphorylation	Cdk1	(22)
Tyrosine 34	Nitration	Peroxynitrite	(102, 137, 141, 163)
Lysine 122	Acetylation, methylation	SIRT3	(129, 150, 168)
Lysine 53	Acetylation	Unknown	(136)
Lysine 89	Acetylation, methylation	Unknown	(136, 150)
Cysteine 196	Glutathionylation	Unknown	(25, 108)
Lysine 68	Methylation	Unknown	(150)
Lysine 202	Methylation	Unknown	(150)
Arginine 197	Methylation	Unknown	(150)
Arginine 216	Methylation	Unknown	(150)

(12). Peroxynitrite is well-established to inactivate MnSOD by nitrating a critical Tyr residue (Tyr 34) at the enzyme active site both *in vitro* and *in vivo* (102, 137, 141, 163). A recent study also reported Tyr nitration of MnSOD and resultant inactivation of the enzyme upon exposure to NO (162). MnSOD protects cells from deleterious effects associated with the overproduction of NO and peroxynitrite in two ways: (i) by reacting with RNS itself; (ii) *via* its active role in the detoxification of ROS, so it prevents the reactions of ROS with NO to form peroxynitrite (163). However, it must be re-emphasized that the reactions of MnSOD with RNS results in impaired enzymatic activity meaning that the nitration/inactivation of MnSOD operates as a positive feedback loop for enhanced mitochondrial peroxynitrite formation, which will eventually lead to the amplification of oxidative stress by allowing the accumulation of $O_2^{\bullet-}$ and trigger apoptosis (138). Besides, peroxynitrite also causes lipid peroxidation, DNA damage and nitration of proteins other than MnSOD. The importance of MnSOD in preventing the nitration of proteins can be better comprehended by looking at the pathologies associated with oxidative stress and overproduction of NO (65, 189). In these pathologic conditions, MnSOD itself is Tyr nitrated and inactivated (103).

Acetylation. The discovery of the mitochondrial localization of at least three sirtuin family of NAD^+ -dependent protein deacetylase enzymes (SIRT3, SIRT4 and SIRT5) suggested the existence of sirtuin substrates in the mitochondria (157). With the addition of several lines of evidence linking SIRT3 to metabolism (157), lysine acetylation has recently emerged as an important post-translational modification to regulate mitochondrial proteins (99). When studies with Sirt3 knockout cells showed increased mitochondrial superoxide levels upon exposure to IR (4), the researchers suspected altered regulations of MnSOD in Sirt3-lacking cells. Additional support came from the studies observing significantly increased MnSOD acetylation and corresponding decrease in the MnSOD superoxide dismutase activity in Sirt3 knockout cells (168). Tao *et al.* further identified Lys 122 of MnSOD as a reversibly acetylated residue (168). These pioneering investigations on the regulation of MnSOD activity by changes in specific lysine acetylation are followed by the reports on SIRT3-mediated regulation of MnSOD activity. Qiu *et al.* found that calorie restriction induces SIRT3-mediated MnSOD activity resulting in reduced oxidative stress (136). In the same study, Lys 53 and Lys 89 were identified as acetylated residues that control enzymatic function. Moreover,

Ozden *et al.* demonstrated that SIRT3 is activated by oxidative stress induced by IR as well as by cellular nutrient status to deacetylase Lys 122 of MnSOD and regulate MnSOD enzymatic activity (129). Taken together, MnSOD acetylation represents an alternative mechanism of post-translational regulation of MnSOD, which may contribute to the overall MnSOD activity in stress response.

Glutathionylation. Glutathionylation is the post-translational modification of protein Cys residues by the addition of glutathione. It is promoted by oxidative and nitrosative stress, although, some also occur in unstressed cells (31). Human Cu/ZnSOD can be glutathionylated at Cys 111, resulting in a reduction in its activity and promoting its aggregation (181). Iron superoxide dismutase (FeSOD) from a eubacterium species was also shown to be glutathionylated at Cys 57, which did not result in a significant change in its enzymatic activity; however, this modification protected the enzyme from Tyr nitration *via* peroxynitrite and was enhanced upon cell exposure to oxidative stress (26). In mammalian cells, protein glutathionylation is involved in redox signaling and the defense against oxidative stress. In mitochondria, the high concentration of glutathione has been related to the presence of several glutathionylated proteins (83). The known sensitivity of human mitochondrial MnSOD to reducing agents (108) has been associated with a highly reactive cysteine residue (Cys 196) exposed on the surface of the tetrameric enzyme. So far, glutathionylation of MnSOD was only found for rat recombinant MnSOD grown in *E. coli* using mass spectrometry analysis (25). On the basis of high similarity between rat and human MnSOD, the candidate cysteine residue, Cys 196, was identified as the target of glutathionylation (25). Accumulating evidence suggests glutathionylation as an important mechanism of cellular response to oxidative damage and redox signaling (133). Future research is needed to elucidate the biological function of glutathionylation of MnSOD as well as other mitochondrial targets *in vivo* and the possible role of such post-translational modification in enzyme activity and cellular outcomes.

Methylation. A recent study with mass spectrometry revealed a complex pattern of lysine and arginine methylation of MnSOD in both quiescent and proliferating mouse embryonic fibroblast (MEF) cells (150). Methylation of MnSOD was detected at lysines 68, 89, 122, and 202; and arginines 197 and 216. Lysine 68 was shown to be dimethylated during quiescence and monomethylated during proliferation states

of fibroblasts. Lysines 89 and lysine 202, on the other hand, were monomethylated during quiescence and unmethylated in proliferating cells. Arginine 197 was shown to be dimethylated, whereas arginine 216 was monomethylated in quiescent cells; and the arginine 197 dimethylation did not change between quiescent and proliferative growth states. These results indicated a dynamic regulation of MnSOD *via* methylation during cell cycle. In the same study, mutagenesis analysis determined that lysine 89 methylation is important for MnSOD activity, while lysine 202 methylation is not. Further computational modeling simulations based on the mass spectrometry data demonstrated that lysine and arginine methylation of MnSOD may increase the accessibility of the active site to superoxide during quiescence in MEF cells.

Metal incorporation. Insertion of the catalytic metals into the SOD is necessary for full activation of the enzymes and occurs post-translationally. After the precursor polypeptide MnSOD enters the mitochondria matrix, the mitochondria targeting sequence is cleaved off and manganese is subsequently inserted into the MnSOD. Two membrane transporters have been studied as facilitators of manganese insertion to MnSOD in yeast: (i) Smf2p, a manganese transporter critical for cell surface uptake of manganese (101); (ii) Mtm1p, a member of the mitochondrial carrier family of transporters residing in the mitochondrial inner membrane and important for the exchange of solutes between mitochondria and cytosol (100). Studies with Smf2 mutants showed accumulation of an iron-substituted form of MnSOD in the mitochondria (192). Monomethylamine permease (Mtmp1) mutants exhibited lost MnSOD activity, albeit accumulation of high levels of mitochondrial manganese occurred (100). Iron incorporated MnSOD was also detected in Mtmp1 mutants (100). Biological significance of iron misincorporated MnSOD is not fully understood, but several indications have emerged from *in vitro* metal substitution studies (191). Iron-substituted MnSOD is inactive but showed more thermal stability than the native MnSOD and a H₂O₂-mediated •OH (hydroxyl free radical) generating activity (191), pointing to a multitude of disadvantages for cells: a loss of enzymatic activity, a gain of radical generating ability and increased stability of iron-misoncorporated MnSOD. Considering iron excess in the mitochondria has been linked to various diseases (105), it would not be surprising to see the presence of iron-misincorporated MnSOD in these diseases.

The regulation of MnSOD via mitochondria-localized p53

p53 is a homotetrameric transcription factor with well-established roles in tumor suppression *via* induction of apoptosis or cell cycle arrest by post-translational and transcriptional mechanisms (97, 194). Diverse biological actions of p53 in regulating cellular functions is cell type- and stimulus-dependent, complicated, and beyond the scope of this review. However, this review will address the evidence linking mitochondrial p53 to oxidative stress and MnSOD.

Mitochondrial translocation of p53 induces apoptosis by promoting changes in mitochondrial membrane potential, cytochrome c release and caspase activation (111, 126). p53 exerts its apoptosis-inducing role by binding to multiple targets on the outer membrane of the mitochondria, such

as B-cell lymphoma (Bcl)-2/Bcl-XL (111), p53-regulated apoptosis inducing protein 1 (p53AIP1) (126) and Bcl-2-associated X protein (Bax) (28). Studies by Zhao *et al.* showed that p53 also localizes to the matrix of the mitochondria (195). In this same study, p53 was found to physically interact with MnSOD and inhibit its superoxide scavenging activity. This was the first study to show a direct role for mitochondrial p53 in the regulation of MnSOD activity and oxidative stress. Importantly, the mitochondrial translocation of p53 can be enhanced by radiation insult in wild type p53-bearing tumor cells (120, 175). Analysis of downstream effects and possible modulators of p53-MnSOD interaction will largely improve our knowledge of the molecular mechanisms of cancer progression and may provide novel therapeutic targets for cancer treatment. Among the upstream effectors of p53 is CyclinB1/Cdk1 complex, which can phosphorylate p53 at Ser315 (16). Establishment of mitochondrial localization of both p53 and CyclinB1/Cdk1 complex (120) suggested that this phosphorylation may also occur in the mitochondria. Nantajit *et al.* showed that mitochondrial Cdk1 interacted with and phosphorylated mitochondrial p53 at Ser315 in colon cancer cells upon exposure to IR (120). Interestingly, the phosphorylation of p53 at Ser315 by mitochondrial Cdk1 resulted in increased survival of the cells, suggesting an antiapoptotic role for mitochondrial phospho-p53. Further elucidation of the role of mitochondrial matrix p53 and the effect of Ser315 phosphorylation on proapoptotic functions of p53 is needed. However, we propose that the enzymatic regulation of MnSOD *via* p53 and Cdk1 induces two competing mechanisms and results in two opposite outcomes potentially depending on the severity of the oxidative stress and the resultant cellular damage: (i) proapoptotic response in case of severe injury *via* p53/MnSOD interaction and (ii) antiapoptotic response in case of minor fixable injury *via* Cdk1-p53-MnSOD pathway. Nevertheless, our current knowledge presents that the interaction of p53 with MnSOD in the mitochondria inhibits MnSOD scavenging activity and induces apoptosis, while the Cdk1-mediated phosphorylation of p53 in the mitochondria inhibits apoptosis, possibly by interfering with p53-MnSOD interaction. The studies demonstrating mitochondrial localization of p53 and CyclinB1/Cdk1 in oxidative stress conditions, together with our recently published findings that Cdk1 can directly phosphorylate and enhance MnSOD activity (22), may point to a Cdk1-p53-MnSOD regulatory mechanism in oxidative stress response (Fig. 4).

All in all, identification of MnSOD as a direct target of p53 is of great importance for tumor biology and holds enormous potential for clinical applications. p53-mediated regulation of MnSOD at the transcriptional level has also been implicated and will be addressed in the next section of this review.

Transcriptional/post-transcriptional regulation of MnSOD

MnSOD regulation is complex and occurs at both pre- and post-translational levels. Here we will briefly discuss some transcriptional/post-transcriptional mechanisms of MnSOD regulation (Fig. 5). The examples include changes in the rate of MnSOD protein synthesis (70), mRNA stability (29), and promoter activity (127) as well as regulation of chromatin accessibility *via* the methylation of CpG islands upstream of the Sod2 gene (84) and histone modifications (73). Of the




FIG. 4. Cdk1-p53-mediated MnSOD regulation. p53 interacts with MnSOD in the mitochondria and inhibits its superoxide scavenging activity, which results in accumulation of ROS and finally apoptosis. Recent findings showing that Cdk1-mediated phosphorylation of p53 at Ser 315 in the mitochondria results in antiapoptotic response supports a model, where both p53 and MnSOD are regulated by Cdk1 *via* phosphorylation. According to this model, phosphorylation of p53 by Cdk1 in the mitochondria releases MnSOD, which then becomes a free substrate for Cdk1. Phosphorylation of MnSOD at Ser 106 by Cdk1, enhances its superoxide dismutase activity, resulting in reduced ROS levels and apoptosis.

factors regulating MnSOD expression, we will address nuclear factor-kappaB (NF- κ B), p53, mTOR and mitogen-activated protein kinases (MAPKs) (ERK and p38).

The connection between NF- κ B and MnSOD has been observed in several studies on radioprotection compounds, where radioprotective thiols are shown to control MnSOD gene expression through NF- κ B activation (32, 60, 117). In mammals, NF- κ B family of transcription factors consists of five members: RelA (p65), RelB, c-Rel, p50/p105 (NF- κ B1) and p52/p100 (NF- κ B2). Although different combinations of homo- and heterodimers can be formed (172), the heterodimer of p50 and p65 is the most abundant form of NF- κ B (119). NF- κ B binding sites were found to be located in the promoter region (174, 190) as well as within the intronic enhancer element of the Sod2 gene (91). NF- κ B is actively involved in the regulation of MnSOD expression in cells induced with various stimuli, including radiation (47, 64), thiol-containing drugs (117) and tumor necrosis factor, tumor necrosis factor alpha (TNF α) (32, 60). Moreover, NF- κ B-induced MnSOD has been related to radioresistant phenotype of normal and cancer cells (64, 117, 124), pointing to MnSOD as a key NF- κ B effector gene in radioadaptive resistance.

Likewise, there are p53 binding regions at 328 and 2032 bp upstream of the transcriptional start site of the Sod2 gene, mutations of which were shown to result in altered MnSOD expression (36, 37, 85). Apart from direct binding to the gene, p53 has been suggested to repress Sod2 gene expression by a number of mechanisms, including interfering with transcription initiation (85), blocking the access of gene activators at the promoter region by forming an inhibitory complex with them (48) and protein-protein interactions (96). In contrast, p53 can induce Sod2 gene expression (36, 96). The opposite

effects of p53 on Sod2 expression is proposed to be cell type-, species- and concentration-dependent such that low concentrations of p53 increases MnSOD expression jointly with other transcription factors, such as NF- κ B; and high concentrations of p53 suppresses MnSOD expression (37, 91). Interestingly, mutations of the NF- κ B binding sites within the intronic enhancer element of the MnSOD gene inhibited p53-induced MnSOD gene transcription; and knocking down p53 *via* siRNA similarly resulted in reduced MnSOD gene transcription through p53 (91).

MnSOD upregulation through ROS-driven ERK-dependent activation of p53 has also been indicated (96). According to this study, ROS induced the translocation of ERK into the nucleus, where it interacted with and phosphorylated p53 at Ser15, resulting in the activation of p53 and subsequent increase in MnSOD expression. Involvement of another MAPK, p38, was also implicated in regulation of MnSOD expression (167, 184). A recent study with epithelial stem cells proposed that inhibition of mammalian target of rapamycin (mTOR) mediated the increased expression of MnSOD, resulting in the suppression of oxidative stress caused by the accumulation of ROS after radiation in normal human oral keratinocytes but not in head and neck squamous cell carcinoma (HNSCC) cells (86). Interestingly, the inhibition of mTOR did not increase the MnSOD mRNA levels, indicating that translational (such as increased rate of translation) or post-translational (such as enhanced protein stability due to protein modifications) mechanisms, rather than transcriptional events, may account for the increased MnSOD expression. Nonetheless, these results showed that mTOR inhibitors may reduce oxidative stress by inducing MnSOD overexpression and provided strong clinical implications for anticancer treatment.




FIG. 5. MnSOD-mediated adaptive response through transcriptional regulation. Oxidative stresses, such as ionizing radiation, can activate several cytoplasmic factors, including NF- κ B, p53, mTOR, ERK and p38 MAPKs, which then translocate into the nucleus and induce the expression of Sod2 gene. The enhanced MnSOD expression, in turn, results in reduced ROS levels in the mitochondria (orange arrows) and induces other transcription factors and transcriptional regulators that modulate the expression of several MnSOD-regulated genes with various cellular functions, such as cell cycle, apoptosis and DNA repair (green arrows). Coherent activity of a number of these MnSOD-overexpression-driven genes dictates the adaptive response of the cell to the oxidative stress. ERK, extracellular regulated kinase; MAPK, mitogen-activated protein kinase; mTOR, mammalian target of rapamycin; NF- κ B, nuclear factor-kappaB.





FIG. 6. A pie chart showing the percentages of putative binding partners of MnSOD in human skin keratinocytes upon LDIR [adapted from (42)]. These results indicate that MnSOD may act as a signaling mediator to activate the cellular adaptive response induced by LDIR. LDIR, low dose ionizing radiation.

Finally, the concept of MnSOD overexpression raises the questions regarding the levels of H_2O_2 that will accumulate as a result of such overexpression. However, mitochondria seem to co-regulate MnSOD expression with mitochondrial H_2O_2 -removing capacity to prevent extreme fluctuations in redox status (90). The essence of maintaining optimal MnSOD protein levels is also supported by the findings that even though MnSOD mRNA levels were greatly elevated following the induction of gene expression, the resultant MnSOD protein levels were not as high (90). These observations implied a tight control over the expression of MnSOD protein despite the availability of mRNA for translation.

MnSOD-regulated gene expression

MnSOD has been shown to increase the expression of a number of genes participating in diverse functions related to radiation-induced adaptive responses (64, 118). These genes include MET, α -catenin, p21, v-myc myelocytomatosis viral oncogene homolog (Myc), 14-3-3 zeta, CyclinA, and CyclinB1. A more recent study with MnSOD-overexpressing HeLa cells identified MnSOD overexpression-driven expression of several other transcripts (79). This new set of genes includes GADD45, RAS homolog, tumor necrosis factor receptor, NF- κ B, glucocorticoid receptor, heat shock proteins, cyclinD1, survivin and ataxin1. A similar study in MnSOD-overexpressing adult *Drosophila* revealed numerous cellular signaling pathways induced by MnSOD overexpression (30). Among these are both pro- and antiapoptotic genes as well as members of NF- κ B, Jun N-terminal kinase (JNK), ERK, Janus kinase (JAK)/signal transducer and activator of transcription (STAT), cell cycle, and ubiquitin-mediated degradation pathways. These studies significantly contributed to the understanding of the downstream signaling molecules involved in the MnSOD-induced cell radiation-resistance phenotype. Regulation of a variety of cellular functions by MnSOD may encourage studies to determine specific targets to enhance radiation tolerance in normal cells (Fig. 5). Likewise, further studies to fully elucidate the mechanisms of MnSOD-mediated radioprotection may provide novel therapeutic targets to modulate tumor radiosensitivity during anticancer therapy as antioxidants, such as MnSOD are currently investigated for cancer treatment.

MnSOD interactome

Many hemostatic functions of cells, including the maintenance of the mitochondrial membrane (1, 2), activation of TNF signaling (115) and inhibition of aggressive phenotype of cancer cells (81, 94), can be induced by MnSOD under varied oxidative conditions (43, 46, 122). However, the exact MnSOD-mediated signaling network regulating the oxidative stress-induced adaptive response remains elusive. A recent study carried out in human skin keratinocytes (HK18) aimed to determine the signaling network associated with the MnSOD-induced radiation protection (42). In this study, a MnSOD-interacting protein profile was established in LDIR-treated HK18 cells. Analysis of the profiles of MnSOD-interacting partners before and after LDIR identified different patterns of MnSOD protein-protein interactions, suggesting a shift in the MnSOD-interacting partners following radiation exposure. Interestingly, many of the MnSOD-interacting proteins detected in this study are known to have functions related

to mitochondrial regulation of cell metabolism, apoptosis and DNA repair (Fig. 6). These results led to the assumption that the role of MnSOD in the adaptive response is not only due to its dismutase activity, but also relates to its interactions/communications with many cellular/mitochondrial proteins, an interesting topic worthy of further investigation.

Innovation

Extensive communications between the mitochondria and cytoplasm regulate the activity of mitochondrial antioxidant MnSOD under oxidative stress. Proteins, such as cyclins, Cdks and p53 are imported into the mitochondria to alter MnSOD activity *via* post-translational modifications. Recent findings suggest phosphorylation of human MnSOD in mitochondria as a novel post-translational modification that enhances its superoxide scavenging activity in oxidative stress conditions. The regulation of mitochondrial import of MnSOD itself as well as of the proteins that modulate MnSOD activity was broadly discussed in this Forum review.

Conclusion

The mitochondrion imports and processes the vast majority of its proteins, including its structural elements and metabolic pathways from cytosol. This suggests that the regulation of MPI is key to the proper functioning of the mitochondria and of the cells. Here we discussed the specifics of mitochondrial protein import and its regulation, especially under oxidizing conditions. The mitochondrial superoxide dismutase MnSOD, the primary defense agent of mitochondria against oxidants, is of particular interest since its dysfunction is indicated in many pathological conditions. The mitochondrial import and processing of MnSOD itself as well as the mitochondrial influx of several proteins that regulate MnSOD enzymatic activity under oxidative insult reveals the importance of mitochondria, specifically its protein import apparatus, in cellular response to genotoxic environment. Along with the post-translational regulation of MnSOD activity, the transcriptional regulation of Sod2 gene and MnSOD-interactome profile in oxidative stress present a unique coordinated network that governs the adaptive response. Further studies will unfold additional regulatory mechanisms of this pivotal antioxidant enzyme and will instigate novel strategies to fight against diseases associated with defective MnSOD activity.

Acknowledgments

The authors would like to thank Dominik J. Green for his help in presentation of MnSOD protein structure and discussion on specific residues of MnSOD. The authors acknowledge Gayle Woloschak, David Grdina, Douglas Spitz, and Daret St. Clair for their communication and suggestions to the research in the author's lab. We would like to acknowledge the support of National Institutes of Health Grants RO1 CA133402 and RO1 CA152313, and the Department of Energy Office of Science Grant DE-SC0001271.

References

1. Ahmed KM, Fan M, Nantajit D, Cao N, and Li JJ. Cyclin D1 in low-dose radiation-induced adaptive resistance. *Oncogene* 27: 6738–6748, 2008.

2. Ahmed KM and Li JJ. NF-kappa B-mediated adaptive resistance to ionizing radiation. *Free Radic Biol Med* 44: 1–13, 2008.
3. Ahmed KM, Nantajit D, Fan M, Murley JS, Grdina DJ, and Li JJ. Coactivation of ATM/ERK/NF-kappaB in the low-dose radiation-induced radioadaptive response in human skin keratinocytes. *Free Radic Biol Med* 46: 1543–1550, 2009.
4. Ahn BH, Kim HS, Song S, Lee IH, Liu J, Vassilopoulos A, Deng CX, and Finkel T. A role for the mitochondrial deacetylase Sirt3 in regulating energy homeostasis. *Proc Natl Acad Sci U S A* 105: 14447–14452, 2008.
5. Aitken A, Collinge DB, van Heusden BP, Isobe T, Roseboom PH, Rosenfeld G, and Soll J. 14-3-3 proteins: a highly conserved, widespread family of eukaryotic proteins. *Trends Biochem Sci* 17: 498–501, 1992.
6. Alonso A, Sasin J, Bottini N, Friedberg I, Osterman A, Godzik A, Hunter T, Dixon J, and Mustelin T. Protein tyrosine phosphatases in the human genome. *Cell* 117: 699–711, 2004.
7. Archambaud C, Nahori MA, Pizarro-Cerda J, Cossart P, and Dussurget O. Control of *Listeria* superoxide dismutase by phosphorylation. *J Biol Chem* 281: 31812–31822, 2006.
8. Aulak KS, Miyagi M, Yan L, West KA, Massillon D, Crabb JW, and Stuehr DJ. Proteomic method identifies proteins nitrated *in vivo* during inflammatory challenge. *Proc Natl Acad Sci U S A* 98: 12056–12061, 2001.
9. Azarashvili TS, Tynnela J, Odinkova IV, Grigorjev PA, Baumann M, Evtodienko YV, and Saris NE. Phosphorylation of a peptide related to subunit c of the F0F1-ATPase/ATP synthase and relationship to permeability transition pore opening in mitochondria. *J Bioenerg Biomembr* 34: 279–284, 2002.
10. Azzam EI, Jay-Gerin JP, and Pain D. Ionizing radiation-induced metabolic oxidative stress and prolonged cell injury. *Cancer Lett* 327: 48–60, 2012.
11. Barra D, Schinina ME, Simmaco M, Bannister JV, Bannister WH, Rotilio G, and Bossa F. The primary structure of human liver manganese superoxide dismutase. *J Biol Chem* 259: 12595–12601, 1984.
12. Beckman JS, Beckman TW, Chen J, Marshall PA, and Freeman BA. Apparent hydroxyl radical production by peroxynitrite: implications for endothelial injury from nitric oxide and superoxide. *Proc Natl Acad Sci U S A* 87: 1620–1624, 1990.
13. Bhattacharjee D. Role of radioadaptation on radiation-induced thymic lymphoma in mice. *Mutat Res* 358: 231–235, 1996.
14. Bijur GN and Jope RS. Glycogen synthase kinase-3 beta is highly activated in nuclei and mitochondria. *Neuroreport* 14: 2415–2419, 2003.
15. Bijur GN and Jope RS. Rapid accumulation of Akt in mitochondria following phosphatidylinositol 3-kinase activation. *J Neurochem* 87: 1427–1435, 2003.
16. Bischoff JR, Friedman PN, Marshak DR, Prives C, and Beach D. Human p53 is phosphorylated by p60-cdc2 and cyclin B-cdc2. *Proc Natl Acad Sci U S A* 87: 4766–4770, 1990.
17. Boerner JL, Demory ML, Silva C, and Parsons SJ. Phosphorylation of Y845 on the epidermal growth factor receptor mediates binding to the mitochondrial protein cytochrome c oxidase subunit II. *Mol Cell Biol* 24: 7059–7071, 2004.
18. Boonstra J and Post JA. Molecular events associated with reactive oxygen species and cell cycle progression in mammalian cells. *Gene* 337: 1–13, 2004.
19. Borgstahl GE, Parge HE, Hickey MJ, Beyer WF, Jr., Hallewell RA, and Tainer JA. The structure of human mitochondrial manganese superoxide dismutase reveals a novel tetrameric interface of two 4-helix bundles. *Cell* 71: 107–118, 1992.
20. Bustamante E and Pedersen PL. High aerobic glycolysis of rat hepatoma cells in culture: role of mitochondrial hexokinase. *Proc Natl Acad Sci U S A* 74: 3735–3739, 1977.
21. Bykova NV, Egsgaard H, and Moller IM. Identification of 14 new phosphoproteins involved in important plant mitochondrial processes. *FEBS Lett* 540: 141–146, 2003.
22. Candas D, Fan M, Nantajit D, Vaughan AT, Murley J, Woloschak G, Grdina DJ, and Li JJ. CyclinB1/Cdk1 phosphorylates mitochondrial antioxidant MnSOD in cell adaptive response to radiation stress. *J Mol Cell Biol* 2013 [Epub ahead of print]; DOI: 10.1093/jmcb/mjs062
23. Cardone L, Carlucci A, Affaitati A, Livigni A, DeCristofaro T, Garbi C, Varrone S, Ullrich A, Gottesman ME, Avvedimento EV, and Feliciello A. Mitochondrial AKAP121 binds and targets protein tyrosine phosphatase D1, a novel positive regulator of src signaling. *Mol Cell Biol* 24: 4613–4626, 2004.
24. Cartwright P, Beilharz T, Hansen P, Garrett J, and Lithgow T. Mft52, an acid-bristle protein in the cytosol that delivers precursor proteins to yeast mitochondria. *J Biol Chem* 272: 5320–5325, 1997.
25. Castellano I, Cecere F, De Vendittis A, Cotugno R, Chambrey A, Di Maro A, Michniewicz A, Parlato G, Masullo M, Avvedimento EV, De Vendittis E, and Ruocco MR. Rat mitochondrial manganese superoxide dismutase: amino acid positions involved in covalent modifications, activity, and heat stability. *Biopolymers* 91: 1215–1226, 2009.
26. Castellano I, Ruocco MR, Cecere F, Di Maro A, Chambrey A, Michniewicz A, Parlato G, Masullo M, and De Vendittis E. Glutathionylation of the iron superoxide dismutase from the psychrophilic eubacterium *Pseudoalteromonas haloplanktis*. *Biochim Biophys Acta* 1784: 816–826, 2008.
27. Chang TS, Jeong W, Choi SY, Yu S, Kang SW, and Rhee SG. Regulation of peroxiredoxin I activity by Cdc2-mediated phosphorylation. *J Biol Chem* 277: 25370–25376, 2002.
28. Chipuk JE, Kuwana T, Bouchier-Hayes L, Droin NM, Newmeyer DD, Schuler M, and Green DR. Direct activation of Bax by p53 mediates mitochondrial membrane permeabilization and apoptosis. *Science* 303: 1010–1014, 2004.
29. Clerch LB and Massaro D. Tolerance of rats to hyperoxia. Lung antioxidant enzyme gene expression. *J Clin Invest* 91: 499–508, 1993.
30. Curtis C, Landis GN, Folk D, Wehr NB, Hoe N, Waskar M, Abdueva D, Skvortsov D, Ford D, Luu A, Badrinath A, Levine RL, Bradley TJ, Tavare S, and Tower J. Transcriptional profiling of MnSOD-mediated lifespan extension in *Drosophila* reveals a species-general network of aging and metabolic genes. *Genome Biol* 8: R262, 2007.
31. Dalle-Donne I, Rossi R, Colombo G, Giustarini D, and Milzani A. Protein S-glutathionylation: a regulatory device from bacteria to humans. *Trends Biochem Sci* 34: 85–96, 2009.
32. Das KC, Lewis-Molock Y, and White CW. Thiol modulation of TNF alpha and IL-1 induced MnSOD gene expression and activation of NF-kappa B. *Mol Cell Biochem* 148: 45–57, 1995.
33. de Diego-Otero Y, Romero-Zerbo Y, el Bekay R, Decara J, Sanchez L, Rodriguez-de Fonseca F, and del Arco

- Herrera I. Alpha-tocopherol protects against oxidative stress in the fragile X knockout mouse: an experimental therapeutic approach for the Fmr1 deficiency. *Neuropsychopharmacology* 34: 1011–1026, 2009.
34. Dekker PJ, Martin F, Maarse AC, Bomer U, Muller H, Guiard B, Meijer M, Rassow J, and Pfanner N. The Tim core complex defines the number of mitochondrial translocation contact sites and can hold arrested preproteins in the absence of matrix Hsp70-Tim44. *EMBO J* 16: 5408–5419, 1997.
 35. Desai BN, Myers BR, and Schreiber SL. FKBP12-rapamycin-associated protein associates with mitochondria and senses osmotic stress via mitochondrial dysfunction. *Proc Natl Acad Sci U S A* 99: 4319–4324, 2002.
 36. Dhar SK, Xu Y, and St Clair DK. Nuclear factor kappaB and specificity protein 1-dependent p53-mediated bidirectional regulation of the human manganese superoxide dismutase gene. *J Biol Chem* 285: 9835–9846, 2010.
 37. Drane P, Bravard A, Bouvard V, and May E. Reciprocal down-regulation of p53 and SOD2 gene expression-implication in p53 mediated apoptosis. *Oncogene* 20: 430–439, 2001.
 38. Droge W. Free radicals in the physiological control of cell function. *Physiol Rev* 82: 47–95, 2002.
 39. Dudley P, Wood CK, Pratt JR, and Moore AL. Developmental regulation of the plant mitochondrial matrix located HSP70 chaperone and its role in protein import. *FEBS Lett* 417: 321–324, 1997.
 40. Duttaroy A, Paul A, Kundu M, and Belton A. A Sod2 null mutation confers severely reduced adult life span in *Drosophila*. *Genetics* 165: 2295–2299, 2003.
 41. Eiserich JP, Patel RP, and O'Donnell VB. Pathophysiology of nitric oxide and related species: free radical reactions and modification of biomolecules. *Mol Aspects Med* 19: 221–357, 1998.
 42. Eldridge A, Fan M, Woloschak G, Grdina DJ, Chromy BA, and Li JJ. Manganese superoxide dismutase interacts with a large scale of cellular and mitochondrial proteins in low-dose radiation-induced adaptive radioprotection. *Free Radic Biol Med* 53: 1838–1847, 2012.
 43. Epperly MW, Sikora C, Defilippi S, Bray J, Koe G, Liggitt D, Luketich JD, and Greenberger JS. Plasmid/liposome transfer of the human manganese superoxide dismutase transgene prevents ionizing irradiation-induced apoptosis in human esophagus organ explant culture. *Int J Cancer* 90: 128–137, 2000.
 44. Epperly MW, Sikora CA, DeFilippi SJ, Gretton JA, Zhan Q, Kufe DW, and Greenberger JS. Manganese superoxide dismutase (SOD2) inhibits radiation-induced apoptosis by stabilization of the mitochondrial membrane. *Radiat Res* 157: 568–577, 2002.
 45. Evans MD, Dizdaroglu M, and Cooke MS. Oxidative DNA damage and disease: induction, repair and significance. *Mutat Res* 567: 1–61, 2004.
 46. Fan J, Li Y, Levy RM, Fan JJ, Hackam DJ, Vodovotz Y, Yang H, Tracey KJ, Billiar TR, and Wilson MA. Hemorrhagic shock induces NAD(P)H oxidase activation in neutrophils: role of HMGB1-TLR4 signaling. *J Immunol* 178: 6573–6580, 2007.
 47. Fan M, Ahmed KM, Coleman MC, Spitz DR, and Li JJ. Nuclear factor-kappaB and manganese superoxide dismutase mediate adaptive radioresistance in low-dose irradiated mouse skin epithelial cells. *Cancer Res* 67: 3220–3228, 2007.
 48. Farmer G, Friedlander P, Colgan J, Manley JL, and Prives C. Transcriptional repression by p53 involves molecular interactions distinct from those with the TATA box binding protein. *Nucleic Acids Res* 24: 4281–4288, 1996.
 49. Feinendegen LE, Bond VP, Sondhaus CA, and Muehlen-siepen H. Radiation effects induced by low doses in complex tissue and their relation to cellular adaptive responses. *Mutat Res* 358: 199–205, 1996.
 50. Feng Y, Ariza ME, Goulet AC, Shi J, and Nelson MA. Death-signal-induced relocalization of cyclin-dependent kinase 11 to mitochondria. *Biochem J* 392: 65–73, 2005.
 51. Folz RJ and Crapo JD. Extracellular superoxide dismutase (SOD3): tissue-specific expression, genomic characterization, and computer-assisted sequence analysis of the human EC SOD gene. *Genomics* 22: 162–171, 1994.
 52. Forman HJ, Fukuto JM, and Torres M. Redox signaling: thiol chemistry defines which reactive oxygen and nitrogen species can act as second messengers. *Am J Physiol Cell Physiol* 287: C246–C256, 2004.
 53. Freeman BA and Crapo JD. Hyperoxia increases oxygen radical production in rat lungs and lung mitochondria. *J Biol Chem* 256: 10986–10992, 1981.
 54. Frei B, Winterhalter KH, and Richter C. Menadione- (2-methyl-1,4-naphthoquinone-) dependent enzymatic redox cycling and calcium release by mitochondria. *Biochemistry* 25: 4438–4443, 1986.
 55. Fridovich I. The biology of oxygen radicals. *Science* 201: 875–880, 1978.
 56. Fridovich I. Superoxide radical and superoxide dismutases. *Annu Rev Biochem* 64: 97–112, 1995.
 57. Funfschilling U and Rospert S. Nascent polypeptide-associated complex stimulates protein import into yeast mitochondria. *Mol Biol Cell* 10: 3289–3299, 1999.
 58. Gius D and Spitz DR. Redox signaling in cancer biology. *Antioxid Redox Signal* 8: 1249–1252, 2006.
 59. Gonzalez FA, Raden DL, and Davis RJ. Identification of substrate recognition determinants for human ERK1 and ERK2 protein kinases. *J Biol Chem* 266: 22159–22163, 1991.
 60. Grdina DJ, Murley JS, Kataoka Y, and Calvin DP. Differential activation of nuclear transcription factor kappaB, gene expression, and proteins by amifostine's free thiol in human microvascular endothelial and glioma cells. *Semin Radiat Oncol* 12: 103–111, 2002.
 61. Gregory EM and Fridovich I. Oxygen toxicity and the superoxide dismutase. *J Bacteriol* 114: 1193–1197, 1973.
 62. Groves JT. Peroxynitrite: reactive, invasive and enigmatic. *Curr Opin Chem Biol* 3: 226–235, 1999.
 63. Gudi R, Bowker-Kinley MM, Kedishvili NY, Zhao Y, and Popov KM. Diversity of the pyruvate dehydrogenase kinase gene family in humans. *J Biol Chem* 270: 28989–28994, 1995.
 64. Guo G, Yan-Sanders Y, Lyn-Cook BD, Wang T, Tamae D, Ogi J, Khaletskiy A, Li Z, Weydert C, Longmate JA, Huang TT, Spitz DR, Oberley LW, and Li JJ. Manganese superoxide dismutase-mediated gene expression in radiation-induced adaptive responses. *Mol Cell Biol* 23: 2362–2378, 2003.
 65. Guo W, Adachi T, Matsui R, Xu S, Jiang B, Zou MH, Kirber M, Lieberthal W, and Cohen A. Quantitative assessment of tyrosine nitration of manganese superoxide dismutase in angiotensin II-infused rat kidney. *Am J Physiol Heart Circ Physiol* 285: H1396–H1403, 2003.
 66. Guo Y, Cheong N, Zhang Z, De Rose R, Deng Y, Farber SA, Fernandes-Alnemri T, and Alnemri ES. Tim50, a

- component of the mitochondrial translocator, regulates mitochondrial integrity and cell death. *J Biol Chem* 279: 24813–24825, 2004.
67. Hachiya N, Alam R, Sakasegawa Y, Sakaguchi M, Mihara K, and Omura T. A mitochondrial import factor purified from rat liver cytosol is an ATP-dependent conformational modulator for precursor proteins. *EMBO J* 12: 1579–1586, 1993.
 68. Halliwell B. Oxidative stress and cancer: have we moved forward? *Biochem J* 401: 1–11, 2007.
 69. Halliwell B and Gutteridge JMC. *Free Radicals in Biology and Medicine*. Oxford: Clarendon Press, 1989.
 70. Hass MA and Massaro D. Regulation of the synthesis of superoxide dismutases in rat lungs during oxidant and hyperthermic stresses. *J Biol Chem* 263: 776–781, 1988.
 71. Hawlitschek G, Schneider H, Schmidt B, Tropschug M, Hartl FU, and Neupert W. Mitochondrial protein import: identification of processing peptidase and of PEP, a processing enhancing protein. *Cell* 53: 795–806, 1988.
 72. He H, Chen M, Scheffler NK, Gibson BW, Spremulli LL, and Gottlieb RA. Phosphorylation of mitochondrial elongation factor Tu in ischemic myocardium: basis for chloramphenicol-mediated cardioprotection. *Circ Res* 89: 461–467, 2001.
 73. Hitchler MJ, Oberley LW, and Domann FE. Epigenetic silencing of SOD2 by histone modifications in human breast cancer cells. *Free Radic Biol Med* 45: 1573–1580, 2008.
 74. Holley AK, Bakthavatchalu V, Velez-Roman JM, and St Clair DK. Manganese superoxide dismutase: guardian of the powerhouse. *Int J Mol Sci* 12: 7114–7162, 2011.
 75. Holley AK, Xu Y, St Clair DK, and St Clair WH. RelB regulates manganese superoxide dismutase gene and resistance to ionizing radiation of prostate cancer cells. *Ann NY Acad Sci* 1201: 129–136, 2010.
 76. Holness MJ and Sugden MC. Regulation of pyruvate dehydrogenase complex activity by reversible phosphorylation. *Biochem Soc Trans* 31: 1143–1151, 2003.
 77. Hopper RK, Carroll S, Aponte AM, Johnson DT, French S, Shen RF, Witzmann FA, Harris RA, and Balaban RS. Mitochondrial matrix phosphoproteome: effect of extra mitochondrial calcium. *Biochemistry* 45: 2524–2536, 2006.
 78. Horbinski C and Chu CT. Kinase signaling cascades in the mitochondrion: a matter of life or death. *Free Radic Biol Med* 38: 2–11, 2005.
 79. Hosoki A, Yonekura S, Zhao QL, Wei ZL, Takasaki I, Tabuchi Y, Wang LL, Hasuike S, Nomura T, Tachibana A, Hashiguchi K, Yonei S, Kondo T, and Zhang-Akiyama QM. Mitochondria-targeted superoxide dismutase (SOD2) regulates radiation resistance and radiation stress response in HeLa cells. *J Radiat Res* 53: 58–71, 2012.
 80. Houten SM and Wanders RJ. A general introduction to the biochemistry of mitochondrial fatty acid beta-oxidation. *J Inherit Metab Dis* 33: 469–477, 2010.
 81. Huang P, Feng L, Oldham EA, Keating MJ, and Plunkett W. Superoxide dismutase as a target for the selective killing of cancer cells. *Nature* 407: 390–395, 2000.
 82. Hunt CR, Sim JE, Sullivan SJ, Featherstone T, Golden W, Von Kapp-Herr C, Hock RA, Gomez RA, Parsian AJ, and Spitz DR. Genomic instability and catalase gene amplification induced by chronic exposure to oxidative stress. *Cancer Res* 58: 3986–3992, 1998.
 83. Hurd TR, Costa NJ, Dahm CC, Beer SM, Brown SE, Filipovska A, and Murphy MP. Glutathionylation of mitochondrial proteins. *Antioxid Redox Signal* 7: 999–1010, 2005.
 84. Hurt EM, Thomas SB, Peng B, and Farrar WL. Molecular consequences of SOD2 expression in epigenetically silenced pancreatic carcinoma cell lines. *Br J Cancer* 97: 1116–1123, 2007.
 85. Hussain SP, Amstad P, He P, Robles A, Lupold S, Kaneko I, Ichimiya M, Sengupta S, Mechanic L, Okamura S, Hofseth LJ, Moake M, Nagashima M, Forrester KS, and Harris CC. p53-induced up-regulation of MnSOD and GPx but not catalase increases oxidative stress and apoptosis. *Cancer Res* 64: 2350–2356, 2004.
 86. Iglesias-Bartolome R, Patel V, Cotrim A, Leelahavanichkul K, Molinolo AA, Mitchell JB, and Gutkind JS. mTOR Inhibition Prevents Epithelial Stem Cell Senescence and Protects from Radiation-Induced Mucositis. *Cell Stem Cell* 11: 401–414, 2012.
 87. Kang BP, Urbonas A, Baddoo A, Baskin S, Malhotra A, and Meggs LG. IGF-1 inhibits the mitochondrial apoptosis program in mesangial cells exposed to high glucose. *Am J Physiol Renal Physiol* 285: F1013–F1024, 2003.
 88. Kennedy G, Spence VA, McLaren M, Hill A, Underwood C, and Belch JJ. Oxidative stress levels are raised in chronic fatigue syndrome and are associated with clinical symptoms. *Free Radic Biol Med* 39: 584–589, 2005.
 89. Kim A, Murphy MP, and Oberley TD. Mitochondrial redox state regulates transcription of the nuclear-encoded mitochondrial protein manganese superoxide dismutase: a proposed adaptive response to mitochondrial redox imbalance. *Free Radic Biol Med* 38: 644–654, 2005.
 90. Kim A, Zhong W, and Oberley TD. Reversible modulation of cell cycle kinetics in NIH/3T3 mouse fibroblasts by inducible overexpression of mitochondrial manganese superoxide dismutase. *Antioxid Redox Signal* 6: 489–500, 2004.
 91. Kinningham KK and St Clair DK. Overexpression of manganese superoxide dismutase selectively modulates the activity of Jun-associated transcription factors in fibrosarcoma cells. *Cancer Res* 57: 5265–5271, 1997.
 92. Koehler CM, Beverly KN, and Leverich EP. Redox pathways of the mitochondrion. *Antioxid Redox Signal* 8: 813–822, 2006.
 93. Kumar S, Bharti A, Mishra NC, Raina D, Kharbanda S, Saxena S, and Kufe D. Targeting of the c-Abl tyrosine kinase to mitochondria in the necrotic cell death response to oxidative stress. *J Biol Chem* 276: 17281–17285, 2001.
 94. Li JJ, Oberley LW, St Clair DK, Ridnour LA, and Oberley TD. Phenotypic changes induced in human breast cancer cells by overexpression of manganese-containing superoxide dismutase. *Oncogene* 10: 1989–2000, 1995.
 95. This reference has been deleted.
 96. Li Z, Shi K, Guan L, Cao T, Jiang Q, Yang Y, and Xu C. ROS leads to MnSOD upregulation through ERK2 translocation and p53 activation in selenite-induced apoptosis of NB4 cells. *FEBS Lett* 584: 2291–2297, 2010.
 97. Lin T, Chao C, Saito S, Mazur SJ, Murphy ME, Appella E, and Xu Y. p53 induces differentiation of mouse embryonic stem cells by suppressing Nanog expression. *Nat Cell Biol* 7: 165–171, 2005.
 98. Liu SZ, Cai L, and Sun SQ. Induction of a cytogenetic adaptive response by exposure of rabbits to very low dose-rate gamma-radiation. *Int J Radiat Biol* 62: 187–190, 1992.
 99. Lombard DB, Alt FW, Cheng HL, Bunkenborg J, Streeper RS, Mostoslavsky R, Kim J, Yancopoulos G, Valenzuela D, Murphy A, Yang Y, Chen Y, Hirschey MD, Bronson RT, Haigis M, Guarente LP, Farese RV, Jr., Weissman S, Verdin E, and Schwer B. Mammalian Sir2 homolog SIRT3

- regulates global mitochondrial lysine acetylation. *Mol Cell Biol* 27: 8807–8814, 2007.
100. Luk E, Carroll M, Baker M, and Culotta VC. Manganese activation of superoxide dismutase 2 in *Saccharomyces cerevisiae* requires MTM1, a member of the mitochondrial carrier family. *Proc Natl Acad Sci U S A* 100: 10353–10357, 2003.
 101. Luk EE and Culotta VC. Manganese superoxide dismutase in *Saccharomyces cerevisiae* acquires its metal co-factor through a pathway involving the Nramp metal transporter, Smf2p. *J Biol Chem* 276: 47556–47562, 2001.
 102. MacMillan-Crow LA, Crow JP, and Thompson JA. Peroxynitrite-mediated inactivation of manganese superoxide dismutase involves nitration and oxidation of critical tyrosine residues. *Biochemistry* 37: 1613–1622, 1998.
 103. Macmillan-Crow LA and Cruthirds DL. Invited review: manganese superoxide dismutase in disease. *Free Radic Res* 34: 325–336, 2001.
 104. Maiorino M, Scapin M, Ursini F, Biasolo M, Bosello V, and Flohe L. Distinct promoters determine alternative transcription of gpx-4 into phospholipid-hydroperoxide glutathione peroxidase variants. *J Biol Chem* 278: 34286–34290, 2003.
 105. Mancuso M, Davidzon G, Kurlan RM, Tawil R, Bonilla E, Di Mauro S, and Powers JM. Hereditary ferritinopathy: a novel mutation, its cellular pathology, and pathogenetic insights. *J Neuropathol Exp Neurol* 64: 280–294, 2005.
 106. Martin J, Mahlke K, and Pfanner N. Role of an energized inner membrane in mitochondrial protein import. Delta psi drives the movement of presequences. *J Biol Chem* 266: 18051–18057, 1991.
 107. Martin RC, Li Y, Liu Q, Jensen NS, Barker DF, Doll MA, and Hein DW. Manganese superoxide dismutase V16A single-nucleotide polymorphism in the mitochondrial targeting sequence is associated with reduced enzymatic activity in cryopreserved human hepatocytes. *DNA Cell Biol* 28: 3–7, 2009.
 108. Matsuda Y, Higashiyama S, Kijima Y, Suzuki K, Kawano K, Akiyama M, Kawata S, Tarui S, Deutsch HF, and Taniguchi N. Human liver manganese superoxide dismutase. Purification and crystallization, subunit association and sulfhydryl reactivity. *Eur J Biochem* 194: 713–720, 1990.
 109. May T and Soll J. 14-3-3 proteins form a guidance complex with chloroplast precursor proteins in plants. *Plant Cell* 12: 53–64, 2000.
 110. Mettler FA. *Medical Effects of Ionizing Radiation*. Philadelphia: Saunders, 1985, p. 304.
 111. Mihara M, Erster S, Zaika A, Petrenko O, Chittenden T, Pancoska P, and Moll UM. p53 has a direct apoptogenic role at the mitochondria. *Mol Cell* 11: 577–590, 2003.
 112. Morgan DO. Cyclin-dependent kinases: engines, clocks, and microprocessors. *Annu Rev Cell Dev Biol* 13: 261–291, 1997.
 113. Morgan DO. *The Cell Cycle: Principles of Control*. London: New Science Press, Ltd. 2007.
 114. Murakami K, Tanase S, Morino Y, and Mori M. Pre-sequence binding factor-dependent and -independent import of proteins into mitochondria. *J Biol Chem* 267: 13119–13122, 1992.
 115. Murley JS, Baker KL, Miller RC, Darga TE, Weichselbaum RR, and Grdina DJ. SOD2-mediated adaptive responses induced by low-dose ionizing radiation via TNF signaling and amifostine. *Free Radic Biol Med* 51: 1918–1925, 2011.
 116. Murley JS, Kataoka Y, Baker KL, Diamond AM, Morgan WF, and Grdina DJ. Manganese superoxide dismutase (SOD2)-mediated delayed radioprotection induced by the free thiol form of amifostine and tumor necrosis factor alpha. *Radiat Res* 167: 465–474, 2007.
 117. Murley JS, Kataoka Y, Cao D, Li JJ, Oberley LW, and Grdina DJ. Delayed radioprotection by NF-kappaB-mediated induction of Sod2 (MnSOD) in SA-NH tumor cells after exposure to clinically used thiol-containing drugs. *Radiat Res* 162: 536–546, 2004.
 118. Murley JS, Kataoka Y, Weydert CJ, Oberley LW, and Grdina DJ. Delayed radioprotection by nuclear transcription factor kappaB-mediated induction of manganese superoxide dismutase in human microvascular endothelial cells after exposure to the free radical scavenger WR1065. *Free Radic Biol Med* 40: 1004–1016, 2006.
 119. Nabel GJ and Verma IM. Proposed NF-kappa B/I kappa B family nomenclature. *Genes Dev* 7: 2063, 1993.
 120. Nantajit D, Fan M, Duru N, Wen Y, Reed JC, and Li JJ. Cyclin B1/Cdk1 phosphorylation of mitochondrial p53 induces anti-apoptotic response. *PLoS One* 5: e12341, 2010.
 121. Nulton-Persson AC and Szweda LI. Modulation of mitochondrial function by hydrogen peroxide. *J Biol Chem* 276: 23357–23361, 2001.
 122. Oberley LW. Mechanism of the tumor suppressive effect of MnSOD overexpression. *Biomed Pharmacother* 59: 143–148, 2005.
 123. Oberley LW, Lindgren LA, Baker SA, and Stevens RH. Superoxide ion as the cause of the oxygen effect. *Radiat Res* 68: 320–328, 1976.
 124. Oberley LW, St Clair DK, Autor AP, and Oberley TD. Increase in manganese superoxide dismutase activity in the mouse heart after X-irradiation. *Arch Biochem Biophys* 254: 69–80, 1987.
 125. Oberley TD, Verwiebe E, Zhong W, Kang SW, and Rhee SG. Localization of the thioredoxin system in normal rat kidney. *Free Radic Biol Med* 30: 412–424, 2001.
 126. Oda K, Arakawa H, Tanaka T, Matsuda K, Tanikawa C, Mori T, Nishimori H, Tamai K, Tokino T, Nakamura Y, and Taya Y. p53AIP1, a potential mediator of p53-dependent apoptosis, and its regulation by Ser-46-phosphorylated p53. *Cell* 102: 849–862, 2000.
 127. Ohman T, Parish G, and Jackson RM. Hypoxic modulation of manganese superoxide dismutase promoter activity and gene expression in lung epithelial cells. *Am J Respir Cell Mol Biol* 21: 119–127, 1999.
 128. Okado-Matsumoto A and Fridovich I. Subcellular distribution of superoxide dismutases (SOD) in rat liver: Cu,Zn-SOD in mitochondria. *J Biol Chem* 276: 38388–38393, 2001.
 129. Ozden O, Park SH, Kim HS, Jiang H, Coleman MC, Spitz DR, and Gius D. Acetylation of MnSOD directs enzymatic activity responding to cellular nutrient status or oxidative stress. *Aging (Albany, NY)* 3: 102–107, 2011.
 130. Ozeki M, Tamae D, Hou DX, Wang T, Lebon T, Spitz DR, and Li JJ. Response of cyclin B1 to ionizing radiation: regulation by NF-kappaB and mitochondrial antioxidant enzyme MnSOD. *Anticancer Res* 24: 2657–2663, 2004.
 131. Pagliarini DJ and Dixon JE. Mitochondrial modulation: reversible phosphorylation takes center stage? *Trends Biochem Sci* 31: 26–34, 2006.
 132. Pagliarini DJ, Wiley SE, Kimple ME, Dixon JR, Kelly P, Worby CA, Casey PJ, and Dixon JE. Involvement of a mitochondrial phosphatase in the regulation of ATP production and insulin secretion in pancreatic beta cells. *Mol Cell* 19: 197–207, 2005.
 133. Pan S and Berk BC. Glutathiolation regulates tumor necrosis factor-alpha-induced caspase-3 cleavage and

- apoptosis: key role for glutaredoxin in the death pathway. *Circ Res* 100: 213–219, 2007.
134. Pandey BN, Gordon DM, De Toledo SM, Pain D, and Azzam EI. Normal human fibroblasts exposed to high- or low-dose ionizing radiation: differential effects on mitochondrial protein import and membrane potential. *Antioxid Redox Signal* 8: 1253–1261, 2006.
135. Pfanner N and Geissler A. Versatility of the mitochondrial protein import machinery. *Nat Rev Mol Cell Biol* 2: 339–349, 2001.
136. Qiu X, Brown K, Hirsche MD, Verdin E, and Chen D. Calorie restriction reduces oxidative stress by SIRT3-mediated SOD2 activation. *Cell Metab* 12: 662–667, 2010.
137. Quijano C, Hernandez-Saavedra D, Castro L, McCord JM, Freeman BA, and Radi R. Reaction of peroxynitrite with Mn-superoxide dismutase. Role of the metal center in decomposition kinetics and nitration. *J Biol Chem* 276: 11631–11638, 2001.
138. Radi R, Cassina A, Hodara R, Quijano C, and Castro L. Peroxynitrite reactions and formation in mitochondria. *Free Radic Biol Med* 33: 1451–1464, 2002.
139. Ramond A, Godin-Ribuot D, Ribaut C, Totoson P, Koritchneva I, Cachot S, Levy P, and Joyeux-Faure M. Oxidative stress mediates cardiac infarction aggravation induced by intermittent hypoxia. *Fundam Clin Pharmacol* 2011 [Epub ahead of print]; DOI: 10.1111/j.1472-8206.2011.01015.x.
140. Ravindranath SD and Fridovich I. Isolation and characterization of a manganese-containing superoxide dismutase from yeast. *J Biol Chem* 250: 6107–6112, 1975.
141. Redondo-Horcajo M, Romero N, Martinez-Acedo P, Martinez-Ruiz A, Quijano C, Lourenco CF, Movilla N, Enriquez JA, Rodriguez-Pascual F, Rial E, Radi R, Vazquez J, and Lamas S. Cyclosporine A-induced nitration of tyrosine 34 MnSOD in endothelial cells: role of mitochondrial superoxide. *Cardiovasc Res* 87: 356–365, 2010.
142. Rhee SG, Kang SW, Chang TS, Jeong W, and Kim K. Peroxiredoxin, a novel family of peroxidases. *IUBMB Life* 52: 35–41, 2001.
143. Ridnour LA, Sim JE, Choi J, Dickinson DA, Forman HJ, Ahmad IM, Coleman MC, Hunt CR, Goswami PC, and Spitz DR. Nitric oxide-induced resistance to hydrogen peroxide stress is a glutamate cysteine ligase activity-dependent process. *Free Radic Biol Med* 38: 1361–1371, 2005.
144. Robinson KM and Lemire BD. Covalent attachment of FAD to the yeast succinate dehydrogenase flavoprotein requires import into mitochondria, presequence removal, and folding. *J Biol Chem* 271: 4055–4060, 1996.
145. Roise D, Horvath SJ, Tomich JM, Richards JH, and Schatz G. A chemically synthesized pre-sequence of an imported mitochondrial protein can form an amphiphilic helix and perturb natural and artificial phospholipid bilayers. *EMBO J* 5: 1327–1334, 1986.
146. Roise D and Schatz G. Mitochondrial presequences. *J Biol Chem* 263: 4509–4511, 1988.
147. Rosini P, De Chiara G, Bonini P, Lucibello M, Marcocci ME, Garaci E, Cozzolino F, and Torcia M. Nerve growth factor-dependent survival of CESS B cell line is mediated by increased expression and decreased degradation of MAPK phosphatase 1. *J Biol Chem* 279: 14016–14023, 2004.
148. Salvi M, Brunati AM, and Toninello A. Tyrosine phosphorylation in mitochondria: a new frontier in mitochondrial signaling. *Free Radic Biol Med* 38: 1267–1277, 2005.
149. Salvi M, Stringaro A, Brunati AM, Agostinelli E, Arancia G, Clari G, and Toninello A. Tyrosine phosphatase activity in mitochondria: presence of Shp-2 phosphatase in mitochondria. *Cell Mol Life Sci* 61: 2393–2404, 2004.
150. Sarsour EH, Kalen AL, Xiao Z, Veenstra TD, Chaudhuri L, Venkataraman S, Reigan P, Buettner GR, and Goswami PC. Manganese Superoxide Dismutase Regulates a Metabolic Switch during the Mammalian Cell Cycle. *Cancer Res* 72: 3807–3816, 2012.
151. Schatz G. Mitochondria: beyond oxidative phosphorylation. *Biochim Biophys Acta* 1271: 123–126, 1995.
152. Schatz G. The protein import system of mitochondria. *J Biol Chem* 271: 31763–31766, 1996.
153. Scheffler IE. Mitochondria. New York: John Wiley & Sons, Inc., 2002, pp. 345–351.
154. Schwoch G, Trinczek B, and Bode C. Localization of catalytic and regulatory subunits of cyclic AMP-dependent protein kinases in mitochondria from various rat tissues. *Biochem J* 270: 181–188, 1990.
155. Seo MS, Kang SW, Kim K, Baines IC, Lee TH, and Rhee SG. Identification of a new type of mammalian peroxiredoxin that forms an intramolecular disulfide as a reaction intermediate. *J Biol Chem* 275: 20346–20354, 2000.
156. Shadley JD and Wolff S. Very low doses of X-rays can cause human lymphocytes to become less susceptible to ionizing radiation. *Mutagenesis* 2: 95–96, 1987.
157. Shi T, Wang F, Stieren E, and Tong Q. SIRT3, a mitochondrial sirtuin deacetylase, regulates mitochondrial function and thermogenesis in brown adipocytes. *J Biol Chem* 280: 13560–13567, 2005.
158. Shimoda-Matsubayashi S, Matsumine H, Kobayashi T, Nakagawa-Hattori Y, Shimizu Y, and Mizuno Y. Structural dimorphism in the mitochondrial targeting sequence in the human manganese superoxide dismutase gene. A predictive evidence for conformational change to influence mitochondrial transport and a study of allelic association in Parkinson's disease. *Biochem Biophys Res Commun* 226: 561–565, 1996.
159. Slot JW, Geuze HJ, Freeman BA, and Crapo JD. Intracellular localization of the copper-zinc and manganese superoxide dismutases in rat liver parenchymal cells. *Lab Invest* 55: 363–371, 1986.
160. Spitz DR and Sullivan SJ. The generation of stable oxidative stress-resistant phenotypes in Chinese hamster fibroblasts chronically exposed to hydrogen peroxide or hyperoxia. *Methods Mol Biol* 610: 183–199, 2010.
161. St Clair DK, Wan XS, Oberley TD, Muse KE, and St Clair WH. Suppression of radiation-induced neoplastic transformation by overexpression of mitochondrial superoxide dismutase. *Mol Carcinog* 6: 238–242, 1992.
162. Stojanovic S, Stanic D, Nikolic M, Raicevic S, Spasic M, and Niketic V. Manganese superoxide dismutase (MnSOD) catalyzes NO-dependent tyrosine residue nitration. *J Serb Chem Soc* 70: 601–608, 2005.
163. Stroupe ME, DiDonato M, and Tainer JA. Manganese superoxide dismutase. In: *Handbook of Metalloproteins*, edited by Messerschmidt A, Huber R, Poulos T, and Wieghardt K. Chichester, UK: John Wiley & Sons, Ltd., 2006.
164. Sui D and Wilson JE. Structural determinants for the intracellular localization of the isozymes of mammalian hexokinase: intracellular localization of fusion constructs incorporating structural elements from the hexokinase isozymes and the green fluorescent protein. *Arch Biochem Biophys* 345: 111–125, 1997.
165. Sutton A, Imbert A, Igoudjil A, Descatoire V, Cazanave S, Pessayre D, and Degoul F. The manganese superoxide

- dismutase Ala16Val dimorphism modulates both mitochondrial import and mRNA stability. *Pharmacogenet Genomics* 15: 311–319, 2005.
166. Sutton A, Khoury H, Prip-Buus C, Capanec C, Pessayre D, and Degoul F. The Ala16Val genetic dimorphism modulates the import of human manganese superoxide dismutase into rat liver mitochondria. *Pharmacogenetics* 13: 145–157, 2003.
 167. Takada Y, Hachiya M, Park SH, Osawa Y, Ozawa T, and Akashi M. Role of reactive oxygen species in cells overexpressing manganese superoxide dismutase: mechanism for induction of radioresistance. *Mol Cancer Res* 1: 137–146, 2002.
 168. Tao R, Coleman MC, Pennington JD, Ozden O, Park SH, Jiang H, Kim HS, Flynn CR, Hill S, Hayes McDonald W, Olivier AK, Spitz DR, and Gius D. Sirt3-mediated deacetylation of evolutionarily conserved lysine 122 regulates MnSOD activity in response to stress. *Mol Cell* 40: 893–904, 2010.
 169. Tsan MF. Superoxide dismutase and pulmonary oxygen toxicity: lessons from transgenic and knockout mice (Review). *Int J Mol Med* 7: 13–19, 2001.
 170. Ubersax JA, Woodbury EL, Quang PN, Paraz M, Blethrow JD, Shah K, Shokat KM, and Morgan DO. Targets of the cyclin-dependent kinase Cdk1. *Nature* 425: 859–864, 2003.
 171. Van Remmen H, Ikeno Y, Hamilton M, Pahlavani M, Wolf N, Thorpe SR, Alderson NL, Baynes JW, Epstein CJ, Huang TT, Nelson J, Strong R, and Richardson A. Life-long reduction in MnSOD activity results in increased DNA damage and higher incidence of cancer but does not accelerate aging. *Physiol Genomics* 16: 29–37, 2003.
 172. Verma IM, Stevenson JK, Schwarz EM, Van Antwerp D, and Miyamoto S. Rel/NF-kappa B/I kappa B family: intimate tales of association and dissociation. *Genes Dev* 9: 2723–2735, 1995.
 173. Voisin S, Watson DC, Tessier L, Ding W, Foote S, Bhatia S, Kelly JF, and Young NM. The cytoplasmic phosphoproteome of the Gram-negative bacterium *Campylobacter jejuni*: evidence for modification by unidentified protein kinases. *Proteomics* 7: 4338–4348, 2007.
 174. Wan XS, Devalaraja MN, and St Clair DK. Molecular structure and organization of the human manganese superoxide dismutase gene. *DNA Cell Biol* 13: 1127–1136, 1994.
 175. Wang F, Liu J, Robbins D, Morris K, Sit A, Liu YY, and Zhao Y. Mutant p53 exhibits trivial effects on mitochondrial functions which can be reactivated by ellipticine in lymphoma cells. *Apoptosis* 16: 301–310, 2011.
 176. Wang HG, Rapp UR, and Reed JC. Bcl-2 targets the protein kinase Raf-1 to mitochondria. *Cell* 87: 629–638, 1996.
 177. Wang X, Martindale JL, Liu Y, and Holbrook NJ. The cellular response to oxidative stress: influences of mitogen-activated protein kinase signalling pathways on cell survival. *Biochem J* 333 (Pt 2): 291–300, 1998.
 178. Waris G and Ahsan H. Reactive oxygen species: role in the development of cancer and various chronic conditions. *J Carcinog* 5: 14, 2006.
 179. Weisiger RA and Fridovich I. Mitochondrial superoxide dismutase. Site of synthesis and intramitochondrial localization. *J Biol Chem* 248: 4793–4796, 1973.
 180. Wiedemann N, Frazier AE, and Pfanner N. The protein import machinery of mitochondria. *J Biol Chem* 279: 14473–14476, 2004.
 181. Wilcox KC, Zhou L, Jordon JK, Huang Y, Yu Y, Redler RL, Chen X, Caplow M, and Dokholyan NV. Modifications of superoxide dismutase (SOD1) in human erythrocytes: a possible role in amyotrophic lateral sclerosis. *J Biol Chem* 284: 13940–13947, 2009.
 182. Williams MD, Van Remmen H, Conrad CC, Huang TT, Epstein CJ, and Richardson A. Increased oxidative damage is correlated to altered mitochondrial function in heterozygous manganese superoxide dismutase knockout mice. *J Biol Chem* 273: 28510–28515, 1998.
 183. Wispe JR, Clark JC, Burhans MS, Kropp KE, Korfhagen TR, and Whitsett JA. Synthesis and processing of the precursor for human manganese-superoxide dismutase. *Biochim Biophys Acta* 994: 30–36, 1989.
 184. Wong GH. Protective roles of cytokines against radiation: induction of mitochondrial MnSOD. *Biochim Biophys Acta* 1271: 205–209, 1995.
 185. Wright G and Reichenbecher V. The effects of superoxide and the peripheral benzodiazepine receptor ligands on the mitochondrial processing of manganese-dependent superoxide dismutase. *Exp Cell Res* 246: 443–450, 1999.
 186. Wright G, Reichenbecher V, Green T, Wright GL, and Wang S. Paraquat inhibits the processing of human manganese-dependent superoxide dismutase by SF-9 insect cell mitochondria. *Exp Cell Res* 234: 78–84, 1997.
 187. Wright G, Terada K, Yano M, Sergeev I, and Mori M. Oxidative stress inhibits the mitochondrial import of preproteins and leads to their degradation. *Exp Cell Res* 263: 107–117, 2001.
 188. Wynn RM, Kato M, Machius M, Chuang JL, Li J, Tomchick DR, and Chuang DT. Molecular mechanism for regulation of the human mitochondrial branched-chain alpha-ketoacid dehydrogenase complex by phosphorylation. *Structure* 12: 2185–2196, 2004.
 189. Xu S, Ying J, Jiang B, Guo W, Adachi T, Sharov V, Lazar H, Menzoian J, Knyushko TV, Bigelow D, Schoneich C, and Cohen RA. Detection of sequence-specific tyrosine nitration of manganese SOD and SERCA in cardiovascular disease and aging. *Am J Physiol Heart Circ Physiol* 290: H2220–H2227, 2006.
 190. Xu Y, Kiningham KK, Devalaraja MN, Yeh CC, Majima H, Kasarskis EJ, and St Clair DK. An intronic NF-kappaB element is essential for induction of the human manganese superoxide dismutase gene by tumor necrosis factor-alpha and interleukin-1beta. *DNA Cell Biol* 18: 709–722, 1999.
 191. Yamakura F, Kobayashi K, Furukawa S, and Suzuki Y. *In vitro* preparation of iron-substituted human manganese superoxide dismutase: possible toxic properties for mitochondria. *Free Radic Biol Med* 43: 423–430, 2007.
 192. Yang M, Cobine PA, Molik S, Naranuntarat A, Lill R, Winge DR, and Culotta VC. The effects of mitochondrial iron homeostasis on cofactor specificity of superoxide dismutase 2. *EMBO J* 25: 1775–1783, 2006.
 193. Young JC, Hoogenraad NJ, and Hartl FU. Molecular chaperones Hsp90 and Hsp70 deliver preproteins to the mitochondrial import receptor Tom70. *Cell* 112: 41–50, 2003.
 194. Zacchi P, Gostissa M, Uchida T, Salvagno C, Avolio F, Volinia S, Ronai Z, Blandino G, Schneider C, and Del Sal G. The prolyl isomerase Pin1 reveals a mechanism to control p53 functions after genotoxic insults. *Nature* 419: 853–857, 2002.
 195. Zhao Y, Chaiswing L, Velez JM, Batinic-Haberle I, Colburn NH, Oberley TD, and St Clair DK. p53 translocation to mitochondria precedes its nuclear translocation and targets

mitochondrial oxidative defense protein-manganese superoxide dismutase. *Cancer Res* 65: 3745–3750, 2005.

196. Zorov DB, Filburn CR, Klotz LO, Zweier JL, and Sollott SJ. Reactive oxygen species (ROS)-induced ROS release: a new phenomenon accompanying induction of the mitochondrial permeability transition in cardiac myocytes. *J Exp Med* 192: 1001–1014, 2000.

Address correspondence to:

Dr. Jian Jian Li

Department of Radiation Oncology

University of California Davis

2700 Stockton Blvd.

Sacramento, CA 95817

E-mail: jijli@ucdavis.edu

Date of first submission to ARS Central, March 13, 2013; date of acceptance, April 14, 2013.

Abbreviations Used

Bax = Bcl-2-associated X protein
 Bcl = B-cell lymphoma
 Bcl-XL = B-cell lymphoma-extra large
 Cdk = cyclin-dependent kinase
 Cu/ZnSOD = copper/zinc superoxide dismutase
 Cys = cysteine
 ECSOD = extracellular superoxide dismutase
 ERK = extracellular regulated kinase
 FeSOD = iron superoxide dismutase
 G2 = growth two
 GADD = growth arrest and DNA damage
 H₂O₂ = hydrogen peroxide

HK = hexokinase
 HNSCC = head and neck squamous cell carcinoma
 IR = ionizing radiation
 JAK = Janus kinase
 JNK = Jun N-terminal kinase
 LDIR = low dose ionizing radiation
 Leu = leucine
 M = mitosis
 MAPK = mitogen-activated protein kinase
 MET = hepatocellular growth factor receptor
 MnSOD = manganese superoxide dismutase
 MPI = mitochondrial protein influx
 Mtmp1 = monomethylamine permease
 mTOR = mammalian target of rapamycin
 Myc = v-myc myelocytomatosis viral oncogene homolog
 NF- κ B = nuclear factor-kappaB
 NO = nitric oxide
 OH = hydroxyl
 p53AIP = p53-regulated apoptosis inducing protein
 PDK = pyruvate dehydrogenase kinase
 Pro = proline
 Prx = peroxiredoxin
 RNS = reactive nitrogen species
 ROS = reactive oxygen species
 Ser/Thr = serine/threonine
 SIRT = sirtuin
 STAT = signal transducer and activator of transcription
 TIM = translocase of the inner membrane
 TNF α = tumor necrosis factor alpha
 TOM = translocase of the outer membrane
 Tyr = tyrosine