UC Berkeley

UC Berkeley Previously Published Works

Title

Assessing the aerodynamic diameters of taxon-specific fungal bioaerosols by quantitative PCR and next-generation DNA sequencing

Permalink

<https://escholarship.org/uc/item/7sn332j0>

Authors

Yamamoto, N
Nazaroff, WW
Peccia, J

Publication Date

2014-12-01

DOI

10.1016/j.jaerosci.2014.08.007

Peer reviewed

Assessing the aerodynamic diameters of
taxon-specific fungal bioaerosols by quantita-
tive PCR and next-generation DNA sequencing

Naomichi Yamamoto, William W Nazaroff,
Jordan Peccia

PII: S0021-8502(14)00132-3
DOI: <http://dx.doi.org/10.1016/j.jaerosci.2014.08.007>
Reference: AS4818

To appear in: *Journal of Aerosol Science*

Received date: 14 June 2014
Revised date: 25 August 2014
Accepted date: 25 August 2014

Cite this article as: Naomichi Yamamoto, William W Nazaroff, Jordan Peccia, Assessing the aerodynamic diameters of taxon-specific fungal bioaerosols by quantitative PCR and next-generation DNA sequencing, *Journal of Aerosol Science*, <http://dx.doi.org/10.1016/j.jaerosci.2014.08.007>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting galley proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

**Assessing the aerodynamic diameters of taxon-specific fungal
bioaerosols by quantitative PCR and next-generation DNA
sequencing**

Submitted to:

Journal of Aerosol Science

Naomichi Yamamoto ^a, William W Nazaroff ^b, Jordan Peccia ^{c, *}

- a. Department of Environmental Health, Graduate School of Public Health, Seoul National University, 1 Gwanak-ro, Gwanak-gu, Seoul 151-742, Korea
- b. Department of Civil and Environmental Engineering, University of California, Berkeley, CA 94720-1710, USA
- c. Department of Chemical and Environmental Engineering, Yale University, New Haven, CT 06520, USA

* Corresponding author.

Tel: +1 203 432 4385

Fax: +1 203 432 4387

E-mail address: jordan.peccia@yale.edu (J. Peccia)

Abstract

Aerodynamic diameter is an important determinant of the physical processes that act upon airborne fungi. Processes include gravitational settling, respiratory deposition, penetration into buildings, resuspension from surfaces into air, and long-range transport. This study combined next-generation DNA sequencing (NGS) with quantitative PCR (qPCR) to evaluate diverse, taxon-specific, fungal aerodynamic diameters from bioaerosol samples. The accuracy of the method was demonstrated by comparing geometric mean aerodynamic diameters of selected taxa produced by the NGS-based method to the diameters produced by taxon-specific qPCR ($r = 0.996$). Geometric means (d_g) and geometric standard deviations (σ_g) of aerodynamic diameters were characterized for more than 50 fungal taxa, spanning 55 genera, 9 classes, and 2 phyla. The results reported in this study demonstrate the robust nature of this method, provide novel insights into aerodynamic properties of diverse airborne fungal species, and potentially enable a better accounting of taxon-specific fungal fate and exposure both in indoor air and in the atmosphere.

Keywords: Fungi, Internal transcribed spacer (ITS), Pyrosequencing, Aerodynamic diameter, Quantitative PCR (qPCR), Bioaerosols

1 **1. Introduction**

2 Aerodynamic diameter exerts significant influence over the important physical
3 processes that act upon airborne biological particles. These processes include but are not
4 limited to ice nucleation, gravitational settling, respiratory deposition, penetration into
5 buildings, resuspension into air, and long-range transport (Ariya *et al.*, 2009; Nazaroff,
6 2004; Prospero *et al.*, 2005; Riley *et al.*, 2002; Thatcher & Layton, 1995; Yeh *et al.*,
7 1996). A particle's aerodynamic diameter is the diameter of a sphere with a density of 1
8 g/cm³ that has the same settling velocity as the particle of interest. Traditionally, the
9 aerodynamic diameters of airborne fungi have been characterized by means of
10 enumerating culturable organisms that were captured on multistage cascade impactors
11 (Madelin & Johnson, 1992; McCartney *et al.*, 1993; Reponen, 1995) or by
12 time-of-flight (TOF)-based aerodynamic particle sizing (Han *et al.*, 2011; Madelin &
13 Johnson, 1992; Reponen *et al.*, 1996). These methods have important limitations for
14 characterizing fungal aerodynamic diameter in indoor air or in the outdoor atmosphere.
15 Impactor data derived from culturing is restricted to species that can be readily
16 identified, and cannot account for nonviable fungal spores and fragments, or fungi that
17 are not culturable under the given conditions (Peccia & Hernandez, 2006; Reponen,
18 1995). TOF-based methods do not allow for the identification of fungal taxa or the
19 discrimination of fungal particles from nonfungal particles. Thus, their use is limited to
20 fungal aerosol studies that start from pure cultures.

21 Fungi are remarkably diverse with an estimated 1.5 million species (Bass &
22 Richards, 2011; Blackwell, 2011; Hawksworth, 2001). Recent airborne fungal diversity
23 analyses based on fungal barcoding via next-generation DNA sequencing (NGS) has
24 resulted in the identification of hundreds to thousands of different fungal taxa in outdoor

aerosol samples (Adams *et al.*, 2013; Dannemiller *et al.*, 2014; Yamamoto *et al.*, 2012). When coupled with size-resolved sampling, the NGS/fungal barcoding technique may enable approaches to determine the aerodynamic diameters of bioaerosols associated with specific fungal taxa. However, NGS-based data provides taxon results as a relative abundance per size bin. Such data must be transformed into absolute concentrations to assess the distribution of aerodynamic diameters for the different phyla, genera, or species that are identified.

The goal of this study was to demonstrate an approach using NGS/fungal barcoding and universal qPCR data to characterize taxon-specific fungal aerodynamic diameters in environmental aerosol samples. Relative abundance results were coupled with universal qPCR data from particle-size-fractionated outdoor air samples collected at a site in the northeastern United States. Absolute concentrations of each fungal taxon in each size fraction were obtained by multiplying universal fungal qPCR results by the NGS-based relative abundance data (Dannemiller *et al.*, 2014). Geometric mean aerodynamic diameters and geometric standard deviations for bioaerosols associated with specific taxa were computed. These values were then compared to geometric mean aerodynamic diameters determined by taxon-specific qPCR. This study provides, for the first time, a robust approach for quantifying geometric means and geometric standard deviations of taxon-specific fungal aerodynamic diameters for aerosol particles based on NGS and qPCR data, broadening the potential to determine the aerodynamic diameters for fungal taxa in air.

2. Experimental

2.1. Seasonal fungal study

The study utilized data and samples from a seasonal sampling campaign conducted in New Haven, Connecticut, USA (41°18'29"N 72°55'43"W) in 2009-2011 (Yamamoto *et al.*, 2012). Airborne fungi were collected at a continuous sampling rate of 28.3 l min⁻¹ for ~ 4-week sampling periods in each of four seasons. We sampled onto glass fiber filter substrates using an eight-stage non-viable Andersen sampler with aerodynamic diameter (d_a) cutoffs of 0.4, 0.7, 1.1, 2.1, 3.3, 4.7, 5.8, and 9.0 μ m. Glass fiber substrates were used to minimize particle bounce (Hu, 1971). The data used here from the previous study included the taxonomic libraries for the different impactor stages for the four different seasons and taxon-specific qPCR data for *Alternaria alternata*, *Aspergillus fumigatus*, *Cladosporium cladosporioides*, *Epicoccum nigrum*, and *Penicillium chrysogenum*. To estimate the absolute concentration for specific taxa, total fungal qPCR was carried out in the present study using DNA extracts of the previously collected seasonal samples. Below, brief descriptions of the methods performed in the prior study are provided along with a more in-depth description of the universal qPCR methods and estimation of aerodynamic diameters for specific taxa.

2.2. Universal fungal qPCR

Total fungal concentrations were measured by universal fungal qPCR with primers FF2 (5'-GGTTCTATTTTGGTTGGTTTCTA-3') and FR1 (5'-CTCTCAATCTGTCAATCCTTATT-3') (Zhou *et al.*, 2000). Reaction mixtures totaling 50 μ L included template DNA (2 μ L), 1 $\times$ SYBR Green Master Mix (FastStart Universal SYBR Green Master (ROX); Roche Applied Science) and 0.3 μ M of each

primer. A real-time PCR system (ABI 7500 Fast Real-time PCR System; Applied Biosystems) was used with these thermal conditions: 50 °C for 2 min, 95 °C for 15 min of initial denaturation and 45 cycles of 95 °C for 15 s of dissociation and 60 °C for 1 min of annealing and extension. Threshold cycles were determined by the ABI 7500 auto function. The results were calibrated against an *Aspergillus fumigatus* standard. Thus, reported universal fungal qPCR results are based on *A. fumigatus* spore equivalents (SE) per m³ as previously described (Dannemiller *et al.*, 2014; Hospodsky *et al.*, 2010; Lang-Yona *et al.*, 2012).

2.3. Species-specific fungal qPCR

Fungal aerodynamic diameters assessed by the NGS-based method were compared with those characterized by taxon-specific qPCR. Fungal genera of *Alternaria*, *Aspergillus*, *Cladosporium*, *Epicoccum*, and *Penicillium* were selected for this comparison. *Aspergillus* and *Penicillium* produce small unicellular asexual spores, whereas *Alternaria* and *Epicoccum* produce large multicellular dictyospores. *Cladosporium* spp. produces both unicellular and multicellular spores. These taxa were selected to cover a wide range of fungal spore size. Concentrations of *Alternaria alternata*, *Aspergillus fumigatus*, *Cladosporium cladosporioides*, *Epicoccum nigrum*, and *Penicillium chrysogenum* were quantified by the reported species-specific qPCR assays (Haugland *et al.*, 2004; Meklin *et al.*, 2004), and their geometric means (d_g) and geometric standard deviations of aerodynamic diameters, calculated according to the method described below, were 10.6 µm (1.55), 3.88 µm (1.25), 4.62 µm (1.46), 11.0 µm (1.61) and 3.89 µm (1.66), respectively. These species-specific d_g values were compared with the corresponding genus-specific aerodynamic diameters as determined by NGS.

The genus-specific values were used for NGS since the numbers of sequences were small for *A. alternata*, *A. fumigatus*, and *P. chrysogenum* (< 15 sequences) if the results were analyzed at the species level.

2.4. Next-generation DNA sequencing and taxonomic assignment

The well-established methods used for the NGS/barcoding methodology are described elsewhere (Yamamoto *et al.*, 2012). Briefly, the internal transcribed spacer (ITS) region of fungal DNA is targeted as a barcode marker for identification (Schoch *et al.*, 2012). The extracted DNA was amplified for the ITS sequences circumscribed by universal fungal primers ITS1F (5'-CTTGGTCATTTAGAGGAAGTAA-3') and ITS4 (5'-TCCTCCGCTTATTGATATGC-3') (Larena *et al.*, 1999; Manter & Vivanco, 2007). The purified amplicons were normalized and sequenced on the 454 GS FLX Titanium Platform (454 Life Sciences) at the Yale University Center for Genome Analysis. The samples with $d_a < 2.1 \mu\text{m}$ were not included for the sequencing analyses owing to no or weak PCR amplification. Overall, 15,326 ITS sequences were produced. Two phyla, 19 classes, and 558 genera were identified, and the relative abundances were calculated at the phylum, class, and genus ranks (Yamamoto *et al.*, 2012). Taxa with at least 40 ITS sequences detected were included for subsequent analyses of particle size distributions.

2.5. Evaluating aerodynamic diameters

A schematic diagram of a method to assess fungal aerodynamic diameters from the bioaerosol samples is shown in Fig. 1. NGS provides relative abundance fractions of each taxon within a sample (Fig. 1A), whereas the universal fungal qPCR measures absolute concentrations of total fungi (Fig. 1B). Thus, absolute

concentrations of each taxon (N_{taxon}) are determined by the following equation:

$$N_{\text{taxon}} = F_{\text{taxon}} \times N_{\text{total}} \quad (1)$$

where F_{taxon} is the relative abundance fraction of a taxon obtained by NGS and N_{total} is the absolute concentration (SE m^{-3}) of total fungi measured by the universal fungal qPCR. The absolute concentrations of each taxon were calculated for each particle size range (Fig. 1C), and geometric means (d_g) and geometric standard deviation (σ_g) of aerodynamic diameters were calculated for each taxon (Fig. 1D).

As reported in the results section, the taxon-specific d_g values ranged widely from $< 2.1 \mu\text{m}$ to $11.8 \mu\text{m}$. For some taxa, particle size distributions were left- or right-truncated, which potentially results in inaccurate d_g and σ_g estimations if traditional forward-calculation methods are used (Yamamoto *et al.*, 2012). Here we used a best-fit d_g and σ_g method which has the advantage of not requiring knowledge of the upper size limit on the largest bin, and does not utilize any *a priori* assumptions about the distribution within each size bin. In this method, we executed a search procedure to solve for the best-fit values of d_g and σ_g under the assumption that the sampled size distributions of taxon-specific fungal DNA were lognormal. The procedure aimed to minimize the residual between predicted and measured particle size distributions (Fig. 2) by using the least-squares method. To avoid the risk of settling on a local minimum, all possible combinations of d_g (0.4 to $15 \mu\text{m}$ in steps of $0.01 \mu\text{m}$) and σ_g (1.01 to 3 in steps of 0.01) were tested. The computations were executed using Excel Visual Basic for Applications ver. 7.0, which can be downloaded at <https://sourceforge.net/projects/gmcalculator/>.

For most taxa, we used the seasonally averaged particle size distributions as the input for computing values of d_g and σ_g . For some taxa, the largest relative

1 abundances were in the winter; however, the absolute fungal DNA concentrations were
 2 the lowest in the winter. In some of these cases, we found a high residual when
 3 computing the geometric parameters for annual averaged data. Thus, in cases in which
 4 the residual between the predicted and measured particle size distributions was greater
 5 than 20% for the annual average data and the greatest relative abundance was observed
 6 in the winter, only the winter data were used to compute d_g and σ_g . The distributional
 7 parameters for these taxa were excluded from the final reporting if the residuals also
 8 were greater than 20% using the winter data. Some taxa showed a residual greater than
 9 20%, and the greatest relative abundance was not found in the winter. However, none
 10 of these taxa was found to show a residual smaller than 20% even when selecting only
 11 the most abundant season.

13 **2.6. Count median diameter (CMD) estimation**

14 The DNA-based methods provide particle size distributions of airborne fungi
 15 based on the third moment of particle size distribution (i.e., corresponding to mass
 16 distributions). To allow for comparisons with culture-based literature data that reflect
 17 the first moment of particle size distributions (i.e., corresponding to count distributions),
 18 we also estimated count median diameters (CMD) for geometric means of fungal
 19 aerodynamic diameters characterized by the NGS-based method and for species-specific
 20 qPCR. To estimate CMD, the Hatch-Choate equation was used (Hinds, 1999):

$$21 \quad \text{CMD} = d_g \exp(-3 \ln^2 \sigma_g) \quad (2)$$

22

23

24

3. Results

3.1. Comparing the NGS-based method and species-specific qPCR

Fig. 3 shows the relationship of fungal aerodynamic diameters characterized by the NGS-based method and by species-specific qPCR. Only the winter data were used for *Aspergillus* and *Penicillium* since large residuals between the measured and predicted particle size distributions (>20%) were observed for the annual averages. The d_g values by the species-specific qPCR were 10.6, 3.88, 4.62, 11.0 and 3.89 μm for *Alternaria*, *Aspergillus*, *Cladosporium*, *Epicoccum*, and *Penicillium*, respectively. The NGS-based method produced the respective d_g values of 10.6, 5.16, 5.52, 11.8, and 5.07 μm . Although different taxonomic ranks were used for the comparison, a strong correlation was observed between the two methods (Pearson's $r = 0.996$, $p < 0.001$). The σ_g values were also consistent between these two methods (Table 1), substantiating the capability of the NGS-based method to determine, with reasonable accuracy, taxon-specific fungal aerodynamic diameters.

3.2. Comparing aerodynamic diameters of different methods

Table 1 shows the values of d_g and σ_g of selected fungal genera obtained by different methods. The d_g values measured by NGS and species-specific qPCR were consistently greater than those reported from culture-based methods. To estimate the d_g values based on the moment of count distributions, count median diameters (CMD) were estimated for aerodynamic diameters obtained by the NGS- and species-specific qPCR-based methods by Eq. (2). The resulting CMD values were correlated with, but smaller than the original d_g values (Pearson's $r = 0.772$, $p < 0.001$) and more similar to the d_g values characterized by the growth-based method (Table 1).

3.3. Taxon-specific particle size distributions

Two phyla, 19 classes, and 558 genera were identified (Yamamoto *et al.*, 2012), of which 2 phyla, 9 classes, and 55 genera were found to have (a) more than 40 ITS sequences, and (b) residual errors between predicted and measured size distributions smaller than 20%. At the phylum rank, geometric means and standard deviations of aerodynamic diameters were 7.68 μm (1.73) for *Ascomycota* and 4.44 μm (1.43) for *Basidiomycota* (Fig. 1D) when data for all seasons were pooled. The season-specific values for *Ascomycota* were 8.28 μm (1.86), 5.75 μm (1.73), 8.59 μm (1.45), and 7.55 μm (1.69) in the spring, summer, fall, and winter, respectively, whereas the respective values for *Basidiomycota* were 4.85 μm (1.38), 3.61 μm (1.37), 5.06 μm (1.41), and 4.42 μm (1.24).

Fig. 4 illustrates particle size-resolved taxonomic compositions of airborne fungi. The class *Agaricomycetes* of the phylum *Basidiomycota* represented the largest fraction (45%), whereas the class *Dothideomycetes* of the phylum *Ascomycota* accounted for the second largest fraction (35%). The dominance of these two classes was consistent throughout each season. Their geometric mean diameters varied seasonally with the smallest values observed in the summer. The season-specific d_g and σ_g values for *Agaricomycetes* were 4.78 μm (1.34), 3.59 μm (1.37), 4.88 μm (1.37), and 4.38 μm (1.23) in the spring, summer, fall, and winter, respectively, whereas the respective values for *Dothideomycetes* were 8.03 μm (2.06), 6.08 μm (1.78), 9.22 μm (1.43), and 10.4 μm (1.84).

Tables 2 and 3 list d_g , σ_g , and the estimated CMD of aerodynamic diameters of the 9 most abundant classes and 55 most abundant genera, respectively. The d_g values

varied substantially across the fungal taxa, ranging from $< 2.1 \mu\text{m}$ for *Antrodia* to $11.8 \mu\text{m}$ for *Epicoccum*. Large d_g values were observed for the *Ascomycota* genera of *Alternaria* ($10.6 \mu\text{m}$), *Epicoccum* ($11.8 \mu\text{m}$), *Leptosphaerulina* ($9.55 \mu\text{m}$), and *Monilinia* ($9.68 \mu\text{m}$). The d_g values smaller than $3.3 \mu\text{m}$ were observed for the *Basidiomycota* genera of *Antrodia* ($< 2.1 \mu\text{m}$), *Phlebia* ($3.04 \mu\text{m}$) *Sistotrema* ($2.26 \mu\text{m}$) and *Wallemia* ($3.01 \mu\text{m}$).

4. Discussion

Although the aerodynamic diameter is an important microbial feature that influences fungal aerosol source emissions, deposition rates and environmental fate, and human exposure, this parameter has not been well characterized in prior studies for relevant fungal taxa in an environmental setting. The present study demonstrates an approach for determining the aerodynamic diameters of a broad diversity of fungal taxa (2 phyla, 9 classes, and 55 genera) suspended in bioaerosols. The method, applied here to atmospheric samples, is also appropriate for indoor air studies.

The reported NGS-based method produced fungal particle-size distributions that were highly consistent with those characterized by taxon-specific qPCR (Fig. 3). Given that taxon-specific qPCR has established accuracy as a reference method (Haugland *et al.*, 2004; Meklin *et al.*, 2004), the results indicate the capability of the NGS-based method to accurately assess fungal aerodynamic diameters.

Application of this approach using size-resolved relative abundance data from prior sampling campaigns revealed a diversity of aerodynamic diameters among taxa. Taxon-dependent d_g values are expected owing to fungal physiology, physical spore sizes, method of spore release, and environmental fate and transport that are unique to

1 each fungal group. At the phylum rank, average geometric means of aerodynamic
2 diameters were 7.68 μm for *Ascomycota* and 4.44 μm for *Basidiomycota* (Fig. 1). The
3 dominant classes of the *Ascomycota* and *Basidiomycota* phyla were *Dothideomycetes*
4 and *Agaricomycetes*, respectively (Fig. 4). Large proportions of *Dothideomycetes* and
5 *Agaricomycetes* in outdoor air were also reported in a previous sequencing-based study
6 (Fröhlich-Nowoisky *et al.*, 2009). The four most abundant genera of the class
7 *Dothideomycetes* and their d_g values were *Leptosphaerulina* (9.55 μm), *Epicoccum*
8 (11.8 μm), *Cladosporium* (5.52 μm), and *Alternaria* (10.6 μm) (Table 3). These fungi
9 produce large multicellular dictyospores with reported spore sizes of 24–36 $\times$ 10–14,
10 15–25, 3–11 $\times$ 2–5, 18–83 $\times$ 7–18 μm , respectively (Cole & Samson, 1984; Mitkowski &
11 Browning, 2004). Meanwhile, the four most abundant genera of the class
12 *Agaricomycetes* and their d_g values were *Peniophora* (4.41 μm), *Exidia* (5.70 μm),
13 *Stereum* (4.13 μm), and *Trametes* (3.34 μm) (Table 2). Their reported spore sizes are
14 6.5–8 $\times$ 3–3.5, 2–4 $\times$ 1, 2.5 $\times$ 6–7, and 6.6–9.2 $\times$ 2.4–3 μm , respectively (Burt, 1920; Ingold,
15 1995; Li & Cui, 2010; Whelden, 1936).

16 As described in the results section, seasonal variations in the aerodynamic
17 diameters have been observed, with the smallest values found in the summer. The
18 observed tendency was consistent with results using species-specific qPCR in our
19 previous study (Yamamoto *et al.*, 2012). Though the mechanisms are unknown, the
20 smaller spores might be produced during summer owing to higher temperature
21 (Phillips, 1982). The finding may be clinically relevant as changes in spore sizes can
22 affect inhalability and respirability of allergenic and pathogenic airborne fungal spores
23 (Reponen, 1995).

24 Comparing the aerodynamic diameters of fungal bioaerosols with spore sizes

1 derived in a prior culture-based environmental study reveals that the geometric mean
 2 aerodynamic diameters estimated from the qPCR and NGS methods are greater (Table
 3 1). A possible cause of this finding is the agglomeration of fungal spores in the
 4 atmosphere or in indoor air (Heikkila *et al.*, 1988; Lacey, 1991). Using culture-based
 5 methods, a single aerosol dispersal unit that contains multiple spores may develop into
 6 only one colony and thus result in one identifiable fungal count per dispersal unit. NGS
 7 and qPCR methods, in contrast, quantify the multiple spores in the above dispersal unit,
 8 thus assigning a value greater than one fungal count to this larger, aggregate particle.
 9 Consequently, the moments of particle size distributions are different, and molecular
 10 techniques may produce a larger d_g value than culture-based techniques. In addition,
 11 comparing the qPCR- and NGS-based d_g values or the culture based d_g values from
 12 environmental studies (Table 1) with pure culture TOF-based data (*Aspergillus*
 13 *fumigatus* $d_g = 2.15 \mu\text{m}$, *Penicillium chrysogenum* $d_g = 2.8 \mu\text{m}$, and *Cladosporium*
 14 *cladosporioides* $d_g = 1.8 \mu\text{m}$) (Madelin & Johnson, 1992; Reponen *et al.*, 1996)
 15 reinforces a finding that many fungal spores sampled from the atmosphere or indoor
 16 air are not in the form of single isolated spores. The difference may also be attributable
 17 to different environmental samples. In each environment, the sizes of airborne fungal
 18 spores or DNA may vary by attachment to other abiotic particulate matter (Lighthart,
 19 1997; Yamaguchi *et al.*, 2012). Finally, the existence of nonculturable fungal fragments
 20 that produce an NGS/qPCR signal might further differentially impact the observed
 21 particle size statistics of fungal aerosols (Peccia & Hernandez, 2006).

22 Limitations in the present method for determining taxon-specific size
 23 characteristics center upon assumptions made in converting relative abundance values
 24 to absolute concentrations. The accuracy of this taxon-specific concentration

estimation has previously been described (Dannemiller *et al.*, 2014). While the NGS/qPCR methods for determining taxon-specific concentration are strongly correlated with taxon-specific qPCR results from the same sample (Pearson's $r = 0.996$ $p < 0.001$), systematic biases have been observed in the conversions of a quantity of a reference fungal strain into absolute quantities of different fungal taxa (Dannemiller *et al.*, 2014). Using a single strain of *A. fumigatus* for universal qPCR calibration, while necessary, was expected to cause biases owing to taxon-dependent variations in the numbers of rDNA copies per fungal genome (Maleszka & Clarkwalker, 1993; Rooney & Ward, 2005).

Notably, unlike taxon-specific concentration calculation, determination of particle size metrics such as d_g and σ_g by the NGS/qPCR method appear to be less sensitive to a bias associated with copy number variation of fungal ITS. Indeed, a strong correlation between the NGS/qPCR method and taxon-specific qPCR for quantifying the d_g values has been observed (Fig. 2). This outcome is expected due to the nature of the d_g and σ_g calculations, which use the relative proportion of absolute concentrations of each fungal taxon quantified across each particle size bin. Though strain-dependent variation has been reported in the numbers of rDNA copies in *A. fumigatus* (Herrera *et al.*, 2009), our findings indicate that particle size is not a major metric systematically influenced by variations in the numbers of ITS copies.

Additional future benefits can be anticipated from ongoing improvements in fungal databases, NGS identification accuracy, and a better understanding of the number of ITS genes in the genomes of a diversity of fungal species (Yamamoto & Bibby, 2014; Yamamoto *et al.*, 2014).

5. Conclusions

Aerodynamic diameter is of central importance for determining physical processes that influence airborne particles, with implications for fungal ecology, human exposure, plant pathogen transport, and climate. Traditionally, aerodynamic diameters of airborne fungi have been studied based on TOF- or growth-based techniques. Taxon-specific qPCR, culturing, or the use of TOF techniques significantly limits the extent of taxon-specific aerodynamic diameters that can be determined in environmental aerosols. The present study used a new approach combining NGS and universal fungal qPCR to evaluate aerodynamic diameters of multiple fungal taxa. The method characterized particle-size distributions of 55 specific fungal genera with a single set of NGS and universal fungal qPCR data, providing important information about the aerodynamic properties of diverse airborne fungal DNA. This new method expands the scope of fungal genera bioaerosol sizes. By avoiding the culture-based underestimation of fungal spore in aggregate, the method also results in larger mean particle sizes than previously reported by culture-based analysis.

Acknowledgements

Primary funding for this project was provided by the Alfred P Sloan foundation. N.Y. is supported by the BK 21 PLUS from the Ministry of Education and National Research Foundation of Korea.

References

- Adams, R.I., Miletto, M., Taylor, J.W., & Bruns, T.D. (2013). Dispersal in microbes: Fungi in indoor air are dominated by outdoor air and show dispersal limitation at short distances. *ISME Journal*, **7**, 1262-1273.
- Ariya, P.A., Sun, J., Eltouny, N.A., Hudson, E.D., Hayes, C.T., & Kos, G. (2009). Physical and chemical characterization of bioaerosols - Implications for nucleation processes. *International Reviews in Physical Chemistry*, **28**, 1-32.
- Bass, D., & Richards, T.A. (2011). Three reasons to re-evaluate fungal diversity 'on Earth and in the ocean'. *Fungal Biology Reviews*, **25**, 159-164.
- Blackwell, M. (2011). The Fungi: 1, 2, 3 ... 5.1 million species? *American Journal of Botany*, **98**, 426-438.
- Burt, E.A. (1920). The *Thelephoraceae* of North America. XII. *Stereum*. *Annals of the Missouri Botanical Garden*, **7**, 81-249.
- Cole, G.T., & Samson, R.A. (1984). The conidia. In Al-Doory, Y. & Domson, J.F. (Eds) *Mould allergy* (pp. 66-104). Philadelphia: Lea & Fibiger.
- Dannemiller, K., Lang-Yona, N., Yamamoto, N., Rudich, Y., & Peccia, J. (2014). Combining real-time PCR and next-generation DNA sequencing to provide quantitative comparisons of fungal aerosol populations. *Atmospheric Environment*, **84**, 113-121.
- Fröhlich-Nowoisky, J., Pickersgill, D.A., Després, V.R., & Pöschl, U. (2009). High diversity of fungi in air particulate matter. *Proceedings of the National Academy of Sciences of the United States of America*, **106**, 12814-12819.
- Han, T., Nazarenko, Y., Liroy, P.J., & Mainelis, G. (2011). Collection efficiencies of an electrostatic sampler with superhydrophobic surface for fungal bioaerosols. *Indoor*

- 1 *Air*, **21**, 110-120.
- 2 Haugland, R.A., Varma, M., Wymer, L.J., & Vesper, S.J. (2004). Quantitative PCR
- 3 analysis of selected *Aspergillus*, *Penicillium* and *Paecilomyces* species. *Systematic*
- 4 *and Applied Microbiology*, **27**, 198-210.
- 5 Hawksworth, D.L. (2001). The magnitude of fungal diversity: The 1.5 million species
- 6 estimate revisited. *Mycological Research*, **105**, 1422-1432.
- 7 Heikkila, P., Kotimaa, M., Tuomi, T., Salmi, T., & Louhelainen, K. (1988).
- 8 Identification and counting of fungal spores by scanning electron-microscope.
- 9 *Annals of Occupational Hygiene*, **32**, 241-248.
- 10 Herrera, M.L., Vallor, A.C., Gelfond, J.A., Patterson, T.F., & Wickes, B.L. (2009).
- 11 Strain-dependent variation in 18S ribosomal DNA copy numbers in *Aspergillus*
- 12 *fumigatus*. *Journal of Clinical Microbiology*, **47**, 1325-1332.
- 13 Hinds, W.C. (1999) *Aerosol technology: Properties, behavior, and measurement of*
- 14 *airborne particles* (2nd ed.). New York: John Wiley & Sons, Inc. (Chapter 4).
- 15 Hospodsky, D., Yamamoto, N., & Peccia, J. (2010). Accuracy, precision, and method
- 16 detection limits of quantitative PCR for airborne bacteria and fungi. *Applied and*
- 17 *Environmental Microbiology*, **76**, 7004-7012.
- 18 Hu, J. N.-H., (1971). Communication. An improved impactor for aerosol studies
- 19 —modified Andersen sampler. *Environmental Science & Technology*, **5**, 251-253.
- 20 Ingold, C.T. (1995). Types of reproductive cell in *Exidia recisa* and *Sirobasidium*
- 21 *intermediae*. *Mycological Research*, **99**, 1187-1190.
- 22 Lacey, J. (1991). Aggregation of spores and its effect on aerodynamic behavior. *Grana*,
- 23 **30**, 437-445.
- 24 Lang-Yona, N., Dannemiller, K., Yamamoto, N., Burshtein, N., Peccia, J., Yarden, O.,

- 1 & Rudich, Y. (2012). Annual distribution of allergenic fungal spores in atmospheric
2 particulate matter in the Eastern Mediterranean; a comparative study between
3 ergosterol and quantitative PCR analysis. *Atmospheric Chemistry and Physics*, **12**,
4 2681-2690.
- 5 Larena, I., Salazar, O., Gonzalez, V., Julian, M.C., & Rubio, V. (1999). Design of a
6 primer for ribosomal DNA internal transcribed spacer with enhanced specificity for
7 ascomycetes. *Journal of Biotechnology*, **75**, 187-194.
- 8 Li, H.-J., & Cui, B.-K. (2010). A new *Trametes* species from Southwest China.
9 *Mycotaxon*, **113**, 263-267.
- 10 Lighthart, B. (1997). The ecology of bacteria in the alfresco atmosphere. *FEMS*
11 *Microbiology Ecology*, **23**, 263-274.
- 12 Madelin, T.M., & Johnson, H.E. (1992). Fungal and actinomycete spore aerosols
13 measured at different humidities with an aerodynamic particle sizer. *Journal of*
14 *Applied Bacteriology*, **72**, 400-409.
- 15 Maleszka, R., & Clarkwalker, G.D. (1993). Yeasts have a 4-fold variation in ribosomal
16 DNA copy number. *Yeast*, **9**, 53-58.
- 17 Manter, D.K., & Vivanco, J.M. (2007). Use of the ITS primers, ITS1F and ITS4, to
18 characterize fungal abundance and diversity in mixed-template samples by qPCR
19 and length heterogeneity analysis. *Journal of Microbiological Methods*, **71**, 7-14.
- 20 McCartney, H.A., Schmechel, D., & Lacey, M.E. (1993). Aerodynamic diameter of
21 conidia of *Alternaria* species. *Plant Pathology*, **42**, 280-286.
- 22 Meklin, T., Haugland, R.A., Reponen, T., Varma, M., Lummus, Z., Bernstein, D.,
23 Wymer, L.J., & Vesper, S.J. (2004). Quantitative PCR analysis of house dust can
24 reveal abnormal mold conditions. *Journal of Environmental Monitoring*, **6**, 615-620.

- 1 Mitkowski, N.A., & Browning, M. (2004). *Leptosphaerulina australis* associated with
2 intensively managed stands of *Poa annua* and *Agrostis palustris*. *Canadian Journal*
3 *of Plant Pathology*, **26**, 193-198.
- 4 Nazaroff, W.W. (2004). Indoor particle dynamics. *Indoor Air*, **14**, 175-183.
- 5 Peccia, J., & Hernandez, M. (2006). Incorporating polymerase chain reaction-based
6 identification, population characterization, and quantification of microorganisms
7 into aerosol science: A review. *Atmospheric Environment*, **40**, 3941-3961.
- 8 Phillips, D.J. (1982). Changes in conidia of *Monilinia fructicola* in response to
9 incubation temperature. *Phytopathology*, **72**, 1281-1283.
- 10 Prospero, J.M., Blades, E., Mathison, G., & Naidu, R. (2005). Interhemispheric
11 transport of viable fungi and bacteria from Africa to the Caribbean with soil dust.
12 *Aerobiologia*, **21**, 1-19.
- 13 Reponen, T. (1995). Aerodynamic diameters and respiratory deposition estimates of
14 viable fungal particles in mold problem dwellings. *Aerosol Science and Technology*,
15 **22**, 11-23.
- 16 Reponen, T., Willeke, K., Ulevicius, V., Reponen, A., & Grinshpun, S.A. (1996). Effect
17 of relative humidity on the aerodynamic diameter and respiratory deposition of
18 fungal spores. *Atmospheric Environment*, **30**, 3967-3974.
- 19 Riley, W.J., McKone, T.E., Lai, A.C.K., & Nazaroff, W.W. (2002). Indoor particulate
20 matter of outdoor origin: Importance of size-dependent removal mechanisms.
21 *Environmental Science & Technology*, **36**, 200-207.
- 22 Rooney, A.P., & Ward, T.J. (2005). Evolution of a large ribosomal RNA multigene
23 family in filamentous fungi: Birth and death of a concerted evolution paradigm.
24 *Proceedings of the National Academy of Sciences of the United States of America*,

- 1 **102**, 5084-5089.
- 2 Schoch, C.L., Seifert, K.A., Huhndorf, S., Robert, V., Spouge, J.L., Levesque, C.A., &
- 3 Chen, W. (2012). Nuclear ribosomal internal transcribed spacer (ITS) region as a
- 4 universal DNA barcode marker for *Fungi*. *Proceedings of the National Academy of*
- 5 *Sciences of the United States of America*, **109**, 6241-6246.
- 6 Thatcher, T.L., & Layton, D.W. (1995). Deposition, resuspension, and penetration of
- 7 particles within a residence. *Atmospheric Environment*, **29**, 1487-1497.
- 8 Whelden, R.M. (1936). A comparative study of basidia and cystidia in *Peniophora*
- 9 *livida*. *American Journal of Botany*, **23**, 539-545.
- 10 Yamaguchi, N., Ichijo, T., Sakotani, A., Baba, T., & Nasu, M. (2012). Global
- 11 dispersion of bacterial cells on Asian dust. *Scientific Reports*, **2**, 525.
- 12 Yamamoto, N., & Bibby, K. (2014). Clustering of fungal community internal
- 13 transcribed spacer (ITS) sequence data obscures taxonomic diversity. *Environmental*
- 14 *Microbiology*, doi:10.1111/1462-2920.12390.
- 15 Yamamoto, N., Bibby, K., Qian, J., Hospodsky, D., Rismani-Yazdi, H., Nazaroff, W.W.,
- 16 & Peccia, J. (2012). Particle-size distributions and seasonal diversity of allergenic
- 17 and pathogenic fungi in outdoor air. *ISME Journal*, **6**, 1801-1811.
- 18 Yamamoto, N., Dannemiller, K., Bibby, K., & Peccia, J. (2014). Identification accuracy
- 19 and diversity reproducibility associated with ITS-based fungal taxonomic library
- 20 preparation. *Environmental Microbiology*, doi:10.1111/1462-2920.12338.
- 21 Yeh, H.C., Cuddihy, R.G., Phalen, R.F., & Chang, I.Y. (1996). Comparisons of
- 22 calculated respiratory tract deposition of particles based on the proposed NCRP
- 23 model and the new ICRP66 model. *Aerosol Science and Technology*, **25**, 134-140.
- 24 Zhou, G., Whong, W.Z., Ong, T., & Chen, B. (2000). Development of a fungus-specific

PCR assay for detecting low-level fungi in an indoor environment. *Molecular and Cellular Probes*, **14**, 339-348.

Table 1

Geometric means (d_g , μm) and geometric standard deviations (σ_g) of aerodynamic diameters of airborne fungi obtained by different methods. ^a

Genera	NGS			Species-specific qPCR			Culture		Microscopy
	d_g	σ_g	Estimated CMD	d_g	σ_g	Estimated CMD	d_g	σ_g	
<i>Aspergillus</i>	5.16	1.20	4.67	3.88	1.25	3.34	1.8 ^b	1.5 ^b	2–3.5 ^c
<i>Penicillium</i>	5.07	1.20	4.59	3.89	1.66	1.80	2.3 ^b	1.2 ^b	2.5 ^d
<i>Cladosporium</i>	5.52	1.58	2.95	4.62	1.46	3.01	2.8 ^b	1.1 ^b	3–11×2–5 ^c
<i>Alternaria</i>	10.6	1.45	7.00	10.6	1.55	5.97	n.a.	n.a.	18–83×7–18 ^c
<i>Epicoccum</i>	11.8	1.54	6.76	11.0	1.61	5.59	n.a.	n.a.	15–25 ^c

^a Count median diameters (CMD, μm) are estimated for the NGS- and species-specific qPCR-based methods by Eq. (2). The microscopy-based sizes of fungal spores are also listed. For the qPCR and microscopy data, the values specific for the species *Aspergillus fumigatus*, *Penicillium chrysogenum*, *Cladosporium cladosporioides*, *Alternaria alternata*, and *Epicoccum nigrum* are shown. For the NGS and culture data, the genus-specific values are listed. Abbreviations: n.a., not available in the literature.

^b Reponen (1995).

^c Cole & Samson (1984).

^d Madelin & Johnson (1992).

Table 2

Geometric means (d_g) and geometric standard deviation (σ_g) of fungal aerodynamic diameters evaluated by the NGS-based method, showing the seasonally averaged data for the 9 most abundant classes.

Phylum	Class	d_g (μm)	σ_g	Residual ^a (%)	Estimated CMD ^b (μm)
<i>Ascomycota</i>	<i>Dothideomycetes</i>	7.94	1.80	4.7	2.82
	<i>Eurotiomycetes</i>	5.38	1.49	13.8	3.34
	<i>Lecanoromycetes</i>	9.71	1.62	2.4	4.83
	<i>Leotiomycetes</i>	7.55	1.51	4.3	4.54
	<i>Sordariomycetes</i>	6.16	1.60	5.1	3.18
<i>Basidiomycota</i>	<i>Agaricomycetes</i>	4.35	1.40	4.0	3.10
	<i>Tremellomycetes</i>	6.98	1.69	4.1	3.06
	<i>Microbotryomycetes</i>	6.19	1.55	2.0	3.48
	<i>Wallemiomycetes</i>	3.12	1.42	5.9	2.16

^a Residual is the square root of the sum of the squared deviations between the predicted and measured concentrations of each size bin of the particle size distribution.

^b Count median diameters (CMD) are estimated by Eq. (2).

1 **Table 3**

2 Geometric mean (d_g) and geometric standard deviation (σ_g) of fungal aerodynamic
 3 diameters evaluated by the NGS-based method, showing the seasonally averaged data
 4 for the 55 most abundant genera.

Phylum	Class	Genus	d_g (μm)	σ_g	Residual ^a (%)	Estimated CMD ^b (μm)
<i>Ascomycota</i>	<i>Dothideomycetes</i>	<i>Alternaria</i>	10.6	1.45	1.1	7.00
		<i>Cladosporium</i>	5.52	1.58	2.2	2.95
		<i>Epicoccum</i>	11.8	1.54	2.2	6.76
		<i>Eudarluka</i>	4.99	1.71	19.0	2.10
		<i>Leptosphaerulina</i>	9.55	1.58	7.2	5.10
		<i>Lophiostoma</i>	6.00	1.44	11.3	4.03
		<i>Mycosphaerella</i>	5.26	1.47	15.5	3.37
		<i>Phaeothecoidea</i>	8.12	1.52	7.3	4.80
		<i>Ramularia</i>	4.89	1.28	10.2	4.07
		<i>Teratosphaeria</i>	9.02	1.67	2.4	4.10
	<i>Eurotiomycetes</i>	<i>Aspergillus</i> ^c	5.16	1.20	1.1	4.67
		<i>Eurotium</i>	4.36	1.14	3.4	4.14
		<i>Penicillium</i> ^c	5.07	1.20	4.8	4.59
		<i>Phaeococcomyces</i>	8.26	1.37	4.6	6.14
	<i>Leotiomycetes</i>	<i>Allantophomopsis</i>	8.16	1.48	12.6	5.15
		<i>Botryotinia</i>	8.16	1.38	0.8	5.98
		<i>Monilinia</i>	9.68	1.45	3.4	6.40
		<i>Trimmatostroma</i>	7.34	1.56	18.6	4.06
	<i>Sordariomycetes</i>	<i>Biscogniauxia</i> ^c	5.02	1.12	0.2	4.83
		<i>Colletotrichum</i>	3.55	1.32	9.0	2.82
		<i>Daldinia</i>	7.68	1.35	7.1	5.86
		<i>Diatrype</i>	7.70	1.48	2.6	4.86
		<i>Eutypa</i>	7.05	1.14	4.0	6.70
		<i>Eutypella</i>	6.23	1.41	3.3	4.37
	<i>Incertae sedis</i>	<i>Microcyclospora</i>	9.77	1.49	1.3	6.06
<i>Basidiomycota</i>	<i>Agaricomycetes</i>	<i>Antrodia</i>	<2.1	n.d.	2.4	n.d.
		<i>Cerrena</i>	4.35	1.21	5.0	3.90
		<i>Coprinellus</i>	5.38	1.21	7.4	4.82
		<i>Cortinarius</i>	5.61	1.22	6.3	4.98
		<i>Cylindrobasidium</i>	5.95	1.33	3.6	4.66
		<i>Daedaleopsis</i>	3.74	1.16	3.4	3.50
		<i>Exidia</i>	5.70	1.19	13.1	5.21
		<i>Ganoderma</i>	5.13	1.15	2.2	4.84
		<i>Hymenochaete</i>	3.66	1.22	3.0	3.25
		<i>Hyphoderma</i>	4.86	1.16	11.3	4.55
		<i>Lycoperdon</i>	4.38	1.16	3.6	4.10
		<i>Mycena</i>	6.75	1.28	6.9	5.62
		<i>Oxyporus</i>	4.00	1.16	6.8	3.74
		<i>Panellus</i>	3.52	1.34	16.9	2.72
		<i>Peniophora</i>	4.41	1.27	7.7	3.72
		<i>Perenniporia</i>	4.17	1.25	7.1	3.59
		<i>Phlebia</i>	3.04	1.46	5.1	1.98
		<i>Piptoporus</i>	2.50	2.07	15.5	0.51
		<i>Pleurotus</i>	5.21	1.33	9.6	4.08

	<i>Polyporus</i>	5.02	1.42	12.3	3.47
	<i>Schizophyllum</i>	3.96	1.39	2.8	2.86
	<i>Sebacina</i>	5.44	1.17	10.2	5.05
	<i>Sistotrema</i>	2.26	2.24	5.2	0.32
	<i>Stereum</i>	4.13	1.32	9.7	3.28
	<i>Trametes</i>	3.34	1.46	3.8	2.17
	<i>Trichaptum</i>	3.85	1.34	3.4	2.98
<i>Tremellomycetes</i>	<i>Cryptococcus</i>	7.71	1.65	4.2	3.63
	<i>Dioszegia</i>	5.55	1.44	10.9	3.72
	<i>Hannaella</i>	9.47	1.75	5.1	3.70
<i>Wallemiomycetes</i>	<i>Wallemia</i>	3.01	1.46	7.0	1.96

^a Residual is the square root of the sum of the squared deviations between the predicted and measured concentrations of each size bin of the particle size distribution.

^b Count median diameters (CMD) are estimated by Eq. (2).

^c The values are based on the winter data alone.

Abbreviation: n.d., not determined owing to the left-truncated particle size distributions.

Figure Legends

Fig. 1. Geometric means (d_g) and geometric standard deviations (σ_g) of taxon-specific fungal aerodynamic diameters as assessed by the next generation DNA sequencing (NGS)-based method. Relative abundances of each fungal taxon were determined by NGS (A), whereas particle size distributions of total airborne fungi were obtained by universal fungal qPCR (B). The NGS-derived relative abundances of each fungal taxon were multiplied by the qPCR-derived particle size distributions of airborne total fungi (C) and the resulting taxon-specific particle size distributions were produced to compute d_g and σ_g for each fungal taxon (D). The values shown are for the phyla *Ascomycota* and *Basidiomycota*, with data pooled for all four seasons.

Fig. 2. Comparison between the measured and predicted particle size distributions. The values of geometric means (d_g) and geometric standard deviation (σ_g) of aerodynamic

diameters were obtained by an analysis that minimized the residual between the predicted and measured particle size distributions. The particle size distributions shown are for *Cladosporium*.

Fig. 3. Comparison of geometric mean of aerodynamic diameters of selected fungal taxa characterized by the species-specific qPCR and the NGS-based method. Each datapoint is an average for four seasons, with the exception of *Aspergillus* and *Penicillium* where only winter data were use due to the large residuals (> 20%) observed for annual averages.

Fig. 4. Particle size-resolved taxonomic compositions of airborne fungi in New Haven, Connecticut, USA in 2009-2011.

Highlights (max 85 characters for each of 3 to 5 bullet points)

- Study combined NGS and qPCR to evaluate the aerodynamic diameters of fungal taxa
- Aerodynamic diameters determined for >50 fungal genera in atmospheric bioaerosols
- Good agreement obtained between diameters estimated by NGS and taxon-specific qPCR