

UC Davis

UC Davis Previously Published Works

Title

Plant responses to heterogeneous salinity: agronomic relevance and research priorities

Permalink

<https://escholarship.org/uc/item/7t32v7cc>

Journal

Annals of Botany, 129(5)

ISSN

0305-7364

Authors

Valenzuela, Francisco Jose
Reineke, Daniela
Leventini, Dante
et al.

Publication Date

2022-04-13

DOI

10.1093/aob/mcac022

Peer reviewed

INVITED REVIEW

Plant responses to heterogeneous salinity: agronomic relevance and research priorities

Francisco Jose Valenzuela¹, Daniela Reineke¹, Dante Leventini¹, Christopher Cody Lee Chen¹, Edward G. Barrett-Lennard^{2,3}, Timothy D. Colmer^{4,5}, Ian C. Dodd⁶, Sergey Shabala^{7,8,*}, Patrick Brown^{1,*} and Nadia Bazihizina^{9,*}

¹Department of Plant Sciences, University of California, Davis, CA, USA, ²Land Management Group, Agriculture Discipline, College of Science, Health, Engineering and Education, Murdoch University, WA, Australia, ³Department of Primary Industries and Regional Development, South Perth, WA, Australia, ⁴UWA School of Agriculture and Environment, The University of Western Australia, Crawley, WA, Australia, ⁵Institute of Agriculture, The University of Western Australia, Crawley, WA, Australia, ⁶The Lancaster Environment Centre, Lancaster University, Lancaster, UK, ⁷Tasmanian Institute of Agriculture, University of Tasmania, Hobart, TAS 7001, Australia, ⁸International Research Centre for Environmental Membrane Biology, Foshan University, Foshan 528000, China, and ⁹Department of Agriculture, Food, Environment and Forestry, University of Florence, Florence, Italy

* For correspondence. E-mail nadia.bazihizina@unifi.it or phbrown@ucdavis.edu

Received: 18 November 2021 Returned for revision: 2 February 2022 Editorial decision: 8 February 2022 Accepted: 14 February 2022
Electronically published: 16 February 2022

- **Background** Soil salinity, in both natural and managed environments, is highly heterogeneous, and understanding how plants respond to this spatiotemporal heterogeneity is increasingly important for sustainable agriculture in the era of global climate change. While the vast majority of research on crop response to salinity utilizes homogeneous saline conditions, a much smaller, but important, effort has been made in the past decade to understand plant molecular and physiological responses to heterogeneous salinity mainly by using split-root studies. These studies have begun to unravel how plants compensate for water/nutrient deprivation and limit salt stress by optimizing root-foraging in the most favourable parts of the soil.
- **Scope** This paper provides an overview of the patterns of salinity heterogeneity in rain-fed and irrigated systems. We then discuss results from split-root studies and the recent progress in understanding the physiological and molecular mechanisms regulating plant responses to heterogeneous root-zone salinity and nutrient conditions. We focus on mechanisms by which plants (salt/nutrient sensing, root-shoot signalling and water uptake) could optimize the use of less-saline patches within the root-zone, thereby enhancing growth under heterogeneous soil salinity conditions. Finally, we place these findings in the context of defining future research priorities, possible irrigation management and crop breeding opportunities to improve productivity from salt-affected lands.

Key words: Irrigation, nutrient heterogeneity, phytohormones, root foraging, root-to-shoot signalling, salt sensing, stomatal conductance, water uptake.

INTRODUCTION

The world's population is expected to increase to 9.9 billion by 2050 (Liu *et al.*, 2020a), prompting a need to double annual food production within the next 30 years (Razzaq *et al.*, 2021). To achieve this production goal, agriculture will inevitably expand further into marginal lands (Pancaldi and Trindade, 2020; Ahmadzai *et al.*, 2021; Khanna *et al.*, 2021; Razzaq *et al.*, 2021), which often suffer from poor soil structure and low fertility (Mantovani *et al.*, 2015; Shukla *et al.*, 2017). Many of the marginal areas are also affected by soil salinity.

Soil salinity, both naturally occurring (i.e. primary salinization) and as a consequence of human activities (i.e. secondary salinization), is a threat to agriculture and a major limitation to food production. The salinization of agricultural land commonly

occurs as a result of using irrigation water containing elevated levels of ions (e.g. Na⁺ and Cl⁻) without adequate, periodic leaching of the accumulated salts from the soil. Secondary dry-land salinization can also occur in non-irrigated areas due to changes in the hydrological balance of a landscape and rising water tables (e.g. as a result of replacing deep-rooted vegetation with annual crop and pasture species, which results in a higher proportion of incoming rain entering the groundwater; Pannell and Ewing, 2006; McFarlane *et al.*, 2016). The problem is often exacerbated by decreases in soil permeability caused by sodicity (i.e. when the accumulation of Na⁺ exceeds that of other cations) and over-exploitation of groundwater, which exhaust high-quality water resources, resulting in water extraction from less favourable groundwater that may be brackish or

saline (Ruto *et al.*, 2021). Increases in salinity can also be expected in low-lying coastal areas associated with sea-level rise due to climate change and salinization of groundwater due to salt water intrusion into depleted aquifers (Vellinga and Barrett-Lennard, 2021).

While the vast majority of research on crop responses to salinity has been conducted under homogeneous saline conditions, root-zones of plants in both natural and managed environments can commonly experience spatial and temporal heterogeneity in soil salinity (Yakir and Yechieli, 1995; Bleby *et al.*, 1997; Davidson *et al.*, 1996; Silvestri *et al.*, 2005; Xing *et al.*, 2019; Zhang *et al.*, 2019). The nature of soils and irrigation practice, crop type and phenology, climate-type and seasonal weather, and the duration of crop exposure, together determine the extent and impact of salinity and its heterogeneity on plant growth and crop productivity (Bazihizina *et al.*, 2012a; Northey *et al.*, 2006; Bogunovic *et al.*, 2017; Xing *et al.*, 2019; Xu *et al.*, 2019). Despite this complexity, most experiments on the impacts of salinity on plants have imposed homogeneous root-zone salinity, which does not represent saline agricultural settings. The extent of the temporal heterogeneity in soil salinity in the field is illustrated within an irrigated wheat trial in China (Fig. 1A) and a rain-fed wheat trial in Western Australia (Fig. 1B). These patterns of saline heterogeneity can be contrasted with the near homogeneous conditions commonly imposed in controlled-environment research trials (Fig. 1C). Since complex multi-faceted traits (developmental, physiological, anatomical, morphological and biochemical) are involved in plant tolerance of salinity, this raises the question of whether trials conducted under near-uniform soil (or root-zone) salinity are indeed optimal for identifying and selecting traits of most value to increasing plant tolerance to the common reality of heterogeneous salinity.

The generic guidelines used to predict crop response to soil and water salinity, produced under near homogeneous conditions, are generally described by crop yield curves that consist of a threshold value at which salinity induced damage first occurs, and a linear percentage yield reduction with every increment in the electrical conductivity of the saturated soil extracts (EC_e) thereafter (Maas and Hoffman, 1977; Maas and Grattan, 1999). Such static diagnostic criteria do not reflect soil salinity under realistic field conditions that are highly spatially and temporally heterogeneous (Chen *et al.*, 2019). Thus, the prevailing standard of describing plant response to salinity is both inadequate and generally overestimates crop response (Tanji & Kielen, 2002), is not relevant nor easily interpreted under field reality, and may not adequately inform irrigation practice, crop selection or salinity mitigation strategies (Ayars, 2012). In the following, we summarize the pattern of occurrence of salinity heterogeneity in rain-fed and irrigated systems, and discuss results from studies of plant responses to heterogeneous root-zone salinity. We then discuss the mechanistic understanding of root physiological and morphological adaptations to heterogeneous conditions, and place these findings in the context of defining future research priorities and possible management and crop breeding opportunities to improve productivity in saline lands.

SOIL SALINITY HETEROGENEITY

In naturally saline environments, within the rooting zone of a single plant, non-saline patches can coexist with nearby saline

ones, ranging from a few millimolar to several times seawater (Bazihizina *et al.*, 2012a). The magnitude of this heterogeneity varies in time and space depending upon soil parent material, landscape position (Aldabaa *et al.*, 2015), soil physical and chemical characteristics (e.g. texture, sodicity and alkalinity; Hillel, 1980; Robbins *et al.*, 1980), surface runoff and subsurface lateral flow of water, intrusion by saline groundwaters or seawater (Tiggeloven *et al.*, 2020; Choukr-Allah, 2021), and root water extraction (Heuperman, 1995; Barrett-Lennard and Malcolm, 2000; Alharby *et al.*, 2014, 2018). Climatic conditions also affect the temporal and spatial heterogeneity of soil salinity, with rainfall leading to soil leaching events while droughts and heatwaves concentrate solutes depending on soil features and topography, and differences in radiation and resulting evaporation due to aspect and slope (Schwantes *et al.*, 2018).

Irrigation also dramatically influences soil salinity (Figs 2 and 3). Irrigation-induced heterogeneity can commonly result in differences in soil EC_e greater than 10-fold (Bernstein *et al.*, 1955; Bernstein and Fireman, 1957; Bernstein, 1975). An extremely heterogeneous distribution of salinity in irrigated systems makes it difficult to design a soil sampling regime to determine the truly effective root-zone salinity (Bernstein *et al.*, 1955). This effect is highly relevant to modern drip and micro-irrigated agriculture (see Box 1 for definition of terms) in arid regions (Fig. 2), which are the most common irrigation strategies in many areas of the world. While micro-irrigation is generally considered a valuable way to improve water use efficiency and allows controlled fertigation strategies, these systems may complicate salinity management, generating highly non-uniform salt and disparate nutrient deposition patterns below the irrigation emitter (Bar-Yosef, 1999). These salt/nutrient deposition patterns below the micro-irrigation emitter directly impact root growth, root activity, and nutrient and salt movement in the soil within the root zone, with effects strongly determined by crop placement, soil preparation, irrigation design and management.

Bar-Yosef (1999) further discussed the risk of salt accumulation in the root-zone under drip irrigation, suggesting that salts are not efficiently displaced to the periphery of the wetted soil volume as might occur under a full surface irrigation system. Under drip irrigation, salts can accumulate in the wetting front after several irrigation cycles but this wetting front will shrink and swell with subsequent irrigation events and root water consumption. Varying the frequency and volumes of irrigation events can manipulate this salt displacement and represents a management strategy. Salts can also accumulate at the upper margin of the wetted soil volume (close to the soil surface) due to capillarity driven by soil evaporation. This effect is particularly marked with buried drip irrigation systems where the depth of irrigation tubes, shape of the furrow, plant and environmental water use patterns (Fig. 3) influence the ultimate salt distribution.

Based upon current understanding, irrigation system placement and operation could theoretically be managed to ensure that the deposition of salinity is largely restricted to the outer margins of the wetted root-zone, thereby providing a zone of lower inner salinity with abundant plant root activity. A better understanding of soil processes and plant responses under heterogeneous conditions may therefore allow us to mitigate the adverse effects of salinity (Lycoskoufis *et al.*, 2005). Although




FIG. 1. Temporal variations in soil salinity measured in irrigated and rain-fed wheat in saline land compared with the typical experimental setup used to assess salt tolerance. In (A) the crop was irrigated with water diverted from the Yellow River that had an average electrical conductivity of 0.75 ds m^{-1} . Crops were planted in a field with shallow saline groundwater. The electrical conductivity (EC) and depth of the groundwater varied from 0.5 to 3 ds m^{-1} and 80 to 200 cm , respectively. Irrigation events are indicated with black arrows. Red arrows indicate the crop harvest (date of crop harvest is assumed based on the maturation days generally required for spring wheat). Data modified from Xu *et al.* (2013). In (B) field trials to evaluate the salinity tolerance of wheat accessions under rain-fed conditions were conducted on saline sites in Western Australia. Data modified from Setter *et al.* (2016). (C) Diagram showing a typical experimental protocol used to assess salt tolerance in irrigated sand culture in pots, where the salinity of the soil solution is increased gradually to achieve the desired concentration, which then remains constant throughout the experimental period (Hussain *et al.*, 2021). After an initial trial in hydroponics to evaluate salt tolerance at the seedlings stage, the protocol shown in (C) was used to screen the salt tolerance in different wheat germplasm grown in sand irrigated with a saline Hoagland solution. Depending on the irrigation schedule (not indicated), the salinity of the soil solution is expected to have varied depending on the evapo-transpiration and the decline in water content in the pots.

very few field experiments have tested this theory that drip irrigation can be optimized to minimize the impacts of salinity, several studies of split-root plants showed greater plant growth under heterogeneous salinities than uniform salinity, at the same

average root-zone salinity (Sonneveld and Voegt, 1990; Zekri and Parsons, 1990; Flores *et al.*, 2002; Mulholland *et al.*, 2002; Tabatabaei *et al.*, 2004; Attia *et al.*, 2009; Kong *et al.*, 2012; Sun *et al.*, 2016; Zhang *et al.*, 2019). This suggests that




FIG. 2. Irrigation induced heterogeneity in root-zone salinity. (A) Typical salt accumulation patterns in surface soils for various methods of water application. Salinity ranges from low (unshaded) to high (darkened). Arrows indicate the direction of soil water flow. Reproduced with permission from Ayers and Westcot (1985). (B, C) Examples of sloping bed irrigation style and their impact on localized salt deposition around sloped furrow irrigation. Reproduced with permission from Zaman et al. (2018).

further developing these irrigation strategies should be fruitful. Understanding the nature of plant responses to heterogeneous salinity is therefore essential to develop and implement improved irrigation practices for saline systems. In particular, this opens an excellent opportunity to improve production by manipulating the heterogeneity in the salinity of the soil solution, thereby harnessing the abilities of plants to make optimum use of less-saline patches within root-zones.

ROOT RESPONSES TO HETEROGENEOUS SOILS

Under naturally occurring and agriculturally induced salinity, plant growth is affected by the salinity of the soil solution or the ratio of salt (of which the EC_e or $EC_{1.5}$ are measures) and the water content of the soil. Soil salinities vary on spatial scales of micrometres to metres, and on temporal scales ranging from seconds to seasonal changes (Bazihizina et al., 2012a; Rellán-Álvarez et al., 2016; Dinneny, 2019). Thus, roots of a single plant will be exposed to a range of soil water salinity levels that vary temporally and spatially, with differential effects depending upon the stage of plant growth. Nevertheless, while heterogeneous salinities typically occur in salt-affected soils, experiments have almost exclusively imposed homogeneous salinity or highly manipulated experimental conditions such as split-root systems, which expose a portion of a root system to salinity while the remainder receives non-saline conditions. Although split-root experiments may not adequately mimic a complex field condition, these have provided valuable insights by demonstrating how plant responses to heterogeneous conditions differ markedly from those of homogeneous saline conditions.

Split-root experiments indicate a more nuanced plant response to saline environments than commonly recognized. As summarized in Bazihizina et al. (2012a), key features of plants exposed to heterogeneous salinities are: (1) shoot water potentials are determined by the salinity level of the low-salinity zone, (2) water uptake occurs predominantly from the low-salinity medium and (3) greater maintenance of shoot growth even when a large proportion of the root system is exposed to high NaCl concentrations that would greatly inhibit growth if applied uniformly to the roots. Transcriptome profiling of plants exposed for 6–9 h to heterogeneous salinities indicated that improved performance under heterogeneous conditions compared to uniform salinities is related to the rapid activation of salt resistance genes and crosstalk between the non-saline and high-saline root sides (Kong et al., 2016; Xiong et al., 2018; Zhang et al., 2019). This suggests that roots operate as the central hub that control: (1) how stress is perceived, (2) long-distance communication with the shoots and (3) the integration of long-distance systemic signals with local root-based ones. Furthermore, salinity heterogeneity is inevitably linked to temporal and spatial variation in the distribution and biological availability of water, essential nutrients and soil pH (Li et al. 2011; Feng et al., 2018; Zhang et al., 2019), with the latter having a major impact on root membrane potential, thus affecting both a plant's ability to acquire essential nutrients and exclude toxic Na^+ and Cl^- ions (Babourina et al., 2001; Zhang et al., 2018). Thus, responses at the root level also play a critical role in: (4) how plants compensate for water/nutrient deprivation and limit salt stress by optimizing root-foraging in the most favourable part of the soil. Understanding how roots respond to heterogeneous salinities is therefore of utmost importance and is needed to develop




FIG. 3. Depiction of salt accumulation patterns from subsurface irrigation in a lettuce crop in the Santa Maria Valley (California, USA). While a low EC zone is formed beneath the emitter, salts can accumulate above and require rain or other surface irrigation to percolate salts beneath the emitter for leaching. Soil types were a clay loam and a fine sand. Reproduced with permission from Hanson and Bendixen (1995).

Box 1. Irrigation system terminology

Term explanation

Surface Irrigation

All formats of irrigation which rely on gravity (rather than pressurized conveyance systems) to distribute water across a field. Examples include flood and furrow irrigation. It is important to note that while these systems are collectively referred to as surface irrigation, they are not the only irrigation systems which apply water to the soil surface; overhead sprinkler, micro-sprinkler and surface drip also irrigate the soil surface but are not generally categorized as traditional 'surface irrigation'.

Drip

Highly local, small volumes of water emanating from points along a pressurized plastic pipe ('drip line') which is installed either on the soil surface (surface drip) or buried beneath the crop ('subsurface drip'/'subirrigation') with very little loss to evaporation.

Micro-irrigation

Micro-sprinkler

Any small, stationary sprinklers which are installed near the soil's surface and irrigate the soil's surface without applying water to the canopy. Micro-sprinkler irrigation is more targeted than overhead sprinkler irrigation, particularly with orchard crops which are widely spaced and have high canopies which can lose large fractions of overhead irrigated water to evaporation.

efficiently resources are captured. These are fundamentally controlled at variable spatial scales, from the single cell to the entire organ.

Salinity sensing

Local patches of high salinity are sensed in individual cells, and then integrated into organ-scale processes. After salinity increases, plants experience multiple constraints ranging from reduced water availability, disturbance to cytosolic ion homeostasis, and dramatic increases in reactive oxygen species (ROS) accumulation. The emerging picture suggests that more than one sensory mechanism may operate in the same cell at the same time, with some common downstream signalling pathway(s) (Shabala *et al.*, 2015; Feng *et al.*, 2018; Niu *et al.*, 2018; Fichman and Mittler, 2020, 2021; Peck and Mittler, 2020).

Calcium and ROS signals are amongst the first signals commonly evoked upon biotic and abiotic stressors. Ca^{2+} and ROS signals are established second messengers involved in most (local) stress responses, and increasing evidence suggests that these act in tandem, interacting and amplifying each other during root salt sensing (Dodd *et al.*, 2010; Choi *et al.*, 2014; Shabala *et al.*, 2015; Evans *et al.*, 2016; Niu *et al.*, 2018; Pottosin and Zepeda-Jazo, 2018). Several molecular components underlying Ca^{2+} and ROS signalling (including *MOCA1*, *OSCA1* and *RBOHs*) have been identified and are currently being considered as potential salt sensors (Yuan *et al.*, 2014; Jiang *et al.*, 2019; Liu *et al.*, 2020b). Interestingly, local salt stress at the root apex triggers immediate cytosolic Ca^{2+} increases at the point of application, leading to propagation of

management strategies to optimize resource use and crop productivity in saline soils. Different processes enable roots to integrate fluctuating soil conditions into appropriate developmental and physiological responses that ultimately determine how

a *TPC1*- ('two-pore channel 1') dependent Ca^{2+} wave to distal shoot tissues, passing through cortical and endodermal cell layers (Choi et al., 2014). By combining experimental analyses and mathematical modelling, Evans et al. (2016) also clearly linked the $[\text{Ca}^{2+}]_{\text{cyt}}$ wave triggered by a localized salt application with systemic ROS waves. Additional salt sensors (extensively reviewed in Shabala et al., 2015, 2016; Byrt et al., 2018; Rui and Dinneny, 2020; Gigli-Bisceglia et al., 2020) include: (1) cell walls, and in particular the salt-induced alterations in cell wall integrity and composition that are sensed by the receptor-like kinase, FERONIA (*FER*) (Feng et al., 2018); (2) mechanosensory channels and transporters (e.g. *OSCA1*, *MSLs*, *MCA*s) that sense the mechanical force exerted on the plasma membrane due to the osmotic component of salinity and translate hydraulic cues into chemical signals (Yuan et al., 2014; Yoshimura et al., 2021); and (3) Na^+ transport systems and proteins with regulatory Na^+ binding sites (e.g. *MOCA1*; Jiang et al., 2019).

While it is becoming increasingly clear that plant cells sense and respond to salinity stress by activating multiple sensing networks, much of our knowledge on root salt sensing and signalling has utilized uniform conditions, with no such studies attempted for heterogeneous salinities. Such experiments will generate valuable information on how salt sensing at the single cell level is integrated into organ-scale processes, revealing how the signal propagates and its effects on root system architecture, developmental trade-offs and root plasticity.

Root foraging

Scaling up to the whole root level, the root system is a highly dynamic physical network that enables a plant to forage for resources and rapidly explore favourable soil patches. Under spatially heterogeneous soil salinities, preferential root growth can occur in the least (or non-) saline compartment, compensating to different degrees for root growth inhibition in the saline patches (Bazihizina et al., 2009, 2012b; Feng et al., 2017; Sun et al., 2016; Xiong et al., 2018; Table 1, Fig. 4). A split-root experiment that closely examined root morphology under heterogeneous salinities revealed that compensatory root growth in

the non-saline areas was associated with increased lateral root growth, which doubled compared to plants with both root halves in non-saline conditions (Feng et al., 2017). However, root proliferation in the non-saline compartment does not always occur, with several studies showing no differences, or even a decline, in root growth compared to measurements under uniform non-saline conditions (see references in Table 1). This highlights the complexity of interpreting how heterogeneous conditions alter root growth, because responses depend on timescale, salt concentration and species sensitivity to salinity.

To understand root foraging it will be necessary to determine whether heterogeneous salinities (and the associated variability in water availability and nutrient distribution) affect root anatomical features, in addition to any effects on root morphology. The section below considers suberin deposition in root cell walls. Even when heterogeneous salinities do not alter root architecture, it remains possible that traits that reduce the metabolic cost of soil exploration, such as cortical cell enlargement and cortical senescence, could be beneficial. This has yet to be tested. Nevertheless, the anatomical traits that reduce the metabolic cost of root soil exploration are currently considered an advantage in water-, nitrogen- and O_2 -limited soils as these improve water and nutrient uptake per unit investment in roots (Lynch, 2018, 2019; Schneider and Lynch, 2020; Colombi et al., 2022). This topic therefore merits greater research efforts to identify key root traits that maximize soil resource capture under heterogeneous salinity.

Water uptake

Irrespective of environmental heterogeneity within the root-zone, plant water uptake is essential to maintain photosynthesis. Typically, water uptake from the non-saline side of the root system increases significantly, which is not always accompanied by increased root biomass (Fig. 4A). Roots can dynamically alter their water transport capacity to acclimate to the ever-changing soil conditions and rapidly explore favourable soil patches. Under heterogeneous salinity, preferential water uptake from the regions with the least negative water potentials are mediated by changes in root hydraulic conductivity that

TABLE 1. Root distribution in different species under horizontally heterogeneous salinities as percentage of dry mass in control plants with low or no-salt media

Species	Salinity (mM NaCl)	Duration	Root biomass (% control)		Ratio L : H	Reference
			L	H		
<i>Atriplex nummularia</i>	10/500	3 weeks	88	87	1.0	Bazihizina et al. (2012b)
<i>Atriplex nummularia</i>	10/1500	3 weeks	153	32	4.8	Bazihizina et al. (2012b)
<i>Hibiscus moscheutos</i>	0/200	2 months	72	57	1.3	Feng et al. (2021)
<i>Hibiscus moscheutos</i>	0/200	2 months	79	20	4.0	Feng et al. (2021)
<i>Medicago sativa</i>	0/200	15 d	140	56	2.5	Xiong et al. (2018)
<i>Medicago sativa</i>	50/200	15 d	131	61	2.1	Xiong et al. (2018)
<i>Medicago sativa</i>	0/75	9 d	99	90	1.1	Sun et al. (2016)
<i>Medicago sativa</i>	0/150	9 d	113	82	1.4	Sun et al. (2016)
<i>Medicago sativa</i>	0/225	9 d	118	56	2.1	Sun et al. (2016)
<i>Medicago sativa</i>	75/150	9 d	82	60	1.4	Sun et al. (2016)
<i>Medicago sativa</i>	75/250	9 d	72	55	1.3	Sun et al. (2016)
<i>Sorghum bicolor</i>	0/200	2 weeks	100	49	2.0	Zhang et al. (2019)




FIG. 4. Taking advantage of the patch. Plants use the most accessible water source under spatially and temporally variable salinity. The datasets in (A) show water uptake and root growth in the non-halophytic almond rootstock ‘Nemaguard’ (*Prunus persica* × *P. davidiana*) and the halophyte *Atriplex nummularia*, after 28 and 21 d of heterogeneous salinity treatment respectively. The data shown for almond rootstock and *Atriplex nummularia* are modified from Valenzuela (2018) and Bazihizina et al. (2009), respectively. Values are mean ($n = 4$) ± SE and different letters indicate a significant difference ($P < 0.05$) between treatments and root sides. The dataset in (B) shows temporal changes in soil water potentials and number of new roots in *Melaleuca halmaturorum* in response to fluctuating saline groundwater. The data shown in the upper two panels refer to soil water potentials and new roots observed at the end of summer, while those shown in the bottom two panels refer to soil water potentials and new roots at the end of winter. The red asterisks indicate the location of main water uptake (based on the uptake of stable isotope data). The figure is modified from Mensforth and Walker (1996).

occur within hours of salt exposure (Kong et al., 2016). These are achieved through changes in the abundance or activity of water channel proteins named aquaporins that facilitate water diffusion across cell membranes (Maurel et al., 2008; Gambetta et al., 2017; Kong et al., 2017; Maurel and Nacry, 2020). The activity of aquaporins is regulated at many levels, including altered transcription levels, channel gating between an open/closed state by various mechanisms including phosphorylation, pH or Ca^{2+} , and changed cellular trafficking (Maurel et al., 2008; Gambetta et al., 2017; Maurel and Nacry, 2020). Under heterogeneous conditions, increased water uptake from the non-saline roots has largely been attributed to changes in aquaporin expression levels (Kong et al., 2017). After applying 200 mM NaCl to one root half in split-root cotton seedlings (with 0 mM NaCl to the other half), gene expression profiling revealed several aquaporin genes were up-regulated within 3 h in the non-salinized root half, resulting in 16 % higher root hydraulic conductivity when measured against NaCl-free controls (Kong et al., 2017). By contrast, both root hydraulic conductivity and most of the differentially expressed aquaporin genes

were largely inhibited in the high-salinity side (Kong et al., 2017).

The deposition of hydrophobic lignin and suberin in the cell walls of the exo- and endodermis also alters root hydraulic conductivity and restricts the free diffusion of solutes and water, including restricting entry of Na^+ and Cl^- from the soil into the vascular stream with high root-zone salinity (Krishnamurthy et al., 2011; Barberon et al., 2016; Wang et al., 2019; Cui et al., 2021). Accordingly, dynamic regulation of root hydraulic conductivity under heterogeneous salinities was also associated with altered expression of genes associated with cutin, suberin and wax biosynthesis in the salinized root portions (Xiong et al., 2020). This could potentially explain decreased endodermal and exodermal permeabilities, which limit water and solute transport from the highly saline areas.

The ability of plants to acquire and transport water from the roots to the leaves also depends on root anatomy and architecture, and the combined hydraulic conductivities among root types and along the root length (Meunier et al., 2017; Ahmed et al., 2018). Thus, over the longer term (days), increases in new

root growth and altered root architecture (i.e. root proliferation and increased lateral root formation) and anatomy may have a more significant effect than localized changes in root hydraulic conductivity at the single root level. Nevertheless, our understanding of the timescale and concentration-dependent drivers of the long- and short-term responses of roots to localized salinity is inadequate. As highlighted in the following sections and above, responses are expected to become increasingly complex when heterogeneous salinity interacts with other environmental factors, such as heterogeneous nutrients as discussed below, and their impacts on plant nutrient and water acquisition.

Phytohormone root–shoot communication

Heterogeneous salinity can induce variable degrees of stomatal closure, with stomatal conductance similar to uniform salinity in some studies (Lycoskoufis *et al.*, 2005; Fig. 5). However, most split-root studies indicate greater plant water use under heterogeneous than uniform salinity, at the same average root-zone salinity. This is mostly because plant water uptake from the non-salinized part of the root system substantially increases, even exceeding water uptake from roots of non-salinized plants (see section above). Long-distance signalling *in planta* is implicated in regulating these plant water relationships under heterogeneous salinity by modulating root hydraulic conductivity and stomatal conductance.

Although leaf water status is regarded as an important regulator of stomatal responses (Christmann *et al.*, 2007), it

is generally determined by the non-salinized part of the root-zone under heterogeneous salinities (Bazihizina *et al.*, 2009, 2012a, b; Feng *et al.*, 2021). Considerable stomatal closure of these plants (Fig. 5) suggests non-hydraulic mechanisms of stomatal closure. Homogeneous salinity induced multiple phytohormonal changes in salinized roots, according to the duration of exposure, with phytohormones such as abscisic acid (ABA), auxin and cytokinins (Albacete *et al.*, 2008), and their crosstalk, mediating the balance between growth and salinity stress responses (Yu *et al.*, 2020). The same applies to heterogeneous salinities. In cotton grown with heterogeneous salinity (0/200 mM NaCl), 200 mM NaCl induced only transient (within 3–12 h of treatment) increases in root ABA concentration. Root ABA levels were similar to controls after 24 h, presumably as sustained up-regulation of ABA catabolism (*CYP707A*) genes influenced root ABA concentrations more than concurrent up-regulation of ABA biosynthesis (*NCED*) genes (Kong *et al.*, 2016). Paradoxically, root ABA concentrations of the non-salinized roots exceeded those of salinized roots throughout the experiment, despite a limited and transient (3–6 h) up-regulation of (*NCED*) genes, implying considerable ABA transport into these non-salinized roots. Further studies are needed to elucidate the source of this additional ABA, since the shoot can regulate root ABA concentration (Manzi *et al.*, 2015; McAdam *et al.*, 2016), which in turn upregulates root hydraulic conductance (Thompson *et al.*, 2007).

Heterogeneous salinity also altered the concentrations of other phytohormones in the non-salinized portion of split-root cotton plants: with indole acetic acid, isopentenyladenine and




FIG. 5. Stomatal conductance of salinized plants (expressed as a percentage of non-salinized controls) exposed to heterogeneous (hollow symbols) and homogeneous (filled symbols) salinity at the same average root-zone salinity. Original papers were from: *Solanum lycopersicum* (Wang *et al.*, 2021), with plants grown with uniform (0 and 50 mM NaCl) and heterogeneous salinity (17/85 mM NaCl); *Hibiscus moscheutos* (Feng *et al.*, 2021), with plants grown with uniform (0 and 200 mM NaCl) and heterogeneous salinity (0/400 mM NaCl); *Sorghum bicolor* (Zhang *et al.*, 2019), with plants grown with uniform (0 and 100 mM NaCl) and heterogeneous salinity (0/200 mM NaCl); *Lycium chinense* (Feng *et al.*, 2017), with plants grown with uniform (0 and 170 mM NaCl) and heterogeneous salinity (0/340 mM NaCl); *Gossypium hirsutum* (Kong *et al.*, 2012), with plants grown with uniform (0 and 100 mM NaCl) and heterogeneous salinity (0/200 mM NaCl); and *Atriplex nummularia* (Bazihizina *et al.*, 2009), with plants grown with uniform (10 and 230 mM NaCl) and heterogeneous salinity (10/450 mM NaCl).

zeatin riboside concentrations increasing compared to their concentrations in plants that were not exposed to salinity (Kong et al., 2016). In this case, increased root cytokinin concentrations correlated with increased expression of *IPT* genes, which were maximal 3 h after salinizing the other part of the root system. Measuring root water potential in a transpiring plant (Adeoye and Rawlins, 1981) may help determine whether this was a transient response to altered root water relationships in the non-salinized roots. Such measurements (along with root gene expression) are required in girdled and non-girdled plants (since girdling at the root–shoot junction blocks phloem transport to the roots) to determine whether local root water relationships and/or a cumulative message from other parts of the plant regulate gene expression. Under heterogeneous salinity, Na^+ accumulation in the non-salinized portion of the root system doubled compared to roots from non-salinized controls. Such Na^+ accumulation depended on phloem transport from the salinized roots, as girdling prevented Na^+ transport to these roots (Kong et al., 2012). Whether girdling eliminates changes in root phytohormone concentration in non-salinized roots, when the other part of the root system is exposed to salinity, needs to be addressed.

Irrespective of whether changes in root phytohormone concentration occur, it is uncertain whether they actually affect shoot phytohormone concentrations and physiological responses, since root-to-shoot signalling under heterogeneous soil conditions depends on relative sap flow from different parts of the root system (Dodd et al., 2008). Under heterogeneous salinity, changes in root phytohormone concentration in the salinized root system may have little impact on shoot physiology since these roots contribute relatively little to total transpirational flow (Kong et al., 2012). Interestingly, changes in root phytohormone concentration in the non-salinized roots may have a greater influence on shoot physiology, since these roots contribute most of the total water flux. Grafting techniques allow the relative contribution of different parts of the root system to root phytohormone export to be evaluated (Dodd et al., 2008), but to date this has only been attempted in plants exposed to different soil moisture levels and such experiments should be applied to plants with heterogeneous root-zone salinity.

HETEROGENEOUS SALINITY AND NUTRIENT DISTRIBUTION: THE MISSING LINK?

In both natural and managed systems, a variable distribution of nutrients (and salinity) in soils is inevitable. This is associated with differential ion mobility and solubility, localized decomposition of organic matter, or applying soluble nutrients through a fertigation system. Furthermore, considerable evidence suggests that nutrient ‘patches’ can influence root foraging, lateral root formation and root hair formation. Thus, the mutual effects of salinity and nutrient heterogeneity are relevant.

Root physiological responses to nutrient heterogeneity in fertigated crops

Though probably a common occurrence in drip irrigated crops, very few studies have simultaneously varied both salinity

and nutrient distribution. The following discussion first considers experiments with only nutrient heterogeneity, before discussing the integration of nutrients with salinity heterogeneity.

In tomato, preferential nitrate (NO_3^-) uptake was found to occur from areas of the root-zone with higher (1.6- to 3.3-fold greater, with 10 ds m^{-1} being the highest EC) electrical conductivity (or more negative osmotic potential) generated by locally high nutrient concentrations (Sonneveld and Voogt, 1990), suggesting a local response of roots exposed to high concentrations probably due to their enhanced NO_3^- uptake kinetics. Mathematical simulations of nutrient uptake under heterogeneous conditions of NO_3^- and phosphate (PO_4^{3-}) using the Barber–Cushman model found a greater impact of soil heterogeneity and root plasticity, with NO_3^- uptake increasing 7–20 times under heterogeneous conditions (Jackson and Caldwell, 1996). Root proliferation and increased uptake kinetics from the enriched root-zones accounted for up to 75 % of NO_3^- supply of a plant and over 50 % of PO_4^{3-} acquired from enriched soil patches. Simulations demonstrated that plants lacking plasticity of root growth or uptake always acquired less nutrients under heterogeneous NO_3^- and PO_4^{3-} distributions.

In a split-root solution culture experiment on *Lolium multiflorum*, less than 24 h after depriving NO_3^- from half the root volume, net NO_3^- influx to roots in the nitrate-rich area increased, with root growth increments observed only after 1 week (Lainé et al., 1998). *Brassica napus* responded similarly (Lainé et al., 1995). This rapid variation in NO_3^- uptake was strongly associated with altered root hydraulic conductivities, with a sudden increase in NO_3^- concentration around the roots almost simultaneously increasing root hydraulic conductivity and preferential water uptake from the nitrate-rich patch (Gorska et al., 2008). Split-root experiments applying NO_3^- to a portion of the root system demonstrated a localized and reversible response, with N starvation on one side of the root system leading to compensatory and enhanced NO_3^- uptake in the other root portion (Tabata et al., 2014).

Heterogeneous NO_3^- distribution to split-root *Acer rubrum* and *Betula papyrifera* plants demonstrated a species-dependent response, with two-fold more fine roots measured for *B. papyrifera* in the high NO_3^- portion than *A. rubrum*, yet similar total NO_3^- uptake rate (Gloser et al., 2008). Under heterogeneous conditions, *A. rubrum* had smaller leaves and N deficiency symptoms in the shoot portion directly above the nutrient-deficient root portion, while *B. papyrifera* had regular leaves with no visible deficiency symptoms. Vascular system architecture may explain this differential response (Orians and Jones, 2001). In species with sectorized vascular systems (e.g. *A. rubrum*), in which contiguous and largely exclusive vascular traces occur from a specific root to a specific branch, N deficiencies occurring in isolated parts of the canopy reflect the nutritional status of the specific root that feeds that branch. In contrast, other species (*B. papyrifera*) have an integrated vascular system allowing nutrient transfer from an individual root to the canopy as a whole, avoiding the consequences of patchy nutritional deficiencies.

Root morphological responses to nutrient heterogeneity

In *Betula pendula*, dry matter allocation to roots can be modified in three different ways when the availability of

mineral nutrients is limited: (1) increased root growth in N-, P- or S-limited soils; (2) decreased root growth when K^+ , Mg^{2+} and Mn^{2+} is limited; and (3) no effect on root growth when Ca^{2+} , Fe^{2+} and Zn^+ are limited (Ericsson, 1995). Root growth plasticity in patchy soil enhances the ability of plants to fill the soil volume rich in nutrients and was the most important trait influencing species success (Hodge, 2006; Rajaniemi, 2007). The ability of a plant to ‘find’ the nutrient-rich patch is essential if morphological/physiological root responses are to be expressed. For instance, while nutrient (N) heterogeneity in *Lolium perenne* did not lead to preferential root growth in the nutrient-rich soil patches (suggesting the patch was not explored to any greater extent than the bulk soil), there were overall increases in specific root length (length/biomass) and root elongation throughout the entire soil profile compared to the uniform N treatment (Nakamura et al., 2008). This suggests that the overall plant N deficiency induced root elongation and

not the patchiness per se. The differential response of roots to nutritional patchiness is probably a consequence of complex nutrient-specific signal transduction pathways (López-Bucio et al., 2003).

Impacts of simultaneous salinity and nutrient heterogeneity

To investigate the effects of heterogeneous root salinity and nutrient conditions, several split-root tomato experiments were conducted (Fig. 6; Valenzuela et al., 2021). Water uptake from the saline root-zone dramatically decreased within 8 h of treatment (Fig. 6A, B) in contrast to the non-saline root-zone, with a more pronounced effect when nutrients were provided only to the non-salinized root-zone (Fig. 6A, B). This reduction in water uptake did not correlate with decreased root growth (which was maintained during Days 1–3), with the saline root-zone only




FIG. 6. Daily measurement of the percentage of water consumption per root side in a split-root system under hydroponics. The vertical dashed line shows the time when salinity application was initiated. Dots represent treatments applied to side A and triangles to side B. The saline agent was NaCl with a concentration of 50 mM. Bar graphs, at the right, show the root biomass allocation for each treatment. In these experiments, tomato (*Lycopersicon esculentum*) seedlings were grown in a hydroponic split-root method for 9 d under heterogeneous saline and nutritional conditions applied separately and in combination. Root activity was monitored by directly measuring root uptake of water and nutrients, biomass allocation and total nutrient uptake. The short experimental time frame and modest salinity concentrations were selected to minimize ionic toxicity and plant growth effects. Reproduced with permission from Valenzuela et al. (2021).

showing significantly less root growth towards the end of the experiment (Day 9). The rapidity and consistency of decreased water uptake by roots in the saline zone, from treatment imposition through to Day 9, suggests that a primary physiological response (possibly due to reduced aquaporin activity) was followed by a morphological response.

To further explore the role of heterogeneous nutrient provision on root activity, complete nutrient solutions were selectively depleted of either N or K⁺ in the non-saline root half while the other root half received a saline, complete nutrient solution (Fig. 6C, D). These treatments provoked a ‘two-phase-response’. Immediately upon treatment application, the saline conditions given to one side of the roots dominated, immediately decreasing water uptake of those roots. Subsequently, water uptake from the saline-treated, nutrient-supplied roots proportionally increased, probably in response to the nutrient deficiency induced by the omission of the nutrient on the non-saline side. This effect was marked when K⁺ was only present in the saline root half and slight in the case of N. The presence of K⁺ in the nutrient solution was the most important determinant of root activity even when coinciding with salinity, resulting in a notably higher shoot tissue Na⁺ and Cl⁻ concentration when the sole source of K⁺ was to the saline root volume (Valenzuela et al., 2021).

This experiment and others described herein suggests that interpreting root responses to heterogeneous conditions depends markedly on context, time, salinity concentrations and plant nutrient status:

- Immediate (within hours) reductions in water uptake in salt-exposed root-zones commonly occur and are expected to be determined by the relative difference in salinity between root parts. Subsequently, relative nutrient availability affects root activity, with responses to K⁺ depletion perceived within days. In the longer term, ionic stress and changes in relative root growth may further alter relative root activity in each root-zone.
- Plant responses to the relative distribution of nutrients in the root-zone probably also depend on plant nutrient status and the absolute concentrations of nutrients and salinity present in each root-zone. Thus, a K⁺-replete plant, provided low levels of K⁺ in the non-saline root-zone, or the presence of extreme salinity (>100 mM), would probably diminish the dramatic response seen here (Fig. 6).
- All split-root systems or otherwise manipulated heterogeneous root-zone experiments do not reflect the complexity of natural ecosystems, where soil heterogeneity is probably significantly more complex in space and time. However, certain agricultural conditions may closely resemble split-root studies, such as in substrate hydroponic systems, raised bed vegetable production and micro-irrigated arid zone crops.

While these simple split-root experimental approaches cannot explain all potential nutrient interactions, they do illustrate the rapidity and plasticity of plant responses and the importance of considering nutrients when studying heterogeneous salinity. Furthermore, this raises several important questions on the signalling pathways underlying root system architecture and functions under heterogeneous saline conditions: (1) What is the relative importance of the signal(s) under heterogeneous saline conditions (salt ions vs. nutrients vs. water vs. hormones)? (2) How does variation in salt tolerance affect this response

(e.g. halophytes vs. non-halophytes)? (3) How do the local conditions (salinity vs. nutrient vs. water availability) and whole plant status (e.g. shoot Na⁺, Cl⁻ and/or nutrient concentrations) modulate the response? Interestingly nutrient availability alters the endodermal specific ABA signalling in roots that modulates lateral root formation and root system architecture in response to salinity stress (Duan et al., 2013). As demonstrated above, this suggests that the ‘nutrient signal’ might eventually override, or at least affect, the ‘salinity signal’ in regulating root growth and functions under heterogeneous conditions.

IMPLICATIONS FOR CROP MANAGEMENT

When considering crop responses to salinity, two parameters define salt tolerance: (1) the threshold salinity that causes the initial significant reduction in the maximum expected yield, and (2) the rate of yield decline as salinity increases beyond the threshold (i.e. slope; Maas and Hoffman, 1977; Maas and Grattan, 1999). However, the important underlying assumption of the threshold model by Maas and Hoffman (1977) is that steady-state conditions exist in the root-zone, with water content and salt concentration remaining constant in time and space. Since these conditions rarely exist in crop root-zones in the field (e.g. Fig. 1), conclusions from steady-state analyses can be questionable (Letey and Feng, 2007). This has critical repercussions when defining the salinity experienced by the roots when identifying breeding targets and soil management practices.

Re-defining soil salinity

One valuable tool in categorizing and quantifying genetic variation in salt tolerance has been to define crop relative yield responses in terms of threshold salinities up to which yields are unaffected and linear decreases in relative yield with increasing salinity thereafter (cf. Maas and Hoffman, 1977, and their successors). However, it is critical to recognize that these relationships have generally always been presented in terms of variation in parameters such as EC_e (the electrical conductivity of the soil saturation extract) or more occasionally in terms of variation in EC_{1:5} (the electrical conductivity of a 1 : 5 soil : water slurry) that relate to the salinity of the soil. However, it is not the salinity of the soil (a parameter that does not account for variations in soil water content) that affects plant growth but the salinity of the soil solution, and thus the ratio of salt to water in the soil. This means that the salinity stress on a plant can be doubled by doubling the salt concentration in a soil or by halving the water concentration of the soil. Furthermore, as soils become drier, plant growth becomes affected by the increasingly negative matrix potentials (Ψ_m values) that develop in soils because of the adhesion of water by soil pores.

This view profoundly affects the whole idea of the heterogeneity of salinity stress in soils, because heterogeneity arises because of variable: (1) leaching effects of irrigation or rainfall on salt concentrations in soil, (2) hydrating effects of irrigation or rainfall on soil water contents, (3) effects of surface soil evaporation increasing salt concentrations by capillarity and decreasing water contents in the soil, and/or (4) water extraction

rates of roots and the ion uptake/exclusion capacity, which over time also influence ion and water abundances near the roots.

One variable that captures variation in both salt and water concentrations in soil is solute potential (Ψ_s ; units MPa). For soils salinized with NaCl, this can be calculated as:

$$\Psi_s = -22.75 \times EC_{1:5}/W \quad (1)$$

where the $EC_{1:5}$ of the bulk soil is in units of $ds\ m^{-1}$, and soil water content of the bulk soil (W) has units of % dry mass.

Water potential of the soil (Ψ_{soil}) can be calculated (Slatyer, 1967) as:

$$\Psi_{soil} = \Psi_m + \Psi_s \quad (2)$$

Furthermore, a leaf's transpiration rate (E_t) can be related to its water potential (Ψ_{leaf}), the water potential of the soil (Ψ_{soil}) and the resistance to flow (R) as follows (Nulsen and Thurtell, 1980):

$$E_t = (\Psi_{soil} - \Psi_{leaf})/R \quad (3)$$

While theoretically sound, plants change these simple mathematical relationships. First, salt accumulates in the root-zone making Ψ_s more negative than can be calculated using the $EC_{1:5}$ and W measurements of the bulk soil. Passioura and Freere (1967) define the variable U as the factor by which Ψ_s at the root surface is more negative than in the bulk soil. Experimental approaches suggest that U can be around 2 in well-hydrated soils (e.g. Sinha and Singh 1974, 1976), but modelling approaches suggest that U could increase to values around 10 as the soil becomes drier (Passioura and Freere, 1967). Given this, eqn (2) can be modified to:

$$\Psi_{soil} = \Psi_m + U\Psi_s \quad (4)$$

Second, the resistance of water flow to the surface of the root increases as the soil pores around the root become depleted of water (Stirzaker and Passioura, 1996). This increases the variable R in eqn (3).

Given this background, how should heterogeneity of salinity in the soil solution in the root-zone be viewed? Unfortunately, very few of the critical experiments have been done under uniform conditions in the root-zone, let alone variable ones. To our knowledge, no split-root experiments have ever attempted to compare the effects of different (or even the same) Ψ_{soil} values by manipulating the salt and water concentrations on each side of the root-zone of a plant, even though the components of Ψ_{soil} (Ψ_m , Ψ_s and U) can all be determined experimentally. Maintaining such treatments is technically challenging, requiring new experimental protocols to be developed. In such experiments, there could be considerable rewards by comparing plants of different salt tolerance.

Opportunities to better manage irrigated agriculture

All irrigation water introduces salts to the system (Hanson and Bendixen, 1995) and in regions with high evapotranspiration and low rainfall, traditional salinity management emphasizes deliberate leaching of salts away from the root-zone while avoiding elevation of the water table to prevent damage to crops (Hopmans et al., 2021). Leaching is usually achieved by

applying irrigation water in excess of crop evapo-transpirational demands. The fraction of applied water that drains below the root-zone is referred to as the 'leaching fraction' and this value is used to coarsely gauge the extent of leaching (Hanson et al., 2009). Larger leaching fractions generally result in larger zones with a low soil water salinity but may necessitate disposal of large volumes of saline drainage water and may cause additional salinization through capillary rise of saline water by raising the water table (Grismer et al., 1988; Corwin, 2021), as well as environmental impacts of drainage water disposal.

Designing the appropriate leaching fractions needed to avoid yield loss is context-specific and will depend on the crop, soil texture, climate, irrigation system and irrigation schedule, and the salinity of irrigation water being used (Ayers and Westcot, 1985; Hanson and Bendixen, 1995; Assouline et al., 2015). Ayers and Westcot (1985) developed a simple approach to calculate the leaching requirement based on salt mass balance calculations. This approach estimates the leaching fraction required to keep the average root-zone salinity below the salinity threshold of the crop, assuming a specific root distribution and a strictly vertical, continual water flow. Approaches like this neglect the spatial non-uniformity of irrigation water application as well as the temporal dynamics of irrigation and water uptake during the season (Letey et al., 2011) and assume that the average root-zone salinity determines the impact of salinity on the crop (Letey and Feng, 2007).

While the physical principles underlying salinity management have not changed since Ayers and Westcot developed these leaching guidelines, management goals have shifted over time to better recognize environmental impacts of nutrient and salinity losses and develop more advanced micro-irrigation and fertigation systems. This has given rise to both new challenges and new opportunities in managing salinity.

Challenge 1: Managing salinity under micro-irrigation systems. Spatial patterns of salt accumulation are diverse and differ by irrigation system (Riaz et al., 2018; Wallender and Tanji, 2011), with each irrigation system having specific challenges to salinity management. In the simplest case, flood irrigation applies water uniformly across the whole surface (although local topography and soil heterogeneity can cause spatially heterogeneous infiltration). In this case, salinity distribution is approximately uniform in the horizontal direction, but a salinity gradient exists vertically (Figs 2 and 3). Assuming sufficient leaching, salinity increases with depth in these systems (Ayers and Westcot, 1985) and uniform leaching of salts below the root-zone causes the salinity within it to be relatively homogeneous.

In contrast, applying water to only part of the surface causes strong horizontal salinity heterogeneity, as in furrow irrigation and more advanced micro-irrigation systems. Micro-irrigation aims to target water application to the root-zone, thereby improving water use efficiency by applying less water to regions with low root density and providing an opportunity to deliver water at a rate which matches crop demand. Flood and overhead sprinkler irrigation manage soil moisture and salt content at the field scale, while micro-irrigation approaches management at the root-zone scale. Targeted water application results in targeted leaching, with micro-irrigation leaching salts in zones which are rich with plant roots, while flood irrigation

requires additional water to also leach salts from field zones between plants with low root density, making micro-irrigation more efficient than furrow/sprinkler irrigation for managing salinity (Hanson *et al.*, 2009). When drip and furrow irrigation were compared, drip irrigation sustained higher yields of salt-sensitive crops compared to furrow irrigation when saline groundwater is shallow, while using less water than furrow irrigation (Hanson *et al.*, 2009).

The economic incentive to install micro-irrigation systems is context-dependent, with the advantage of micro-irrigation over conventional irrigation becoming less clear when growing salt-tolerant crops or when irrigation water is abundant. Despite its potential to accumulate salts in the root-zone, even subsurface drip can have advantages over salinity management with traditional irrigation. While higher tomato yields justified the expense of installing a subsurface drip irrigation system in California, the same was not true of cotton, which remained lucrative with furrow irrigation (Hoffmann and Johnsson, 2000; Hanson *et al.*, 2009), as such salt-tolerant crops tend to tolerate flood irrigation without yield loss provided that irrigation is applied pre-planting to avoid stand establishment losses (Ayars *et al.*, 1993; Hanson *et al.*, 2009).

In drip irrigation systems with strongly localized water application, salt is not only leached downwards, but significant lateral water movement away from the drip emitter also leaches salt horizontally (Raine *et al.*, 2007) resulting in salt accumulation in the fringes of the wetted volume (Fig. 2A). This leads to a strongly heterogeneous small-scale salt distribution where soil salinity levels in the top 20 cm can vary by a factor of more than five within only 40 cm of horizontal distance (e.g. May and Hanson, 2006). Although the extent of horizontal salt movement depends on the soil texture and can be partially controlled by emitter spacing, under micro-irrigation, salts concentrated between emitters near the surface generally have little opportunity to intrude into the root-zone without precipitation, due to surface evaporation and irrigation (Hanson and Bendixen, 1995; Hanson and May, 2011). It is therefore recommended that crops be arranged close to emitters where salinity is low and that new lines be installed as close as possible to where old lines existed to avoid the need for pre-season reclamation leaching (Hanson and May, 2011).

Subsurface drip irrigation results in a different pattern of water flow and salinity accumulation. While water application at the soil surface causes salts to leach downward and outward from the water source, subsurface irrigation causes resident and irrigated salts to flow upward through advection and accumulate above the dripline where plants are present (Hanson and Bendixen, 1995; Hopmans *et al.*, 2021). This accumulation pattern antagonizes the establishment of many row crops because germination is relatively sensitive to salt stress (Bernstein *et al.*, 1955). Such production systems rely on pre-season rain, sprinkler or surface irrigation to leach salts below the drip line where they may be leached downward by subsurface irrigation (Hanson and Bendixen, 1995). Shallow installation of subsurface drip lines is advantageous where sufficient pre-season rains are present as irrigating the soil surface may be avoided altogether (Hanson and Bendixen, 1995). This issue can be mechanically managed in processing tomato by adding soil to planting beds (Hanson *et al.*, 1995), followed by irrigation to

accumulate salts into the uppermost zone of the bed, which is subsequently removed and placed in the furrow between rows, where very little horizontal salt movement occurs (Hanson and Bendixen, 1995).

The strong localization of water application in drip irrigation questions the applicability of historical steady-state leaching models to micro-irrigation systems (Letey and Feng, 2007). These models insufficiently account for the highly local nature of micro-irrigation and underestimate both the local leaching fraction experienced by plants and the tolerable EC of irrigation water (Wallender and Tanji, 2011). Adequate management of heterogeneous salinity patterns and localized leaching under drip or micro-sprinkler may allow sustainable crop production in soils that would otherwise be deemed too saline for that species. Using transient models such as the HYDRUS model has been suggested as an alternative (Letey *et al.*, 2011). These models account for localized application of water and changes in flow rates over time by explicitly simulating two-dimensional (or even three-dimensional) water and solute transport in the root-zone by numerically solving mechanistic models. However, although these models are very strong in depicting physical transport processes, they often oversimplify the description of plant physiological processes governing water and solute uptake. For example, the HYDRUS model neglects that the distribution of water uptake is also affected by nutrient concentrations. Moreover, even if it was possible to perfectly simulate the water, nutrient and salinity dynamics for a given scenario, it would still be unclear how the calculated heterogeneous salinity distribution would translate into plant performance. Incorporating current knowledge of plant responses to heterogeneous conditions (both salinity and nutrient) might make these models more suitable for evaluating salinity management practices.

Challenge 2: How to simultaneously optimize N efficiency and minimize the impact of salinity. The necessity of a leaching fraction for long-term salinity management is coupled with the issue of nutrient loss, especially for nitrate (NO_3^-), which exhibits similar leaching potential as Cl^- . Any practice designed to remove Na^+ or Cl^- from the root-zone probably also leaches NO_3^- (Assouline *et al.*, 2015; Vaughan and Letey, 2015; Libutti and Monteleone, 2017). Although a common problem, few studies have addressed the integrated nature of salinity and nutrient management (Libutti and Monteleone, 2017). While NO_3^- and Cl^- are subject to very similar transport mechanisms and rates in the soil, their distribution in the soil can nevertheless be quite different, and high Na^+ and Cl^- concentrations do not necessarily coincide with high NO_3^- concentrations. This is because: (1) in contrast to Na^+ and Cl^- , NO_3^- is preferentially taken up by plant roots; and (2) nitrogen fertilizer is deliberately added to the irrigation water during fertigation and is to some degree independent of water (and therefore salt) application. Understanding crop nitrogen demands and responses to spatially localized nutrients and salinity may help manage fertigation systems to achieve the simultaneous goal of salinity leaching and minimal nitrate loss.

By providing nutrients through fertigation in a manner (rate, duration and timing during a fertigation event) that retains nutrients in the low-salinity zone adjacent to the drip-emitter, roots can avoid exploring the saline fringes of the wetted zones,

thus reducing salt exposure. HYDRUS-based modelling suggests that high-frequency applications of small amounts of nitrate, timed toward the end of a fertigation event, can help retain NO_3^- in the root-zone adjacent to the irrigation source while allowing salt (i.e. Na^+ and Cl^-) to be leached to the peripheral root-zone. Scheduling low but frequent NO_3^- applications, attuned to crop demand, allows the crop to take up most of the NO_3^- before it passes through the low-salinity zone into the saline fringes. Figure 7 simulates continuous NO_3^- application and a scenario which applies NO_3^- only every 10 d, while the total amount of NO_3^- applied is the same for both simulations. High-frequency applications of NO_3^- using drip irrigation increased N uptake efficiency in some cases (e.g. Scholberg et al., 2002; Quiñones et al., 2007).

Breeding targets

Challenging as they are, modern irrigation systems provide some leeway for controlling water and nutrient supply, to match plant demands. Nevertheless, genetic approaches to enhance salt tolerance are also needed.

Avenue 1: Breeding for root traits that facilitate water and nutrient uptake. Do specific root traits facilitate resource foraging

(water and nutrients) under heterogeneous soil salinities? With the current (or lack of) knowledge it is very difficult to address this question. As advocated for other marginal environments (Lynch, 2018, 2019; Schneider and Lynch, 2020; Colombi et al., 2022), root architectural traits and anatomical plasticity that reduce the metabolic cost of soil exploration might be beneficial in saline environments. Since heterogeneous salt distribution is probably associated with non-uniform water and nutrient distribution, careful consideration is necessary. For example, under saline conditions plants modify their root system architecture to reduce salt uptake (Julkowska et al., 2014) by reducing the length and density of root hairs and thus the overall absorption surface area (Shabala et al., 2003). However, marginal soils can also be highly deficient in phosphorus (P), and root hairs would be critical to allow root exploration beyond the root depletion zones and acquire P (and also other nutrients and water) from impoverished soil (Lynch, 2018, 2019; Rongsawat et al., 2021). Our ability to understand how plants could resolve such dilemma, and identify which root traits might be more favourable under heterogeneous salinities, is hindered by the simplicity of the experimental systems employed to date.

Avenue 2: Breeding for tissue tolerance. Traditionally, crop breeding for salinity tolerance has targeted Na^+ exclusion traits (Genc et al., 2010; Munns et al., 2012; Wu et al., 2019). This




FIG. 7. Simulated spatial distributions of salinity and nitrate following a growing season with an equal amount of nitrate applied (A) continuously and (B) once every 10 d for 8 h. The density of black dots represents the concentration of nitrate in the soil and the isolines indicate volumetric water content (---). This simulation accounts for plant uptake of nitrate and water over the growing season. The simulation was done using the software HYDRUS 2D (Šimunek et al., 2012) assuming a constant transpiration rate of 8 mm d^{-1} and no surface evaporation over a period of 75 d (D. Reineke and P. H. Brown, pers. comm.).

strategy comes with a caveat of a progressive build-up of Na⁺ in a root-zone (Liu *et al.*, 2020a), thus further exacerbating heterogeneity of Na⁺ distribution profiles in the rhizosphere and affecting water uptake and ultimately growth (Alharby *et al.*, 2014, 2018). Furthermore, this strategy requires a heavy reliance on energetically expensive *de novo* synthesis of compatible solutes for osmotic adjustment (Munns *et al.*, 2020). A viable alternative may be to target crop halophytism, e.g. a set of anatomical and physiological traits that allow plants to include significant amounts of Na⁺ in their tissues, without compromising their metabolic activity (Flowers and Colmer, 2015; Munns *et al.*, 2016). Amongst key traits conferring crop halophytism, vacuolar Na⁺ sequestration, ROS desensitization, tissue succulence and salt deposition in trichomes are considered as promising targets in breeding programmes (Liu *et al.*, 2020a).

Avenue 3: Understanding the nature of root to shoot signals. Plant biomass is ultimately proportional to the amount of CO₂ assimilated by the shoot that, in turn, is determined by the efficiency of stomata in balancing CO₂ gain and water loss via leaf transpiration. Root-borne signals play a critical role in coordinating plant gas exchange and optimizing plant water use efficiency. The signalling between roots and shoots integrates various signals (from electrical and hydraulic signals, Ca²⁺ and ROS waves to hormones, peptides and RNA; Gilroy *et al.*, 2016; Shabala *et al.*, 2016; Li *et al.*, 2021) that ultimately determine a plant's ability to adapt to saline conditions. Stress-induced elevations in ROS levels is accelerated in halophytes compared to glycophytes (Ellouzi *et al.*, 2011), and NADPH oxidase-mediated root-borne ROS signals induce early stomatal closure in salt-tolerant species (Niu *et al.*, 2018). Although shoot ABA levels increase within 30 min of salinity exposure, the magnitude of this increase appears to be species-specific (Geilfus *et al.*, 2015; Hedrich and Shabala, 2018). While xylem sap ABA concentrations also increase (Albacete *et al.*, 2008), grafting experiments with ABA-deficient mutants indicate this is shoot-mediated (Li *et al.*, 2018). Nevertheless, grafting wild-type tomato scions onto an ABA-overproducing rootstock enhanced salt tolerance, even if it was difficult to establish consistent evidence of root-to-shoot ABA signalling (Martínez-Andújar *et al.*, 2021). As discussed above, it remains uncertain how plants exposed to heterogeneous salinity integrate signals from exposed and non-exposed roots to regulate stomatal aperture. Since a multitude of signals interact during root-to-shoot communication, it is critical to understand how they confer plant stress tolerance.

Avenue 4: Understanding how water and ion transport are coupled. Another emerging topic is the possibility of water and ion flow coupling by aquaporins. Initially described as water and neutral solute channels, aquaporins can also transport ions across various cellular membranes (Byrt *et al.*, 2017; Qiu *et al.*, 2020). This discovery challenges current concepts that water and solutes move across membranes via separate pathways and may account for situations where water movement into the xylem goes against an apparent water potential gradient (Wegner, 2017; Tyerman *et al.*, 2021). Such coupling may be especially crucial for roots exposed to heterogeneous salinity as transcriptional changes are probably too slow to account for the highly dynamic external ionic environment. In this context, a phosphorylation-dependent switch between ion

and water permeation in aquaporins (and, specifically, PIP2;1) might enable plant cells to rapidly adjust to altered ionic conditions in the rhizosphere and optimize ion transport at minimal energy cost. This option implies that plants that rely on Na⁺ accumulation for osmotic adjustment and thus water uptake under hypersaline soil conditions can tolerate tissue Na⁺ loads, thereby avoiding cytotoxicity, requiring that halophytic traits be incorporated into modern elite varieties.

CONCLUDING REMARKS

There are several constraints in conducting experiments or genotypic selection of plants under conditions that do not reflect real agricultural conditions of heterogeneous soil environments. Homogeneous conditions limit more complex interactions between salinity prevalence and plant responses such as exclusion of saline ions, nutrient and water uptake, root architecture, or adjustment to varying pH within the root-zone. As highlighted above, fundamental questions remain on plant responses to heterogeneous salinities and how this is affected by the associated variations in water and nutrient distribution. We call for a greater focus on understanding plant responses to heterogeneous soil salinity, which should be considered as the next frontier for salinity research and land management. Understanding responses to heterogeneous saline conditions holds significant promise for identifying new breeding targets for crop salt tolerance and adequate management practices of saline environments, which will accelerate the implementation of solutions to improve the productive use of saline land.

ACKNOWLEDGMENTS

This review was instigated from the workshop on 'Plant responses to heterogeneous salinity' held at the University of Western Australia, 12–16 December 2016, and we thank all participants for their contributions and discussion.

FUNDING

E.B.L., T.C., I.C.D., P.B. and N.B. thank the University of Western Australia for funding the workshop. I.C.D. thanks the European Union (Project 773903 – SHui) for supporting ongoing work on split-root plants. N.B. was a recipient of the Marie Curie Fellowship funded by the European Union (grant 700001).

LITERATURE CITED

- Adeoye KB, Rawlins SL. 1981. A split-root technique for measuring root water potential. *Plant Physiology* **68**: 44–47.
- Ahmadzai H, Tutundjian S, Elouafi I. 2021. Policies for sustainable agriculture and livelihood in marginal lands: a review. *Sustainability* **13**: 8692.
- Ahmed MA, Zarebanadkouki M, Meunier F, Javaux M, Kaestner A, Carminati A. 2018. Root type matters: measurement of water uptake by seminal, crown, and lateral roots in maize. *Journal of Experimental Botany* **69**: 1199199–1191206.
- Albacete A, Ghanem ME, Martínez-Andújar C, *et al.* 2008. Hormonal changes in relation to biomass partitioning and shoot growth

- impairment in salinized tomato (*Solanum lycopersicum* L.) plants. *Journal of Experimental Botany* **59**: 4119–4131.
- Aldabaa AAA, Weindorf DC, Chakraborty S, Sharma A, Li B. 2015.** Combination of proximal and remote sensing methods for rapid soil salinity quantification. *Geoderma* **239–240**: 34–46.
- Alharby HF, Colmer TD, Barrett-Lennard EG. 2014.** Salt accumulation and depletion in the root-zone of the halophyte *Atriplex nummularia* Lindl.: influence of salinity, leaf area and plant water use. *Plant and Soil* **382**: 31–41.
- Alharby HF, Colmer TD, Barrett-Lennard EG. 2018.** Salinization of the soil solution decreases the further accumulation of salt in the root zone of the halophyte *Atriplex nummularia* Lindl. growing above shallow saline groundwater. *Plant, Cell & Environment* **41**: 99–110.
- Assouline S, Russo D, Silber A, Or D. 2015.** Balancing water scarcity and quality for sustainable irrigated agriculture. *Journal of the American Water Resources Association* **51**: 3419–3436.
- Attia H, Nouaïli S, Soltani A, Lachaâl M. 2009.** Comparison of the responses to NaCl stress of two pea cultivars using split-root system. *Scientia Horticulturae* **123**: 164–169.
- Ayars JE. 2012.** On-farm irrigation and drainage practices. In: Wallender WW, Tanji K, eds. *Agricultural salinity assessment and management*. ASCE manual and reports on engineering practices, 2nd ed. Reston, VA: American Society of Civil Engineers, 511–538.
- Ayars JE, Hutmacher RB, Schoneman RA, Vail SS, Pflaum T. 1993.** Long term use of saline water for irrigation. *Irrigation Science* **14**: 27–34.
- Ayers RS, Westcot DW. 1985.** *Water quality for agriculture*. Rome: FAO Irrigation and Drainage, Paper 29, Food and Agriculture Organization.
- Babourina O, Hawkins B, Lew RR, Newman I, Shabala S. 2001.** K⁺ transport by *Arabidopsis* root hairs at low pH. *Australian Journal of Plant Physiology* **28**: 635–641.
- Barberon M, Vermeer JEM, De Bellis D, et al. 2016.** Adaptation of root function by nutrient-induced plasticity of endodermal differentiation. *Cell* **164**: 447–459.
- Barrett-Lennard EG, Malcolm CV. 2000.** Increased concentrations of chloride beneath stands of saltbushes (*Atriplex* species) suggest substantial use of groundwater. *Australian Journal of Experimental Agriculture* **39**: 949–955.
- Bar-Yosef B. 1999.** *Advances in fertigation*. In: Sparks DLBTAiA, ed. New York: Academic Press.
- Bazihizina N, Barrett-Lennard EG, Colmer TD. 2012a.** Plant growth and physiology under heterogeneous salinity. *Plant and Soil* **354**: 1–19.
- Bazihizina N, Barrett-Lennard EG, Colmer TD. 2012b.** Plant responses to heterogeneous salinity: growth of the halophyte *Atriplex nummularia* is determined by the root-weighted mean salinity of the root zone. *Journal of Experimental Botany* **63**: 6347–6358.
- Bazihizina N, Colmer TD, Barrett-Lennard EG. 2009.** Response to non-uniform salinity in the root zone of the halophyte *Atriplex nummularia*: growth, photosynthesis, water relations and tissue ion concentrations. *Annals of Botany* **104**: 737–745.
- Bernstein L. 1975.** Effects of salinity and sodicity on plant growth. *Annual Review of Phytopathology* **13**: 295–312.
- Bernstein L, Fireman M. 1957.** Laboratory studies on salt distribution in furrow-irrigated soil with special reference to the pre-emergence period. *Soil Science* **83**: 249–264.
- Bernstein L, MacKenzie AJ, Krantz BA. 1955.** The interaction of salinity and planting practice on the germination of irrigated row crops. *Soil Science Society of America Journal* **19**: 240–243.
- Bleby TM, Aucote M, Kennett-Smith AK, Walker GR, Schachtman DP. 1997.** Seasonal water use characteristics of tall wheatgrass (*Agropyron elongatum* (Host) Beauv) in a saline environment. *Plant Cell and Environment* **20**: 1361–1371.
- Bogunovic I, Trevisani S, Seput M, Juzbasic D, Durdevic B. 2017.** Short-range and regional spatial variability of soil chemical properties in an agro-ecosystem in eastern Croatia. *CATENA* **154**: 50–62.
- Byrt CS, Munns R, Burton RA, Gilliham M, Wege S. 2018.** Root cell wall solutions for crop plants in saline soils. *Plant Science* **269**: 47–55.
- Byrt CS, Zhao M, Kourghi M, et al. 2017.** Non-selective cation channel activity of aquaporin *AtPIP2;1* regulated by Ca²⁺ and pH. *Plant, Cell & Environment* **40**: 802–815.
- Chen S, Wang Z, Guo X, et al. 2019.** Effects of vertically heterogeneous soil salinity on tomato photosynthesis and related physiological parameters. *Scientia Horticulturae* **249**: 120–130.
- Choi W-G, Toyota M, Kim S-H, Hilleary R, Gilroy S. 2014.** Salt stress-induced Ca²⁺ waves are associated with rapid, long-distance root-to-shoot signaling in plants. *Proceedings of the National Academy of Sciences* **111**: 6497–6502.
- Choukr-Allah R. 2021.** Use and management of saline water for irrigation in the Near East and North Africa (NENA) region. In: Negacz K, Vellinga P, Barrett-Lennard E, Choukr-Allah R, Elzenga T, eds. *Future of sustainable agriculture in saline environments*. Boca Raton: CRC Press.
- Christmann A, Weiler EW, Steudle E, Grill E. 2007.** A hydraulic signal in root-to-shoot signalling of water shortage. *The Plant Journal* **52**: 167–174.
- Colombi T, Chakrawal A, Herrmann AM. 2022.** Carbon supply–consumption balance in plant roots: effects of carbon use efficiency and root anatomical plasticity. *New Phytologist* **233**: 1542–1547.
- Corwin DL. 2021.** Climate change impacts on soil salinity in agricultural areas. *European Journal of Soil Science* **72**: 842–862.
- Cui B, Liu R, Flowers TJ, Song J. 2021.** Casparian bands and suberin lamellae: Key targets for breeding salt tolerant crops? *Environmental and Experimental Botany* **191**: 104600.
- Davidson NJ, Galloway R, Lazarescu G. 1996.** Growth of *Atriplex amnicola* on salt-affected soils in Western Australia. *Journal of Applied Ecology* **33**: 1257–1266.
- Dinneny JR. 2019.** Developmental responses to water and salinity in root systems. *Annual Review of Cell and Developmental Biology* **35**: 239–257.
- Dodd AN, Kudla J, Sanders D. 2010.** The language of calcium signaling. *Annual Review of Plant Biology* **61**: 593–620.
- Dodd IC, Egea G, Davies WJ. 2008.** Accounting for sap flow from different parts of the root system improves the prediction of xylem ABA concentration in plants grown with heterogeneous soil moisture. *Journal of Experimental Botany* **59**: 4083–4093.
- Duan L, Dietrich D, Ng CH, et al. 2013.** Endodermal ABA signaling promotes lateral root quiescence during salt stress in *Arabidopsis* seedlings. *The Plant Cell* **25**: 324–341.
- Ellouzi H, Ben Hamed K, Cela J, Munné-Bosch S, Abdely C. 2011.** Early effects of salt stress on the physiological and oxidative status of *Cakile maritima* (halophyte) and *Arabidopsis thaliana* (glycophyte). *Physiologia Plantarum* **142**: 128–143.
- Ericsson T. 1995.** Growth and shoot: root ratio of seedlings in relation to nutrient availability. *Plant and Soil* **168–169**: 205–214.
- Evans MJ, Choi W-G, Gilroy S, Morris RJ. 2016.** A ROS-assisted calcium wave dependent on the *ATRBOHD* NADPH oxidase and *TPC1* cation channel propagates the systemic response to salt stress. *Plant Physiology* **171**: 1771–1784.
- Feng W, Kita D, Peaucelle A, et al. 2018.** The FERONIA receptor kinase maintains cell-wall integrity during salt stress through Ca²⁺ signaling. *Current Biology* **28**: 666–675.e5.
- Feng X, An P, Guo K, Li X, Liu X, Zhang X. 2017.** Growth, root compensation and ion distribution in *Lycium chinense* under heterogeneous salinity stress. *Scientia Horticulturae* **226**: 24–32.
- Feng X, Hussain T, Guo K, An P, Liu X. 2021.** Physiological, morphological and anatomical responses of *Hibiscus moscheutos* to non-uniform salinity stress. *Environmental and Experimental Botany* **182**: 104301.
- Fichman Y, Mittler R. 2020.** Rapid systemic signaling during abiotic and biotic stresses: is the ROS wave master of all trades? *The Plant Journal* **102**: 887–896.
- Fichman Y, Mittler R. 2021.** Integration of electric, calcium, reactive oxygen species and hydraulic signals during rapid systemic signaling in plants. *The Plant Journal* **107**: 7–20.
- Flores P, Angeles Botella M, Martínez V, Cerdá A. 2002.** Response to salinity of tomato seedlings with a split-root system: nitrate uptake and reduction. *Journal of Plant Nutrition* **25**: 177–187.
- Flowers TJ, Colmer TD. 2015.** Plant salt tolerance: adaptations in halophytes. *Annals of Botany* **115**: 327–331.
- Gambetta GA, Knipfer T, Fricke W, McElrone AJ. 2017.** Aquaporins and root water uptake. In: Chaumont F, Tyerman SD, eds. *Plant aquaporins: from transport to signaling*. Cham: Springer International Publishing.
- Geilfus CM, Mithöfer A, Ludwig-Müller J, Zörb C, Muehling Karl H. 2015.** Chloride-inducible transient apoplastic alkalizations induce stomata closure by controlling abscisic acid distribution between leaf apoplast and guard cells in salt-stressed *Vicia faba*. *New Phytologist* **208**: 803–816.
- Genç Y, Oldach K, Verbyla AP, et al. 2010.** Sodium exclusion QTL associated with improved seedling growth in bread wheat under salinity stress. *Theoretical and Applied Genetics* **121**: 877–894.

- Gigli-Bisceglia N, Engelsdorf T, Hamann T. 2020. Plant cell wall integrity maintenance in model plants and crop species-relevant cell wall components and underlying guiding principles. *Cellular and Molecular Life Sciences* 77: 2049–2077.
- Gilroy S, Bialasek M, Suzuki N, et al. 2016. ROS, calcium, and electric signals: key mediators of rapid systemic signaling in plants. *Plant Physiology* 171: 1606.
- Gloser V, Libera K, Orians CM. 2008. Contrasting below- and aboveground responses of two deciduous trees to patchy nitrate availability. *Tree Physiology* 28: 37–44.
- Gorska A, Ye Q, Holbrook NM, Zwieniecki MA. 2008. Nitrate control of root hydraulic properties in plants: translating local information to whole plant response. *Plant Physiology* 148: 1159–1167.
- Grismer M, Gates T, Hanson B. 1988. Irrigation and drainage strategies in salinity problem areas. *California Agriculture* 42: 23–24.
- Hanson B, May D. 2011. *Drip irrigation salinity management for row crops*. Publication 8447, University of California, 1–13.
- Hanson BR, Bendixen WE. 1995. Drip irrigation controls soil salinity under row crops. *California Agriculture* 49: 19–23.
- Hanson BR, May D, Simnek J, Hopmans JW, Hutmacher RB. 2009. Drip irrigation provides the salinity control needed for profitable irrigation of tomatoes in the San Joaquin Valley. *California Agriculture* 63: 131–136.
- Hedrich R, Shabala S. 2018. Stomata in a saline world. *Current Opinion in Plant Biology* 46: 87–95.
- Heuperman AF. 1995. *Salt and water dynamics beneath a tree plantation growing on a shallow watertable*. Technical Report. Tatura, Victoria: Institute of Sustainable Irrigated Agriculture.
- Hillel D. 1980. *Fundamental of soil physics*. New York: Academic Press.
- Hodge A. 2006. Plastic plants and patchy soils. *Journal of Experimental Botany* 57: 401–411.
- Hoffmann M, Johnsson H. 2000. Nitrogen leaching from agricultural land in Sweden: model calculated effects of measures to reduce leaching loads. *Ambio* 29: 67–73.
- Hopmans JW, Qureshi AS, Kisekka I, et al. 2021. Critical knowledge gaps and research priorities in global soil salinity. *Advances in Agronomy* 169: 1–191.
- Hussain N, Ghaffar A, Zafar ZU, et al. 2021. Identification of novel source of salt tolerance in local bread wheat germplasm using morpho-physiological and biochemical attributes. *Scientific Reports* 11: 10854.
- Jackson RB, Caldwell MM. 1996. Integrating resource heterogeneity and plant plasticity: modelling nitrate and phosphate uptake in a patchy soil environment. *Journal of Ecology* 84: 891–903.
- Jiang Z, Zhou X, Tao M, et al. 2019. Plant cell-surface GIPC sphingolipids sense salt to trigger Ca²⁺ influx. *Nature* 572: 341–346.
- Julkowska MM, Hoefsloot HCJ, Mol S, et al. 2014. Capturing Arabidopsis root architecture dynamics with root-fit reveals diversity in responses to salinity. *Plant Physiology* 166: 1387–1402.
- Khanna M, Chen L, Basso B, et al. 2021. Redefining marginal land for bio-energy crop production. *GCB Bioenergy* 13: 1590–1609.
- Kong X, Luo Z, Dong H, Eneji AE, Li W. 2012. Effects of non-uniform root zone salinity on water use, Na⁺ recirculation, and Na⁺ and H⁺ flux in cotton. *Journal of Experimental Botany* 63: 2105–2116.
- Kong X, Luo Z, Dong H, Eneji AE, Li W. 2016. H₂O₂ and ABA signaling are responsible for the increased Na⁺ efflux and water uptake in *Gossypium hirsutum* L. roots in the non-saline side under non-uniform root zone salinity. *Journal of Experimental Botany* 67: 2247–2261.
- Kong X, Luo Z, Dong H, Li W, Chen Y. 2017. Non-uniform salinity in the root zone alleviates salt damage by increasing sodium, water and nutrient transport genes expression in cotton. *Scientific Reports* 7: 2879.
- Krishnamurthy P, Ranathunge K, Nayak S, Schreiber L, Mathew MK. 2011. Root apoplastic barriers block Na⁺ transport to shoots in rice (*Oryza sativa* L.). *Journal of Experimental Botany* 62: 4215–4228.
- Lainé P, Ourry A, Boucaud J. 1995. Shoot control of nitrate uptake rates by roots of *Brassica napus* L.: effects of localized nitrate supply. *Planta* 196: 77–83.
- Lainé P, Ourry A, Boucaud J, Salette J. 1998. Effects of a localized supply of nitrate on NO₃⁻ uptake rate and growth of roots in *Lolium multiflorum* Lam. *Plant and Soil* 202: 61–67.
- Letey J, Feng GL. 2007. Dynamic versus steady-state approaches to evaluate irrigation management of saline waters. *Agricultural Water Management* 91: 1–10.
- Letey J, Hoffman GJ, Hopmans JW, et al. 2011. Evaluation of soil salinity leaching requirement guidelines. *Agricultural Water Management* 98: 502–506.
- Li C, Li Y, Ma J. 2011. Spatial heterogeneity of soil chemical properties at fine scales induced by *Haloxylon ammodendron* (*Chenopodiaceae*) plants in a sandy desert. *Ecological Research* 26: 385–394.
- Li H, Testerink C, Zhang Y. 2021. How roots and shoots communicate through stressful times. *Trends in Plant Science* 26: 940–952.
- Li W, de Ollas C, Dodd IC. 2018. Long-distance ABA transport can mediate distal tissue responses by affecting local ABA concentrations. *Journal of Integrative Plant Biology* 60: 16–33.
- Libutti A, Monteleone M. 2017. Soil vs. groundwater: The quality dilemma. Managing nitrogen leaching and salinity control under irrigated agriculture in Mediterranean conditions. *Agricultural Water Management* 186: 40–50.
- Liu M, Pan T, Allakhverdiev SI, Yu M, Shabala S. 2020a. Crop halophytism: an environmentally sustainable solution for global food security. *Trends in Plant Science* 25: 630–634.
- Liu M, Yu H, Ouyang B, et al. 2020b. NADPH oxidases and the evolution of plant salinity tolerance. *Plant, Cell & Environment* 43: 2957–2968.
- López-Bucio J, Cruz-Ramírez A, Herrera-Estrella L. 2003. The role of nutrient availability in regulating root architecture. *Current Opinion in Plant Biology* 6: 280–287.
- Lycoskoufis IH, Savvas D, Mavrogianopoulos G. 2005. Growth, gas exchange, and nutrient status in pepper (*Capsicum annuum* L.) grown in recirculating nutrient solution as affected by salinity imposed to half of the root system. *Scientia Horticulturae* 106: 147–161.
- Lynch JP. 2018. Rightsizing root phenotypes for drought resistance. *Journal of Experimental Botany* 69: 3279–3292.
- Lynch JP. 2019. Root phenotypes for improved nutrient capture: an under-exploited opportunity for global agriculture. *New Phytologist* 223: 548–564.
- Maas EV, Grattan SR. 1999. Crop yields as affected by salinity. In: Skaggs RW, van Schilfegaarde J, eds. *Agricultural drainage agronomy monograph*. Madison: ASA.
- Maas EV, Hoffman GJ. 1977. Crop salt tolerance – current assessment. *Journal of the Irrigation and Drainage Division* 103: 115–134.
- Mantovani D, Veste M, Bohm C, Vignudelli M, Freese D. 2015. Spatial and temporal variation of drought impact on black locust (*Robinia pseudoacacia* L.) water status and growth. *iForest – Biogeosciences and Forestry* 8: 743–747.
- Manzi M, Lado J, Rodrigo MJ, Zacarias L, Arbona V, Gómez-Cadenas A. 2015. Root ABA accumulation in long-term water-stressed plants is sustained by hormone transport from aerial organs. *Plant and Cell Physiology* 56: 2457–2466.
- Martínez-Andújar C, MartíMartínez-Andúñez-Pérez A, Albacete A, et al. 2021. Overproduction of ABA in rootstocks alleviates salinity stress in tomato shoots. *Plant, Cell & Environment* 44: 2966–2986.
- Maurel C, Nacry P. 2020. Root architecture and hydraulics converge for acclimation to changing water availability. *Nature Plants* 6: 744–749.
- Maurel C, Verdoucq L, Luu D-T, Santoni V. 2008. Plant aquaporins: membrane channels with multiple integrated functions. *Annual Review of Plant Biology* 59: 595–624.
- May D, Hanson B. 2006. Is drip irrigation sustainable in the salt affected soil of the San Joaquin Valley of California? *WIT Transactions on Ecology and the Environment* 96.
- McAdam SAM, Brodribb TJ, Ross JJ. 2016. Shoot-derived abscisic acid promotes root growth. *Plant, Cell & Environment* 39: 652–659.
- McFarlane DJ, George RJ, Barrett-Lennard EG, Gilfedder M. 2016. Salinity in dryland agricultural systems: challenges and opportunities. In: Farooq M, Siddique K, eds. *Innovations in dryland agriculture*. Cham: Springer International Publishing.
- Mensforth LJ, Walker GR. 1996. Root dynamics of *Melaleuca halmaturorum* in response to fluctuating saline groundwater. *Plant and Soil* 184: 75–84.
- Meunier F, Couvreur V, Draye X, Vanderborcht J, Javaux M. 2017. Towards quantitative root hydraulic phenotyping: novel mathematical functions to calculate plant-scale hydraulic parameters from root system functional and structural traits. *Journal of Mathematical Biology* 75: 1133–1170.
- Mulholland BJ, Fussell M, Edmondson RN, et al. 2002. The effect of split-root salinity stress on tomato leaf expansion, fruit yield and quality. *The Journal of Horticultural Science and Biotechnology* 77: 509–519.
- Munns R, Day DA, Fricke W, et al. 2020. Energy costs of salt tolerance in crop plants. *New Phytologist* 225: 1072–1090.
- Munns R, James RA, Gilliam M, Flowers TJ, Colmer TD. 2016. Tissue tolerance: an essential but elusive trait for salt-tolerant crops. *Functional Plant Biology* 43: 1103–1113.

- Munns R, James RA, Xu B, et al. 2012. Wheat grain yield on saline soils is improved by an ancestral Na⁺ transporter gene. *Nature Biotechnology* 30: 360–364.
- Nakamura R, Kachi N, Suzuki J-I. 2008. Root growth and plant biomass in *Lolium perenne* exploring a nutrient-rich patch in soil. *Journal of Plant Research* 121: 547–557.
- Niu M, Huang Y, Sun S, et al. 2018. Root respiratory burst oxidase homologue-dependent H₂O₂ production confers salt tolerance on a grafted cucumber by controlling Na⁺ exclusion and stomatal closure. *Journal of Experimental Botany* 69: 3465–3476.
- Northey JE, Christen EW, Ayars JE, Jankowski J. 2006. Occurrence and measurement of salinity stratification in shallow groundwater in the Murrumbidgee Irrigation Area, south-eastern Australia. *Agricultural Water Management* 81: 23–40.
- Nulsen RA, Thurtell GW. 1980. Effects of osmotica around the roots on water uptake by maize plants. *Functional Plant Biology* 7: 27–34.
- Orians CM, Jones CG. 2001. Plants as resource mosaics: a functional model for predicting patterns of within-plant resource heterogeneity to consumers based on vascular architecture and local environmental variability. *Oikos* 94: 493–504.
- Pancaldi F, Trindade LM. 2020. Marginal lands to grow novel bio-based crops: a plant breeding perspective. *Frontiers in Plant Science* 11: 227.
- Pannell DJ, Ewing MA. 2006. Managing secondary dryland salinity: options and challenges. *Agricultural Water Management* 80: 41–56.
- Passioura JB, Frere MH. 1967. Numerical analysis of the convection and diffusion of solutes to roots. *Soil Research* 5: 149–159.
- Peck S, Mittler R. 2020. Plant signaling in biotic and abiotic stress. *Journal of Experimental Botany* 71: 1649–1651.
- Pottosin I, Zepeda-Jazo I. 2018. Powering the plasma membrane Ca²⁺-ROS self-amplifying loop. *Journal of Experimental Botany* 69: 3317–3320.
- Qiu J, McGaughey SA, Groszmann M, Tyerman SD, Byrt CS. 2020. Phosphorylation influences water and ion channel function of AtPIP2;1. *Plant, Cell & Environment* 43: 2428–2442.
- Quiñones A, Martínez-Alcántara B, Legaz F. 2007. Influence of irrigation system and fertilization management on seasonal distribution of N in the soil profile and on N-uptake by citrus trees. *Agriculture, Ecosystems and Environment* 122: 399–409.
- Raine SR, Meyer WS, Rassam DW, Hutson JL, Cook FJ. 2007. Soil–water and solute movement under precision irrigation: knowledge gaps for managing sustainable root zones. *Irrigation Science* 26: 91–100.
- Rajaniemi TK. 2007. Root foraging traits and competitive ability in heterogeneous soils. *Oecologia* 153: 145–152.
- Razzaq A, Wani SH, Saleem F, Yu M, Zhou M, Shabala S. 2021. Rewilding crops for climate resilience: economic analysis and *de novo* domestication strategies. *Journal of Experimental Botany* 72: 6123–6139.
- Reyllán-Álvarez R, Lobet G, Dinneny JR. 2016. Environmental control of root system biology. *Annual Review of Plant Biology* 67: 619–642.
- Riaz M, Arif MS, Ashraf MA, et al. 2018. A comprehensive review on rice responses and tolerance to salt stress. In: Hasanuzzaman M, Fujita M, Nahar K, Biswas JK, eds. *Advances in rice research for abiotic stress tolerance*. Sawston: Woodhead Publishing, 133–158.
- Robbins CW, Wagenet RJ, Jurinak JJ. 1980. A combined salt transport-chemical equilibrium model for calcareous and gypsiferous soils. *Soil Science Society of America Journal* 44: 1191–1194.
- Rongsawat T, Peltier J-B, Boyer J-C, Véry A-A, Sentenac H. 2021. Looking for root hairs to overcome poor soils. *Trends in Plant Science* 26: 83–94.
- Rui Y, Dinneny JR. 2020. A wall with integrity: surveillance and maintenance of the plant cell wall under stress. *New Phytologist* 225: 1428–1439.
- Ruto E, Tzemi D, Gould I, Bosworth G. 2021. Economic impact of soil salinization and the potential for saline agriculture. In: Negacz K, Vellinga P, Barrett-Lennard E, Choukr-Allah R, Elzenga T, eds. *Future of sustainable agriculture in saline environments*. Boca Raton: CRC Press.
- Schneider HM, Lynch JP. 2020. Should root plasticity be a crop breeding target? *Frontiers in Plant Science* 11: 546.
- Scholberg JMS, Parsons LR, Wheaton TA, McNeal BL, Morgan KT. 2002. Soil temperature, nitrogen concentration, and residence time affect nitrogen uptake efficiency in Citrus. *Journal of Environmental Quality* 31: 759–768.
- Schwantes AM, Parolari AJ, Swenson JJ, et al. 2018. Accounting for landscape heterogeneity improves spatial predictions of tree vulnerability to drought. *New Phytologist* 220: 132–146.
- Setter TL, Waters I, Stefanova K, Munns R, Barrett-Lennard EG. 2016. Salt tolerance, date of flowering and rain affect the productivity of wheat and barley on rainfed saline land. *Field Crops Research* 194: 31–42.
- Shabala S, Bose J, Fuglsang AT, Pottosin I. 2016. On a quest for stress tolerance genes: membrane transporters in sensing and adapting to hostile soils. *Journal of Experimental Botany* 67: 1015–1031.
- Shabala S, Shabala L, Volkenburgh EV. 2003. Effect of calcium on root development and root ion fluxes in salinised barley seedlings. *Functional Plant Biology* 30: 507–514.
- Shabala S, Wu H, Bose J. 2015. Salt stress sensing and early signalling events in plant roots: current knowledge and hypothesis. *Plant Science* 241: 109–119.
- Shukla AK, Sinha NK, Tiwari PK, et al. 2017. Spatial distribution and management zones for sulphur and micronutrients in Shivalik Himalayan region of India. *Land Degradation & Development* 28: 959–969.
- Silvestri S, Defina A, Marani M. 2005. Tidal regime, salinity and salt marsh plant zonation. *Estuarine, Coastal and Shelf Science* 62: 119–130.
- Simunek J, Van Genuchten MT, Sejna M. 2012. HYDRUS: Model use, calibration, and validation. *Transactions of the ASABE* 55: 1263–1274.
- Sinha BK, Singh NT. 1974. Effect of transpiration rate on salt accumulation around corn roots in a saline soil. *Agronomy Journal* 66: 557–560.
- Sinha BK, Singh NT. 1976. Salt distribution around roots of wheat under different transpiration rates. *Plant and Soil* 44: 141–147.
- Slatyer RO. 1967. *Plant water relationships*. London: Academic Press.
- Sonneveld C, Voogt W. 1990. Response of tomatoes (*Lycopersicon esculentum*) to an unequal distribution of nutrients in the root environment. *Plant and Soil* 124: 251–256.
- Stirzaker RJ, Passioura JB. 1996. The water relations of the root–soil interface. *Plant, Cell & Environment* 19: 201–208.
- Sun J, Yang G, Zhang W, Zhang Y. 2016. Effects of heterogeneous salinity on growth, water uptake, and tissue ion concentrations of alfalfa. *Plant and Soil* 408: 211–226.
- Tabata R, Sumida K, Yoshii T, Ohyama K, Shinohara H, Matsubayashi Y. 2014. Perception of root-derived peptides by shoot LRR-RKs mediates systemic N-demand signaling. *Science* 346: 343–346.
- Tabatabaei SJ, Gregory P, Hadley P, Ho L. 2004. Use of unequal salinity in the root zone to improve yield and quality in hydroponically grown tomato. *Acta Horticulturae* 648: 47–54.
- Tanji KK, Kielen NC. 2002. *Agricultural drainage water management in arid and semi-arid areas*. Rome: Food and Agriculture Organization of the United Nations.
- Thompson AJ, Mulholland BJ, Jackson AC, et al. 2007. Regulation and manipulation of ABA biosynthesis in roots. *Plant, Cell & Environment* 30: 67–78.
- Tiggeloven T, de Moel H, Winsemius HC, et al. 2020. Global-scale benefit–cost analysis of coastal flood adaptation to different flood risk drivers using structural measures. *Natural Hazards and Earth System Sciences* 20: 1025–1044.
- Tyerman SD, McGaughey SA, Qiu J, Yool AJ, Byrt CS. 2021. Adaptable and multifunctional ion-conducting aquaporins. *Annual Review of Plant Biology* 72: 703–736.
- Valenzuela F. 2018. *Effects of nonuniform root-zone salinity on Na⁺, Cl⁻ and nitrate uptake in irrigated almond*. PhD Dissertation, The University of California Davis.
- Valenzuela F, Fuentes A, Brown PH. 2021. Heterogeneous saline and nutritional conditions in the root-zone and its effect in water and nutrient uptake. *Acta Horticulturae* 1333: 59–68.
- Vaughan P, Letey J. 2015. Irrigation water amount and salinity dictate nitrogen requirement. *Agricultural Water Management* 157: 6–11.
- Vellinga P, Barrett-Lennard E. 2021. Saline agriculture as a way to adapt to sea level rise. In: Negacz K, Vellinga P, Barrett-Lennard E, Choukr-Allah R, Elzenga T, eds. *Future of sustainable agriculture in saline environments*. Boca Raton: CRC press.
- Wallender WW, Tanji KK. 2011. *Agricultural salinity assessment and management*. Reston: American Society of Civil Engineers (ASCE).
- Wang P, Calvo-Polanco M, Reyt G, et al. 2019. Surveillance of cell wall diffusion barrier integrity modulates water and solute transport in plants. *Scientific Reports* 9: 4227.
- Wang W, Cai L, Long Z, Zhang X, Zhao F. 2021. Effects of non-uniform salt stress on growth, yield, and quality of tomato. *Soil Science and Plant Nutrition* 67: 545–556.
- Wegner LH. 2017. Cotransport of water and solutes in plant membranes: The molecular basis, and physiological functions. *AIMS Biophysics* 4: 192–209.

- Wu H, Shabala L, Zhou M, et al. 2019.** Root vacuolar Na⁺ sequestration but not exclusion from uptake correlates with barley salt tolerance. *The Plant Journal* **100**: 55–67.
- Xing X, Du W, Ma X. 2019.** Field-scale distribution and heterogeneity of soil salinity in the mulched-drip-irrigation cotton field. *Archives of Agronomy and Soil Science* **65**: 1248–1261.
- Xiong X, Liu N, Wei Y-Q, et al. 2018.** Effects of non-uniform root zone salinity on growth, ion regulation, and antioxidant defense system in two alfalfa cultivars. *Plant Physiology and Biochemistry* **132**: 434–444.
- Xiong X, Wei Y-Q, Chen J-H, Liu N, Zhang Y-J. 2020.** Transcriptome analysis of genes and pathways associated with salt tolerance in alfalfa under non-uniform salt stress. *Plant Physiology and Biochemistry* **151**: 323–333.
- Xu L, Du H, Zhang X. 2019.** . Spatial distribution characteristics of soil salinity and moisture and its influence on agricultural irrigation in the Ili River Valley, China. *Sustainability*, **11**: 7142.
- Xu X, Huang G, Sun C, et al. 2013.** Assessing the effects of water table depth on water use, soil salinity and wheat yield: Searching for a target depth for irrigated areas in the upper Yellow River basin. *Agricultural Water Management* **125**: 46–60.
- Yakir D, Yechieli Y. 1995.** Plant invasion of newly exposed hypersaline Dead Sea shores. *Nature* **374**: 803–805.
- Yoshimura K, Iida K, Iida H. 2021.** MCAs in Arabidopsis are Ca²⁺-permeable mechanosensitive channels inherently sensitive to membrane tension. *Nature Communications* **12**: 6074.
- Yu Z, Duan X, Luo L, Dai S, Ding Z, Xia G. 2020.** How plant hormones mediate salt stress responses. *Trends in Plant Science* **25**: 1117–1130.
- Yuan F, Yang H, Xue Y, et al. 2014.** OSCA1 mediates osmotic-stress-evoked Ca²⁺ increases vital for osmosensing in Arabidopsis. *Nature* **514**: 367–371.
- Zaman M, Shahid S, Heng L. 2018.** Irrigation systems and zones of salinity development. In: Zaman M, Shahid S, Heng L, eds. *Guideline for salinity assessment, mitigation and adaptation using nuclear and related techniques*. Cham: SpringerOpen.
- Zekri M, Parsons LR. 1990.** Response of split-root sour orange seedlings to NaCl and polyethylene-glycol stresses. *Journal of Experimental Botany* **41**: 35–40.
- Zhang XC, Chen LM, Wu HH, Liu LL, Wan XC. 2018.** Root plasma membrane H⁺-ATPase is involved in low pH-inhibited nitrogen accumulation in tea plants (*Camellia sinensis* L.). *Plant Growth Regulation* **86**: 423–432.
- Zhang H, Wang R, Wang H, et al. 2019.** Heterogeneous root zone salinity mitigates salt injury to *Sorghum bicolor* (L.) Moench in a split-root system. *PLoS One* **14**: e0227020.

