

UCLA

UCLA Previously Published Works

Title

Structure-neurotoxicity relationships of amyloid β -protein oligomers

Permalink

<https://escholarship.org/uc/item/7tj7v363>

Journal

Proceedings of the National Academy of Sciences of the United States of America, 106(35)

ISSN

0027-8424

Authors

Ono, Kenjiro
Condron, Margaret M
Teplow, David B

Publication Date

2009-09-01

DOI

10.1073/pnas.0905127106

Peer reviewed

Structure–neurotoxicity relationships of amyloid β -protein oligomers

Kenjiro Ono^{a,b}, Margaret M. Condrón^a, and David B. Teplow^{a,1}

^aDepartment of Neurology, David Geffen School of Medicine, and Molecular Biology Institute and Brain Research Institute, University of California, Los Angeles, CA 90095; and ^bDepartment of Neurology and Neurobiology and Aging, Kanazawa University Graduate School of Medical Science, Kanazawa 920-8640, Japan

Edited by H. Eugene Stanley, Boston University, Boston, MA, and approved July 6, 2009 (received for review May 11, 2009)

Amyloid β -protein ($A\beta$) oligomers may be the proximate neurotoxins in Alzheimer's disease (AD). "Oligomer" is an ill-defined term because many kinds have been reported and they often exist in rapid equilibrium with monomers and higher-order assemblies. We report here results of studies in which specific oligomers have been stabilized structurally, fractionated in pure form, and then studied by using a combination of CD spectroscopy, Thioflavin T fluorescence, EM, atomic force microscopy (AFM), and neurotoxicity assays. $A\beta$ monomers were largely unstructured, but oligomers exhibited order-dependent increases in β -sheet content. EM and AFM data suggest that dimerization and subsequent monomer addition are processes in which significant and asymmetric monomer conformational changes occur. Oligomer secondary structure and order correlated directly with fibril nucleation activity. Neurotoxic activity increased disproportionately (order dependence >1) with oligomer order. The structure–activity correlations reported here significantly extend our understanding of the conformational dynamics, structure, and relative toxicity of pure $A\beta$ oligomers of specific order.

Alzheimer's disease | toxicity

Alzheimer's disease (AD) is the most common form of late-life dementia. Current estimates of AD incidence are >24 million worldwide, a number that is expected to double every 20 years, reaching 81 million in 2040 (1). AD is a slowly progressive disorder with insidious onset and progressive impairment of episodic memory and executive function coupled with aphasia, apraxia, and agnosia (1).

The amyloid β -protein ($A\beta$) appears to play an essential role in the pathogenesis of AD. $A\beta$ is produced throughout life through posttranslational processing of the $A\beta$ precursor (APP). Familial forms of AD increase $A\beta$ production or the propensity of $A\beta$ to aggregate (2). The "amyloid cascade hypothesis" proposes that assemblies of $A\beta$ initiate a process leading to neuronal dysfunction and cell death (2). The most potent neurotoxic assemblies appear to be oligomeric, rather than fibrillar, in nature (3, 4). For example, oligomers extracted from AD brain potentially impair synapse structure and function (5). The smallest of these oligomers appears to be dimeric (5). However, a systematic correlation of oligomer structure and neurotoxic potency has not been reported. This correlation is critical for the targeting and design of disease-modifying therapeutic agents.

Efforts to establish rigorous structure–toxicity correlations have been hindered by the complex, dynamic equilibria displayed by $A\beta$ (for recent reviews, see refs. 6 and 7). To enable determination of the oligomer frequency distribution, we have used in situ chemical cross-linking to prevent oligomer dissociation or growth (8–10). Oligomers thus stabilized can be visualized and quantified by SDS/PAGE. However, in theory, the method also could be used to produce pure populations of oligomers of defined order[†], enabling the biophysical and biological studies necessary to establish structure–toxicity correla-

tions. We report here that this goal is attainable in practice and discuss insights into the structural biology of $A\beta$ thus obtained.

Results

Preparation and Characterization of $A\beta$ 40 Oligomers of Defined Order. To study specific $A\beta$ oligomers, we used the technique of photo-induced cross-linking of unmodified proteins (PICUP) (10) to chemically stabilize oligomers. PICUP has been shown to rapidly (<1 s) and efficiently (80–90%) cross-link native $A\beta$ oligomers, allowing accurate determination of the oligomer size distribution using SDS/PAGE (10). We found, as reported (9), that cross-linked $A\beta$ 40 comprises a mixture of monomers and oligomers of order 2–5 (Fig. 1, lane 1).

To obtain pure oligomer populations, we developed a method to extract the protein component from a gel band and remove the accompanying SDS (see *Materials and Methods*). We performed this procedure first on an entire gel lane in which cross-linked $A\beta$ had been electrophoresed. When the material thus isolated was electrophoresed on a second SDS gel, the oligomer distribution was identical, within experimental error, to that of the original cross-linked preparation (Fig. 1, lane 2). Proteins extracted in this manner thus "ran true," as determined by SDS/PAGE behavior[‡]. We next purified monomers through tetramers by using this method (Fig. 1, lanes 3–6). Highly pure preparations were obtained. Percentage purities determined by densitometric analysis of the monomer, dimer, trimer, and tetramer preparations were 99.9%, 97.0%, 94.3%, and 99.6%. The low abundance ($<5\%$) of pentamers and higher-order oligomers precluded their isolation and study.

To determine whether the purified, cross-linked species self-associated to produce Thioflavin T (ThT)-positive assemblies, we incubated them in F12K medium at 37 °C and monitored β -sheet content periodically. In addition to the pure oligomers, we studied unfractionated, cross-linked $A\beta$. All of the samples displayed low initial ThT fluorescence intensities that did not change significantly over a 2-day period (data not shown). These data show that cross-linking prevents the continued evolution of the β -sheet structures that are characteristic of $A\beta$ assembly.

To determine whether dissociation of the purified oligomers

Author contributions: K.O. and D.B.T. designed research; K.O. performed research; M.M.C. contributed new reagents/analytic tools; D.B.T. analyzed data; and K.O. and D.B.T. wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission.

¹To whom correspondence should be addressed. E-mail: dteplow@ucla.edu.

[†]In this context, "order" refers to the number of monomers comprising a specific assembly.

[‡]Similar experiments done using $A\beta$ 42 revealed that some, but not all, of the oligomers formed were stable upon re-electrophoresis. For the purpose of structure–activity studies, the partial instability of the repurified $A\beta$ 42 oligomers precludes structure–activity studies. This is an important difference between $A\beta$ 40 and $A\beta$ 42. This difference requires further investigation to develop the means to eliminate the instability of the $A\beta$ 42 oligomers and to understand this fundamental difference in biochemical behavior, which likely correlates with known differences in pathophysiologic behavior.

This article contains supporting information online at www.pnas.org/cgi/content/full/0905127106/DCSupplemental.




Fig. 1. Stability of purified oligomers. $A\beta$ samples were subjected to PICUP and SDS/PAGE. Individual gel bands were stained with Coomassie blue, excised, and then extracted under alkaline conditions. The extracts then were reconstituted in F12K medium and analyzed by SDS/PAGE and silver staining. Lane 1, cross-linked $A\beta$ immediately after PICUP (cross-linking control). Lane 2, cross-linked $A\beta$ subjected to the entire protocol, but with all bands pooled together (control for unfractionated $A\beta$ subjected to alkaline extraction). Lane 3, monomer band. Lane 4, dimer band. Lane 5, trimer band. Lane 6, tetramer band. Lane 7, a “band-equivalent”-sized piece of gel (“no protein” control). The data are representative of results from each of three independent experiments.

occurred during incubation in culture medium, an important question for interpretation of cytotoxicity assays, aliquots of pure, cross-linked monomers, dimers, and trimers were removed from the F12K medium at the end of the incubation (2 days) and were analyzed by SDS/PAGE (Fig. S1). Densitometry then was done to quantify the relative amounts of each band in each lane. In this experiment, before incubation, monomers, dimers, and trimers displayed purities of 91.2%, 93.5%, and 100%, respectively. After incubation, the purities were 80.0%, 89.7%, and 94.5%. The decreases in purity reflected small increases (4–12%) in the intensity of oligomers of order $2n$, where n was the initial order of the purified oligomer. The similarity in oligomer states for each of the “before and after” samples means that experimental differences observed among the pure oligomer populations in subsequent experiments must be caused primarily by oligomer order and not postincubation alterations thereof.

Oligomer Conformation. Studies have revealed that $A\beta$ exists predominately as a statistical coil (SC) if prepared under conditions designed to prevent its folding and self-assembly (7, 11). To determine whether initial $A\beta$ oligomerization involves changes in peptide secondary structure, each purified oligomer population was studied by using CD (Fig. 2). Uncross-linked $A\beta$ (Fig. 2A) and purified monomer (Fig. 2B) produced initial spectra characteristic of SCs. The major feature of these spectra was a large magnitude minimum centered at ≈ 200 nm. In contrast, the major spectral feature of fibrils was a large magnitude minimum centered at ≈ 216 nm, indicative of β -sheet structure (Fig. 2A). The major feature of the spectrum produced by cross-linked, unfractionated $A\beta$ was a minimum at ≈ 205 nm, a wavelength between those of the minima of $A\beta$ monomers and fibrils. This finding indicates that oligomerization is accompanied by an increase in structural order⁸. Spectra from dimers, trimers, and tetramers all displayed single minima, the wavelengths of which (≈ 210 , ≈ 212 , and ≈ 214 nm, respectively) correlated directly with oligomer order (Fig. 2B). The tetramer wavelength minimum approached that of fibrils (≈ 216 nm). Deconvolution of the spectra (Table 1) revealed that the sum of the percentages of β -sheet and SC remained almost constant ($\approx 90\%$) among monomers and oligomers, but that β -sheet content increased from $\approx 25\%$ in the monomer to $\approx 45\%$ in the

⁸“Order,” in this context, refers to secondary and tertiary structure, not to “oligomer order,” the number of monomers in an assembly.




Fig. 2. Secondary structure dynamics of $A\beta$ assemblies. Uncross-linked (UnXL), cross-linked (XL), or fibrillar (Fib) $A\beta$ (A) or isolated oligomers [monomer (Mo), dimer (Di), trimer (Tr), and tetramer (Te)] (B) were prepared in 10 mM PBS, pH 7.4, and then monitored immediately by CD. Data are representative of three independent experiments.

tetramer (Fig. S2). The largest relative change in secondary structure content was between monomer and dimer. Cross-correlation of the SC and β -sheet structural elements yielded $r^2 = 0.995$, suggesting that the increased β -sheet could have formed directly from disordered elements. However, it also is possible that an SC \rightarrow α -helix \rightarrow β -sheet transition could occur.

$A\beta$ Assembly Morphology. We next determined the morphologies of the oligomers by using EM (Fig. S3 and Table 1). Uncross-linked $A\beta$ and purified monomers displayed irregular thread-like and globular shapes that had average diameters of 1.15–1.30 nm. As oligomer order increased, the relative amounts of thread-like structures decreased and the amounts of quasi-spherical structures increased (compare Fig. S3 C and F). Diameter increased progressively with oligomer order, from 1.78 nm for dimers to 11.00 nm for tetramers. Unfractionated, cross-linked species had the largest average diameter, 12.39 nm, but this value was not significantly different from that of tetramers.

In prior studies of $A\beta$ assembly, we have found that EM and atomic force microscopy (AFM) sometimes reveal different features in a single population, which is not surprising considering the differences in substrates (carbon-coated formvar versus mica) to which assemblies must adhere and in visualization physics (electron transmission versus physical displacement). Therefore, in addition to EM, we studied oligomer morphology by using AFM (Fig. S4 and Table 1). The rank order of heights

Table 1. Characteristics of A β assemblies

Sample	α -Helix*	β -Sheet*	SC*	Diameter [†]	Height [‡]	EC ₅₀ [§]
Uncross-linked	9.4	25.2	65.4	1.15 \pm 0.17 (79)	0.14 \pm 0.01 (135)	102.5 \pm 5.6
Cross-linked	11.1	33.6	55.3	12.39 \pm 2.13 (55)	2.93 \pm 0.40 (56)	43.0 \pm 2.7
Pure monomer	8.7	24.0	67.3	1.30 \pm 0.13 (127)	0.24 \pm 0.01 (178)	67.3 \pm 8.7
Pure dimer	10.5	38.6	50.9	1.78 \pm 0.23 (116)	0.53 \pm 0.03 (165)	41.6 \pm 3.9
Pure trimer	10.3	40.8	48.9	7.22 \pm 1.12 (64)	0.94 \pm 0.13 (67)	24.5 \pm 1.9
Pure tetramer	12.7	44.9	42.4	11.00 \pm 2.08 (39)	1.51 \pm 0.30 (38)	20.5 \pm 0.4
Fibrils	ND	57.0	43.0	ND	ND	57.6 \pm 2.2

ND, not determined.

*CD data are expressed as percent of each secondary structure element.

[†]Mean diameter \pm SE, in nm, is listed for (*n*) A β assemblies visualized by EM.

[‡]Mean height \pm SE, in nm, is listed for (*n*) A β assemblies visualized by AFM.

[§]Effective concentration (EC₅₀) is the concentration of a particular A β assembly that produced a level of toxicity in MTT assays that was half maximal. EC₅₀ [mean concentration (μ M) \pm SE] values were calculated after sigmoidal curve fitting of the data shown in Fig. S5, using GraphPad Prism software (version 4.0a).

was identical to that for diameter. The smallest assemblies were in the uncross-linked population, and the largest were in the unfractionated, cross-linked population. Analysis of the diameter and height data revealed that these parameters were correlated ($r^2 > 0.95$). EM and AFM thus revealed the same order dependency of assembly geometry, although the absolute values of the respective diameters and heights were unequal.

Nucleation of Fibril Formation. A β fibril assembly is a nucleation-dependent elongation process (12–14). To determine the effects of specific oligomers on this process, we used ThT to monitor temporal changes in β -sheet content in samples of A β to which were added preformed fibrils, unfractionated oligomer mixtures, or pure monomers, dimers, trimers, or tetramers. ThT fluorescence is not a measure of fibril content *per se*, but because β -sheet formation correlates with amyloid fibril formation, ThT fluorescence is a useful surrogate marker (15–17). The assay also provides the means to monitor assembly kinetics.

A β displayed a sigmoidal process curve characterized by a \approx 1-day lag time, a \approx 4-day period of increasing ThT binding, and a binding plateau occurring after \approx 5 d (Fig. 3A), results consistent with a nucleation-dependent polymerization process (18). The addition of preformed fibrils (nuclei) eliminated the lag period and produced a rapid increase in ThT binding that reached maximal levels after \approx 4 h. Similar effects were observed after addition of unfractionated, cross-linked A β , but the rate of increase of ThT fluorescence was lower (maximal signal occurred after \approx 2 days). These data showed that stable oligomers function as fibril nuclei.

To differentiate effects among the oligomers, the experiments were repeated with the pure cross-linked monomer and oligomer preparations (Fig. 3B). Monomers had no effect, as would be expected because monomers cannot nucleate fibril formation (12, 13). However, each stabilized oligomer nucleated assembly growth. Dimers displayed a process curve similar to that of unfractionated, cross-linked oligomers (compare Fig. 3A and B), demonstrating that they nucleate assembly significantly. However, substantially greater effects were observed with trimers and tetramers. Initial rates of ThT signal increase for these latter oligomers were \approx 75% that of fibrils (Fig. 3A), showing that these oligomeric assemblies were almost as efficient as preformed fibrils in nucleating assembly. If one defines $t_{1/2}$ as the time at which half maximal ThT binding is observed, $t_{1/2}$ values ranged from 1 h for fibril seeds to 65 h for monomers (Fig. 3, arrows). In contrast, dimers displayed $t_{1/2} = 10$ h and trimers and tetramers had identical (within experimental error) $t_{1/2} = 2$ h. Dimers thus can act as seeds, and trimers and tetramers are almost as active as preformed fibrils themselves.

It is important to recognize that the nominal total A β mono-

mer concentration in each experiment was identical. This means that the molar concentration of each assembly differed. If one calculates effectiveness relative to that of the monomer (relative activity; Table S1), the data suggest that dimers are at least an order of magnitude more effective in nucleating fibril formation than are monomers (which must assemble *de novo* to produce nuclei). Trimers and tetramers are at least 100-fold more effective. Fibrils are more effective than oligomers, but fibril-specific activity cannot be determined accurately because fibril structure is not static, but is represented by a quaternary structure frequency distribution that itself comprises dynamic lower-order structural equilibria.

Biological Activities of Oligomers. To establish structure–activity relationships, we determined the effects of specific oligomers on cellular metabolism and survival by using 3-[4,5-dimethylthiazol-2-yl]-2,5-diphenyltetrazolium bromide (MTT) (Table 1 and Fig. S5) and lactate dehydrogenase (LDH) (Fig. 4) assays (19, 20), respectively. For the MTT experiments, differentiated PC12 cells were treated with oligomers of varying concentration and then the concentration at which a 50% effect was observed (EC₅₀) was determined for each (see *Materials and Methods*). Uncross-linked A β , cross-linked unfractionated A β , and A β fibrils yielded EC₅₀ values of 102.5 \pm 5.6, 43.0 \pm 2.7, and 57.6 \pm 2.2 μ M (mean \pm SE), respectively (Fig. S5A). Both cross-linked oligomeric A β and fibrillar A β thus were significantly more toxic than unassembled A β .

To determine the contributions of each oligomer species to that of the cross-linked mixture, EC₅₀ values were determined for each pure population. As observed in the biophysical studies discussed above, EC₅₀ was proportional (in this case, inversely) to oligomer order (monomer = 67.3 \pm 8.7 μ M, dimer = 41.6 \pm 3.9 μ M, trimer = 24.5 \pm 1.9 μ M, and tetramer = 20.5 \pm 0.4 μ M) (Fig. S5B).

We next determined the cytotoxic ability of each assembly by measuring LDH release from differentiated PC12 cells (Fig. 4). Uncross-linked A β produced modest toxicity, averaging \approx 35%. Unfractionated, cross-linked A β was significantly ($P < 0.001$) more toxic, producing almost 80% LDH release, and this level of release was significantly ($P < 0.05$) greater than that of fibrils (\approx 66%). Purified monomers were slightly more toxic than was A β alone (\approx 44% versus \approx 35%), although this toxicity difference was not significant statistically. However, dimers were substantially and significantly ($P < 0.001$) more toxic and their toxicity was equal, within experimental error, to that of fibrils. These data show that toxicity was directly related to oligomer order. This was a consistent trend that was statistically significant between each assembly pair, except trimers and tetramers. In summary, the MTT and LDH assays produced identical rank

tions is a prerequisite for establishing rigorous structure–activity correlations leading to knowledge-based therapeutic drug design. Here, we have reported results of studies using pure (94.3–99.6%), low-order, oligomer populations of constant quaternary structure and restricted conformational complexity. Structural studies revealed that a consistent feature of A β assembly space was a correlation between oligomer structure and order. This correlation extended not only from an intuitively obvious relationship between size (determined by EM and AFM) and oligomer order, but to the level of secondary structure, fibril nucleation activity, and neurotoxicity. Interestingly, and potentially importantly, nonlinear correlations were observed (see below).

Nascent A β in vivo and in vitro is significantly unstructured (6). In vivo, this is a consequence of the necessity for peptide α -helix denaturation during β -secretase and γ -secretase function (29). In vitro, strongly denaturing conditions are used before experiments to eliminate preexistent aggregates and other amyloid-like assemblies and conformers (7). In contrast, protofibrils and fibrils are structures exhibiting high β -sheet content ($\geq 50\%$) (30). Given this continuum in secondary structure from SC to extended β -sheet, we began the studies reported here by asking the question, where do specific oligomers lie within the conformational continuum?

We observed a direct relationship between oligomer and structural (secondary) order (Table 1 and Fig. S2). The relative magnitudes of conformational change between each oligomer are informative. The data suggest that the monomer \rightarrow dimer transition involves substantial secondary structure change, significantly more than in the higher-order transitions. Our data do not reveal whether this change occurs in the monomer before its dimerization, within a nascent dimer, or both. It is interesting that fibril elongation also has been found to involve structural rearrangement in monomers and these rearrangements were suggested to occur in the incoming monomer or the monomers comprising the fibril ends themselves (31).

Examination of the quaternary structure data in Table 1 suggests, as did the results from CD analyses, that the monomer \rightarrow dimer transition is qualitatively different from higher-order transitions. In both the EM and AFM experiments, the dimer \rightarrow trimer \rightarrow tetramer transitions involved increases in assembly size that significantly exceeded that in the monomer \rightarrow dimer transition. This finding suggests that monomer addition to dimeric and higher-order assemblies is associated with tertiary conformational changes producing relatively extended conformers. For example, in the AFM experiments, each monomer added to a dimer increases assembly height by ≈ 0.5 nm. This relationship [$h = 0.5(o - 1)$], where h is height and o is order) predicts $h = 0$ for monomer, which is impossible. Alternatively, if monomer height were constant (≈ 0.25 nm), then the maximum height increase per monomer addition would be 0.25 nm. This relationship ($h = 0.25 o$) holds for monomer and dimer, but not for higher-order oligomers, again suggesting oligomerization-linked conformational change. Taken together, the data show a nonlinear dependence of oligomer geometry on oligomer order.

A fundamental question in any nucleated assembly process is whether particular assemblies can act as nuclei. Many oligomeric assemblies are intermediates in fibril assembly (see Table S1 in ref. 6). What has not been clear is whether the smallest of these oligomers, in pure form, function as fibril “seeds,” i.e., can eliminate the lag phase in fibril elongation process. We found here that purified oligomers as small as dimers seed fibril formation. However, they are not as effective as trimers or tetramers, which have an effectiveness closer to that of fibrils than that of monomers.

An important aspect of the analysis of the nucleation data was the consideration of number average nucleation activity, i.e.,

what is the quotient of activity/number of oligomers? This is an important question in the A β system, because experiments usually are done, and analyzed, at a constant total A β concentration, without consideration for the number of moles of each assembly present. When the nucleation data were analyzed in this manner, the relative activity of dimers was an order of magnitude higher than that of monomers, and trimer and tetramer activity was two orders of magnitude higher.

Once static and dynamic features of the purified oligomers were determined, we next sought to correlate these features with biological activities. This structure–activity correlation is central to investigation of the pathobiology of oligomer-mediated neurotoxicity in vivo. We observed an order dependence of toxicity (Table S2) that correlated with the dependencies observed in the prior structural studies, namely tetramers \geq trimers $>$ dimers \geq fibrils $>$ monomers $>$ uncross-linked A β . Analysis of the molar toxicity of each oligomer, determined similarly to the molar nucleation activity, showed that dimers are 3-fold more toxic than monomers and that the toxicity of trimers and tetramers are up to ≈ 13 -fold more toxic.

Recently, Chimon et al. (32) reported results of structure–toxicity studies of nonfibrillar A β assemblies. These studies focused on a high-order ($n = 150$) spherical assembly that had substantial β -sheet content and a fibril-like β -strand organization, I_β . They suggested that I_β was an immediate precursor to fibrils and showed that its toxicity was significantly greater than monomers but significantly less than fibrils. Our results show that the lowest-order ($n = 2$ –4) A β oligomers, which evidence suggests are the most clinically relevant, are less organized structurally than I_β and likely are the link between the SC state of the monomer and that in I_β and fibrils.

Insights and Implications

Our results provide significant insights into the biophysical and pathobiological behavior of A β , and importantly, into strategies for developing therapeutics for AD. The “specific activity” of A β assemblies depends nonlinearly on oligomer order. In fact, dimers were ≈ 3 -fold more toxic than monomers, and tetramers were ≈ 13 -fold more toxic. Are tetramers the “most toxic” assembly? We do not know. Determination of a complete toxicity/order correlation is not possible because of the progressive decrease in occurrence frequency of higher-order oligomers, which precludes their isolation and study.

A broader consideration of the pathobiology of AD suggests that determination of a toxicity/oligomer-order frequency distribution may not be necessary. Fibrils, the most toxic assemblies on a molar basis, accumulate in plaques, a process now thought to be beneficial by sequestering these toxic entities (2). Monomers have very low toxic activity. Only when A β self-associates does toxicity rise substantially. How likely to occur are these associations? Because the occurrence frequency has a negative exponential dependence on oligomer order, and because initial assembly can nucleate rapid A β polymerization, the steady-state levels of any oligomer must be relatively low. This logic suggests that a strategy targeting low-order oligomers, e.g., dimers through hexamers, is reasonable because these targets likely contribute most substantially to AD neurotoxicity (according to Eq. 1). The larger of these assemblies is significantly more toxic than the smaller, but the concentration of these highly toxic larger oligomers will be lower (assuming they do not exist in covalently associated form).

How should one target low-order oligomers? Two points in the oligomerization process may be particularly attractive (Fig. 5). The first point is the conformational organization of the monomer during its dimerization. Therapeutic agents could stabilize a disordered state of the monomer or destabilize the dimer state, resulting in complete blockage of oligomerization and fibril formation. The second point would be the stabilization of the less extended form of the A β monomer within the dimer or in incoming monomers

