UC Santa Barbara

UC Santa Barbara Previously Published Works

Title

A copper(I)-arene complex with an unsupported η^6 interaction.

Permalink

<https://escholarship.org/uc/item/7v41p19v>

Journal

Angewandte Chemie (International ed. in English), 54(10)

ISSN

1433-7851

Authors

Wright, Ashley M
Irving, Benjamin J
Wu, Guang
et al.

Publication Date

2015-03-01

DOI

10.1002/anie.201410948

Peer reviewed

Structural and theoretical investigation into a copper(I)–arene complex with an unsupported η^6 interaction**

Ashley M. Wright, Benjamin J. Irving, Guang Wu, Anthony J. H. M. Meijer, Trevor W. Hayton*

Abstract. Addition of PR_3 ($\text{R} = \text{Ph}$ or OPh) to $[\text{Cu}(\eta^2\text{-Me}_6\text{C}_6)_2][\text{PF}_6]$ results in formation of $[(\eta^6\text{-Me}_6\text{C}_6)\text{Cu}(\text{PR}_3)][\text{PF}_6]$, the first copper arene complexes to feature an unsupported η^6 arene interaction. A DFT analysis reveals that the preference for the η^6 binding mode is enforced by the steric clash between the methyl groups of the arene ligand and the phenyl rings of the phosphine co-ligand.

The synthesis of the first metal–arene complex $\text{Cr}(\eta^6\text{-C}_6\text{H}_6)_2$, nearly 100 years ago by Hein and co-workers,^[1] was an important milestone in the development of organometallic chemistry.^[2–4] Since its discovery, metal–arene complexes have been isolated for most of the transition metal elements,^[5–10] and these species have found use in a wide variety of applications. For example, the binding of an arene to a metal activates the ring toward nucleophilic addition, and also renders the ring protons more acidic, making these species synthetically useful.^[11] In addition, metal–arene complexes have found utility as both catalyst pre-cursors and common starting materials.^[12, 13] These points are well illustrated by the chemistry of the coinage metals. For example, Cu–arene species are thought to play a role in copper(I)–catalysed cross-coupling reactions.^[14] Cu–arene complexes are also pre-cursors for catalytic alkoxylations and aminations,^[15] and are known to mediate the reactivity of Cu(I) with oxygen.^[16, 17]

Group 11 metal–arene complexes strongly prefer the η^2 binding mode.^[12, 18–23] This binding mode is favoured as it lessens the interaction between the filled nd orbitals with the filled arene π -orbitals.^[24] Consistent with this analysis, the only reported Cu arene complexes that contain the η^6 binding mode feature tethered arene rings and long Cu–C_{cent} distances (~ 2.5 – 3.0 Å) indicative of weak interactions (Chart 1).^[25, 26] Interestingly, though, the η^6 configuration is proposed to be slightly more stable than the η^2 configuration for the “naked” $[\text{Cu}(\text{C}_6\text{H}_6)]^+$ fragment in the gas phase, but the difference between the η^6 and η^2 modes is less than 1 kJ mol^{−1}.^[27] Inclusion of a counterion, such as a halide or pseudo-halide, to the Cu coordination sphere was calculated to increase the stability of the η^2 mode over η^6 ,^[28] which led to the proposal that the unsupported η^6 –arene copper interaction is unlikely to be observed in the condensed state.^[28]

Chart 1. Known examples of $\text{Cu}(\eta^6\text{-arene})$ interactions.

Herein, we report the synthesis and structural characterization of the first Cu(I)–arene complexes that feature unsupported η^6 interactions, demonstrating that complexes of this type are, in fact, isolable.

We previously reported the synthesis of the copper–arene sandwich complex, $[\text{Cu}(\eta^2\text{-Me}_3\text{C}_6\text{H}_3)_2][\text{PF}_6]$, which was prepared by oxidation of Cu metal with $[\text{NO}][\text{PF}_6]$ in the presence of mesitylene.^[29] The hexamethylbenzene analogue, $[\text{Cu}(\eta^2\text{-Me}_6\text{C}_6)_2][\text{PF}_6]$ (**1**), was prepared in a similar fashion. As determined by X-ray crystallography, complex **1** features two Me_6C_6 ligands coordinated to Cu via η^2 interactions. The Cu–C bond lengths are 2.092(2) and 2.192(2) Å, and are comparable with those previously reported for η^2 –arene,^[13, 22, 30] alkyne,^[31] and alkene^[32] complexes of copper.

With **1** in hand, we postulated that substitution of one Me_6C_6 ligand with a $2e^-$ σ -donor ligand would result in a η^2 to η^6 ring slip to generate a pseudo-tetrahedral $18e^-$ Cu(I) complex. Gratifyingly, addition of 1 equiv of PR_3 ($\text{R} = \text{Ph}$, OPh) to a CH_2Cl_2 solution of **1** results in formation of the half-sandwich complexes $[(\eta^6\text{-Me}_6\text{C}_6)\text{Cu}(\text{PR}_3)][\text{PF}_6]$ ($\text{R} = \text{2, Ph; 3, OPh}$) (eq. 1). Complexes **2** and **3** can be crystallized from CH_2Cl_2 /hexanes at -25°C to give colourless blocks in moderate to good yields (**2**: 40%; **3**: 78%).

Figure 1. ORTEP drawing of $[(\eta^6\text{-Me}_6\text{C}_6)\text{Cu}(\text{PPh}_3)][\text{PF}_6]$ (**2**) (left) and $[(\eta^6\text{-Me}_6\text{C}_6)\text{Cu}(\text{P}(\text{OPh})_3)][\text{PF}_6]$ (**3**) (right). Hydrogen atoms and a PF_6 anion have been omitted for clarity.

Single crystals of **2** and **3** suitable for X-ray crystallography were grown from a dilute CH_2Cl_2 /hexane solution at -25°C . Both complexes feature a copper(I) centre bound by a $\eta^6\text{-Me}_6\text{C}_6$ ligand and a single PR_3 ligand, in an overall pseudo-tetrahedral geometry (Fig. 1). The η^6 coordination mode of the Me_6C_6 ligand is exemplified by the narrow range of Cu–C_{aryl} bond lengths [**2**: 2.284(5)–2.293(5) Å; **3**: 2.253(1)–2.300(1) Å] and short Cu–C_{cent} distances [**2**: 1.800(3) Å; **3**: 1.775(6) Å]. Notably, these Cu–C_{cent} distances are significantly shorter than those previously reported for the tethered Cu(I)– η^6 –arene interactions (Chart 1).^[25, 26] For instance, Mascall and co-workers reported a Cu(I) cyclophane complex that features a Cu–C_{cent} distance of 2.97 Å.^[25] Similarly, Zhang and co-

workers reported a Cu(I) complex, supported by 9,10-bis{N,N-propyl-N-(diphenylphosphino)aminomethyl}anthracene, which features a Cu–C_{cent} distance of 2.50 Å.^[26] Finally, the average C–C bond length of the Me₆C₆ ligand in **2** is 1.411 Å (range = 1.395(7) – 1.423(7) Å), and is comparable with those of unbound Me₆C₆, suggesting that there is little net charge transfer in the Cu–arene interaction.^[9] Complex **3** features comparable C–C bond lengths for its Me₆C₆ ligand (1.416(1) – 1.421(2) Å). The Cu–P bond lengths in **2** (2.158(2) Å) and **3** (2.1242(4) Å) are similar to those of (η⁵-Cp)Cu(PPh₃) (2.116(2) Å).^[33, 34]

The binding of an arene to a metal can result in a significant shift of its ¹³C resonances.^[35] However, only small shifts are observed for the aryl carbons in **2** (131.2 ppm) vs. those of unbound Me₆C₆ (132.1 ppm). A slight down field shift of the aryl carbon is observed for **3** (149.2 ppm). Additionally, the methyl carbon resonances in **2** and **3** do not differ significantly from those observed for unbound Me₆C₆ (**2**: 19.06 ppm; **3**: 16.07 ppm; free Me₆C₆: 16.71 ppm). Overall, these data are consistent with a relatively weak Cu–arene interaction.

To gain further insight into the bonding interactions in **2** and **3**, DFT calculations were performed. For comparison, the “naked” Cu–arene complexes, [Cu(C₆H₆)]⁺ and [Cu(C₆Me₆)]⁺, were also studied, as were their C₆H₆-bearing sister molecules, [(C₆H₆)Cu(PPh₃)]⁺ and [(C₆H₆)Cu(P(OPh)₃)]⁺. The optimized coordinates for all calculated species are given in the Supporting Information. The energy differences between the η⁶ and η² binding modes for the naked complexes and **2** and **3** are tabulated in Table 1. For the “naked” complexes the calculations reveal that the η² structures are generally more stable (apart from [Cu(C₆Me₆)]⁺ in the gas phase), in accordance with earlier calculations.^[27, 28] Structurally, the two η⁶ complexes feature similar geometries; however, the Cu⁺ ion in [Cu(η⁶-C₆Me₆)]⁺ is closer to the arene than in [Cu(η⁶-C₆H₆)]⁺ (Cu–C_{cent} = 1.66 Å and 1.73 Å, respectively). The two η² structures are also similar, but the distance between Cu⁺ and the C=C bond has increased to 1.89 Å and 1.92 Å for [Cu(η²-C₆Me₆)]⁺ and [Cu(η²-C₆H₆)]⁺, respectively.

Interestingly, the addition of PPh₃ or P(OPh)₃ to the [Cu(arene)]⁺ motif results in a reversal of the favored arene binding mode. The η⁶ motif becomes favoured by 8.78 and 13.83 kJ mol^{−1} for **2** and **3** (in the gas phase), respectively (Table 1). Inclusion of solvent (CH₂Cl₂) in the calculation results in a slight destabilization of η⁶ binding mode relative to the η² mode, which we attribute to the attenuation of the electrostatic interaction due to the change in dielectric constant from vacuum to CH₂Cl₂, but the η⁶ mode is still favored. We attribute the lower stability of the η² mode to the steric clash between the CH₃ groups on Me₆C₆ and the [Cu(PR₃)]⁺ moiety. The larger ΔE(η⁶ – η²) values calculated for compound **3**, which features the phenyl groups on the P(OPh)₃ moiety pointing towards the methyl groups of the Me₆C₆ ligand (see Table 1, Fig. S18), also support the notion that sterics dictate the binding mode. Moreover, we calculate that [(C₆H₆)Cu(PPh₃)]⁺, which features the smaller C₆H₆ ligand, prefers the η² motif, whereas [(C₆H₆)CuP(OPh)₃]⁺ exhibits a preference for an η¹ motif, consistent with a diminished steric clash between the C₆H₆ ring and the [Cu(PR₃)]⁺ fragment (see SI). The isolation of **1** is also consistent with this argument, as it reveals that a η² binding mode of the Me₆C₆ ligand is preferred in the absence of a phosphine co-ligand. Thus, several lines of evidence support the argument that sterics to play a dominating role in the choice of final binding mode.

Table 1. Calculated relative energies for different binding modes of **2** and **3**.

Complex	ΔE(η ⁶ – η ²) / kJ mol ^{−1}	
	Gas phase	CH ₂ Cl ₂
[Cu(C ₆ H ₆)] ⁺	9.79	17.6 ^[a]
[Cu(C ₆ Me ₆)] ⁺	−3.99	4.51
2 / PPh ₃ ^[c]	−8.78	−7.19 ^[b]
3 / P(OPh) ₃ ^[c]	−13.83	−12.16

[a] The η⁶ structure is not a stationary point, so was calculated as a restricted optimization at a fixed C_{6v} geometry [b] The η⁶ structure in this case is actually slightly off centre, i.e. η⁴. [c] The η² binding mode was calculated using a restricted optimization.

Inspection of the molecular orbitals (both for B3LYP and for a Hartree-Fock calculation, based on the DFT-optimized geometry) does not reveal a clear molecular orbital picture for the Cu–arene interaction. Moreover, the effect of solvent on the relative stabilities of the η² and η⁶ binding modes, along with the importance of sterics in determining the hapticity, suggest a significant electrostatic component to the interaction. Thus, to elucidate the bonding interaction further, the complexes were studied using a Mulliken partitioning scheme, permitting the Mulliken charges of the Cu⁺ ions in **2** and **3** to be calculated. The calculated Cu⁺ charges for **2** and **3** are 0.86 and 0.81 (in CH₂Cl₂), respectively (Table 2). Both of these values are close to 1, which supports a largely electrostatic bonding interaction with very little covalency. The electrostatic bonding arrangement is further supported by analysis of the total electron density isosurfaces of **2** and **3** (see SI for details).^[36] We used H₂ (0.24 a.u.) and LiF (0.005 a.u.) as benchmarks for covalent and ionic bonding, respectively (see SI). The calculated values for **2** and **3** (*in vacuo*) are: **2**: η² = 0.07 a.u., η⁶ = 0.05 a.u.; **3**: η² = 0.07 a.u., η⁶ = 0.05 a.u. Notably, these values are similar to that calculated for LiF, which supports the conclusion that arene binding in **2** and **3** is largely electrostatic. These data also suggest that the character of the Cu–arene interaction is insensitive to the choice of PR₃ ligand, but is sensitive to the coordination mode: i.e., the lower electron density cutoff values for the η⁶ coordination mode indicate a less covalent interaction than the η² mode. Similar results are also observed for the ‘naked’ systems. For example, for [Cu(C₆H₆)]⁺, the maximum values of the electron densities are 0.08 a.u. for the η² mode and 0.055 for η⁶ mode. Our calculations also show that the η² motif for the ‘naked’ systems becomes relatively more stable with the introduction of the CH₂Cl₂ solvent into the calculation, which indicates that the binding between naked Cu⁺ and the arene is also largely electrostatic in character, in accordance with previous results.^[27, 28]

Table 2. Mulliken charges on the central Cu⁺ ion for complexes **2** and **3** with a η⁶ bound arene

Complex	Gas phase	CH ₂ Cl ₂
2 / PPh ₃ ^[c]	0.83	0.86
3 / P(OPh) ₃ ^[c]	0.78	0.81

Figure 2. Average M–C_{cent} distances for 18e[−] M(η⁶-C₆R₆) complexes (R = H, closed circles, or Me, open triangles) in the Cambridge Structural Database.^[19] The open circle represents Cr(benzene)L₃-type complexes, where L is not a carbonyl ligand (9 examples).

Finally, the isolation of complexes **2** and **3** allows for a structural comparison with other first row M(η⁶-arene) complexes. A search of the Cambridge Structural Database reveals the existence of 77 18e[−] η⁶-C₆R₆ (R = H or Me) complexes of the first row transition metals (Cr to Ni). Figure 2 depicts the average M–C_{cent} distance for each element for this series of complexes. Most importantly, **2** and **3** feature the longest M–C_{cent} distances, which supports our contention that the Cu-arene interaction is weak and predominantly electrostatic in nature, a consequence of the contracted nature of the 3d¹⁰ shell. This contention is further supported by the observed decrease in M–C_{cent} distances on moving from Cu to Fe. This decrease can be rationalized by the better energy match of the 3d orbitals with the arene π* orbitals as one moves from right to left across the row, which permits a more effective back bonding interaction.

In summary, we have isolated and structurally characterized the first copper complex with an unsupported η⁶ arene interaction. This completes the series of known unsupported [M(η⁶-C₆R₆)]⁺ first row transition metal complexes. These copper arene complexes have much shorter Cu–C_{cent} distances (~1.80 Å) than those previously reported for Cu(η⁶-arene) compounds. Despite this structural difference, density functional theory calculations reveal the Cu⁺-arene interaction is largely electrostatic in nature, while the preference for the η⁶ binding mode is enforced by the steric clash between the methyl groups of the arene ligand and the phenyl rings of the phosphine co-ligand.

[*] Dr. A. M. Wright, Dr. G. Wu, Prof. Dr. T. W. Hayton
Department of Chemistry and Biochemistry
University of California Santa Barbara
Santa Barbara, CA 93106 (USA)
E-mail: hayton@chem.ucsb.edu
Dr. A. J. H. M. Meijer
Department of Chemistry
University of Sheffield
Sheffield, S3 7HF (UK)
E-mail: a.meijer@sheffield.ac.uk
Dr. B. J. Irving
Department of Control Engineering
Faculty of Electrical Engineering
Czech Technical University in Prague,

Karlovo náměstí 13, 121 35 Prague 2, Czech Republic
E-mail: irvinben@fel.cvut.cz

[**] Research at UCSB supported by the University of California, Santa Barbara and the National Science Foundation (CHE 1059097). This publication was also supported by the European social fund within the framework of realizing the project “Support of inter-sectoral mobility and quality enhancement of research teams at Czech Technical University in Prague”, CZ.1.07/2.3.00/30.0034.

Received: ((will be filled in by the editorial staff))
Published online on ((will be filled in by the editorial staff))

Keywords: Copper • metal arene • π complex • bonding analysis • organometallics

References

- [1] F. Hein, *Chem. Ber.* **1919**, 52, 195-196.
- [2] E. O. Fischer, W. Hafner, *Z. Naturforsch. Part B* **1955**, 10, 665-668.
- [3] G. Wilkinson, M. Rosenblum, M. C. Whiting, R. B. Woodward, *J. Am. Chem. Soc.* **1952**, 74, 2125-2126.
- [4] P. Powell, *Principles of Organometallic Chemistry*, 2nd ed., Chapman and Hall, New York, **1988**.
- [5] H. G. Smith, R. E. Rundle, *J. Am. Chem. Soc.* **1958**, 80, 5075-5080.
- [6] R. K. McMullan, T. F. Koetzle, C. J. Fritchie, Jr, *Acta Crystall. Sec. B* **1997**, 53, 645-653.
- [7] E. L. Muetterties, J. R. Bleeke, E. J. Wucherer, T. Albright, *Chem. Rev.* **1982**, 82, 499-525.
- [8] M. N. Bochkarev, *Chem. Rev.* **2002**, 102, 2089-2118.
- [9] S. M. Hubig, S. V. Lindeman, J. K. Kochi, *Coord. Chem. Rev.* **2000**, 200–202, 831-873.
- [10] G. Pampaloni, *Coord. Chem. Rev.* **2010**, 254, 402-419.
- [11] R. H. Crabtree, *The Organometallic Chemistry of the Transition Metals*, 3rd ed., John Wiley & Sons, Inc., New York, NY, **2001**.
- [12] M. B. Dines, P. H. Bird, *J. Chem. Soc., Chem. Comm.* **1973**, 12-12.
- [13] G. Santiso-Quinones, A. Higelin, J. Schaefer, R. Brückner, C. Knapp, I. Krossing, *Chem. Eur. J.* **2009**, 15, 6663-6677.
- [14] D. Ma, Y. Zhang, J. Yao, S. Wu, F. Tao, *J. Am. Chem. Soc.* **1998**, 120, 12459-12467.
- [15] R. T. Gephart, C. L. McMullin, N. G. Sapiezynski, E. S. Jang, M. J. B. Aguila, T. R. Cundari, T. H. Warren, *J. Am. Chem. Soc.* **2012**, 134, 17350-17353.
- [16] T. Osako, Y. Tachi, M. Doe, M. Shiro, K. Ohkubo, S. Fukuzumi, S. Itoh, *Chem. Eur. J.* **2004**, 10, 237-246.
- [17] T. Osako, Y. Tachi, M. Taki, S. Fukuzumi, S. Itoh, *Inorg. Chem.* **2001**, 40, 6604-6609.
- [18] R. W. Turner, E. L. Amma, *J. Am. Chem. Soc.* **1966**, 88, 1877-1882.
- [19] *Cambridge Structural Database*, 6.16 ed., **2014**.
- [20] R. R. Conry, *Chem. Commun.* **1998**, 555-556.

- [21] R. R. Conry, W. S. Striejewske, A. A. Tipton, *Inorg. Chem.* **1999**, *38*, 2833-2843.
- [22] A. M. Dattelbaum, J. D. Martin, *Inorg. Chem.* **1999**, *38*, 6200-6205.
- [23] S. Y. Lee, S. J. Na, H. Y. Kwon, B. Y. Lee, S. O. Kang, *Organometallics* **2004**, *23*, 5382-5385.
- [24] F. Meyer, F. A. Khan, P. B. Armentrout, *J. Am. Chem. Soc.* **1995**, *117*, 9740-9748.
- [25] M. Mascal, J.-L. Kerdelhué, A. J. Blake, P. A. Cooke, *Angew. Chem. Int. Ed.* **1999**, *38*, 1968-1971.
- [26] F.-B. Xu, Q.-S. Li, L.-Z. Wu, X.-B. Leng, Z.-C. Li, X.-S. Zeng, Y. L. Chow, Z.-Z. Zhang, *Organometallics* **2003**, *22*, 633-640.
- [27] T. K. Dargel, R. H. Hertwig, W. Koch, *Mol. Phys.* **1999**, *96*, 583-591.
- [28] S.-L. Zhang, L. Liu, Y. Fu, Q.-X. Guo, *J. Mol. Struct.* **2005**, *757*, 37-46.
- [29] A. M. Wright, G. Wu, T. W. Hayton, *J. Am. Chem. Soc.* **2010**, *132*, 14336-14337.
- [30] D. S. Laitar, C. J. N. Mathison, W. M. Davis, J. P. Sadighi, *Inorg. Chem.* **2003**, *42*, 7354-7356.
- [31] A. Das, C. Dash, M. A. Celik, M. Yousufuddin, G. Frenking, H. V. R. Dias, *Organometallics* **2013**, *32*, 3135-3144.
- [32] G. Santiso-Quinones, A. Reisinger, J. Slattery, I. Krossing, *Chem. Commun.* **2007**, 5046-5048.
- [33] F. A. Cotton, J. Takats, *J. Am. Chem. Soc.* **1970**, *92*, 2353-2358.
- [34] J. A. Tang, B. D. Ellis, T. H. Warren, J. V. Hanna, C. L. B. Macdonald, R. W. Schurko, *J. Am. Chem. Soc.* **2007**, *129*, 13049-13065.
- [35] J. A. Chudek, G. Hunter, R. L. MacKay, P. Kremminger, K. Schlogl, W. Weissensteiner, *J. Chem. Soc., Dalton Trans.* **1990**, 2001-2005.
- [36] B. J. Irving, F. Y. Naumkin, *Phys. Chem. Chem. Phys.* **2014**, *16*, 7697-7709.
-

Copper arenes

Ashley M. Wright†, Benjamin J. Irving‡,
Guang Wu†, Anthony J. H. M. Meijer‡,
Trevor W. Hayton†* ————— Page
– Page

**Structural and theoretical
investigation into a copper(I)–arene
complex with an unsupported η^6
interaction**

Addition of PR_3 ($\text{R} = \text{Ph}$ or OPh) to $[\text{Cu}(\eta^2\text{-Me}_6\text{C}_6)_2][\text{PF}_6]$ results in formation of $[(\eta^6\text{-Me}_6\text{C}_6)\text{Cu}(\text{PR}_3)][\text{PF}_6]$, the first copper arene complexes to feature an unsupported η^6 arene interaction. A DFT analysis reveals that the preference for the η^6 binding mode is enforced by the steric clash between the methyl groups of the arene ligand and the phenyl rings of the phosphine co-ligand.