Lawrence Berkeley National Laboratory

Recent Work

Title

THE CRYSTAL STRUCTURE OF MANGANESE DICHLORIDE TETRAHYDRATE

Permalink

<https://escholarship.org/uc/item/7vf7p79j>

Authors

Zalkin, A.
Forrester, J.D.
Templeton, D.H.

Publication Date

1963-10-01

University of California
Ernest O. Lawrence
Radiation Laboratory

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 5545*

**THE CRYSTAL STRUCTURE OF
MANGANESE DICHLORIDE TETRAHYDRATE**

Berkeley, California

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

Submitted to Inorganic Chemistry for publication

UNIVERSITY OF CALIFORNIA
Lawrence Radiation Laboratory
Berkeley, California
Contract No. W-7405-eng-48

THE CRYSTAL STRUCTURE OF MANGANESE DICHLORIDE TETRAHYDRATE

A. Zalkin, J. D. Forrester and D. H. Templeton

October 1963

CONTRIBUTION FROM THE LAWRENCE RADIATION LABORATORY AND DEPARTMENT OF CHEMISTRY,
UNIVERSITY OF CALIFORNIA, BERKELEY, CALIFORNIA

The Crystal Structure of Manganese Dichloride Tetrahydrate¹

BY ALLAN ZALKIN, J. D. FORRESTER, AND DAVID H. TEMPLETON

The stable room temperature form of $\text{MnCl}_2 \cdot 4\text{H}_2\text{O}$ (α -form) has a monoclinic cell $a = 11.186 \text{ \AA}$, $b = 9.513 \text{ \AA}$, $c = 6.186 \text{ \AA}$, and $\beta = 99.74^\circ$. The space group is $P2_1/n$ with four formula units per unit cell and $d_x = 2.03 \text{ g./cc.}$ An X-ray diffraction study of this material yielded the positions of all of the atoms including the hydrogens. The structure consists of discrete octahedra of four oxygen atoms and two chlorine atoms about the manganese, and with the chlorine atoms adjacent to each other. The four Mn-O distances are nearly equidistant being $2.224 \text{ \AA}$, $2.209 \text{ \AA}$, $2.185 \text{ \AA}$, and $2.206 \text{ \AA}$; the two Mn-Cl distances are $2.475 \text{ \AA}$ and $2.500 \text{ \AA}$. Only four of the eight hydrogen atoms make hydrogen bonds. Three of the hydrogen bonds are between oxygen and chlorine and one between two water molecules.

Introduction

Groth² reports two monoclinic crystal modifications of $\text{MnCl}_2 \cdot 4\text{H}_2\text{O}$. One form is metastable at room temperature (Dawson and Williams³), and is isomorphous with $\text{FeCl}_2 \cdot 4\text{H}_2\text{O}$, the structure of which has been described by Penfold and Grigor⁴ with comments on the proton sites by Saffar and Murtz⁵. Our paper deals with the form that is stable at room temperatures, and in keeping with the nomenclature of Dawson and Williams³ we refer to this

modification as the α -form, and to the metastable modification as the β -form. It should be noted here that Groth² in his publication has reversed the naming of these two forms from the above convention.

Delain⁶ has reported the cell dimensions and space group of the α -form, and Gardner⁷ has reported some preliminary results on the proton sites. We became interested in the structure of this crystal after learning of its use as a sample for low temperature magnetic measurements by Professor W. F. Giaque.

Experimental

An aqueous solution of manganous chloride was allowed to evaporate at room temperature and the tetrahydrate crystals grew readily in the form of approximately hexagonal plates. A suitable crystal with maximum dimension of 0.1 mm was mounted and used for collecting the intensity data. Intensities were measured with a General Electric XRD-5 equipped with a goniostat, a scintillation counter and a pulse height discriminator. Mo $K\alpha$ radiation was used and the unit cell dimensions are based on $\lambda = 0.70926 \text{ \AA}$ for Mo $K\alpha_1$. No preliminary X-ray photographs were taken; the crystal was aligned on the goniostat using the cell dimensions reported in Donnay⁸, together with the observations on the direction of extinction of polarized light. The systematic space group absences were carefully checked on the goniostat.

The 848 independent reflections permitted by the space group in the sphere with $\sin\theta/\lambda$ less than 0.538 (2θ less than 45°) were measured with counting times of 10 seconds each; of these, 49 were assigned zero intensity. The maximum intensity counted was 4200 counts/second, which was the 002 reflection.

No corrections were made for either absorption or extinction. The absorption factor μ for this crystal with Mo radiation is 28 cm^{-1} . μ_R is

about 0.14 or less, making the absorption very low.

Calculations were made with an IBM 7094 computer using a full matrix least squares refinement program written by P. K. Gantzell, R. A. Sparks, and K. N. Trueblood with minor changes and Fourier and distance programs written by A. Zalkin. We minimized the function $\sum w(|F_o| - |F_c|)^2 / \sum wF_o^2$.

Atomic scattering factors were taken as the values by Ibers⁹ for Mn^{+2} , Cl^{-1} , neutral O and H. Dispersion corrections¹⁰ of -0.4 and -0.1 electrons were added to the Mn^{+2} and Cl^{-1} scattering factors respectively. The imaginary part of the dispersion correction was neglected.

Results

Unit Cell and Space Group.—The primitive cell is monoclinic, contains four formula units ($MnCl_2 \cdot 4H_2O$) per unit cell and has the dimensions:

$$\begin{aligned} a &= 11.186 \pm 0.006 \text{ \AA}, \quad b = 9.513 \pm 0.005 \text{ \AA}, \quad c = 6.186 \pm 0.002 \text{ \AA}, \\ \beta &= 99.74 \pm 0.04^\circ, \quad V = 648.8 \text{ \AA}^3, \quad a : b : c = 1.1759 : 1 : 0.6503. \end{aligned}$$

These compare with values $a = 11.3$, $b = 9.55$ and $c = 6.15 \text{ \AA}$ and $\beta = 99.63^\circ$ reported by Delain⁶ (after transformation from $P2_1/a$) and the axial ratios of 1.1525 : 1 : 0.6445 with $\beta = 99.42^\circ$ by Groth².

From the X-ray data the calculated density is 2.03 g./cc. which compares with the value of 2.01 g./cc. reported in the Handbook of Chemistry and Physics¹¹.

Reflections $h0l$ are absent unless $h + l = 2n$ and $0k0$ reflections are absent unless $k = 2n$. This is characteristic of the space group $P2_1/n$ (C_{2h}^5) and the success of our structure determination confirms this symmetry.

Determination of the Structure.—Projection data were collected initially and the Patterson functions for three projections $hk0$, $h0l$, and $0kl$ were calculated from the observed intensities after correction for Lorentz and polarization effects. From these three projections, the positions of the Mn and the two independent Cl atoms required by the four formula units per

unit cell were obtained directly. These three atoms were all in 4 - fold general positions:

$$4(e) \pm (\underline{x}, \underline{y}, \underline{z}; 1/2 + \underline{x}, 1/2 - \underline{y}, 1/2 + \underline{z}).$$

When all of the three dimensional data were available, a three dimensional Fourier was calculated with phases based on the positions of the Mn and two Cl atoms. In this Fourier, all four independent oxygen atoms were found in general positions.

The parameters of these 7 sets of atoms were refined by least squares using all the data, each with unit weight, and giving each atom an isotropic temperature factor of the form $\exp(-B\lambda^{-2}\sin^2\theta)$.

After 8 cycles of refinement, the conventional reliability factor or

$R = \sum ||F_o| - |F_c|| / \sum |F_o|$ decreased to 0.083. Correction of several blunders in the data taking and in the card punching and the introduction of anisotropic temperature factors of the form $\exp(-\beta_{11}h^2 - \beta_{22}k^2 - \beta_{33}l^2 - 2\beta_{12}hk - 2\beta_{13}hl - 2\beta_{23}kl)$ for each of the 7 atoms reduced R to 0.047 after four more cycles of refinement.

An electron density difference function with all atoms subtracted out with the exception of hydrogen was calculated using the results of this anisotropic refinement and using the terms for which $\sin\theta/\lambda$ less than 0.48. Eight out of the ten highest peaks in this function corresponded to reasonable positions for the eight independent hydrogen atoms in general positions. Four of these hydrogen atoms form hydrogen bonds, but the other four atoms have no suitable environment for forming bonds and remain unbonded.

Four cycles of least squares refinement with all eight hydrogen atoms each having isotropic temperature factors and the remaining seven heavier atoms having anisotropic temperature factors resulted in an R factor of 0.041. The temperature factors of the four bonded hydrogen atoms remained normal, but those of two of the unbonded atoms, attached to O(3), had

higher values.

After four more cycles of refinement, the final R value was 0.039 .

The final positional parameters and thermal parameters are listed in Table I and Table II respectively. The observed and calculated structure factors are listed in Table III.

TABLE I.

FINAL POSITIONAL PARAMETERS AND STANDARD DEVIATIONS FOR $\text{MnCl}_2 \cdot 4\text{H}_2\text{O}$

Atom	<u>x</u>	<u>y</u>	<u>z</u>	$\sigma(\underline{x})$	$\sigma(\underline{y})$	$\sigma(\underline{z})$
Mn	0.2329	0.1714	0.9865	0.0001	0.0001	0.0001
Cl(1)	.0610	.3076	.0938	.0001	.0001	.0002
Cl(2)	.3817	.3662	.0355	.0001	.0001	.0002
O(1)	.3010	.1127	.3334	.0004	.0005	.0006
O(2)	.1568	.2280	.6446	.0004	.0005	.0007
O(3)	.1323	.9736	.9590	.0005	.0006	.0011
O(4)	.3695	.0381	.8764	.0004	.0005	.0007
H(1)1 ^a	.39	.15	.37	.01	.01	.01
H(1)2	.30	.02	.35	.01	.01	.01
H(2)1	.08	.20	.62	.01	.01	.01
H(2)2	.19	.21	.56	.01	.01	.02
H(3)1	.11	.95	.87	.01	.01	.02
H(3)2	.10	.99	.01	.02	.02	.03
H(4)1	.43	.09	.81	.01	.01	.02
H(4)2	.35	.95	.86	.01	.01	.01

^a H(1)1 means that it is the 1'th hydrogen of water oxygen O(1).

TABLE II.

THERMAL PARAMETERS ^a AND STANDARD DEVIATIONS.

Atom	<u>B₁₁</u> ^b	<u>B₂₂</u>	<u>B₃₃</u>	<u>B₁₂</u>	<u>B₁₃</u>	<u>B₂₃</u>
Mn	1.98 ± .04	1.81 ± .04	1.50 ± .04	-0.02 ± .03	0.36 ± .03	0.01 ± .03
Cl(1)	2.16 ± .06	2.76 ± .07	2.51 ± .06	0.43 ± .05	0.65 ± .05	-0.29 ± .05
Cl(2)	2.22 ± .06	2.03 ± .06	2.38 ± .06	-0.29 ± .05	0.35 ± .05	0.00 ± .05
O(1)	3.3 ± .2	2.2 ± .2	2.0 ± .2	-0.1 ± .2	0.2 ± .1	0.1 ± .1
O(2)	2.5 ± .2	3.2 ± .2	2.2 ± .2	-0.2 ± .2	0.5 ± .2	0.3 ± .2
O(3)	4.1 ± .2	3.1 ± .2	3.4 ± .3	-1.2 ± .2	1.0 ± .2	-0.6 ± .2
O(4)	3.5 ± .2	2.6 ± .2	3.7 ± .2	0.1 ± .2	1.5 ± .2	-0.1 ± .2
Atom	<u>B</u> ^c	Atom	<u>B</u>	Atom	<u>B</u>	
H(1)1	2 ± 2	H(2)2	5 ± 3	H(4)1	6 ± 3	
H(1)2	3 ± 2	H(3)1	6 ± 5	H(4)2	5 ± 2	
H(2)1	1 ± 1	H(3)2	21 ± 9			

^a Units are Å².^b The anisotropic values $\underline{B}_{ij} = 4\beta_{ij}/\underline{a}_i^* \underline{a}_j^*$ where $\underline{a}_i^*$ is the *i*th reciprocal cell length.^c Isotropic.

TABLE III.

OBSERVED STRUCTURE FACTOR MAGNITUDES (FOB_h) AND CALCULATED STRUCTURE FACTORS (FCAL_h).

(Table submitted as photographic print)

Standard deviations of the parameters of the atoms were calculated assuming that the discrepancies in the structure factors represent random errors. The high standard deviations of the parameters of the hydrogen atoms illustrate that we are at the limit of our data's accuracy in determining these values.

[illegible]

Description of the Structure.—The structure consists of discrete octahedral groups with each manganese atom co-ordinated to two chlorine atoms at an average distance of 2.488 Å and to four oxygen atoms at an average distance of 2.206 Å. Of the eight hydrogen atoms only four are involved in hydrogen bonding which is discussed below. The over-all packing is illustrated in Fig. 1 and the dimensions of the octahedron shown in Table IV. A complete list of distances less than 4.0 Å is shown in Table V; the hydrogen atom distances are excluded.

TABLE IV

DIMENSIONS OF THE OCTAHEDRON AROUND MANGANESE IN $\text{MnCl}_2 \cdot 4\text{H}_2\text{O}$.

Atoms		Distance, Å	
Mn	- Cl(1)	$2.500 \pm .002$	(2.506) ^a
	- Cl(2)	$2.475 \pm .002$	(2.478)
	- O(1)	$2.224 \pm .004$	(2.229)
	- O(2)	$2.209 \pm .004$	(2.215)
	- O(3)	$2.185 \pm .006$	(2.198)
	- O(4)	$2.206 \pm .005$	(2.216)
Atoms		Angles	
Cl(1)	- Mn - Cl(2)	96°	O(1) - Mn - O(2) 177°
	- O(1)	92°	- O(3) 87°
	- O(2)	87°	- O(4) 92°
	- O(3)	93°	O(2) - Mn - O(3) 92°
	- O(4)	174°	- O(4) 90°
Cl(2)	- Mn - O(1)	88°	O(3) - Mn - O(4) 82°
	- O(2)	94°	
	- O(3)	169°	
	- O(4)	89°	

^aDistances in parentheses are values corrected for thermal motion assuming that the O and Cl atoms "ride" on the Mn.

TABLE V

DISTANCES UNDER 4.0 Å WITH STANDARD DEVIATIONS IN $\text{MnCl}_2 \cdot 4\text{H}_2\text{O}$. Å

Atom	Distance, Å	Atom	Distance, Å
Mn - O(1)	2.224 ± .004	O(2) - Mn	2.209 ± .004
- O(2)	2.209 ± .004	- O(4)	3.117 ± .007 <u>b</u>
- O(3)	2.185 ± .006	- O(3)	3.146 ± .008 <u>b</u>
- O(4)	2.206 ± .005	- Cl(1)	3.233 ± .005 <u>b</u>
- Cl(1)	2.500 ± .002	- Cl(2)	3.442 ± .005 <u>b</u>
- Cl(2)	2.475 ± .002	- O(1)	2.926 ± .006 <u>c</u>
Cl(1)- Mn	2.500 ± .002	- Cl(2)	3.166 ± .005 <u>c</u>
- O(2)	3.233 ± .005 <u>b</u>	- Cl(1)	3.477 ± .005
- O(1)	3.391 ± .005 <u>b</u>	O(3) - Mn	2.185 ± .006
- O(3)	3.414 ± .006 <u>b</u>	- O(4)	2.872 ± .008 <u>b</u>
- Cl(2)	3.708 ± .003 <u>b</u>	- O(1)	3.033 ± .008 <u>b</u>
- O(1)	3.169 ± .005 <u>c</u>	- O(2)	3.146 ± .008 <u>b</u>
- O(4)	3.292 ± .005 <u>c</u>	- Cl(1)	3.414 ± .006 <u>b</u>
- O(2)	3.477 ± .005	- Cl(2)	3.815 ± .007
Cl(2)- Mn	2.475 ± .002	O(4) - Mn	2.206 ± .005
- O(1)	3.254 ± .005 <u>b</u>	- O(3)	2.872 ± .008 <u>b</u>
- O(4)	3.285 ± .005 <u>b</u>	- O(2)	3.117 ± .007 <u>b</u>
- O(2)	3.442 ± .005 <u>b</u>	- O(1)	3.185 ± .006 <u>b</u>
- Cl(1)	3.708 ± .003 <u>b</u>	- Cl(2)	3.285 ± .005 <u>b</u>
- O(2)	3.166 ± .005 <u>c</u>	- Cl(1)	3.292 ± .005 <u>c</u>
- O(3)	3.815 ± .007	- O(1)	3.339 ± .006
O(1) - Mn	2.224 ± .004	O(1) - Cl(1)	3.391 ± .005 <u>b</u>
- O(3)	3.033 ± .008 <u>b</u>	- O(2)	2.926 ± .006 <u>c</u>
- O(4)	3.185 ± .006 <u>b</u>	- Cl(1)	3.169 ± .005 <u>c</u>
- Cl(2)	3.254 ± .005 <u>b</u>	- O(4)	3.339 ± .006

TABLE V (continued)

^aHydrogen atom distances are not included.

^bOctahedral edge.

^cHydrogen bond.

The octahedron is only slightly distorted with the primary distortion due to the packing of the two larger chlorine atoms with the four somewhat smaller oxygen atoms. The chlorine atoms are adjacent in each octahedron, and not opposite as in the $\text{FeCl}_2 \cdot 4\text{H}_2\text{O}$ structure⁴ although both structures have discrete octahedra. Also in the $\text{FeCl}_2 \cdot 4\text{H}_2\text{O}$ structure there are two different octahedral Fe-O distances (2.09 Å and 2.59 Å), whereas in the $\text{MnCl}_2 \cdot 4\text{H}_2\text{O}$ all four Mn-O distances are almost equivalent (2.19 Å to 2.22 Å). The densities of the two substances are very similar (2.03 g./cc. for the Mn compound, 1.98 g./cc. for the Fe compound), indicating that the efficiency of packing is also comparable. These comparisons suggest a rather interesting phase transformation if the metastable β -form of $\text{MnCl}_2 \cdot 4\text{H}_2\text{O}$ is indeed isomorphous with $\text{FeCl}_2 \cdot 4\text{H}_2\text{O}$ as is suggested by Groth².

According to Groth² $\text{MnBr}_2 \cdot 4\text{H}_2\text{O}$ has axial cell ratios, and β angle very close to $\text{MnCl}_2 \cdot 4\text{H}_2\text{O}$ and is presumably isostructural with the structure described in this paper.

There is a similarity between the $\text{MnCl}_2 \cdot 4\text{H}_2\text{O}$ structure and that reported by Culot et al¹² for $\text{NaBr} \cdot 2\text{H}_2\text{O}$. Dawson and Williams³ remark that $\text{MnCl}_2 \cdot 4\text{H}_2\text{O}$ and $\text{NaCl} \cdot 2\text{H}_2\text{O}$ are isomorphous, and the relation is further expanded by Groth² who states that $\text{NaCl} \cdot 2\text{H}_2\text{O}$ and $\text{NaBr} \cdot 2\text{H}_2\text{O}$ are isomorphous. If the monoclinic cell for $\text{NaBr} \cdot 2\text{H}_2\text{O}$ is doubled along its a axis and a different setting chosen, the resulting cell has axial ratios and β angle as follows,
 $\underline{a} : \underline{b} : \underline{c} = 1.171 : 1 : 0.653$ with $\beta = 97.3^\circ$; this is quite similar to the values for $\text{MnCl}_2 \cdot 4\text{H}_2\text{O}$ reported here. The $\text{NaBr} \cdot 2\text{H}_2\text{O}$ structure also consists of octahedra with the halide atoms adjacent. These octahedra

however, each share an edge with each of three adjacent octahedra, consistent with one extra anion in the formula, and form sheets in contrast with discrete octahedra in $\text{MnCl}_2 \cdot 4\text{H}_2\text{O}$.

Hydrogen Bonding—All eight independent H atoms were located from the difference function calculated after an anisotropic refinement of the seven heavier atoms. Four of these are in a suitable environment to participate in hydrogen bonding, but the other four have no close neighbors that would indicate such bonding. A representative of each independent H atom is shown in Fig. 1 and appropriate distances for the four bonded hydrogens are indicated in Table V. The four hydrogen bonds are as follows.

O(1)——H(1)1:.....	Cl(1)	3.17 Å
O(2)——H(2)1.....	Cl(2)	3.17 Å
O(2)——H(2)2.....	O(1)	2.93 Å
O(4)——H(4)1.....	Cl(1)	3.29 Å

All of the distances above represent hydrogen bonding between atoms in different octahedra.

The distances calculated for the O-H distances in water from the parameters listed in Table I are .8 and 1.0 Å for O(1); .8 and .9 Å for O(2); .5 and .6 Å for O(3); and .9 and 1.0 Å for O(4). It is because of the limited accuracy of our hydrogen parameters that we have not presented a detailed list of distances to the hydrogen atoms.

References

- (1) Work done under the auspices of the U. S. Atomic Energy Commission.
- (2) P. Groth, "Chemische Krystallographie," Wilhelm Engelmann, Leipzig, 1908, Vol. I.
- (3) H. M. Dawson and P. Williams, Z. physik. Chem., 31, 59 (1899).
- (4) B. R. Penfold and J. A. Grigor, Acta Cryst., 12, 850 (1959).
- (5) Z. M. El Saffar and C. R. K. Murtz, Acta Cryst., 15, 285 (1962).
- (6) C. Delain, Compt. Rend. Acad. Sci. Paris, 238, 1245 (1954).

- (7) W. E. Gardner, Bull. Amer. Phys. Soc., 5, 458 (1960).
- (8) J. D. H. Donnay, "Crystal Data," American Crystallographic Association Monograph, 2nd. Edition, 1963, p. 161.
- (9) J. A. Ibers, "International Tables for X-ray Crystallography," Kynoch Press, Birmingham, England, 1962, Vol. III, p. 210.
- (10) D. H. Templeton, *ibid.*, p. 215.
- (11) "Handbook of Chemistry and Physics," 39th Edition, Chemical Rubber Publishing Co., Ohio, 1957, p. 554.
- (12) J. P. Culot, P. Piret and M. Van Meerssche, Bull. Soc. Franc. Miner. Crist., 85, 282, (1962).

Figure Caption

Fig. 1.—Atomic arrangement in $\text{MnCl}_2 \cdot 4\text{H}_2\text{O}$. y coordinates ($\times 100$) are indicated as numbers inside some of the circles. The small solid circles represent hydrogen atoms, and only one of each type is shown.

MUB-2101