

UC Berkeley

UC Berkeley Previously Published Works

Title

Circadian control of neuroendocrine function: implications for health and disease

Permalink

<https://escholarship.org/uc/item/7z2301n6>

Journal

Current Opinion in Physiology, 5

ISSN

2468-8681

Authors

Gotlieb, Neta

Moeller, Jacob

Kriegsfeld, Lance J

Publication Date

2018-10-01

DOI

10.1016/j.cophys.2018.11.001

Peer reviewed



Published in final edited form as:

Curr Opin Physiol. 2018 October ; 5: 133–140. doi:10.1016/j.cophys.2018.11.001.

Circadian Control of Neuroendocrine Function: Implications for Health and Disease

Neta Gotlieb¹, Jacob Moeller², and Lance J. Kriegsfeld^{1,2,3,*}

¹Department of Psychology, University of California, Berkeley, CA, 94720

²Graduate Group in Endocrinology, University of California, Berkeley, CA 94720

³The Helen Wills Neuroscience Institute, University of California, Berkeley, CA, 94720

Abstract

The circadian timing system orchestrates daily rhythms in physiology and behavior via the suprachiasmatic nucleus (SCN), the master brain clock. Because endocrine secretions have far-reaching influence on the brain and periphery, circadian regulation of hormones is essential for normal functioning and disruptions to circadian timing (e.g., irregular sleep patterns, limited exposure to sunlight, jet lag, nighttime light exposure) have detrimental health consequences. Herein, we provide an overview of circadian timing in three major endocrine axes, the hypothalamo-pituitary-gonadal (HPG), hypothalamo-pituitary-adrenal (HPA) and hypothalamo-pituitary-thyroid (HPT) axes, and then consider the negative health consequences of circadian disruptions in each of these systems. For example, disruptions to HPG axis circadian timing lead to a host of negative reproductive outcomes such as irregular menstrual cycles, low sperm density and increased rates of miscarriages and infertility. Dysregulation of HPA axis timing is associated with obesity and metabolic disease, whereas disruptions to the HPT axis are associated with dysregulated metabolic gene rhythms in the heart. Together, this overview underscores the significance of circadian endocrine rhythms in normal health and disease prevention.

Keywords

biological rhythms; circadian; hormone; reproduction; HPA; HPG; HPT

1. Introduction

Walter Cannon coined the term homeostasis (standing the same) to describe the remarkable precision with which brain and bodily processes are maintained within stable operating parameters to promote optimal health and prevent illness [1]. However, physiological and behavioral needs vary markedly and predictably over the course of the day, necessitating that

*Address correspondence to: Lance J. Kriegsfeld, PhD, Department of Psychology and The Helen Wills Neuroscience Institute, 2121 Berkeley Way, University of California, Berkeley, CA 94720, kriegsfeld@berkeley.edu, Phone: 510-642-5148, Fax: 510-642-5293.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

biological systems adjust correspondingly. The circadian timing system synergizes with homeostatic drive to anticipate changing daily requirements and modify central and peripheral physiology accordingly. Under ideal circumstances, exposure to sunlight during the day and darkness at night optimally entrains (synchronizes) endogenously-generated circadian rhythms to environmental time to temporally coordinate neural and hormonal events for optimal health and functioning. Unfortunately, a major consequence of the modern lifestyle is increased exposure to sun-free environments during the day and artificial lighting at night, resulting in an incongruence between the endogenous circadian timing system and the external environment [2,3]. Such concerns have attracted the attention of the medical community, with the American Medical Association adopting a policy statement on the dangers of light at night for a number of maladies [4].

Hormones are substantial regulators of biological and behavioral events, including sexual motivation and reproduction, feeding and metabolism, sleep and vigilance, and immune function. Given the broad functional implications of hormones and their ability to travel long distances through the bloodstream, rhythms in endocrine secretions have far-reaching consequences for physiology and behavior [5–8]. Likewise, circadian-controlled rhythms in physiology and behavior (e.g., feeding) can influence rhythmic endocrine secretion (e.g., [9]; Figure 1). In the present overview, we consider circadian timing in three major endocrine axes (the hypothalamo-pituitary-gonadal (HPG), hypothalamo-pituitary-adrenal (HPA) and hypothalamo-pituitary-thyroid (HPT) axes) their functional significance, and clinical implications when their circadian timing is disrupted.

2. The Circadian Timing System

The circadian system includes a master brain clock in the suprachiasmatic nucleus (SCN) of the anterior hypothalamus that is synchronized to environmental time via a direct retinal pathway [10]. The SCN has direct access to environmental time cues and uses neural, diffusible and autonomic communication to convey timing information to the whole organism [5,11]. By communicating to hypothalamic neuroendocrine cells and hormone-producing glands, the SCN has widespread influence over the timing of physiology and behavior. At the cellular level, circadian rhythms are generated by an autoregulatory transcription-translation feedback loop consisting of clock genes and their protein products [12]. The core feedback loop begins in the morning with the clock protein, CLOCK, binding to BMAL1 to drive the transcription of the *Period* (*Per1* and *Per2*) and *Cryptochrome* (*Cry1* and *Cry2*) genes. Over the course of the day, *Per* and *Cry* transcripts are translated into their respective proteins that inevitably feed back to the cell nucleus to repress CLOCK:BMAL1-mediated transcription until the next morning when transcription resumes. Circadian timekeeping is a ubiquitous property of cells throughout the brain and body, with virtually all cells exhibiting circadian timekeeping [13]. However, in the absence of light input, the SCN maintains indefinite circadian rhythms at the tissue level due to unique coupling among independent oscillators in this master pacemaker. In contrast, in the absence of master clock communication or other entraining stimuli, extra-SCN brain loci and peripheral organs exhibit loss of rhythmicity after several cycles [14,15]. This loss of rhythmicity in extra-SCN systems results from loss of coupling among cellular oscillators having slightly different periods [16]. Disruptions to this circadian timing through night or rotating shift

work, international travel, irregular sleep patterns, limited exposure to sunlight, and exposure to light pollution and electronic devices at night precipitates a host of illnesses, including obesity and metabolic disease [17,18], breast cancer [19], prostate cancer [20], mental illness [21,22], and reproductive deficits [23,24].

3. The Hypothalamo-Pituitary Gonadal Axis

The HPG axis controls reproduction, including the generation and maintenance of gametes and sexual motivation and behavior. Secretion of hypothalamic gonadotropin-releasing hormone (GnRH) triggers the release of the gonadotropins, luteinizing hormone (LH) and follicle-stimulating hormone (FSH), from the anterior pituitary. In turn, LH and FSH stimulate gonadal sex steroid (i.e., estradiol, progesterone, and testosterone) synthesis and secretion and gamete maturation, respectively. Sex steroids and gonadotropins feed back onto the HPG axis to regulate its activity. Hormones produced by the HPG axis are under strong circadian control [5,6], with male and female reproductive function negatively affected by disruptions to circadian timing [23–25].

3.1 Circadian timing and female reproduction

Converging lines of evidence implicate a critical role for circadian timing in successful female reproduction across mammalian species, including humans (see [23,24,26–28] for comprehensive reviews on this topic). Ovulation, behavioral estrus, fertilization, pregnancy maintenance, and birth each require specific temporal patterns of hormone secretion regulated by the circadian system [29–33]. The negative consequences of chronic circadian disruption for female reproductive health are underscored in studies investigating women or animals with marked circadian deficits. Women with irregular work or sleep cycles, for example, exhibit abnormal menstrual cycles [34,35], reduced fertility [36,37] and increased miscarriage rates [37–39]. In rodents, destruction of the SCN, its neural output, or the genes regulating cellular clock function lead to pronounced abnormalities in ovulation and fecundity [40–44].

Because the majority of functional studies have explored the role of the circadian system in ovulation, the present overview will focus on this aspect of the female reproductive cycle. In spontaneously ovulating mammals, estradiol secretion from maturing ovarian follicles maintains LH at low concentrations through estradiol negative feedback during the follicular phase of the ovulatory cycle. Just prior to ovulation, estradiol negative feedback is suppressed and estradiol acts through positive feedback to stimulate the LH surge that initiates ovulation. Findings across rodent species by our group and others have shown that the SCN acts to coordinate the timing of negative and positive feedback via actions on the inhibitory neuropeptide, gonadotropin-inhibitory hormone (GnIH; also known as RFamide-related peptide 3) and the stimulatory neuropeptide, kisspeptin (reviewed in [45–47]). At the time of the preovulatory LH surge that stimulates ovulation, the SCN suppresses estradiol negative feedback by acting on GnIH neurons and stimulates estradiol positive drive via actions on the kisspeptin and GnRH systems. In addition to these monosynaptic projections to neuroendocrine cells, the SCN may be communicating via autonomic outflow through the PVN to the ovary [48,49]. Finally, circadian timekeeping at the level of GnIH, GnRH and

kisspeptin neurons furthers precision in the balance of these negative and positive and regulators [50–53] (Figure 1).

In addition to actions at the level of the brain, cellular clocks in the ovary have been implicated in follicular growth, hormone synthesis, and ovulation [54], with clock gene expression observed in theca, granulosa, and luteal cells [55]. Additionally, abnormal rhythms of *Per2* are seen in ovary in a mouse model of polycystic ovarian syndrome [56], suggesting abnormal ovarian rhythmicity may contribute to the symptoms of the disease. Using RT-qPCR, rhythms in clock genes are observed in the rat ovary across the ovulatory cycle [57]. Likewise, in rats with a luciferase reporter for *Per1*, circadian rhythms are seen in the ovary *in vitro* with large phase adjustments observed in response to LH and FSH [58]. *In vivo*, rats treated with LH during the subjective night ovulate more frequently and produced more oocytes than animals treated during the subjective day [59]. Together, these results suggest that ovarian timers mediate the sensitivity of the ovary to LH across the day, further establishing an important role for the circadian system in ovulation. The generation of ovarian cell-phenotype-specific clock gene knockout/knockdown mice will help to clarify the specific role of ovarian clocks.

3.2 Circadian timing and male reproductive function

Although the significance of circadian timing in male reproduction has received less attention than that of females, several lines of evidence point to an important role for rhythmicity in males. Men exhibit daily changes in semen quality and sperm numbers [60,61], and those who work irregular shifts exhibit compromised fertility and lower sperm density, motile count, and testosterone [20]. Global knockout of *Bmal1* leads to loss of circadian rhythmicity and infertility in male mice, presumably due to abnormal hormone secretion, small testes and seminal vesicles, and low sperm count [62]. However, testes from these mice still produce viable sperm, suggesting infertility may be due to alterations in behavior in these mice. More recent findings reveal that *Bmal1* knockout males fail to mate with receptive females [63]. Interestingly, this deficit appears to be due to abnormal olfactory processing. Although the vomeronasal organ (VNO) responds appropriately to pheromonal stimulation, hypothalamic targets of the VNO do not respond properly to downstream signaling from the VNO. These findings indicate that, although *Bmal1* knockout mice may be capable of sexual behavior, the motivation to engage in such behavior is abolished due to deficits in olfactory processing. Whether these deficits result from loss of rhythmicity in neural targets of the olfactory system or through pleiotropic effects resulting from the loss of *Bmal1* remains to be determined. Together, these findings point to several factors that may contribute to infertility in *Bmal1* knockout mice, deficits in hormone secretion, reduced sperm production, and reduced sexual motivation.

In mice, clock genes are expressed in the testes, but do not exhibit detectable rhythms [64–66]. In *Clock* mutant mice, *Per1* expression is not altered relative to wild-type mice, indicating that typical CLOCK:BMAL1-mediated transcription may not drive *Period* gene production in mouse testis [66]. Mice lacking *Bmal1* have reduced steroidogenic acute regulatory protein (StAR), the rate-limiting enzyme in steroidogenesis, reduced serum testosterone levels, and elevated LH concentrations [62], pointing to a central role for this

clock gene in normal testicular function. Intriguingly, in Syrian hamster testis, *Per1* and *Bmal1* are expressed rhythmically [67]. This same study identified two *Per1* transcripts in testis that differed from those seen in mice and other hamster organs, with these transcript variants lacking a nuclear localization signal and lacking a putative CRY1-binding domain. Whether or not these hamster transcript variants account for the apparent rhythmicity of testicular clock gene expression in this species remains to be determined.

4. Hypothalamo-Pituitary Adrenal Axis

The HPA axis regulates arousal and energy mobilization under typical conditions and rapidly mobilizes energy from stored sources to facilitate the fight or flight response. Analogous to the HPG axis, the hypothalamic peptide, corticotropin-releasing hormone (CRH), is released into the anterior pituitary blood supply and stimulates the release of adrenocorticotrophic hormone (ACTH). ACTH released into systemic circulation, in turn, acts on the adrenal cortex to stimulate glucocorticoid (GC) (i.e., cortisol and corticosterone) release. GC acts broadly within the brain and body through negative feedback to inhibit its own production. Humans, non-human primates, and rodents exhibit pronounced daily GC rhythms that persist in constant conditions, with GC concentrations rising prior to waking, decreasing throughout the day, and falling in anticipation of sleep [5,68,69]. Like the ovaries, the adrenal glands exhibit rhythms in clock gene expression that likely drive daily changes in responsiveness to ACTH stimulation and stress [70–73]. Rhythms in adrenal GC secretion and adrenal clock gene expression are eliminated by SCN lesions [74,75], suggesting that circadian rhythms in individual cells of the adrenal become uncoupled in the absence of SCN input.

The SCN drives rhythmic secretion of GC through several pathways (Figure 1). The first pathway indirectly targets CRH neurons in the paraventricular nucleus of the hypothalamus (PVN) through SCN arginine vasopressin-ergic projections to an area just below the PVN (the subPVN) and the dorsomedial hypothalamus (DMH) [76–78]. In turn, the subPVN and DMH regulate CRH production. Secondly, the SCN continues through this PVN pathway, sending autonomic outflow through a multisynaptic projection to the adrenal cortex [79]. As removal of the pituitary (and resulting abolition of ACTH secretion) does not alter clock gene rhythmicity in the adrenal cortex [72], it is likely that SCN control of autonomic input to the adrenals is responsible for the coordination/maintenance of adrenal cellular clocks. Finally, feeding influences GC rhythms, underscoring the importance of circadian-controlled behavior in maintaining typical endocrine rhythmicity [9].

GC can act throughout the periphery to set the phase of oscillators in individual peripheral systems [9,80–82]. As a result, disruptions to GC rhythms have far-reaching, negative impact on normal physiology, particularly metabolism (see [83] a review of this topic). Travel in humans and experimental jet lag in rodents increases GC [84,85], contributing to negative health consequences of circadian disruption. In addition, advancing the sleep-wake and light-dark cycles by 8 hours in humans results in elevated nighttime cortisol concentrations, likely contributing to difficulty falling asleep at the new circadian phase. Individuals suffering from depression have abnormal cortisol and ACTH rhythms, with the trough of

both hormones advanced 3 h [86,87]. Whether disruptions to the cortisol rhythm are a cause or consequence of depressive symptoms remains to be determined.

Given the role of GC in energy mobilization and utilization, disruptions to HPA axis rhythmicity are implicated in obesity and metabolic disease. Chronic stress, for example, results in elevated GC concentrations and abnormal GC rhythmicity and is associated with obesity, insulin resistance, dyslipidemia, hypertension, and hyperglycemia [83]. Obese mice and humans exhibit flattened GC rhythms, further pointing to HPA axis dysregulation in obesity [88,89]. People with select polymorphisms in the *Clock* gene are at greater risk for obesity and metabolic disease [90,91]. Furthermore, patients with abnormal HPA axis functioning and receiving cortisol replacement at tonic high levels (rather than mimicking the endogenous rhythm) are at greater risk for cardiovascular and metabolic bone disease [92]. Likewise, *Cry* deficient mice exhibit HPA axis deficits and are vulnerable to obesity and enhanced fat deposition when fed a high fat diet [93]. Similarly, mice deficient in *Clock* or *Bmal1* exhibit abnormal glucose and triglycerides rhythms, and develop obesity, hyperlipidemia and diabetes mellitus [94,95]. Although these disease risks have not been directly linked to disruptions in HPA axis rhythmicity in these knockout mice, it is noteworthy that glucocorticoid excess results in the same negative outcomes across studies [83]. Whereas it has been challenging to specifically link alterations in HPA axis rhythms directly to metabolic outcomes, these converging lines of evidence, along with established functions of GC, suggest that disrupted HPA rhythms contribute to metabolic dysregulation.

4. Hypothalamo-Pituitary Thyroid Axis

The HPT axis is responsible for regulating metabolism. Hypothalamic release of thyrotropin-releasing hormone (TRH) into the anterior pituitary stimulates production and release of thyroid-stimulating hormone (TSH) in the general circulation. TSH stimulates production of thyroid hormones from the thyroid gland. Thyroid hormones are initially produced as thyroglobulin, which is converted primarily to thyroxine (T4). T4 is considered inactive and is further converted into the active thyroid hormone, triiodothyronine (T3) in target tissues. In humans, the HPT axis is under circadian control, with free T3 and TSH being low during the day and high at night [96,97]. In rats, TSH is rhythmic and in antiphase to that of humans (low during the night and high during the day), with rhythmic secretion abolished by SCN lesions [98–100]. As with the adrenals, clock genes (*Per1* and *Bmal1*) are rhythmically expressed in the rat thyroid [101]. Whereas daily rhythms in thyroid hormones are abolished by hypophysectomy, rhythms in clock gene expression are unaffected, suggesting that thyroid clock maintenance is accomplished through SCN autonomic innervation of the thyroid gland. Indeed, retrograde transneuronal tracing from the thyroid gland reveals multisynaptic projections from the SCN [100]. This same study found direct projections to TSH neurons located in the PVN, indicating that, as with the HPA and potentially the HPG axes, the SCN regulates thyroid hormone secretion both via actions on TSH neuroendocrine cells and via autonomic outflow to the thyroid (Figure 1).

Relative to the HPA and HPG axis, less work has focused on the negative effects of disrupted HPT axis rhythms. Given the contribution of circadian disruption to obesity and metabolic disease, it is feasible that disruptions to the HPT axis contribute to these

considers sex difference in the circadian system underscoring the utility of considering sex differences in circadian timing in health and disease disparity.

- [9]. Ikeda Y, Sasaki H, Ohtsu T, Shiraishi T, Tahara Y, Shibata S, Feeding and adrenal entrainment stimuli are both necessary for normal circadian oscillation of peripheral clocks in mice housed under different photoperiods. *Chronobiol Int* 32 (2015) 195–210. [PubMed: 25286135]
- [10]. Morin LP, Allen CN, The circadian visual system, 2005. *Brain Res Rev* 51 (2006) 1–60. [PubMed: 16337005]
- [11]. Buijs RM, Escobar C, Swaab DF, The circadian system and the balance of the autonomic nervous system. *Handb Clin Neurol* 117 (2013) 173–191. [PubMed: 24095125]
- [12]. Takahashi JS, Molecular components of the circadian clock in mammals. *Diabetes Obes Metab* 17 Suppl 1 (2015) 6–11. [PubMed: 26332962]
- [13]. Buijs R, Salgado R, Sabath E, Escobar C, Peripheral circadian oscillators: time and food. *Prog Mol Biol Transl Sci* 119 (2013) 83–103. [PubMed: 23899595]
- [14]. Abe M, Herzog ED, Yamazaki S, Straume M, Tei H, Sakaki Y, Menaker M, Block GD, Circadian rhythms in isolated brain regions. *J Neurosci* 22 (2002) 350–356. [PubMed: 11756518]
- [15]. Yamazaki S, Numano R, Abe M, Hida A, Takahashi R, Ueda M, Block GD, Sakaki Y, Menaker M, Tei H, Resetting central and peripheral circadian oscillators in transgenic rats. *Science* 288 (2000) 682–685 [PubMed: 10784453]
- [16]. Welsh DK, Yoo SH, Liu AC, Takahashi JS, Kay SA, Bioluminescence imaging of individual fibroblasts reveals persistent, independently phased circadian rhythms of clock gene expression. *Curr Biol* 14 (2004) 2289–2295. [PubMed: 15620658]
- [17]. Saulle R, Bernardi M, Chiarini M, Backhaus I, La Torre G, Shift work, overweight and obesity in health professionals: a systematic review and meta-analysis. *Clin Ter* 169 (2018) e189–e197. [PubMed: 30151553]
- [18]. Liu Q, Shi J, Duan P, Liu B, Li T, Wang C, Li H, Yang T, Gan Y, Wang X, et al., Is shift work associated with a higher risk of overweight or obesity? systematic review of observational studies with meta-analysis. *Int J Epidemiol* (2018)* Meta-analysis exploring the impact of shift work/night work on nurses.
- [19]. Ball LJ, Palesh O, Kriegsfeld LJ, The Pathophysiologic Role of Disrupted Circadian and Neuroendocrine Rhythms in Breast Carcinogenesis. *Endocr Rev* 37 (2016) 450–466. [PubMed: 27712099]
- [20]. Deng N, Kohn TP, Lipshultz LI, Pastuszak AW, The Relationship Between Shift Work and Men's Health. *Sex Med Rev* 6 (2018) 446–456. [PubMed: 29371140] * Overview of the literature examining the effects of shift work on men's health. The studies highlighted reveal an association between shift work and hypogonadism, poor semen quality, decreased fertility, and prostate cancer
- [21]. Khan S, Duan P, Yao L, Hou H, Shiftwork-Mediated Disruptions of Circadian Rhythms and Sleep Homeostasis Cause Serious Health Problems. *Int J Genomics* 2018 (2018) 8576890. [PubMed: 29607311] * Review of the negative health consequences associated with shift work.
- [22]. Angerer P, Schmook R, Elfantel I, Li J, Night Work and the Risk of Depression. *Dtsch Arztebl Int* 114 (2017) 404–411. [PubMed: 28669378] * review of the literature and meta-analysis reporting a 42% increased risk of depression among night shift workers.
- [23]. Gamble KL, Resuehr D, Johnson CH, Shift work and circadian dysregulation of reproduction. *Front Endocrinol (Lausanne)* 4 (2013) 92. [PubMed: 23966978]
- [24]. Mahoney MM, Shift work, jet lag, and female reproduction. *Int J Endocrinol* 2010 (2010) 813764. [PubMed: 20224815]
- [25]. Kennaway DJ, Boden MJ, Varcoe TJ, Circadian rhythms and fertility. *Mol Cell Endocrinol* 349 (2012) 56–61. [PubMed: 21872642]
- [26]. Boden MJ, Varcoe TJ, Kennaway DJ, Circadian regulation of reproduction: from gamete to offspring. *Prog Biophys Mol Biol* 113 (2013) 387–397. [PubMed: 23380455]
- [27]. Williams WP, 3rd, Kriegsfeld LJ, Circadian control of neuroendocrine circuits regulating female reproductive function. *Front Endocrinol (Lausanne)* 3 (2012) 60. [PubMed: 22661968]

- [28]. Simonneaux V, Bahougne T, A Multi-Oscillatory Circadian System Times Female reproduction. *Front Endocrinol (Lausanne)* 6 (2015) 157. [PubMed: 26539161]
- [29]. Mong JA, Pfaff DW, Hormonal and genetic influences underlying arousal as it drives sex and aggression in animal and human brains. *Neurobiol Aging* 24 Suppl 1 (2003) S83–88; discussion S91–82. [PubMed: 12829115]
- [30]. McEwen BS, Jones KJ, Pfaff DW, Hormonal control of sexual behavior in the female rat: molecular, cellular and neurochemical studies. *Biol Reprod* 36 (1987) 37–45. [PubMed: 2436675]
- [31]. Blaustein JD, Tetel MJ, Ricciardi KH, Delville Y, Turcotte J, Hypothalamic ovarian steroid hormone-sensitive neurons involved in female sexual behavior. *Psychoneuroendocrinology* 19 (1994) 505–516. [PubMed: 7938350]
- [32]. Kriegsfeld LJ, Silver R, Gore AC, Crews D, Vasoactive intestinal polypeptide contacts on gonadotropin-releasing hormone neurones increase following puberty in female rats. *J Neuroendocrinol* 14 (2002) 685–690. [PubMed: 12213129]
- [33]. Egli M, Bertram R, Sellix M, Freeman M, Rhythmic secretion of prolactin in rats: action of oxytocin coordinated by vasoactive intestinal polypeptide of suprachiasmatic nucleus origin. *Endocrinology* 145 (2004) 3386–3394. [PubMed: 15033917]
- [34]. Wang Y, Gu F, Deng M, Guo L, Lu C, Zhou C, Chen S, Xu Y, Rotating shift work and menstrual characteristics in a cohort of Chinese nurses. *BMC Womens Health* 16 (2016) 24. [PubMed: 27145834]
- [35]. Lawson CC, Whelan EA, Lividoti Hibert E. N., Spiegelman D, Schernhammer ES, Rich-Edwards JW, Rotating shift work and menstrual cycle characteristics. *Epidemiology* 22 (2011) 305–312. [PubMed: 21364464]
- [36]. Ahlborg G, Jr., Axelsson G, Bodin L, Shift work, nitrous oxide exposure and subfertility among Swedish midwives. *Int J Epidemiol* 25 (1996) 783–790. [PubMed: 8921457]
- [37]. Fernandez RC, Marino JL, Varcoe TJ, Davis S, Moran LJ, Rumbold AR, Brown HM, Whitrow MJ, Davies MJ, Moore VM, Fixed or Rotating Night Shift Work Undertaken by Women: Implications for Fertility and Miscarriage. *Semin Reprod Med* 34 (2016) 74–82. [PubMed: 26854708] * Explores the impact of circadian disruption on menstrual cycle irregularities, endometriosis, and miscarriage rates. Significantly, this overview distinguishes between circadian disruption through night work versus rotating shift work.
- [38]. Nurminen T, Shift work and reproductive health. *Scand J Work Environ Health* 24 Suppl 3 (1998) 28–34. [PubMed: 9916814]
- [39]. Lawson CC, Rocheleau CM, Whelan EA, N Lividoti Hibert E, Grajewski B, Spiegelman D, W Rich-Edwards J, Occupational exposures among nurses and risk of spontaneous abortion. *Am J Obstet Gynecol* 206 (2012) 327 e321–328. [PubMed: 22304790]
- [40]. Miller BH, Olson SL, Turek FW, Levine JE, Horton TH, Takahashi JS, Circadian clock mutation disrupts estrous cyclicity and maintenance of pregnancy. *Curr Biol* 14 (2004) 1367–1373. [PubMed: 15296754]
- [41]. Nunez AA, Stephan FK, The effects of hypothalamic knife cuts on drinking rhythms and the estrus cycle of the rat. *Behavioral biology* 20 (1977) 224–234. [PubMed: 561597]
- [42]. Wiegand SJ., Terasawa, Discrete lesions reveal functional heterogeneity of suprachiasmatic structures in regulation of gonadotropin secretion in the female rat. *Neuroendocrinology* 34 (1982) 395–404. [PubMed: 6808412]
- [43]. van der Horst GT, Muijtjens M, Kobayashi K, Takano R, Kanno S, Takao M, de Wit J, Verkerk A, Eker AP, van Leenen D, et al., Mammalian Cry1 and Cry2 are essential for maintenance of circadian rhythms. *Nature* 398 (1999) 627–630. [PubMed: 10217146]
- [44]. Chu A, Zhu L, Blum ID, Mai O, Leliavski A, Fahrenkrug J, Oster H, Boehm U, Storch KF, Global but not gonadotrope-specific disruption of Bmal1 abolishes the luteinizing hormone surge without affecting ovulation. *Endocrinology* 154 (2013) 2924–2935. [PubMed: 23736292]
- [45]. Kriegsfeld LJ, Circadian regulation of kisspeptin in female reproductive functioning. *Adv Exp Med Biol* 784 (2013) 385–410. [PubMed: 23550016]
- [46]. Simonneaux V, Bahougne T, Angelopoulou E, Daily rhythms count for female fertility. *Best Pract Res Clin Endocrinol Metab* 31 (2017) 505–519. [PubMed: 29223284] * Comprehensive review

on the mechanisms by which the circadian system participates in female reproduction, including control by the SCN and extra-SCN/peripheral oscillators.

- [47]. Kriegsfeld LJ, Jennings KJ, Bentley GE, Tsutsui K, Gonadotrophin-inhibitory hormone and its mammalian orthologue RFamide-related peptide-3: Discovery and functional implications for reproduction and stress. *J Neuroendocrinol* 30 (2018) e12597. [PubMed: 29624758] * Overview of the the impact of the inhibitory neuropeptide, RFamide-related peptide-3, on reproductive function and in mediating the response to stressors.
- [48]. Gerendai I, Toth I, Boldogkoi Z, Medveczky I, Halasz B, Neuronal labeling in the rat brain and spinal cord from the ovary using viral transneuronal tracing technique. *Neuroendocrinology* 68 (1998) 244–256. [PubMed: 9772339]
- [49]. Gerendai I, Toth I, Boldogkoi Z, Medveczky I, Halasz B, CNS structures presumably involved in vagal control of ovarian function. *J Auton Nerv Syst* 80 (2000) 40–45. [PubMed: 10742538]
- [50]. Russo KA, La JL, Stephens SB, Poling MC, Padgaonkar NA, Jennings KJ, Piekarski DJ, Kauffman AS, Kriegsfeld LJ, Circadian Control of the Female Reproductive Axis Through Gated Responsiveness of the RFRP-3 System to VIP Signaling. *Endocrinology* 156 (2015) 2608–2618. [PubMed: 25872006]
- [51]. Chassard D, Bur I, Poirel VJ, Mendoza J, Simonneaux V, Evidence for a Putative Circadian Kiss-Clock in the Hypothalamic AVPV in Female Mice. *Endocrinology* 156 (2015) 2999–3011. [PubMed: 25993523]
- [52]. Chappell PE, White RS, Mellon PL, Circadian gene expression regulates pulsatile gonadotropin-releasing hormone (GnRH) secretory patterns in the hypothalamic GnRH-secreting GT1–7 cell line. *J Neurosci* 23 (2003) 11202–11213. [PubMed: 14657179]
- [53]. Zhao S, Kriegsfeld LJ, Daily changes in GT1–7 cell sensitivity to GnRH secretagogues that trigger ovulation. *Neuroendocrinology* 89 (2009) 448–457. [PubMed: 19141986]
- [54]. Sellix MT, Circadian clock function in the mammalian ovary. *J Biol Rhythms* 30 (2015) 7–19. [PubMed: 25367899]
- [55]. Fahrenkrug J, Georg B, Hannibal J, Hindersson P., Gras, Diurnal rhythmicity of the clock genes *Per1* and *Per2* in the rat ovary. *Endocrinology* 147 (2006) 3769–3776. [PubMed: 16675517]
- [56]. Mereness AL, Murphy ZC, Sellix MT, Developmental programming by androgen affects the circadian timing system in female mice. *Biol Reprod* 92 (2015) 88. [PubMed: 25695720]
- [57]. Karman BN, Tischkau SA, Circadian clock gene expression in the ovary: Effects of luteinizing hormone. *Biol Reprod* 75(2006)624–632. [PubMed: 16807384]
- [58]. Yoshikawa T, Sellix M, Pezuk P, Menaker M, Timing of the ovarian circadian clock is regulated by gonadotropins. *Endocrinology* 150 (2009) 4338–4347. [PubMed: 19520783]
- [59]. Sellix MT, Yoshikawa T, Menaker M, A circadian egg timer gates ovulation. *Curr Biol* 20 (2010) R266–267. [PubMed: 20334830]
- [60]. Xie M, Utzinger KS, Blickenstorfer K, Leeners B, Diurnal and seasonal changes in semen quality of men in subfertile partnerships. *Chronobiol Int* (2018) 1–10.* study of semen samples from 7068 men showing circadian and seasonal changes in sperm quality.
- [61]. Cagnacci A, Maxia N, Volpe A, Diurnal variation of semen quality in human males. *Hum Reprod* 14 (1999) 106–109. [PubMed: 10374104]
- [62]. Alvarez JD, Hansen A, Ord T, Bebas P, Chappell PE, Giebultowicz JM, Williams C, Moss S, Sehgal A, The circadian clock protein BMAL1 is necessary for fertility and proper testosterone production in mice. *J Biol Rhythms* 23 (2008) 26–36. [PubMed: 18258755]
- [63]. Schoeller EL, Clark DD, Dey S, Cao NV, Semaan SJ, Chao LW, Kauffman AS, Stowers L, Mellon PL, Bmal1 Is Required for Normal Reproductive Behaviors in Male Mice. *Endocrinology* 157 (2016) 4914–4929. [PubMed: 27704948] ** Identified a role for the circadian clock in male reproduction using Bmal1 knockout mice. Established that a functional circadian clock is necessary for female pheromone detection and sexual motivation.
- [64]. Alvarez JD, Chen D, Storer E, Sehgal A, Non-cyclic and developmental stage-specific expression of circadian clock proteins during murine spermatogenesis. *Biol Reprod* 69 (2003) 81–91. [PubMed: 12606319]

- [65]. Bebas P, Goodall CP, Majewska M, Neumann A, Giebultowicz JM, Chappell PE, Circadian clock and output genes are rhythmically expressed in extratesticular ducts and accessory organs of mice. *FASEB J* 23 (2009) 523–533. [PubMed: 18945877]
- [66]. Morse D, Cermakian N, Brancorsini S., Parvinen, P, Sassone-Corsi, No circadian rhythms in testis:Period1 expression is clock independent and developmentally regulated in the mouse. *Mol Endocrinol* 17 (2003) 141–151. [PubMed: 12511614]
- [67]. Tong Y, Guo H, Brewer JM, Lee H, Lehman MN, Bittman EL, Expression of haPer1 and haBmal1 in Syrian hamsters: heterogeneity of transcripts and oscillations in the periphery. *J Biol Rhythms* 19 (2004) 113–125. [PubMed: 15038851]
- [68]. den Boon FS, Sarabdjitsingh RA. Circadian and ultradian patterns of HPA-axis activity in rodents: Significance for brain functionality. *Best Pract Res Clin Endocrinol Metab* 31 (2017) 445–457. [PubMed: 29223280] * Explores patterns of HPA axis rhythmicity at the circadian and ultradian time scales and considers the importance of these rhythms in gene expression, normal physiology, and stress responsivity.
- [69]. Kalsbeek A, van der Spek R, Lei J, Endert E, Buijs RM, Fliers E, Circadian rhythms in the hypothalamo-pituitary-adrenal (HPA) axis. *Mol Cell Endocrinol* 349 (2012) 20–29. [PubMed: 21782883]
- [70]. Kalsbeek A, Ruiters M, La Fleur SE, Van Heijningen C, Buijs RM, The diurnal modulation of hormonal responses in the rat varies with different stimuli. *J Neuroendocrinol* 15 (2003) 1144–1155 [PubMed: 14636176]
- [71]. Bittman EL, Doherty L, Huang L, Paroskie A, Period gene expression in mouse endocrine tissues. *Am J Physiol Regul Integr Comp Physiol* 285 (2003) R561–569. [PubMed: 12775559]
- [72]. Fahrenkrug J, Hannibal J, Georg B, Diurnal rhythmicity of the canonical clock genes Per1, Per2 and Bmal1 in the rat adrenal gland is unaltered after hypophysectomy. *J Neuroendocrinol* 20 (2008) 323–329. [PubMed: 18208549]
- [73]. Oster H, Damerow S, Kiessling S, Jakubcakova V, Abraham D, Tian J, Hoffmann MW, Eichele G, The circadian rhythm of glucocorticoids is regulated by a gating mechanism residing in the adrenal cortical clock. *Cell Metab* 4 (2006) 163–173. [PubMed: 16890544]
- [74]. Moore RY, Eichler VB, Loss of a circadian adrenal corticosterone rhythm following suprachiasmatic lesions in the rat. *Brain Res* 42 (1972) 201–206. [PubMed: 5047187]
- [75]. Guo H, Brewer JM, Lehman MN, Bittman EL, Suprachiasmatic regulation of circadian rhythms of gene expression in hamster peripheral organs: effects of transplanting the pacemaker. *J Neurosci* 26 (2006) 6406–6412. [PubMed: 16775127]
- [76]. Hermes ML, Ruijter JM, Klop A, Buijs RM, Renaud LP, Vasopressin increases GABAergic inhibition of rat hypothalamic paraventricular nucleus neurons in vitro. *J Neurophysiol* 83 (2000) 705–711. [PubMed: 10669486]
- [77]. Buijs RM, Hermes MH, Kalsbeek A, The suprachiasmatic nucleus-paraventricular nucleus interactions: a bridge to the neuroendocrine and autonomic nervous system. *Prog Brain Res* 119 (1998) 365–382. [PubMed: 10074800]
- [78]. Kalsbeek A, Verhagen LA, Schalij I, Foppen E, Saboureaux M, Bothorel B, Buijs RM, Pevet P, Opposite actions of hypothalamic vasopressin on circadian corticosterone rhythm in nocturnal versus diurnal species. *Eur J Neurosci* 27 (2008) 818–827. [PubMed: 18279365]
- [79]. Buijs RM, Wortel J, Van Heerikhuizen JJ, Feenstra MG, Ter Horst GJ, Romijn HJ, Kalsbeek A, Anatomical and functional demonstration of a multisynaptic suprachiasmatic nucleus adrenal (cortex) pathway. *Eur J Neurosci* 11 (1999) 1535–1544. [PubMed: 10215906]
- [80]. Pezuk P, Mohawk JA, Wang LA, Menaker M, Glucocorticoids as entraining signals for peripheral circadian oscillators. *Endocrinology* 153 (2012) 4775–4783. [PubMed: 22893723]
- [81]. Kamagata M, Ikeda Y, Sasaki H, Hattori Y, Yasuda S, Iwami S, Tsubosaka M, Ishikawa R, Todoh A, Tamura K, et al., Potent synchronization of peripheral circadian clocks by glucocorticoid injections in PER2::LUC-Clock/Clock mice. *Chronobiol Int* 34 (2017) 1067–1082. [PubMed: 28704069] * Study establishing that glucocorticoids can synchronize the peripheral oscillators in mice with a disrupted cellular clock mechanism.

- [82]. Balsalobre A, Brown SA, Marcacci L, Tronche F, Kellendonk C, Reichardt HM, Schutz G, Schibler U, Resetting of circadian time in peripheral tissues by glucocorticoid signaling. *Science* 289 (2000) 2344–2347. [PubMed: 11009419]
- [83]. Nader N, Chrousos GP, Kino T, Interactions of the circadian CLOCK system and the HPA axis. *Trends Endocrinol Metab* 21 (2010) 277–286. [PubMed: 20106676]
- [84]. Doane LD, Kremen WS, Eaves LJ, Eisen SA, Hauger R, Hellhammer D, Levine S, Lupien S, Lyons MJ, Mendoza S, et al., Associations between jet lag and cortisol diurnal rhythms after domestic travel. *Health Psychol* 29 (2010) 117–123. [PubMed: 20230083]
- [85]. Gibson EM, Wang C, Tjho S, Khattar N, Kriegsfeld LJ, Experimental ‘jet lag’ inhibits adult neurogenesis and produces long-term cognitive deficits in female hamsters. *PLoS One* 5 (2010) e15267. [PubMed: 21152025]
- [86]. Linkowski P, Mendlewicz J, Kerkhofs M, Leclercq R, Golstein J, Brasseur M, Copinschi G, Van Cauter E, 24-hour profiles of adrenocorticotropin, cortisol, and growth hormone in major depressive illness: effect of antidepressant treatment. *J Clin Endocrinol Metab* 65 (1987) 141–152. [PubMed: 3034952]
- [87]. Linkowski P, Mendlewicz J, Leclercq R, Brasseur M, Hubain P, Golstein J, Copinschi G, Van Cauter E, The 24-hour profile of adrenocorticotropin and cortisol in major depressive illness. *J Clin Endocrinol Metab* 61 (1985) 429–438. [PubMed: 2991318]
- [88]. Walker CD, Scribner KA, Stern JS, Dallman MF, Obese Zucker (fa/fa) rats exhibit normal target sensitivity to corticosterone and increased drive to adrenocorticotropin during the diurnal trough. *Endocrinology* 131 (1992) 2629–2637. [PubMed: 1332842]
- [89]. Ahima RS, Prabakaran D, Flier JS, Postnatal leptin surge and regulation of circadian rhythm of leptin by feeding. Implications for energy homeostasis and neuroendocrine function. *J Clin Invest* 101 (1998) 1020–1027. [PubMed: 9486972]
- [90]. Scott EM, Carter AM, Grant PJ, Association between polymorphisms in the Clock gene, obesity and the metabolic syndrome in man. *Int J Obes (Lond)* 32 (2008) 658–662. [PubMed: 18071340]
- [91]. Sookoian S, Gemma C, Gianotti TF, Burgueno A, Castano G, Pirola CJ, Genetic variants of Clock transcription factor are associated with individual susceptibility to obesity. *Am J Clin Nutr* 87 (2008) 1606–1615. [PubMed: 18541547]
- [92]. Debono M, Ross RJ, Newell-Price J, Inadequacies of glucocorticoid replacement and improvements by physiological circadian therapy. *Eur J Endocrinol* 160 (2009) 719–729. [PubMed: 19168600]
- [93]. Barclay JL, Shostak A, Leliavski A, Tsang AH, Johren O, Muller-Fielitz H, Landgraf D, Naujokat N, van der Horst GT, Oster H, High-fat diet-induced hyperinsulinemia and tissue-specific insulin resistance in Cry-deficient mice. *Am J Physiol Endocrinol Metab* 304 (2013) E1053–E1063 [PubMed: 23531614]
- [94]. Turek FW, Joshu C, Kohsaka A, Lin E, Ivanova G, McDearmon E, Laposky A, Losee-Olson S, Easton A, Jensen DR, et al., Obesity and metabolic syndrome in circadian Clock mutant mice. *Science* 308 (2005) 1043–1045. [PubMed: 15845877]
- [95]. Rudic RD, McNamara P, Curtis AM, Boston RC, Panda S, Hogenesch JB, Fitzgerald GA, BMAL1 and CLOCK, two essential components of the circadian clock, are involved in glucose homeostasis. *PLoS Biol* 2 (2004) e377. [PubMed: 15523558]
- [96]. Weeke J, Gundersen HJ, Circadian and 30 minutes variations in serum TSH and thyroid hormones in normal subjects. *Acta Endocrinol (Copenh)* 89 (1978) 659–672 [PubMed: 716774]
- [97]. Allan JS, Czeisler CA, Persistence of the circadian thyrotropin rhythm under constant conditions and after light-induced shifts of circadian phase. *J Clin Endocrinol Metab* 79 (1994) 508–512. [PubMed: 8045970]
- [98]. Abe K, Kroning J, Greer MA, Critchlow V, Effects of destruction of the suprachiasmatic nuclei on the circadian rhythms in plasma corticosterone, body temperature, feeding and plasma thyrotropin. *Neuroendocrinology* 29 (1979) 119–131. [PubMed: 503281]
- [99]. Bertani S, Carboni L, Criado A, Michielin F, Mangiarini L, Vicentini E, Circadian profile of peripheral hormone levels in Sprague-Dawley rats and in common marmosets (*Callithrix jacchus*). *In Vivo* 24(2010)827–836. [PubMed: 21164040]

- [100]. Kalsbeek, Fliers E, Franke AN, Wortel J, Buijs RM, Functional connections between the suprachiasmatic nucleus and the thyroid gland as revealed by lesioning and viral tracing techniques in the rat. *Endocrinology* 141 (2000) 3832–3841. [PubMed: 11014240]
- [101]. Fahrenkrug J, Georg B, Hannibal J, Jorgensen HL, Hypophysectomy abolishes rhythms in rat thyroid hormones but not in the thyroid clock. *J Endocrinol* 233 (2017) 209–216. [PubMed: 28348112] * Reports a dissociation between the impact of pituitary removal on thyroid hormone rhythms versus rhythmic clock gene expression in the thyroid. This finding strongly suggests that clock gene rhythms in the thyroid are mediated via autonomic outflow controlled by the SCN.
- [102]. Peliciari-Garcia RA, Bargi-Souza P, Young ME, Nunes MT, Repercussions of hypo and hyperthyroidism on the heart circadian clock. *Chronobiol Int* 35 (2018) 147–159. [PubMed: 29111822]
- [103]. Maroni J, Capri KM, Cushman AV, Monteiro De Pina IK, Chasse MH, Seggio JA, Constant light alters serum hormone levels related to thyroid function in male CD-1 mice. *Chronobiol Int* (2018) 1–8.* A study showing that exposure to constant light disrupts thyroid hormone rhythms and results in dysregulated leptin rhythms.
- [104]. Philippe J, Dibner C, Thyroid circadian timing: roles in physiology and thyroid malignancies. *J Biol Rhythms* 30 (2015) 76–83. [PubMed: 25411240]
- [105]. Mannic T, Meyer P, Triponez F, Pusztaszeri M, Le Martelot G, Mariani O, Schmitter D, Sage D, Philippe J, Dibner C, Circadian clock characteristics are altered in human thyroid malignant nodules. *J Clin Endocrinol Metab* 98 (2013) 4446–4456. [PubMed: 23979949]

Highlights

- The circadian system coordinates physiology and behavior with time of day
- Hormones broadly affect central and peripheral physiology and behavior.
- The circadian system coordinates endocrine timing via direct communication to neuroendocrine cells, autonomic outflow to endocrine glands and through and rhythmic behavior.
- Disruptions to endocrine timing have marked, negative impact on normal physiology and behavior and are associated with a variety of disease states.




Figure 1.

The SCN influences endocrine timing via projections to neuroendocrine cells in the brain, autonomic outflow through initial projections to the PVN, and rhythmic behavioral output. Autonomic outflow from the SCN to the gonads, while likely, has not been specifically examined. Peripheral clocks are found in the gonads, adrenals, and thyroid, keeping their own circadian time coordinated by the SCN. Peripheral clock functioning likely serves to further control hormonal output from these glands. Finally, hormones from these glands feed back to the hypothalamus and pituitary to further regulate their own production and secretion.