

# UCLA

## UCLA Previously Published Works

### Title

Association of Long-term Change and Variability in Glycemia With Risk of Incident Heart Failure Among Patients With Type 2 Diabetes: A Secondary Analysis of the ACCORD Trial

### Permalink

<https://escholarship.org/uc/item/7zp1715b>

### Journal

Diabetes Care, 43(8)

### ISSN

1066-9442

### Authors

Segar, Matthew W  
Patel, Kershaw V  
Vaduganathan, Muthiah  
[et al.](#)

### Publication Date

2020-08-01

### DOI

10.2337/dc19-2541

Peer reviewed



# Association of Long-term Change and Variability in Glycemia With Risk of Incident Heart Failure Among Patients With Type 2 Diabetes: A Secondary Analysis of the ACCORD Trial

*Diabetes Care* 2020;43:1920–1928 | <https://doi.org/10.2337/dc19-2541>

Matthew W. Segar,<sup>1</sup> Kershaw V. Patel,<sup>1</sup>  
Muthiah Vaduganathan,<sup>2</sup>  
Melissa C. Caughey,<sup>3</sup> Javed Butler,<sup>4</sup>  
Gregg C. Fonarow,<sup>5</sup> Justin L. Grodin,<sup>1</sup>  
Darren K. McGuire,<sup>1</sup> and Ambarish Pandey<sup>1</sup>

## OBJECTIVE

To evaluate the associations between long-term change and variability in glycemia with risk of heart failure (HF) among patients with type 2 diabetes mellitus (T2DM).

## RESEARCH DESIGN AND METHODS

Among participants with T2DM enrolled in the Action to Control Cardiovascular Risk in Diabetes (ACCORD) trial, variability in HbA<sub>1c</sub> was assessed from stabilization of HbA<sub>1c</sub> following enrollment (8 months) to 3 years of follow-up as follows: average successive variability (ASV) (average absolute difference between successive values), coefficient of variation (SD/mean), and SD. Participants with HF at baseline or within 3 years of enrollment were excluded. Adjusted Cox models were used to evaluate the association of percent change (from baseline to 3 years of follow-up) and variability in HbA<sub>1c</sub> over the first 3 years of enrollment and subsequent risk of HF.

## RESULTS

The study included 8,576 patients. Over a median follow-up of 6.4 years from the end of variability measurements at year 3, 388 patients had an incident HF hospitalization. Substantial changes in HbA<sub>1c</sub> were significantly associated with higher risk of HF (hazard ratio [HR] for  $\geq 10\%$  decrease 1.32 [95% CI 1.08–1.75] and for  $\geq 10\%$  increase 1.55 [1.19–2.04]; reference  $< 10\%$  change in HbA<sub>1c</sub>). Greater long-term variability in HbA<sub>1c</sub> was significantly associated with higher risk of HF (HR per 1 SD of ASV 1.34 [95% CI 1.17–1.54]) independent of baseline risk factors and interval changes in cardiometabolic parameters. Consistent patterns of association were observed with use of alternative measures of glycemic variability.

## CONCLUSIONS

Substantial long-term changes and variability in HbA<sub>1c</sub> were independently associated with risk of HF among patients with T2DM.

Heart failure (HF) is common and associated with high morbidity and mortality among adults with type 2 diabetes mellitus (T2DM) (1–3). Risk factors for HF among patients with T2DM include high blood pressure (BP), coronary artery disease, abnormal kidney function, obesity, and low cardiorespiratory fitness levels (4,5). Moreover,

<sup>1</sup>Division of Cardiology, Department of Internal Medicine, University of Texas Southwestern Medical Center, Dallas, TX

<sup>2</sup>Brigham and Women's Hospital Heart and Vascular Center, Department of Medicine, Harvard Medical School, Boston, MA

<sup>3</sup>Joint Department of Biomedical Engineering, University of North Carolina and North Carolina State University, Chapel Hill, NC

<sup>4</sup>Department of Medicine, University of Mississippi Medical Center, Jackson, MS

<sup>5</sup>Ahmanson-UCLA Cardiomyopathy Center, Division of Cardiology, Ronald Reagan UCLA Medical Center, Los Angeles, CA

Corresponding author: Ambarish Pandey, [ambarish.pandey@utsouthwestern.edu](mailto:ambarish.pandey@utsouthwestern.edu)

Received 18 December 2019 and accepted 4 May 2020

This article contains supplementary material online at <https://doi.org/10.2337/figshare.12273155>.

M.W.S. and K.V.P. made equal contributions as co-first authors.

© 2020 by the American Diabetes Association. Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered. More information is available at <https://www.diabetesjournals.org/content/license>.

epidemiologic studies have also demonstrated that chronic hyperglycemia in T2DM is associated with higher risk of HF (6–8). However, large-scale randomized controlled trials of intensive glycemetic control among patients with T2DM did not lower the risk of HF (9–11). The null effect of strict glucose control on cardiovascular outcomes has been attributed in part to hypoglycemia, though such a causal link has not been proven (12,13). A single measurement of glycemia may not reflect chronic glucose exposure and its attendant cardiovascular risk. Fluctuation in glycemia has been proposed as an alternative prognostic marker in T2DM. Long-term glycemetic variability is a measure of changes in glucose over time and is associated with adverse cardiovascular events, renal disease, and mortality in patients with T2DM (14,15). Whether long-term change and variability in glycemia are each associated with risk of HF among patients with T2DM is not well established.

The Action to Control Cardiovascular Risk in Diabetes (ACCORD) trial was a multicenter clinical trial that examined the effects of intensive glycemetic control on cardiovascular outcomes among adults with T2DM (9). In ACCORD, glycated hemoglobin (HbA<sub>1c</sub>) and fasting plasma glucose (FPG) were measured at regular intervals and HF hospitalization events were adjudicated as a secondary outcome (9,16). The purpose of the current study was to examine the associations of long-term changes and variability in glycemetic markers, specifically HbA<sub>1c</sub> and FPG, with risk of HF among participants of ACCORD. We hypothesized that substantial changes in HbA<sub>1c</sub> as well as greater long-term glycemetic variability in HbA<sub>1c</sub> on follow-up would be associated with higher risk of HF.

## RESEARCH DESIGN AND METHODS

### Study Population

The current study was performed as a post hoc analysis of ACCORD and the subsequent ACCORD Follow-on (ACCORDION) study. The study design and results have previously been reported and are described in Supplementary Material (9,17,18). The present analysis included participants from the intensive and standard glycemetic control arms of the trial with no history of HF who had at least three repeated measurements of glycemia (HbA<sub>1c</sub> or FPG) after stabilization of the values (8-month visit) following

randomization up to 3 years from enrollment (Supplementary Fig. 1). Patients with an incident HF event within 3 years from enrollment were excluded from the present analysis.

### Measures of Long-term Glycemetic Variability

Long-term glycemetic variability was measured as the variability in HbA<sub>1c</sub> or FPG between visits for each participant and was calculated using repeated measures of HbA<sub>1c</sub> or FPG between 8 months and 3 years of follow-up. The variability assessment was landmarked at 8 months considering the study intervention directly influenced fluctuation in glycemetic markers in the first few months following enrollment (Supplementary Fig. 2). Core laboratory measures of HbA<sub>1c</sub> and FPG were used for calculation of glycemetic variability. The following three established measures of glycemetic variability were used: average successive variability (ASV), SD, and coefficient of variation (CV) (15,19). The ASV was prespecified as the primary exposure variable of interest and was defined as the average absolute difference between successive values. The CV was calculated as the SD divided by the mean value.

### Clinical Covariates

Upon enrollment, patients underwent questionnaires, physical examination, and laboratory measures using a standardized protocol as previously described and detailed in Supplementary Material (9,17). HbA<sub>1c</sub> levels were measured at a core laboratory every 4 months. FPG levels were measured every 4 months for the first 2 years and subsequently every 12 months. Participants randomized to both trial arms achieved lower glucose levels soon after randomization with stabilization of glycemetic levels at 8 months (Supplementary Fig. 2). In the current study, percent change ( $\Delta$ ) in HbA<sub>1c</sub> ( $\Delta$ HbA<sub>1c</sub>) and other cardiometabolic parameters was calculated with use of data collected from the baseline visit until up to 3 years of follow-up.

### Incident HF, Noncardiovascular Death, and Hypoglycemic Events

The primary outcome of interest for the present analysis was incident HF defined as the first hospitalization event for HF or death due to HF. In ACCORD, an independent committee adjudicated HF events

as a prespecified secondary outcome (9,16). Hospitalization for HF was based on documented clinical and radiological evidence. Death due to HF was based on clinical, radiological, or postmortem evidence of HF with absence of an acute ischemic event according to clinical or postmortem evidence. The secondary outcome of interest was defined as an acute ischemic heart disease event, which was a prespecified secondary outcome in ACCORD defined as cardiovascular death, nonfatal myocardial infarction (MI), or unstable angina. Time to event was calculated as the number of months from year 3 of follow-up to occurrence of the first event. Long-term follow-up was recorded in all patients using data from ACCORD and ACCORDION (18). For analyses of all clinical outcomes, patients were censored at the time of their last follow-up.

Death events were adjudicated by a central committee, which was masked to treatment allocation and used predefined criteria. Hypoglycemic events were self-reported by the participant and defined as any episode requiring medical attention in which there was either a documented capillary glucose level <50 mg/dL (<2.8 mmol/L) or prompt recovery following administration of oral carbohydrate, intravenous glucose, or glucagon (20).

### Statistical Analysis

Patients were categorized into long-term HbA<sub>1c</sub> variability groups according to quintiles of ASV. Normality in distribution of patient characteristic variables was assessed by calculating the skewness and plotting a histogram for the vector of values with an overlying normal curve with the same mean and SD. Each continuous variable was confirmed as being normally distributed (skewness statistic between  $-2$  and  $2$ ), and these variables were reported as mean (SD). Categorical variables were reported as percentages across the study groups. Differences between groups were tested using a  $\chi^2$  test for categorical variables and one-way ANOVA for continuous variables. Baseline characteristics were compared across categories of baseline HbA<sub>1c</sub> and  $\Delta$ HbA<sub>1c</sub>. Unadjusted rates of incident HF events were also compared across ASV categories in the overall cohort and across subgroups stratified by categories of baseline HbA<sub>1c</sub> (<7.0% [ $<53$  mmol/mol],

7.0%–8.5% [53–69 mmol/mol], and >8.5% [ $>69$  mmol/mol]) and  $\Delta\text{HbA}_{1c}$  ( $\geq 10\%$  decrease,  $<10\%$  decrease to  $<10\%$  increase, and  $\geq 10\%$  increase). Separate multivariable Cox proportional hazards models were constructed to evaluate the association of categorical and continuous measures of baseline  $\text{HbA}_{1c}$  and  $\Delta\text{HbA}_{1c}$  with risk of HF. Validity of the proportionality assumption was verified using scaled Schoenfeld residuals and by visually examining “log-log” plots. The models were sequentially adjusted with inclusion of the following covariates selected a priori based on biologic plausibility and prior published literature (21): model 1, demographic characteristics (age, sex, race, and level of education), intensive glycemic control treatment group, history of cardiovascular disease (CVD) (MI, stroke, or coronary revascularization), traditional cardiovascular risk factors (systolic BP, BMI, cigarette use, alcohol use, total cholesterol, serum creatinine, LDL cholesterol [LDL-c], and HDL cholesterol [HDL-c]), medication use (angiotensin receptor blocker, ACE inhibitor,  $\beta$ -blocker, loop diuretic, thiazide diuretic, calcium channel blocker, insulin, sulfonylurea, biguanide, meglitinide, and  $\alpha$ -glucosidase inhibitor), and baseline  $\text{HbA}_{1c}$ , and model 2, adjustment for covariates in model 1 +  $\Delta\text{HbA}_{1c}$ ,  $\Delta$ systolic BP,  $\Delta$ BMI, and  $\Delta$ creatinine from enrollment baseline to year 3 + interval MI on follow-up as a time-dependent variable. The associations between measures of variability in  $\text{HbA}_{1c}$  and risk of HF were also assessed using multivariable Cox models that included the covariates described above in models 1 and 2 along with measures of  $\text{HbA}_{1c}$  variability (ASV, CV, and SD in separate models). Multiplicative interaction terms were included in the adjusted Cox models to evaluate whether the association between glycemic variability and risk of HF was modified by baseline  $\text{HbA}_{1c}$  and  $\Delta\text{HbA}_{1c}$ . FPG, an alternative measure of glycemia, was also used to assess the independent associations of glycemic variability and risk of adverse events.

In the subset of the study population free of atherosclerotic CVD at baseline ( $N = 5,822$ ), the association between measures of glycemic variability with the risk of acute ischemic heart disease events was assessed using adjusted Cox models that included covariates described above in model 1 (except

history of CVD) and model 2 (except interval MI).

Additional analyses were performed to evaluate the robustness of the association of  $\Delta\text{HbA}_{1c}$  and variability in  $\text{HbA}_{1c}$  with risk of HF. First, to account for the potential contribution of hypoglycemic events toward the observed association of  $\Delta\text{HbA}_{1c}$  and variability in  $\text{HbA}_{1c}$  with risk of HF, we performed sensitivity analysis excluding individuals with a hypoglycemic event on follow-up ( $N = 1,159$  with hypoglycemic events). Second, to account for seasonal changes in  $\text{HbA}_{1c}$  and the effect of number of  $\text{HbA}_{1c}$  measures used to estimate variability, we performed sensitivity analysis assessing  $\text{HbA}_{1c}$  variability among individuals with higher minimum number of repeated  $\text{HbA}_{1c}$  measures within the first 3 years of follow-up. Third, to account for the prognostic importance of variability in BP (22), BMI, and LDL-c, we additionally adjusted Cox models for variability in these parameters over the first 3 years (from 8 months to 3 years of follow-up). Finally, we performed stratified analyses examining the association of long-term variability in  $\text{HbA}_{1c}$  with risk of HF across intensive glucose control versus standard control groups, across intensive BP control versus standard BP control groups, and among subgroups of participants with versus without history of atherosclerotic CVD. Analyses were performed using R 3.5.1 (R Foundation for Statistical Computing, Vienna, Austria).

## RESULTS

Of the 10,251 patients enrolled in ACCORD, participants were excluded from the present analyses for the following reasons: 492 had a history of HF, 870 were missing at least three  $\text{HbA}_{1c}$  measurements between month 8 and 3 years, and 223 developed HF and an additional 90 died within the first 3 years of enrollment (Supplementary Fig. 1). The current study included 8,576 patients (38.4% of whom were women and 18.3% of whom were black) with a mean  $\pm$  SD baseline  $\text{HbA}_{1c}$  of  $8.3 \pm 1.0\%$  (mean 67 mmol/mol). The median number of  $\text{HbA}_{1c}$  measurements between 8 months and 3 years of follow-up was eight (interquartile range 7–8). The mean  $\text{HbA}_{1c}$  variability, as measured by ASV, was  $0.6 \pm 0.3\%$ . Over a median follow-up of 6.4 (interquartile range 4.0–7.6) years from the end of variability measurements at

year 3, there were 388 (cumulative rate: 4.5%) incident HF events.

The baseline characteristics of participants stratified by quintiles of long-term variability in  $\text{HbA}_{1c}$  are shown in Table 1. The patients with greater long-term glycemic variability more commonly were black, younger, and randomized to standard therapy; used insulin; had a history of CVD; and had higher BMI,  $\text{HbA}_{1c}$ , and total cholesterol at baseline. These patients had more change in BMI and serum creatinine from baseline to 3 years of follow-up. Baseline characteristics of the patients stratified by baseline  $\text{HbA}_{1c}$  and  $\Delta\text{HbA}_{1c}$  from baseline to 3 years of follow-up are shown in Supplementary Tables 1 and 2.

## Baseline $\text{HbA}_{1c}$ , $\Delta\text{HbA}_{1c}$ , and Risk of HF

Baseline  $\text{HbA}_{1c}$  was significantly associated with risk of HF in multivariable-adjusted analysis such that 1 unit higher  $\text{HbA}_{1c}$  was associated with 20% higher risk of incident HF after other baseline confounders were accounted for (hazard ratio [HR] 1.20 [95% CI 1.08–1.33]) (Supplementary Table 3). The association between  $\Delta\text{HbA}_{1c}$  and risk of incident HF was nonlinear with a significantly higher risk noted at either extreme (Fig. 1). Individuals with  $\geq 10\%$  increase or  $\geq 10\%$  decrease in  $\text{HbA}_{1c}$  during the first 3 years of follow-up had higher risk of HF compared with individuals with a more stable  $\text{HbA}_{1c}$  ( $<10\%$  decrease to  $<10\%$  increase) (HR 1.55 [95% CI 1.19–2.04] and 1.32 [1.08–1.75], respectively) (Supplementary Table 3) after adjustment for baseline and longitudinal changes in risk factors on follow-up. In sensitivity analysis excluding individuals with a hypoglycemic event on follow-up, the risk of incident HF was significant among those with a  $>10\%$  increase in  $\text{HbA}_{1c}$  but not among those with  $>10\%$  decrease in  $\text{HbA}_{1c}$  on follow-up (Supplementary Table 4).

## Long-term Variability in $\text{HbA}_{1c}$ and Risk of HF

In unadjusted comparison, the risk of incident HF increased in a graded fashion across increasing quintiles of ASV (quintile 1 [Q1] 3.3% and Q5 6.2%) (Fig. 2). In adjusted analyses, higher measures of long-term variability in  $\text{HbA}_{1c}$ , as measured by ASV, were associated with significantly higher risk of HF independent of baseline risk factors, medication

**Table 1—Baseline characteristics of participants stratified by quintiles of variability in HbA<sub>1c</sub>**

	Quintile 1 (n = 1,748)	Quintile 2 (n = 1,700)	Quintile 3 (n = 1,716)	Quintile 4 (n = 1,702)	Quintile 5 (n = 1,710)	P
HbA <sub>1c</sub> ASV, %	0.2 (0.05)	0.3 (0.03)	0.4 (0.04)	0.6 (0.06)	1.1 (0.44)	<0.001
Age, years	63.4 (6.4)	63.1 (6.6)	62.7 (6.5)	62.5 (6.4)	61.3 (6.4)	<0.001
Female	641 (36.7)	659 (38.8)	645 (37.6)	661 (38.8)	683 (39.9)	0.33
BMI, kg/m <sup>2</sup>	31.9 (5.2)	31.7 (5.3)	32.0 (5.4)	32.3 (5.4)	32.8 (5.5)	<0.001
Intensive glycemic control	1,369 (78.3)	1,077 (63.4)	852 (49.7)	553 (32.5)	415 (24.3)	<0.001
Race						<0.001
Black	222 (12.7)	272 (16.0)	294 (17.1)	350 (20.6)	430 (25.1)	
Hispanic	79 (4.5)	115 (6.8)	97 (5.7)	117 (6.9)	181 (10.6)	
Other	223 (12.8)	232 (13.6)	197 (11.5)	203 (11.9)	169 (9.9)	
White	1,224 (70.0)	1,081 (63.6)	1,128 (65.7)	1,032 (60.6)	930 (54.4)	
Education						0.003
<High school	224 (12.8)	230 (13.5)	215 (12.5)	255 (15.0)	283 (16.6)	
High school (or GED)	465 (26.6)	404 (23.8)	483 (28.2)	454 (26.7)	429 (25.1)	
Some college	552 (31.6)	594 (35.0)	564 (32.9)	555 (32.6)	565 (33.1)	
College	506 (29.0)	471 (27.7)	453 (26.4)	438 (25.7)	431 (25.2)	
Established CVD	499 (28.5)	522 (30.7)	572 (33.3)	576 (33.8)	585 (34.2)	0.001
Alcoholic drinks/week	1.2 (3.0)	1.0 (2.8)	1.0 (2.7)	1.0 (2.8)	0.8 (2.4)	0.001
Current smoker	219 (12.5)	219 (12.9)	227 (13.2)	220 (12.9)	285 (16.7)	0.002
Systolic BP, mmHg	136 (16)	137 (17)	136 (16)	136 (17)	137 (17)	0.46
Serum creatinine, mg/dL	0.90 (0.2)	0.89 (0.2)	0.91 (0.2)	0.91 (0.2)	0.91 (0.2)	0.04
HbA <sub>1c</sub> , % [mmol/mol]	8.0 (0.9) [64]	8.1 (0.9) [65]	8.3 (1.0) [67]	8.4 (1.1) [68]	8.6 (1.2) [70]	<0.001
Total cholesterol, mg/dL	183 (40)	182 (41)	184 (43)	184 (43)	186 (43)	0.03
LDL-c, mg/dL	104 (33)	103 (33)	106 (34)	106 (34)	107 (35)	0.12
HDL-c, mg/dL	42 (11)	42 (12)	42 (12)	42 (12)	42 (11)	0.08
Angiotensin receptor blocker	281 (16.1)	288 (17.0)	320 (18.7)	296 (17.4)	250 (14.7)	0.03
ACE inhibitor	904 (51.8)	911 (53.7)	933 (54.4)	913 (53.8)	963 (56.5)	0.09
β-Blocker	450 (25.8)	464 (27.4)	479 (27.9)	498 (29.3)	507 (29.8)	0.07
Calcium channel blocker	184 (10.5)	206 (12.2)	173 (10.1)	209 (12.3)	211 (12.4)	0.09
Loop diuretic	94 (5.4)	101 (6.0)	121 (7.1)	114 (6.7)	140 (8.2)	0.01
Thiazide diuretic	490 (28.1)	470 (27.7)	502 (29.3)	479 (28.2)	491 (28.8)	0.86
Insulin use	245 (15.2)	368 (23.7)	432 (28.3)	448 (29.5)	448 (29.4)	<0.001
ΔHbA <sub>1c</sub> , %	−0.0 (6.6)	0.3 (9.5)	1.2 (11.5)	1.5 (14.0)	5.1 (22.1)	<0.01
ΔSystolic BP, %	−5.4 (13.0)	−4.9 (13.4)	−4.8 (14.1)	−4.9 (13.7)	−4.5 (14.2)	0.40
ΔSerum creatinine, %	10.4 (20.8)	13.6 (28.8)	13.4 (26.2)	13.9 (24.3)	16.2 (26.9)	<0.001
ΔBMI, %	2.0 (7.4)	2.2 (7.3)	2.3 (7.2)	1.7 (7.2)	2.8 (7.7)	<0.001

Data presented as mean (SD) for continuous variables or n (%) for categorical variables. For HbA<sub>1c</sub>, data in square brackets are means. Long-term variability in HbA<sub>1c</sub> was assessed by ASV from 8 months to 3 years of follow-up. Δ is the percent change in cardiometabolic parameters from enrollment baseline to 3 years of follow-up. GED, General Education Development.

use, and treatment arm (intensive vs. standard glucose control) (HR per 1-SD higher ASV 1.27 [95% CI 1.15–1.40]) (Fig. 3 [model 1]). This association was not attenuated and remained significant after further adjustment for ΔHbA<sub>1c</sub> and changes in other cardiometabolic factors as well as interval MI on follow-up (HR per 1 SD of ASV 1.34 [95% CI 1.17–1.54]) (Fig. 3 [model 2]). Further adjustment for variability in other cardiometabolic parameters (ASV for BP, LDL-c, and BMI) also did not attenuate the association between variability in HbA<sub>1c</sub> and risk of HF (HR per 1 SD higher ASV 1.24 [95% CI 1.12–1.37]) in the most adjusted model.


Similar patterns of association were also observed using other measures of long-term variability in HbA<sub>1c</sub> (CV, SD) (Fig. 3).

The higher risk of incident HF among individuals with greater variability in HbA<sub>1c</sub> (above vs. below median ASV) was consistent across all strata of different baseline HbA<sub>1c</sub> levels ( $P_{\text{interaction}}$  for baseline HbA<sub>1c</sub> \* ASV = 0.29) as well as ΔHbA<sub>1c</sub> categories ( $P_{\text{interaction}}$  for ΔHbA<sub>1c</sub> \* ASV = 0.38) (Supplementary Fig. 3A and B). The association between long-term variability in HbA<sub>1c</sub> and risk of HF was consistent across relevant subgroups including glucose treatment strategies (intensive vs. standard), BP treatment strategies (intensive

vs. standard), and prior history of ischemic event (yes vs. no) (Supplementary Tables 5–7). In sensitivity analysis using a higher minimum number of HbA<sub>1c</sub> measures to estimate variability, the association between long-term variability in HbA<sub>1c</sub> with risk of incident HF was consistent with that observed in the primary analysis (Supplementary Fig. 4).

#### Long-term Variability in FPG and Risk of HF

For further evaluation of the association between long-term glycemic variability and risk of incident HF, an alternative measure of glycemic variability based on



**Figure 1**—Cubic spline demonstrating multivariable-adjusted HRs (95% CI: red dotted line) for incident HF across  $\Delta$ HbA<sub>1c</sub> from baseline to 3 years of follow-up. HR refers to the association of  $\Delta$ HbA<sub>1c</sub> with risk of incident HF. Model included demographic characteristics (age, sex, race, and level of education), intensive glycemic control treatment group, history of CVD (MI, stroke, or coronary revascularization), risk factors (systolic BP, alcohol use, cigarette use, BMI, total cholesterol, serum creatinine, LDL-c, and HDL-c), medication use (angiotensin receptor blocker, ACE inhibitor,  $\beta$ -blocker, loop diuretic, thiazide diuretic, calcium channel blocker, insulin, sulfonylurea, biguanide, meglitinide, and  $\alpha$ -glucosidase inhibitor), baseline HbA<sub>1c</sub>,  $\Delta$ HbA<sub>1c</sub>,  $\Delta$ systolic BP,  $\Delta$ BMI,  $\Delta$ creatinine, and incident MI as a time-dependent variable.

long-term variability in FPG were significantly associated with higher risk of HF (HR per 1-SD higher ASV 1.21 [95% CI 1.07–1.36]) independent of other potential confounders including  $\Delta$ FPG and other cardiometabolic parameters as well as time-updated MI event on follow-up (Fig. 3). Consistent patterns of association were observed with alternative measures of long-term variability in FPG, including CV and SD (Fig. 3). There was no significant interaction between long-term variability in FPG (assessed by ASV) and either baseline FPG ( $P_{\text{interaction}} = 0.72$ ) or  $\Delta$ FPG ( $P_{\text{interaction}} = 0.11$ ) for the risk of HF (Supplementary Fig. 3C and D).

**Glycemic Variability, Hypoglycemic Events, and HF**

The risk of clinical hypoglycemic events was evaluated across categories of long-term glycemic variability. Overall, 13.5% of study participants reported clinically significant hypoglycemic events in the study cohort. The reported rates of hypoglycemic events increased significantly across increasing quintiles of long-term variability in HbA<sub>1c</sub> (assessed by ASV) (Q1 10.8% vs. Q5 15.2%) as well as long-term variability in FPG (Q1 10.4% vs. Q5 19.5%). In adjusted analysis, greater variability in HbA<sub>1c</sub> was significantly associated with greater likelihood of a hypoglycemic event on follow-up (odds ratio 1.23 [95% CI 1.01–1.49]). Furthermore, individuals who developed hypoglycemia on follow-up had higher incidence of HF versus those without reported hypoglycemic event (6.9% [80 of 1,159] vs. 4.2% [308 of 7,417],  $P$  value <0.001). In sensitivity analysis excluding individuals with hypoglycemic events on follow-up ( $N = 7,417$ ), the association between long-term variability in HbA<sub>1c</sub> (assessed by ASV) with higher

risk of HF was significant and consistent with that observed in the primary analysis (HR per 1 SD higher ASV 1.35 [95% CI 1.16–1.56]) in the most adjusted model. Consistent patterns of associations were also observed between long-term variability in FPG and risk of HF in sensitivity analysis excluding individuals with hypoglycemic events on follow-up (HR per 1-SD higher ASV 1.27 [95% CI 1.11–1.45]).


**Long-term Glycemic Variability and Risk of Acute Coronary Ischemic Events**

Among the subset of study participants free of atherosclerotic CVD at baseline, there were 664 (cumulative rate 11.4%) incident acute coronary ischemic events during the study period. In unadjusted comparison, the risk of acute coronary ischemic events increased in a graded fashion across increasing quintiles of HbA<sub>1c</sub> ASV (Q1 8.7% and Q5 15.5%) (Supplementary Fig. 5). In adjusted analyses, higher measures of long-term variability in HbA<sub>1c</sub>, as measured by ASV, were associated with significantly higher risk of acute coronary ischemic events independent of baseline risk factors,  $\Delta$ HbA<sub>1c</sub>, and other cardiometabolic parameters (HR per 1 SD higher ASV 1.25 [95% CI 1.13–1.39]) (Supplementary Fig. 6). Similar patterns of association were also observed using alternative parameters of HbA<sub>1c</sub> variability (CV, SD) as well as using FPG in measures of glycemic variability (Supplementary Fig. 6).

**CONCLUSIONS**


In this post hoc secondary analysis of data from ACCORD, several important findings were observed. First, there was a nonlinear relationship between changes in HbA<sub>1c</sub> during the trial and risk of HF. Individuals with a substantial decrease as well as increase in HbA<sub>1c</sub> had significantly higher risk of HF compared with those with stable HbA<sub>1c</sub> values, independent of other risk factors. Furthermore, the higher risk of HF observed among individuals with a substantial decrease in HbA<sub>1c</sub> on follow-up was largely related to downstream hypoglycemic events. Second, greater long-term variability in HbA<sub>1c</sub> was also significantly associated with higher risk of HF, independent of other risk factors, changes in HbA<sub>1c</sub> and other cardiometabolic parameters, and interval occurrence of MI event. Third, greater long-term variability in FPG,

FPG measurements was examined. The median number of FPG measurements from month 8 to year 3 of follow-up was four. The mean  $\pm$  SD FPG at month 8 (study baseline) was 136  $\pm$  32 mg/dL, and variability, as measured by ASV, was 36  $\pm$  19 mg/dL. A significant graded association was observed between greater long-term variability in FPG and risk of HF such that participants in the highest quintile of ASV for FPG had approximately a twofold higher risk of HF compared with those in the lowest quintile (event rate 6.0% vs. 3.1%, respectively) (Fig. 2B). In adjusted analysis, higher measures of



**Figure 2**—Proportion of participants with incident HF across quintiles of long-term variability in HbA<sub>1c</sub> (A) and FPG (B).





**Figure 3**—Multivariable-adjusted association of long-term variability in HbA<sub>1c</sub> (top) and FPG (bottom) with risk of incident HF. HR data refer to the association of 1 SD higher measure of long-term variability in HbA<sub>1c</sub>/FPG with risk of incident HF. Model 1 included demographic characteristics (age, sex, race, and level of education), intensive glycemic control treatment group, history of CVD (MI, stroke, or coronary revascularization), risk factors (systolic BP, alcohol use, cigarette use, BMI, total cholesterol, serum creatinine, LDL-c, and HDL-c), medication use (angiotensin receptor blocker, ACE inhibitor, β-blocker, loop diuretic, thiazide diuretic, calcium channel blocker, insulin, sulfonylurea, biguanide, meglitinide, and α-glucosidase inhibitor) and baseline HbA<sub>1c</sub>/FPG level. Model 2 included covariates from model 1 plus ΔHbA<sub>1c</sub>/ΔFPG, Δsystolic BP, ΔBMI, Δcreatinine, and incident MI as a time-dependent variable.

another measure of glycemic status, was also significantly associated with higher risk of HF. Finally, greater glycemic variability was also significantly associated with higher risk of acute ischemic cardiac events. Taken together, these findings highlight the prognostic significance of substantial fluctuations in glycemia among patients with T2DM in forecasting future risk of adverse cardiovascular events.

Consistent with our observations, prior studies have demonstrated a significant association between hyperglycemia and risk of HF (6–8). However, the relationship between long-term changes in HbA<sub>1c</sub> and risk of HF is less well established. In a large cohort study from the Kaiser Permanente Health System, Nichols et al. (23) observed that a decline in HbA<sub>1c</sub> in patients with T2DM was associated with lower likelihood of HF on follow-up. In a time-updated analysis from a cohort of patients with T2DM from the U.K., both higher (>10% [ $>86$  mmol/mol]) and lower (<6% [ $<42$  mmol/mol]) HbA<sub>1c</sub> levels on longitudinal follow-up were

associated with higher risk of HF compared with risk in individuals with HbA<sub>1c</sub> in the range of 6–7% (42–53 mmol/mol) (24). However, these studies were limited by evaluation of a referral population, use of administrative data codes to identify HF events, and confounding by indication for HbA<sub>1c</sub> testing. In the present analysis, using data from a large, multicenter, randomized controlled trial that included intensive and standard glycemic control strategies, protocolized and prespecified serial HbA<sub>1c</sub> testing, and clinically adjudicated incident HF outcomes, there was a nonlinear relationship between long-term changes in HbA<sub>1c</sub> and risk of HF, with a higher risk associated with substantial changes in HbA<sub>1c</sub> levels at either extreme.

Several factors may underlie the observed higher risk of HF among individuals with substantial increase or decrease in HbA<sub>1c</sub>. First, we observed that the higher risk of HF among individuals with substantial decrease in HbA<sub>1c</sub> was largely driven by hypoglycemic events and not

observed in individuals without downstream hypoglycemic events. Second, while there was no statistically significant interaction between changes in HbA<sub>1c</sub> levels and HbA<sub>1c</sub> variability for the risk of HF, we observed greater incidence of HF among individuals with substantial increase or decrease in HbA<sub>1c</sub> (vs. those without significant change in HbA<sub>1c</sub>) in the high variability strata. In contrast, among individuals with lesser HbA<sub>1c</sub> variability, the difference in incidence of HF across HbA<sub>1c</sub> change categories was not as contrasting. This suggests that greater variability in HbA<sub>1c</sub> at the extremes of HbA<sub>1c</sub> change distribution may also have contributed to higher risk of HF. Finally, substantial changes in HbA<sub>1c</sub> may also be associated with increased oxidative stress, endothelial dysfunction, and proinflammatory state, which may contribute to the observed higher risk of HF (25,26).

Besides substantial changes in HbA<sub>1c</sub>, greater long-term variabilities in measures of HbA<sub>1c</sub> and FPG on follow-up were each significantly associated with risk of HF. Prior studies have demonstrated associations between long-term variability in glycemic control and risk of adverse clinical outcomes including all-cause mortality and risk of microvascular complications such as nephropathy and retinopathy (15,27). However, the data on associations between long-term variability in glycemic control and risk of adverse cardiovascular outcomes are mixed. Some studies have demonstrated that greater long-term variability in HbA<sub>1c</sub> or FPG is associated with higher risk of adverse cardiovascular outcomes including CV death and nonfatal atherosclerotic events (15,27,28). Furthermore, two studies using large administrative databases in Europe and Republic of Korea have demonstrated a significant association of variability in HbA<sub>1c</sub> or FPG with risk of incident HF (24,29). However, these studies were limited due to nonprotocolized measurement of glycemic levels and use of administrative codes for HF outcome ascertainment. In the study by Kwon et al. (29), patients with a history of diabetes were excluded from the analysis. Other studies have failed to observe a consistent, independent association between glycemic variability and risk of adverse cardiovascular outcomes (19). Secondary analysis of trials comparing usual versus intensive glycemic control in patients with T2DM have demonstrated

a significant association between glycemic variability and risk of adverse CV events only in the intensive control arm—not in the usual care arm (30,31). The current study findings add to the existing literature by demonstrating a robust, significant association between multiple measures of long-term glycemic variability and risk of HF that was consistent across both standard and intensive glycemic control treatment groups. Two separate markers of glycemia were measured according to a standardized procedure. HF events were adjudicated by a committee as part of a prespecified secondary outcome.

The mechanisms underlying the observed associations between long-term variability in HbA<sub>1c</sub> and risk of HF are not well established. One potential mechanism could be the increased incidence of hypoglycemic events among individuals with higher variability in HbA<sub>1c</sub>. Prior studies have demonstrated a strong association between hypoglycemic events and risk of adverse cardiovascular outcomes (13,32). We observed greater likelihood of hypoglycemic events among individuals with greater long-term variability in HbA<sub>1c</sub>. Furthermore, the rates of HF were also higher in individuals with downstream hypoglycemic events. However, the risk of HF associated with greater variability in HbA<sub>1c</sub> was consistent among individuals without a hypoglycemic event on follow-up and across both intensive and standard glucose control arms. Taken together, these observations suggest that the risk of HF associated with higher glycemic variability may be independent of the intensity of glycemic control and associated hypoglycemic events. Among other potential mechanisms, glycemic variability is associated with higher risk of atherosclerotic CV disease outcomes, adverse cardiac remodeling patterns, and systolic dysfunction, all of which are key factors in development of HF (30,33,34). Furthermore, greater glycemic variability is associated with upregulation of stress hormones, activation of inflammation cascade, and downstream oxidative stress and endothelial dysfunction, which may lead to the observed higher risk of HF (25,26). Finally, it is possible that higher glycemic variability may reflect inconsistent compliance with medications, use of steroids, underlying infections, drastic weight changes, or other unmeasured

confounders. Fluctuations in cardiometabolic parameters, including glucose, BMI, and lipid levels, are each associated with adverse CV events and may suggest an underlying systemic process (35–37).

Our study findings have important implications for prevention of HF among patients with T2DM. While epidemiologic studies suggest that hyperglycemia is associated with higher risk of HF, clinical trials examining intensive glycemic control targeting lower HbA<sub>1c</sub> goals among patients with T2DM did not reduce the risk of HF (6–11). It is plausible that the lack of therapeutic benefit of intensive glycemic control on cardiovascular outcomes may be related to greater fluctuations in glycemic control. This is supported by observations from the current study demonstrating increased risk of HF among individuals with substantial changes in HbA<sub>1c</sub> at either extreme as well as greater long-term glycemic variability. The study findings highlight the importance of long-term variability in HbA<sub>1c</sub> to identify individuals with T2DM and adequate glycemic control who are at high risk for developing HF. Future studies are needed to determine whether glycemic variability may meaningfully inform glucose control strategies in patients with T2DM to prevent HF. These findings are particularly relevant since the newer antihyperglycemic therapies such as sodium–glucose cotransporter 2 (SGLT2) inhibitors, which are associated with reduction in the risk of incident HF, have been shown to lower glycemic variability (38). However, the extent to which changes in glycemic variability may mediate the favorable effects of these therapies remains unknown. As many patients with high degree of glycemic variability are treated with insulin, these data suggest that selection of insulins with less day-to-day glycemic variability may lessen future HF risk; however, this hypothesis requires prospective testing (39).

There are several strengths to our analysis that add to the robustness of our study findings. These include the large size of the study cohort with 8,576 participants, protocolized assessments of HbA<sub>1c</sub> at regular intervals, large number of clinically adjudicated cardiovascular events on follow-up, and consistency of the observed associations between greater variability in glycemia and risk of HF across different measures of variability

as well as glycemic control (HbA<sub>1c</sub> and FPG). The current study also has some noteworthy limitations. First, owing to the observational nature of the study, we cannot establish a causal association between glycemic variability and risk of HF and our findings need to be confirmed in future studies. Furthermore, we cannot exclude the possibility of residual or unmeasured confounding given the observational study design of the present analysis. Second, the study population was derived from ACCORD, which included patients with T2DM who had prevalent cardiovascular disease or risk factors. Thus, the findings may not be generalizable to patients with T2DM who are at lower risk for cardiovascular disease outside a clinical trial setting. Third, ACCORD recruited participants until 2005 and there was limited use of contemporary T2DM therapies such as SGLT2 inhibitors and glucagon-like peptide 1 receptor agonists. However, this may have avoided the confounding effects of drug therapy on HF risk, given the reductions in risk of incident HF observed with SGLT2 inhibitors (and, more modestly, with glucagon-like peptide 1 receptor agonists) in T2DM (40,41). Finally, left ventricular ejection fraction and HF subtype data were not available for the present analysis so we cannot evaluate the associations between baseline as well as long-term change and variability in glycemia with risk of incident HF with reduced ejection fraction and HF with preserved ejection fraction.

In conclusion, substantial changes and long-term variability in HbA<sub>1c</sub> were each independently associated with risk of incident HF among patients with T2DM. Future studies are needed to determine whether long-term glycemic variability may be used to guide glycemic control strategies with use of therapies that have less glucose fluctuations to modify the risk of HF among high-risk patients with T2DM.

---

**Funding.** K.V.P. is supported by a National Heart, Lung, and Blood Institute T32 postdoctoral training grant (5T32HL125247-03). M.V. is supported by the KL2/Catalyst Medical Research Investigator Training Award from Harvard Catalyst (National Institutes of Health [NIH]/National Center for Advancing Translational Sciences, UL 1TR002541). M.V. participates on clinical end point committees for studies sponsored by the NIH. J.B. has received research support from the NIH, Patient-Centered Outcomes Research Institute, and the European Union. J.L.G. is supported



by the Texas Health Resources Clinical Scholars Program. A.P. is supported by the Texas Health Resources Clinical Scholars Program.

**Duality of Interest.** M.V. serves on advisory boards for Amgen, AstraZeneca, Baxter Healthcare, Bayer AG, and Boehringer Ingelheim and participates on clinical end point committees for studies sponsored by Novartis. J.B. serves as a consultant for Amgen, Array, AstraZeneca, Bayer AG, Boehringer Ingelheim, Bristol-Myers Squibb, CVRx, G3 Pharmaceuticals, INNOLIFE, Janssen, Luitpold Pharmaceuticals, Medtronic, Merck, Novartis, Relypsa, Stealth Peptides, SC Pharma, Vifor Pharma, and ZS Pharma. G.C.F. reports consulting for Abbott, Amgen, Bayer AG, CHF Solutions, Janssen, Medtronic, and Novartis. J.L.G. reports honoraria for consulting for Pfizer. D.K.M. reports honoraria for trial leadership from AstraZeneca, Sanofi, Janssen, Boehringer Ingelheim, Merck & Co, Pfizer, Novo Nordisk, Lexicon, Eisai, Glaxo-SmithKline, and ESPERION Therapeutics and honoraria for consulting for AstraZeneca, Sanofi, Lilly USA, Boehringer Ingelheim, Merck & Co, Pfizer, Novo Nordisk, Applied Therapeutics, Afimmune, and Metaviant Sciences. A.P. has served on the advisory board of Roche Diagnostics. No other potential conflicts of interest relevant to this article were reported.

**Author Contributions.** M.W.S., K.V.P., and A.P. developed the study concept and design, interpreted the data, and critically revised and drafted the manuscript. M.V., M.C.C., J.B., G.C.F., J.L.G., and D.K.M. contributed to the discussion and critically revised and reviewed the manuscript. M.W.S. and A.P. are the guarantors of this work and, as such, had full access to all the data in the study and take responsibility for the integrity of the data and the accuracy of the data analysis.

**Prior Presentation.** Parts of this study were presented in abstract form at the American Heart Association Scientific Sessions 2019, Philadelphia, PA, 16–18 November 2019.

## References

- Bertoni AG, Hundley WG, Massing MW, Bonds DE, Burke GL, Goff DC Jr. Heart failure prevalence, incidence, and mortality in the elderly with diabetes. *Diabetes Care* 2004;27:699–703
- Nichols GA, Gullion CM, Koro CE, Ephross SA, Brown JB. The incidence of congestive heart failure in type 2 diabetes: an update. *Diabetes Care* 2004;27:1879–1884
- Shah AD, Langenberg C, Rapsomaniki E, et al. Type 2 diabetes and incidence of cardiovascular diseases: a cohort study in 1.9 million people. *Lancet Diabetes Endocrinol* 2015;3:105–113
- Pandey A, Patel KV, Bahnson JL, et al.; Look AHEAD Research Group. Association of intensive lifestyle intervention, fitness, and body mass index with risk of heart failure in overweight or obese adults with type 2 diabetes mellitus: an analysis from the Look AHEAD trial. *Circulation* 2020;141:1295–1306
- Segar MW, Vaduganathan M, Patel KV, et al. Machine learning to predict the risk of incident heart failure hospitalization among patients with diabetes: the WATCH-DM Risk Score. *Diabetes Care* 2019;42:2298–2306
- Iribarren C, Karter AJ, Go AS, et al. Glycemic control and heart failure among adult patients with diabetes. *Circulation* 2001;103:2668–2673
- Thrainsdottir IS, Aspelund T, Thorgeirsson G, et al. The association between glucose abnormalities and heart failure in the population-based Reykjavik study. *Diabetes Care* 2005;28:612–616
- Pazin-Filho A, Kottgen A, Bertoni AG, et al. HbA1c as a risk factor for heart failure in persons with diabetes: the Atherosclerosis Risk in Communities (ARIC) study. *Diabetologia* 2008;51:2197–2204
- Gerstein HC, Miller ME, Byington RP, et al.; Action to Control Cardiovascular Risk in Diabetes Study Group. Effects of intensive glucose lowering in type 2 diabetes. *N Engl J Med* 2008;358:2545–2559
- Patel A, MacMahon S, Chalmers J, et al.; ADVANCE Collaborative Group. Intensive blood glucose control and vascular outcomes in patients with type 2 diabetes. *N Engl J Med* 2008;358:2560–2572
- Duckworth W, Abraira C, Moritz T, et al.; VADT Investigators. Glucose control and vascular complications in veterans with type 2 diabetes. *N Engl J Med* 2009;360:129–139
- Desouza CV, Bolli GB, Fonseca V. Hypoglycemia, diabetes, and cardiovascular events. *Diabetes Care* 2010;33:1389–1394
- Zoungas S, Patel A, Chalmers J, et al.; ADVANCE Collaborative Group. Severe hypoglycemia and risks of vascular events and death. *N Engl J Med* 2010;363:1410–1418
- Ceriello A, Monnier L, Owens D. Glycaemic variability in diabetes: clinical and therapeutic implications. *Lancet Diabetes Endocrinol* 2019;7:221–230
- Gorst C, Kwok CS, Aslam S, et al. Long-term glycemic variability and risk of adverse outcomes: a systematic review and meta-analysis. *Diabetes Care* 2015;38:2354–2369
- Cushman WC, Evans GW, Byington RP, et al.; ACCORD Study Group. Effects of intensive blood-pressure control in type 2 diabetes mellitus. *N Engl J Med* 2010;362:1575–1585
- Buse JB, Bigger JT, Byington RP, et al.; ACCORD Study Group. Action to Control Cardiovascular Risk in Diabetes (ACCORD) trial: design and methods. *Am J Cardiol* 2007;99:21i–33i
- ACCORD Study Group. Nine-year effects of 3.7 years of intensive glycemic control on cardiovascular outcomes. *Diabetes Care* 2016;39:701–708
- Echouffo-Tcheugui JB, Zhao S, Brock G, Matsouaka RA, Kline D, Joseph JJ. Visit-to-visit glycemic variability and risks of cardiovascular events and all-cause mortality: the ALLHAT study. *Diabetes Care* 2019;42:486–493
- Bonds DE, Kurashige EM, Bergenstal R, et al.; ACCORD Study Group. Severe hypoglycemia monitoring and risk management procedures in the Action to Control Cardiovascular Risk in Diabetes (ACCORD) trial. *Am J Cardiol* 2007;99:80i–89i
- Pandey A, Keshvani N, Ayers C, et al. Association of cardiac injury and malignant left ventricular hypertrophy with risk of heart failure in African Americans: the Jackson Heart Study. *JAMA Cardiol* 2019;4:51–58
- Wang J, Shi X, Ma C, et al. Visit-to-visit blood pressure variability is a risk factor for all-cause mortality and cardiovascular disease: a systematic review and meta-analysis. *J Hypertens* 2017;35:10–17
- Nichols GA, Hillier TA, Erbey JR, Brown JB. Congestive heart failure in type 2 diabetes: prevalence, incidence, and risk factors. *Diabetes Care* 2001;24:1614–1619
- Parry HM, Deshmukh H, Levin D, et al. Both high and low HbA1c predict incident heart failure in type 2 diabetes mellitus. *Circ Heart Fail* 2015;8:236–242
- Monnier L, Mas E, Ginet C, et al. Activation of oxidative stress by acute glucose fluctuations compared with sustained chronic hyperglycemia in patients with type 2 diabetes. *JAMA* 2006;295:1681–1687
- Ceriello A, Esposito K, Piconi L, et al. Oscillating glucose is more deleterious to endothelial function and oxidative stress than mean glucose in normal and type 2 diabetic patients. *Diabetes* 2008;57:1349–1354
- Nalysnyk L, Hernandez-Medina M, Krishnarajah G. Glycaemic variability and complications in patients with diabetes mellitus: evidence from a systematic review of the literature. *Diabetes Obes Metab* 2010;12:288–298
- Muggeo M, Verlato G, Bonora E, Zoppini G, Corbellini M, de Marco R. Long-term instability of fasting plasma glucose, a novel predictor of cardiovascular mortality in elderly patients with non-insulin-dependent diabetes mellitus: the Verona Diabetes Study. *Circulation* 1997;96:1750–1754
- Kwon S, Lee SR, Choi EK, et al. Visit-to-visit variability of metabolic parameters and risk of heart failure: a nationwide population-based study. *Int J Cardiol* 2019;293:153–158
- Hirakawa Y, Arima H, Zoungas S, et al. Impact of visit-to-visit glycemic variability on the risks of macrovascular and microvascular events and all-cause mortality in type 2 diabetes: the ADVANCE trial. *Diabetes Care* 2014;37:2359–2365
- Zhou JJ, Schwenke DC, Bahn G, Reaven P; VADT Investigators. Glycemic variation and cardiovascular risk in the Veterans Affairs Diabetes Trial. *Diabetes Care* 2018;41:2187–2194
- Bonds DE, Miller ME, Bergenstal RM, et al. The association between symptomatic, severe hypoglycaemia and mortality in type 2 diabetes: retrospective epidemiological analysis of the ACCORD study. *BMJ* 2010;340:b4909
- Tang X, Zhong J, Zhang H, et al. Visit-to-visit fasting plasma glucose variability is an important risk factor for long-term changes in left cardiac structure and function in patients with type 2 diabetes. *Cardiovasc Diabetol* 2019;18:50
- Gottdiener JS, Arnold AM, Aurigemma GP, et al. Predictors of congestive heart failure in the elderly: the Cardiovascular Health Study. *J Am Coll Cardiol* 2000;35:1628–1637
- Bangalore S, Fayyad R, Laskey R, DeMicco DA, Messerli FH, Waters DD. Body-weight fluctuations and outcomes in coronary disease. *N Engl J Med* 2017;376:1332–1340
- Kim MK, Han K, Kim HS, et al. Cholesterol variability and the risk of mortality, myocardial infarction, and stroke: a nationwide population-based study. *Eur Heart J* 2017;38:3560–3566
- Kim MK, Han K, Park YM, et al. Associations of variability in blood pressure, glucose and cholesterol concentrations, and body mass index with mortality and cardiovascular outcomes in the general population. *Circulation* 2018;138:2627–2637
- Nishimura R, Osonoi T, Kanada S, et al. Effects of luseogliflozin, a sodium-glucose co-transporter 2 inhibitor, on 24-h glucose variability assessed by

continuous glucose monitoring in Japanese patients with type 2 diabetes mellitus: a randomized, double-blind, placebo-controlled, crossover study. *Diabetes Obes Metab* 2015; 17:800–804

39. Marso SP, McGuire DK, Zinman B, et al.; DEVOTE Study Group. Efficacy and safety of

degludec versus glargine in type 2 diabetes. *N Engl J Med* 2017;377:723–732

40. Zelniker TA, Wiviott SD, Raz I, et al. SGLT2 inhibitors for primary and secondary prevention of cardiovascular and renal outcomes in type 2 diabetes: a systematic review and meta-analysis of cardiovascular outcome trials. *Lancet* 2019;393:31–39

41. Kristensen SL, Rørth R, Jhund PS, et al. Cardiovascular, mortality, and kidney outcomes with GLP-1 receptor agonists in patients with type 2 diabetes: a systematic review and meta-analysis of cardiovascular outcome trials. *Lancet Diabetes Endocrinol* 2019;7:776–785