

UC Irvine

Western Journal of Emergency Medicine: Integrating Emergency Care with Population Health

Title

Potentially Avoidable Transfers of Veterans with Mental Health Conditions in the Veterans Health Administration

Permalink

<https://escholarship.org/uc/item/7zs493xc>

Journal

Western Journal of Emergency Medicine: Integrating Emergency Care with Population Health, 20(2)

ISSN

1936-900X

Authors

Mohr, Nicholas M.
Wu, Chaorong
Ward, Michael J.
[et al.](#)

Publication Date

2019

Copyright Information

Copyright 2019 by the author(s). This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

9th Annual National Update on Behavioral Emergencies Conference (NUBE) in Las Vegas, Nevada (December 12-14, 2018)

total number of emergency psychiatry consults, discharge rates, and transfer rates.

Results: The number of consults per day has been increasing by about 13.8% a month over the last few years and is now around 16-17 a day. The service discharges about 45% of the patients consulted to us; and of those requiring admission, about 35% are transferred to other psychiatric unit, with the rest being admitted to UMCNO's 60-bed inpatient psychiatric unit. Looking at the seven months before and after the changes were made, the average LOS has decreased from 15.98 hours to 13.78 hours (a 17% decrease), and the number of hours on saturation decreased from 42.3 hours a month to 19.2 hours (a 55% decrease).

Discussion: While our goal of zero hours on saturation was not met, the data show that by planning for the increase in volume during the weekend with more staff starting Sunday evening to open all 26 beds, we were able to lower saturation hours, which helps throughput in the main ED and throughout the hospital. Furthermore, by increasing the hours of clerks on weekends (who are responsible for transferring patients when our inpatient unit is full), we were able to transfer more patients throughout the weekend than previously. And finally, by integrating our handoff within our EHR, we were able to quickly identify those patients who could potentially be discharged safely and what was needed to ensure that safe discharge. Combined, these efforts lowered the average of LOS in the BHER.

5 Potentially Avoidable Transfers of Veterans with Mental Health Conditions in the Veterans Health Administration

NM Mohr¹, C Wu^{1,2,3}, MJ Ward^{1,4}, CD McNaughton^{5,6}, PJ Kaboli^{1,7}
¹VA Iowa City Healthcare System, Center for Comprehensive Access Delivery Research & Evaluation (CADRE), Iowa City, Iowa; ²University of Iowa Carver College of Medicine, ³Department of Emergency Medicine, ⁴Department of Anesthesia, Iowa City, Iowa; ⁵University of Iowa, Institute for Clinical and Translational Sciences Iowa City, Iowa; ⁶Tennessee Valley Healthcare System, VA Medical Center, Nashville, Tennessee; ⁷Vanderbilt University Medical Center, Department of Emergency Medicine, Nashville, Tennessee; ⁸University of Iowa Carver College of Medicine, Department of Internal Medicine, Iowa City, Iowa

Objective: Over 40% of the 2.4 million emergency department (ED) visits to Veterans Health Administration (VHA) hospitals are from veterans who live in rural areas, a population at increased risk of interfacility transfer. Veterans may undergo interfacility transfer to obtain emergent or urgent access to specialized health services, particularly mental health care. However, such transfers raise questions regarding appropriate use of resources, travel burdens for patients and families, and logistical challenges for ED staff and providers that may delay timely care. We sought to describe ED-based, interfacility transfer

rates within the VHA and to estimate the proportion of potentially avoidable transfers (PAT) of patients with mental health conditions relative to other diseases.

Methods: This observational cohort included all patients who were transferred from a VHA ED to another VHA hospital between 2012 and 2014. We extracted data from Clinical Data Warehouse administrative data. PAT was defined as discharge from the receiving ED without a procedure, or hospital length of stay at the receiving hospital ≤ 1 day without having a procedure performed. We conducted facility-level and diagnosis-level analysis to identify conditions for which an alternative to transfer, such as telehealth access to specialty care, could be developed and implemented in low-volume or rural EDs.

Results: Of 6,131,734 ED visits during the three-year study period, 18,875 (0.3%) were transferred from one VHA ED to another VHA facility. Rural residents were transferred three times as often as urban residents (0.6% vs. 0.2%, $p < 0.001$), and 23.6% of all VHA-to-VHA transfers met the PAT definition. Mental health conditions were the most common reason for interfacility transfer (34% of all interfacility transfers), followed by heart disease (12%). Of transfers that met PAT criteria, 11% were for mental health diagnoses whereas 21% were for heart disease. Geographic analysis suggested that overall PAT proportion ranged across regions from 8-53% with mental health PATs between 2-42%.

Conclusion: VHA interfacility transfer is commonly performed for mental health diagnoses, and there is substantial regional variation in potentially avoidable transfers in a national sample of transfers. A significant proportion of these transfers may be potentially avoidable. Future work should focus on improving capabilities to provide specialty evaluation locally for these conditions, possibly using telehealth solutions. Additional work should also focus on measuring the timeliness of these transfers.

6 Reducing Emergency Department Length of Stay and Wait Times for Psychiatric Patients

JD Richter, D Blackhurst, P Moschella, KM Lommel/ Greenville Health System, South Carolina

Introduction: In the past 20 years there has been a significant decline in the number of inpatient psychiatric beds in the United States, while the number of patients seeking psychiatric treatment in the emergency department (ED) has increased over the same time period. Given the increase in demand for psychiatric services and decrease in availability of inpatient treatment the ED is becoming the de facto place of treatment for the majority of psychiatric crises. Psychiatric patients experience longer lengths of stay (LOS) when compared to non-psychiatric patients,