

Lawrence Berkeley National Laboratory

LBL Publications

Title

A Possible Primary Quantum Conversion Act of Photosynthesis

Permalink

<https://escholarship.org/uc/item/8043t5gf>

Authors

Calvin, M

Barltrop, J A

Publication Date

1952-12-01

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

UCRL- 2024

UNCLASSIFIED

UNIVERSITY OF CALIFORNIA - BERKELEY

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 5545*

RADIATION LABORATORY

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

UCRL-2024
Unclassified-Chemistry Distribution

UNCLASSIFIED

UNIVERSITY OF CALIFORNIA

Radiation Laboratory

Contract No. W-7405-eng-48

A POSSIBLE PRIMARY QUANTUM CONVERSION ACT OF PHOTOSYNTHESIS

M. Calvin and J. A. Barltrop

December 5, 1952


Berkeley, California

A POSSIBLE PRIMARY QUANTUM CONVERSION

ACT OF PHOTOSYNTHESIS¹M. Calvin and J. A. Barltrop²

Radiation Laboratory and Department of Chemistry
 University of California
 Berkeley, California

To account for the observation^{3,4,5} that illumination prevents the appearance of newly assimilated carbon in the compounds of the tricarboxylic acid cycle, it was suggested⁵ that the light shifts the steady-state condition of the thioctic acid-containing coenzyme^{6,7} (protogen, lipoic acid, thioctic acid, P.O.F.) toward the reduced (dithiol) form, in which condition it is incapable of oxidatively decarboxylating pyruvic acid,^{8,9} newly formed from CO₂ to give rise to the acetyl-CoA^{10,11,12} required to bring this carbon into the compounds of the Krebs cycle. We are here reporting some observations leading to the further suggestion that this shift toward the dithiol form is the direct result of the light action and that a biradical formed by dissociation of the disulfide bond in a strained five-membered disulfide containing ring (as in 6,8-thioctic acid and trimethylenedisulfide) is the species in which the quantum absorbed by the plant pigments and stored as electronic excitation in chlorophyll¹³ appears first as chemical bond potential energy; i.e., that a possible primary quantum conversion act of photosynthesis is represented by the equation



Subsequent abstraction of H atoms^{14,15} from a suitable donor by the thiyl free radicals would lead to the dithiol which would be reoxidized ultimately by CO₂. The residual oxidation product of the H donor would lead in the end to molecular oxygen. It is obvious that on both the reductant and oxidant sides the chemical products of the conversion of several quanta will be required to accomplish the reduction of each CO₂ molecule and the generation of each O₂ molecule.

These subsequent reactions, being strictly chemical, may lead to diverse energy rearrangements. For example, the chemical potential of reduced carbon might be converted by oxidative phosphorylation reactions into the energy of phosphoric anhydrides, which in turn, could raise the potential energy of intermediates in the reaction sequences leading to the evolution of molecular oxygen and to the reduction of CO₂.^{16,17}

A value of the dissociation energy for this particular disulfide link lying in the region of 30-40 Kcals. would constitute not only permissive evidence for reaction (I) but positive evidence in its support, since hitherto it has been difficult to suggest any likely primary chemical step capable of usefully absorbing the greater part of the \approx 30-40 Kcal. quantum of electronic excitation available for photosynthesis. Estimates of D(RS-SR) from simple open chain compounds range from 50¹⁸ to 70¹⁹ Kcals. However, the fact that 5,8-thioctic acid is colorless, while 6,8-thioctic acid is yellow,²⁰ suggested that the incorporation of the S-S bond into a 5-membered ring might indeed introduce sufficient strain into it so as to reduce the S-S dissociation energy by as much as 25-30 Kcals., thus bringing it down into the required range. A number of experiments have been performed using the product of the reaction of Na₂S₂ with (CH₂)₃Br₂ (trimethylenedisulfide) as a model substance. Its absorption

spectrum together with that of the two thiocetic acids and n-propyl disulfide is shown in Figure 1. A light induced fading of diphenylpicrylhydrazyl²² at room temperature and dependent upon the presence of the disulfide was demonstrated. A photochemical polymerization of the disulfide by light of wavelength greater than 4,000 Å was sensitized by Zn tetraphenylporphin. These results may be taken to indicate that the dissociation energy of the disulfide bond in this compound is something less than 50 Kcals. and that this dissociation may be brought about by energy transfer from some other molecule in a suitably excited state.

That such energy transfers may take place, and especially efficiently in condensed systems, has been amply demonstrated.^{23,24,25,26,27} Since the grana have the optical properties of a condensed chlorophyll phase,^{28,29,30,31} a quantum absorbed anywhere within that phase is very rapidly transferred among the identical molecules of that phase at the singlet or the triplet levels. Concomitant with this, there occurs a decrease in the probability of emission as fluorescence^{23,24,25,32} and hence an increase in the availability of the quantum for chemical transformation (disulfide fission). In such a system, the high efficiency of energy conversion may be retained even though the ratio of chlorophyll to disulfide molecules be large (10^2-10^3).

References

- (1) The work described in this paper was sponsored by the U. S. Atomic Energy Commission.
- (2) Rockefeller Fellow, 1952-1953, while on leave from Brasenose College and the Dyson Perrins Laboratory, Oxford University, England.
- (3) A. A. Benson and M. Calvin, *J. Exptl. Bot.*, 1, 63 (1950).
- (4) J. W. Weigl, P. M. Warrington and M. Calvin, *J. Am. Chem. Soc.*, 73, 5058 (1951).
- (5) M. Calvin and Peter Massini, *Experientia*, in press.
- (6) G. W. Kidder and V. C. Dewey, *Protozoa*, I, 388, Acad. Press, 1951.
- (7) E. L. Patterson, J. A. Brockman, Jr., F. P. Day, J. V. Pierce, M. E. Macchi, C. E. Hoffman, C. T. O. Long, E. L. R. Stokstad, T. H. Jukes, *J. Am. Chem. Soc.*, 73, 5919 (1951).
- (8) L. J. Reed, B. G. DeBusk, I. C. Gunsalus, G. H. F. Schnakenberg, *J. Am. Chem. Soc.*, 73, 5920 (1951).
- (9) I. C. Gunsalus, L. Struglia, D. J. O'Kane, *J. Biol. Chem.*, 194, 859 (1952).
- (10) J. R. Stern, *Phosphorus Metabolism*, I, Johns Hopkins Press, 1951.
- (11) S. Korke, *Phosphorus Metabolism*, I, Johns Hopkins Press, 1951.
- (12) R. S. Sweet, M. Fuld, K. Cheslock, M. H. Paul, *Phosphorus Metabolism*, I, Johns Hopkins Press, 1951.
- (13) L. N. M. Duysens, *Nature*, 168, 548 (1951).
- (14) A. F. Bickel and E. C. Kooijman, *Nature*, 170, 211 (1952).
- (15) E. F. P. Harris and W. A. Waters, *Nature*, 170, 212 (1952).
- (16) *Phosphorus Metabolism*, I, Johns Hopkins Press, 1951, Sec. V.
- (17) M. Calvin, J. A. Bassham, A. A. Benson and P. Massini, *Ann. Rev. Phys. Chem.*, 3, 215 (1952).
- (18) A. H. Schon, *J. Am. Chem. Soc.*, 74, 4723 (1952).
- (19) J. L. Franklin and H. E. Lumpkin, *J. Am. Chem. Soc.*, 74, 1024 (1952).
- (20) Several milligrams of each of these synthetic²¹ products were obtained through the courtesy of Dr. T. H. Jukes of Lederle Laboratories.

- (21) M. W. Bullock, J. A. Brockman, E. L. Patterson, J. V. Pierce, E. L. R. Stokstad, *J. Am. Chem. Soc.*, 74, 3455 (1952).
- (22) C. E. H. Bawn and S. F. Mellish, *Trans. Farad. Soc.*, 47, 1217 (1951).
- (23) H. Kallman and M. Furst, *Phys. Rev.*, 79, 857 (1950).
- (24) H. Kallman and M. Furst, *Phys. Rev.*, 81, 853 (1951).
- (25) H. Kallman and M. Furst, *Phys. Rev.*, 85, 816 (1952).
- (26) M. M. Moodie and C. Reid, *J. Chem. Phys.*, 20, 1510 (1952).
- (27) C. Reid, *Phys. Rev.*, 88, 422 (1952).
- (28) S. Grancik and K. R. Porter, *Amer. J. Bot.*, 34, 545 (1947).
- (29) M. Calvin and V. Lynch, *Nature*, 169, 455 (1952).
- (30) E. E. Jacobs and A. S. Holt, *J. Chem. Phys.*, 20, 1326 (1952).
- (31) A. Frey-Wyssling and K. Mühlethaler, *Vierteljahrsschrift der Naturf. Ges. Zurich*, 94, 182 (1949).
- (32) D. McClure, private communication.
- (33) We are indebted to Mr. Paul Hayes for the determination of these absorption spectra.

Caption to Figure

Figure 1 - Absorption spectra of several disulfides³³

