

# Influence of time delay and nonlinear diffusion on herbivore outbreak



Gui-Quan Sun<sup>a,b,c,\*</sup>, Amit Chakraborty<sup>c</sup>, Quan-Xing Liu<sup>c</sup>, Zhen Jin<sup>a,b</sup>, Kurt E. Anderson<sup>d</sup>, Bai-Lian Li<sup>c</sup>

<sup>a</sup>Complex Sciences Center, Shanxi University, Taiyuan 030006, PR China

<sup>b</sup>School of Mathematical Sciences, Shanxi University, Taiyuan 030006, PR China

<sup>c</sup>Ecological Complexity and Modeling Laboratory, Department of Botany and Plant Sciences, University of California, Riverside, CA 92521-0124, USA

<sup>d</sup>Department of Biology, University of California, Riverside, CA 92521, USA

## ARTICLE INFO

### Article history:

Received 9 February 2013

Received in revised form 20 August 2013

Accepted 16 September 2013

Available online 26 September 2013

### Keywords:

Herbivore-plant

Time delay

Spatial diffusion

Outbreak

Synchrony

## ABSTRACT

Herbivore outbreaks, a major form of natural disturbance in many ecosystems, often have devastating impacts on their food plants. Understanding those factors permitting herbivore outbreaks to occur is a long-standing issue in conventional studies of plant-herbivore interactions. These studies are largely concerned with the relative importance of intrinsic biological factors and extrinsic environmental variations in determining the degree of herbivore outbreaks. In this paper, we illustrated that how the time delay associated with plant defense responses to herbivore attacks and the spatial diffusion of herbivore jointly promote outbreaks of herbivore population. Using a reaction-diffusion model, we showed that there exists a threshold of time delay in plant-herbivore interactions; when time delay is below the threshold value, there is no herbivore outbreak. However, when time delay is above the threshold value, periodic outbreak of herbivore emerges. Furthermore, the results confirm that during the outbreak period, plants display much lower density than its normal level but higher in the inter-outbreak periods. Our results are supported by empirical findings.

Crown Copyright © 2013 Published by Elsevier B.V. All rights reserved.

## 1. Introduction

Herbivory is an important feeding process. By feeding on different plant parts or materials, herbivores can affect plant growth, transfers of nutrients to the soil surface, and habitat and resource conditions for other organisms (see the review of [58] and references therein). These effects are mostly viewed as beneficial outcomes of plant-herbivore interactions as it exerts positive effects for the maintenance and conservation of plant population structure and composition during non-outbreak periods. However, periodic outbreaks of herbivores can denude or kill plants over many square kilometers and led to changes in plant community structure and composition. This capacity to alter community structure allows herbivores to act as biotic agents of disturbance [69]. Unlike abiotic disturbances, herbivore outbreaks have continuous impact on plant population and hold selectivity for the feeder plant species. As a result, a core issue of herbivory is that a threshold at which it shifts from a normally acting process to a disturbance in terms of intensity, scale and frequency similar to those of fire, storm, drought or flood. In temperate forest, for example, at a normal level of herbivory, defoliating insects consume 5–15 percents of foliage production but it may reach to 100 percents during outbreak periods [50,39,9].

\* Corresponding author at: Complex Sciences Center, Shanxi University, Taiyuan 030006, PR China.

E-mail addresses: [gquansun@126.com](mailto:gquansun@126.com) (G.-Q. Sun), [amitc@ucr.edu](mailto:amitc@ucr.edu) (A. Chakraborty), [q.liu@nioo.knaw.nl](mailto:q.liu@nioo.knaw.nl) (Q.-X. Liu), [jinzhn@263.net](mailto:jinzhn@263.net) (Z. Jin), [kurta@ucr.edu](mailto:kurta@ucr.edu) (K.E. Anderson), [bai-lian.li@ucr.edu](mailto:bai-lian.li@ucr.edu) (B.-L. Li).

Although many herbivore populations exhibit outbreak behavior, the main causes of these phenomena remain unclear [2]. Thus, determining which factors regulate or induce outbreak in herbivore populations continues to be an important issue in the field of both ecological and agricultural research [66,8]. The previous work revealed that herbivore outbreak was caused by interactions with enemies [62,30,41,40,13], inducible defences or physiological stress [14,6,37,65,71,32,67], the case that herbivore population's parents and grandparents experienced in preceding generations [17], environmental forcing [3,15,26] and resource-limited [1,2]. The aim of this paper is to give another mechanism for the outbreak of the herbivore.

While plant-herbivore interactions exhibit cyclic population dynamics, the spatiotemporal outbreak patterns of herbivore population are often explained by those underlying causes of population synchrony in which peak phases of herbivore population cycle are inversely correlated with that of crash phases of plant population cycle. A suite of studies have attempted to link these periodic outbreak patterns with that of the large-scale processes such as climatic variation [46,48]. The other approaches focused on the analysis of the rate of spread of herbivores, seeking traveling wave solutions in the model of plant-herbivore interactions. In this case that rate of spread often depends on distributions and dispersal ranges of both populations [60,32]. Consequently, both spatial and temporal aspects play very important roles in the plant-herbivore population dynamics, which can not be ignored.

On the other hand, biological systems with time delays have been of considerable interest. It means the time between immature and mature [18], the time of pregnancy of the mother [31] and so on [16,49,38]. The interaction of plant and herbivore share some common features with prey-predator model. So delay widely exists in the dynamics of plant-herbivore interaction [10,51]. For example, the delay may arise between herbivore damage and deployment of inducible defenses. Moreover, theoretical analysis pointed out that time lags in reproduction influencing population dynamics [5,57,53,68].

Underwood et al. firstly demonstrated that time delays in inducible defenses and non-linear dispersal can lead to spatial instabilities [67]. However, it was a simple simulation study and thus, studies of the population consequences of time delay and the spatial spread of populations have remained largely unconnected. As a result, in the present paper, our aim is to build a better understanding of how time delay affects the dynamic behavior of herbivores in both space and time, including spatial pattern and herbivore fluctuation. We will analysis a reaction-diffusion model by addressing the following questions: (1) How do time delay influence herbivore cycles; (2) How do time delay and nonlinear diffusion term have effect on the spatial spread of the herbivore?

The paper is organized as follows. In Section 2, we use a reaction-diffusion model with delay to describe the interaction of plant and herbivore. In Section 3, by using both mathematical analysis and numerical simulation, we show that herbivore outbreak can be induced by time delay. Moreover, we show the relationship between wavelength and time delay. Finally, conclusions and discussions are presented in Section 4.

## 2. The model

Since that long time series of the density of both plant and herbivore is needed, it is difficult to identify the causes of the fluctuations empirically [35,1]. Thus, it may provide useful information by constructing mathematical models to explain the phenomenon observation in the real world.

Here, we introduce a simple reaction-diffusion model to analyze the effect of time delays in the induction of plant defenses on the outbreak and spatial distribution of herbivores in a closed, one-dimensional landscape. We model the level that inducible defenses have reached in each location at each time, which is dependent on herbivore density and the level of already induced defenses. We do not explicitly model changes in plant biomass since such changes are potentially uncorrelated with the effects of induction in empirical plant-herbivore systems [29]. Induced defenses increase in response to herbivore densities according to a saturating function such that there is a maximum amount  $\alpha$  by which induction can change between  $t$  and  $t + \Delta t$ ; the shape of the saturation curve is set by  $\theta$ . Induction also exhibits self-limitation, tuned by  $\beta$ , which reduces the potential for increased induction in response to new damage. To reflect delays in the deployment of inducible defenses, we set induction changes at time  $t$  dependent on herbivore densities at  $t - \tau$  time steps previously. Induction decays at a constant rate  $\delta$ , reflecting the breakdown of inducible defenses and plant repair [68]. We assume that the herbivore population grows logistically in the absence of induced defenses and suffers a linear increase in mortality when defenses are activated. Furthermore, we assume that herbivores move away from areas of high induced defenses according to Fokker-Planck nonlinear diffusion, which is dependent on the density of both induction levels and herbivores.

On the basis of the above assumptions, we arrive at the following equations:


$$\frac{\partial I(x, t)}{\partial t} = (\alpha - \beta I(x, t)) \frac{H^\theta(x, t - \tau)}{b^\theta + H^\theta(x, t - \tau)} - \delta I(x, t), \quad (1.a)$$

$$\frac{\partial H}{\partial t} = rH(x, t) \left[ 1 - \frac{H(x, t)}{K} \right] - mI(x, t)H(x, t) + \frac{\partial^2}{\partial x^2} [(D_0 + \chi I(x, t))H(x, t)], \quad (1.b)$$

where  $I(x, t)$  and  $H(x, t)$  represent induced defense and herbivore density in both space and time. The biological meanings of the parameters are given in Table 1.

**Table 1**  
Parameter used in model (1).

Symbol	Comments
$\alpha$	maximum induction rate per herbivore
$\beta$	per-unit reduction of induction rate by self-limitation
$\delta$	per-unit induction decay rate
$b$	half-maximum for herbivore effectiveness of damage
$\theta$	herbivore damage effectiveness shape tuning parameter
$r$	intrinsic rate of herbivore population growth
$K$	herbivore carry-capacity in the absence of inducible defenses
$m$	mortality rate by induction
$D_0$	baseline diffusion rate
$\chi$	sensitivity of diffusion rate to induction
$\tau$	time delay between herbivore damage and deployment of inducible defenses



**Fig. 1.** Plot of  $Q(s)$  as a function of  $s$ , and the expression of  $Q(s)$  is in Eq. (2). Note that there are two pure imaginary solutions of  $Q(s)$ , which have no biological meanings.

### 3. Main results

In this section, we set that  $\alpha = 200$ ,  $\beta = 1$ ,  $\delta = 0.75$ ,  $b = 5$ ,  $\theta = 3$ ,  $r = 1$ ,  $K = 10$ ,  $m = 0.01$ ,  $D_0 = 0.1$ ,  $\chi = 0.1$ , and to see what will occur as the time delay increases. The first step in analyzing the model is to determine the stationary point of the non-spatial model obtained by setting space derivatives equal to zero. It is, naturally, the dynamics in the biologically meaningful region  $I \geq 0, H \geq 0$  that are of interest. By considering the nullclines  $F(I, H) = 0$ ,  $G(I, H) = 0$  (see from Appendix A), and the intersection of these curves in phase space, we have a stationary point  $(0, 0)$  which is a saddle,<sup>1</sup> and  $(I^*, H^*)$ . Additionally,  $I^*$  is the solution of

$$Q(s) = 0.00175s^4 - 0.725s^3 + 112.5s^2 - 7843.75s + 200000 \quad (2)$$


and  $H^* = 10 - 0.1I^*$ . It can be seen from Fig. 1 that there are positive stationary point. One is  $(60, 4)$ , which is locally stable, and the other one does not exist by reason of the negative of  $H^*$ . Thus, from the biological point of view, we are interested to study the stability behavior of the interior equilibrium point  $(60, 4)$ .

From the stability analysis in Appendix B and direct calculations, we know that when  $\tau > \tau_c = 2.54980875$ , Hopf bifurcation of the no-spatial model emerges. On the other hand, the positive equilibrium  $(60, 4)$  is stable when  $\tau < \tau_c$ . This is illustrated by the numerical simulation in Fig. 2 (with initial data  $(70, 2)$ ). Further, when delay passes through the critical value  $\tau_c$ , the positive equilibrium  $(60, 4)$  loses its stability and the system goes into oscillations, see Fig. 3.


Now, we want to see the effect of time delay when spatial terms are added. In Fig. 4, the dispersal relation,<sup>2</sup> which can be obtained from Eq. (B.9) in Appendix B, is given when there is no time delay. It is easy to find that, when there is no time delay, there is no spatial pattern.

<sup>1</sup> The local stability of the stationary point can be checked by Eq. (A.3) in Appendix A.


<sup>2</sup> The dispersion relation can determine how time oscillations  $e^{i\omega t}$  are linked to spatial oscillations  $e^{i\kappa \vec{r}}$  (here,  $\vec{r} = x(y)$  or  $\vec{r} = (x, y)$ ). The dispersion relation can be described by the characteristic value  $-\lambda$ , with respect to wave number  $\kappa$ .



**Fig. 2.** When  $\tau = 1 < \tau_c$ , the positive equilibrium (60,4) is stable. Parameter values are used as:  $\alpha = 200, \beta = 1, \delta = 0.75, b = 5, \theta = 3, r = 1, K = 10, m = 0.01, D_0 = 0$  and  $\chi = 0$ . (A) Time series of the induced defense and herbivore; (B) Phase diagram of induced defense and herbivore.




**Fig. 3.** When  $\tau = 4 > \tau_c$ , Hopf bifurcation occurs and leads periodic solution. Parameter values are used as:  $\alpha = 200, \beta = 1, \delta = 0.75, b = 5, \theta = 3, r = 1, K = 10, m = 0.01, D_0 = 0$  and  $\chi = 0$ . (A) Time series of the induced defense and herbivore; (B) Phase diagram of induced defense and herbivore.




**Fig. 4.** Dispersion relation of the model (1) without time delay. Parameter values are used as:  $\alpha = 200, \beta = 1, \delta = 0.75, b = 5, \theta = 3, r = 1, K = 10, m = 0.01, D_0 = 0.1, \chi = 0.1$  and  $\tau = 0$ . Note that the characteristic value is less than zero, which means there is no pattern.

3.1. Spatial pattern

In the following part, numerical solutions for the system (1) with both time delay and spatial terms in the one-dimensional space will be presented. In practice, the reaction-diffusion system is solved in a discrete domain. The space between the lattice points is defined by the lattice constant  $\Delta x$ . In the discrete system the Laplacian describing diffusion is calculated by using finite differences, i.e., the derivatives are approximated by differences over  $\Delta x$ . For  $\Delta x \rightarrow 0$ , the differences approach



**Fig. 5.** Model solutions with  $\alpha = 200, \beta = 1, \delta = 0.75, b = 5, \theta = 3, r = 1, K = 10, m = 0.01, D_0 = 0.1, \chi = 0.1,$  and  $\tau = 1$ . The initial condition is that  $I(x, 0) = 60,$  and  $H(x, 0) = 4 + 10^{-7}(x - 300)$ .




**Fig. 6.** Model solutions with  $\alpha = 200, \beta = 1, \delta = 0.75, b = 5, \theta = 3, r = 1, K = 10, m = 0.01, D_0 = 0.1, \chi = 0.1,$  and  $\tau = 4$ . The initial condition is the same as in Fig. 5.

the derivatives. The time evolution is also discrete, i.e., the time goes in steps of  $\Delta t$ . The time evolution can be solved by using the Euler method, which means approximating the value of the concentration at the next time step on the basis of the change rate of the concentration at the previous time step. In our calculations, the parameters values were taken to be  $\Delta x = 1, \Delta t = 0.01$  and the space was discretized to  $1 \times 600$  lattices on the one-dimensional space. The periodic boundary condition was used in the simulation. We also find that when we vary the number of mesh points such as  $1 \times 1000$  and  $1 \times 300$ , the results of system (1) have the same dynamical behaviors. At the same time, it was checked that a further decrease of the step values did not lead to any significant modification of the results.


In Fig. 5, we show the solution with respect to the space and time with  $\tau < \tau_c$ . One can see that the solutions are stable and converge to constant. However, as time delay increases as the case  $\tau > \tau_c$ , the model (1) has period solutions which is shown in Fig. 6. As the time delay further increases, the amplitude of the oscillations becomes larger, see in Fig. 7. In a word, time delay plays a constructive role in the pattern formation of the herbivore, i.e., induce periodic pattern.

### 3.2. Synchronization and outbreak

Synchronization is a fundamental phenomenon arising in many biological and physical contexts for which there are two or more coupled oscillating systems. In the classical sense, and dating back at least to Huygens in the 17th century [27], synchronization has been understood as the mutual adjustment of periodic oscillators and the frequency locking that results because of their (often weakly) coupled interaction [63]. Synchrony has also been observed among populations of sympatric species that are not directly linked through trophic interactions, such as herbivorous forest insects [42,25,45,55,33] and allopatric large herbivores [54].



**Fig. 7.** Model solutions with  $\alpha = 200$ ,  $\beta = 1$ ,  $\delta = 0.75$ ,  $b = 5$ ,  $\theta = 3$ ,  $r = 1$ ,  $K = 10$ ,  $m = 0.01$ ,  $D_0 = 0.1$ ,  $\chi = 0.1$ , and  $\tau = 10$ . The initial condition is the same as in Fig. 5.



**Fig. 8.** Synchrony of the induced defense and the herbivore with  $\tau = 2.5$ . Parameter values are used as:  $\alpha = 200$ ,  $\beta = 1$ ,  $\delta = 0.75$ ,  $b = 5$ ,  $\theta = 3$ ,  $r = 1$ ,  $K = 10$ ,  $m = 0.01$ ,  $D_0 = 0.1$  and  $\chi = 0.1$ .


In Figs. 8–10, we show the time series of the induced defense and herbivore as time delay increases. One can see from these three figures, the induced defense and the herbivore are anti-phase synchrony. When the time delay is smaller than but very close to the critical value, both the two population exhibit fluctuation behavior. When it is becoming larger than the critical value, periodic solutions with fixed period emerge, see Figs. 9 and 10.

To well explain the cause of herbivore outbreak, we plot the time series of the two population when  $\tau$  is less than  $\tau_c$  in Fig. 11. When  $\tau$  is small, the density of induced defense and herbivore converge to a constant for a long time. In other words, there is no outbreak for small  $\tau$ . As seen from Fig. 8, when  $\tau$  is larger than the critical values, fluctuation of herbivore is shown. As the time delay further increases, the oscillation aptitude and period becomes larger by comparing with Fig. 9 and 10. We can conclude that herbivore outbreak is induced by time delay. Moreover, Figs. 8–10 show that the density of induced defense is very low in the outbreak period, but high in the inter-outbreak period, which confirm the previous argument [34].


### 3.3. Wavelength

Investigating the relationship between spatial patterns in population densities and environmental heterogeneity is crucial to the understanding of population dynamics and for the management of species in communities. Consequently, determining the wavelength for the spatial pattern is a key issue in the field of spatial ecology [61,70]. In other words, we can see the distribution of the population in the spatial space directly by calculating the wavelength.


In order to see the effect of time delay, we set  $\tau$  as a parameter, and show the corresponding wavelength as it varies in Fig. 12. It can be found that the time delay increase the wavelength of the spatial pattern, and wavelength reaches the maximum values when  $\tau \approx 3.5$ . However, when  $\tau$  further increases, the wavelength is a decreasing function of it. From the



**Fig. 9.** Synchrony of the induced defense and the herbivore, and the parameters are the same as in Fig. 6.



**Fig. 10.** Synchrony of the induced defense and the herbivore and the parameters are the same as in Fig. 7.




**Fig. 11.** Time series of the induced defense and the herbivore with  $\tau = 1$ . Parameter values are used as:  $\alpha = 200$ ,  $\beta = 1$ ,  $\delta = 0.75$ ,  $b = 5$ ,  $\theta = 3$ ,  $r = 1$ ,  $K = 10$ ,  $m = 0.01$ ,  $D_0 = 0.1$  and  $\chi = 0.1$ .

biological point of view, the increase of time delay can largely affect the distance between one peak distribution and the adjacent one.

#### 4. Discussion and conclusion

In this paper, we present a reaction-diffusion model described by interaction of plant and herbivore with delay. The results showed that the time delay can induce the herbivore outbreak (cf. Figs. 8–10). Moreover, herbivore outbreak in our model is characterized by high density of induced defense during the inter-outbreak period, which is consistent with the experimental data [34,36,56,59]. In addition to that, we show that time delay may have great effect on the wavelength of the spatial pattern (cf. Fig. 12). We can see from the figure that there exist a threshold value on time delay that the





**Fig. 12.** An illustration of the variation in pattern wavelength with different  $\tau$ . Parameter values are used as:  $\alpha = 200$ ,  $\beta = 1$ ,  $\delta = 0.75$ ,  $b = 5$ ,  $\theta = 3$ ,  $r = 1$ ,  $K = 10$ ,  $m = 0.01$ ,  $D_0 = 0.1$  and  $\chi = 0.1$ .

wavelength reaches its maximum value. In summary, our results identify some biological features of plant-herbivore dynamics that determines whether the outbreak occurs or not and suggest the further investigation should take into account the effects of both time delay and spatial diffusion.

These above-noted model results are consistent with the empirical findings in birch forest regions, where defoliating insect outbreaks are one of the most common disturbances. Along the Scandinavian mountain chain, some birch forest areas are usually experienced with periodic, synchronous peaks of Geometrid defoliators *Epirrita* (*Oporinia*) *autumnata* and *Operophтера* spp. [64]. Their outbreak cycles are characterized by large amplitude with 1000 to 100,000 fold differences between population lows and peaks [24,28,7], which is clearly larger than in the well known vole cycles with 25–250 fold differences [21]. It has been observed that although the period of the cycle is remarkably stable, the peak densities show large variation even within a single site (see Neuvonen et al., 2001 and references therein). Detailed investigation of these patterns revealed that the time delay associated with inducible plant defenses, which refers to decreasing foliage quality in the years following insect attacks, is one of the major factors behind such cyclic outbreak patterns [24,22,23,47].

An interesting question of interest is whether nonlinear diffusion terms have effect on the herbivore outbreak. To address this issue, we performed extensive computer simulations for different parameter values of  $D$  and  $\chi$  (in total, more than 50 parameter sets were examined). We found that the results nonlinear diffusion just influence the density not the outbreak (for the sake of brevity, we do not provide separated figures).

The present paper provides some new insights into the spatial ecology of interaction of plant and herbivore, but it also leaves many questions open for the future investigations. We would like to emphasize two relevant topics where may be the main subjects of the further research. Firstly, in the model (1), we consider the case that only herbivores diffuse in the space. However, for some plants their weeds also can move in the space caused by purely environmental factors such as wind. In that situation, the model we employ may be changed as

$$\frac{\partial I(x, t)}{\partial t} = (\alpha - \beta I(x, t)) \frac{H^\theta(x, t - \tau)}{b^\theta + H^\theta(x, t - \tau)} - \delta I(x, t) + D_I \frac{\partial^2 I(x, t)}{\partial x^2}, \tag{3.a}$$

$$\frac{\partial H}{\partial t} = rH(x, t) \left[ 1 - \frac{H(x, t)}{K} \right] - mI(x, t)H(x, t) + \frac{\partial^2}{\partial x^2} [(D_0 + \chi I(x, t))H(x, t)], \tag{3.b}$$

where  $D_I$  is the diffusion coefficient of the plant. Secondly, we found that periodic wave solution in the model (1) (cf. Figs. 6 and 7). However, previous work revealed that traveling wave of herbivore may exist by modeling the interaction of plant and herbivore [44]. Thus, we need to check whether model (1) have traveling wave solution, especially how time delay influence such solution.

It also should be noted that, in our model simulations, we have considered all the parameters as constants. However, herbivore populations are sometimes considered to be strongly influenced by abiotic factors such as weather and climatic conditions [3,43,11,12,19]. As a result, all the parameters can show temporal and spatial variations: indeed, some can be both stochastic and show significant seasonal variations [20,52,4]. We hope that our efforts will provide a new starting point for the analysis of more detailed models to understand the outbreak in plant-herbivore dynamics.

**Acknowledgments**

We are grateful for the constructive suggestions of the anonymous referees on our original manuscript.

The research was partially supported by the National Natural Science Foundation of China under Grant Nos. 11301490, 11331009, 11147015 and 11171314, Natural Science Foundation of Shan’Xi Province Grant No. 2012021002-1, the US



National Science Foundation Bio-complexity Program and the University of California Agricultural Experiment Station. KEA was funded in part by a USDA CSREES Award No. 2005-35302-1699.

## Appendix A. Linear stability of the no-spatial model

For sake of simplicity, we rewrite the system (1) as following:

$$\frac{\partial I}{\partial t} = F(I, H), \quad (\text{A.1a})$$

$$\frac{\partial H}{\partial t} = G(I, H) + \frac{\partial^2 P}{\partial x^2}. \quad (\text{A.1b})$$

Denote the constant equilibrium solution as  $(I^*, H^*)$ , i.e.,

$$F(I^*, H^*) = 0, \quad G(I^*, H^*) = 0. \quad (\text{A.2})$$

We are aim to look for the conditions so that  $(I^*, H^*)$  is stable for the no-spatial version of (1) and is unstable for the spatial version of (1). We always assume that  $(I^*, H^*)$  is linearly stable with respect to perturbation of  $I$  and  $H$ , then the eigenvalues of the Jacobian

$$J = \begin{pmatrix} \frac{\partial F}{\partial I} & \frac{\partial F}{\partial H} \\ \frac{\partial G}{\partial I} & \frac{\partial G}{\partial H} \end{pmatrix} \triangleq \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix}$$

at  $(I^*, H^*)$  must have negative real parts, which is equivalent to

$$\text{tr}(J) = a_{11} + a_{22} < 0, \quad \text{Det}(J) = a_{11}a_{22} - a_{12}a_{21} > 0. \quad (\text{A.3})$$

## Appendix B. Dispersal relation

The main goal is to study the instability for the system (1) with the nonlinear diffusion. To deal with the nonlinear terms, we use the method of transformation variable. Define the new variables as  $P = P(I, H)$ . Then we have the following system with three equations:

$$\frac{\partial I}{\partial t} = F(I, H), \quad (\text{B.1a})$$

$$\frac{\partial H}{\partial t} = G(I, H) + \frac{\partial^2 P}{\partial x^2}, \quad (\text{B.1b})$$

$$P = P(I, H). \quad (\text{B.1c})$$

It is easy to see that the instability of  $(I^*, H^*)$  in the system (1) is equivalent to that of  $(I^*, H^*, P^*)$  in the system (B.1) where  $P^* = P(I^*, H^*)$ . Let  $u = I - I^*$ ,  $h = H - H^*$  and  $p = P - P^*$  be a spatial perturbation at the equilibrium  $(I^*, H^*, P^*)$ , then linearizing the reaction-diffusion system (B.1) at  $(I^*, H^*, P^*)$  yields

$$\frac{\partial u}{\partial t} = a_{11}u(x, t) + a_{12}h(x, t - \tau), \quad (\text{B.2a})$$

$$\frac{\partial h}{\partial t} = a_{21}u(x, t) + a_{22}h(x, t) + \frac{\partial^2 p}{\partial x^2}, \quad (\text{B.2b})$$

$$p(x, t) = a_{31}u(x, t) + a_{32}h(x, t), \quad (\text{B.2c})$$

where

$$a_{31} = \frac{\partial P(I, H)}{\partial I} \Big|_{(I^*, H^*)}, \quad a_{32} = \frac{\partial P(I, H)}{\partial H} \Big|_{(I^*, H^*)}. \quad (\text{B.3})$$

In order to examine the linear stability of  $(I^*, H^*, P^*)$  of system (B.1), we linearize the dynamic model (B.2) around the spatially homogeneous fixed point  $(0, 0, 0)$  for small space- and time-dependent fluctuations and expand them in Fourier space:

$$(u, h, p)^T = (A_1, A_2, A_3)^T e^{\lambda t + i k x}, \quad (\text{B.4})$$

which yields

$$A_1 \lambda e^{\lambda t + i k x} = (A_1 a_{11} + A_2 a_{12} e^{-\lambda \tau}) e^{\lambda t + i k x}, \quad (\text{B.5a})$$

$$(A_2\lambda + A_3\kappa^2)e^{\lambda t + i\kappa r} = (A_1a_{21} + A_2a_{22})e^{\lambda t + i\kappa r}, \quad (\text{B.5b})$$

$$A_3e^{\lambda t + i\kappa r} = (A_1a_{31} + A_2a_{32})e^{\lambda t + i\kappa r}. \quad (\text{B.5c})$$

Since  $e^{\lambda t + i\kappa r} \neq 0$ , (B.5) is equivalent to the following linear algebraic equations

$$\begin{pmatrix} \lambda - a_{11} & -a_{12}e^{-\lambda\tau} & 0 \\ -a_{21} & \lambda - a_{22} & \kappa^2 \\ -a_{31} & -a_{32} & 1 \end{pmatrix} \begin{pmatrix} A_1 \\ A_2 \\ A_3 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}. \quad (\text{B.6})$$

Nontrivial solutions to the above equations (B.6) exist if

$$\text{Det} \begin{pmatrix} \lambda - a_{11} & -a_{12}e^{-\lambda\tau} & 0 \\ -a_{21} & \lambda - a_{22} & \kappa^2 \\ -a_{31} & -a_{32} & 1 \end{pmatrix} = 0, \quad (\text{B.7})$$

which equals that

$$\text{Det} \begin{pmatrix} \lambda - a_{11} & -a_{12}e^{-\lambda\tau} \\ -a_{21} + a_{31}\kappa^2 & \lambda - a_{22} + a_{32}\kappa^2 \end{pmatrix} = 0. \quad (\text{B.8})$$

Note that (B.8) can be solved, yielding the characteristic polynomial of the original problem:

$$\lambda^2 - (a_{11} + a_{22} - a_{32}\kappa^2)\lambda + a_{11}a_{22} - a_{12}a_{21}e^{-\lambda\tau} - a_{11}a_{32}\kappa^2 + a_{12}a_{31}e^{-\lambda\tau}\kappa^2 = 0. \quad (\text{B.9})$$

### Appendix C. Critical value of time delay

We firstly pay attention to the model without spatial terms. Thus, the corresponding characteristic equation is that:

$$\lambda^2 - (a_{11} + a_{22})\lambda + a_{11}a_{22} - a_{12}a_{21}e^{-\lambda\tau} = 0. \quad (\text{C.1})$$

We know that  $i\omega$  ( $\omega > 0$ ) is a root of Eq. (C.1) if and only if  $\omega$  satisfies the following equation:

$$-\omega^2 - (a_{11} + a_{22})i\omega + a_{11}a_{22} - a_{12}a_{21}(\cos \omega\tau - i \sin \omega\tau) = 0. \quad (\text{C.2})$$

Separating the real and imaginary parts, one can have

$$(a_{11} + a_{22})\omega = a_{12}a_{21} \sin \omega\tau, \quad (\text{C.3a})$$

$$-\omega^2 + a_{11}a_{22} = a_{12}a_{21} \cos \omega\tau. \quad (\text{C.3b})$$

By squaring and adding the two parts of Eq. (C.3), it follows that

$$\omega^4 + (a_{11}^2 + a_{22}^2)\omega^2 + a_{11}^2a_{22}^2 - a_{12}^2a_{21}^2 = 0. \quad (\text{C.4})$$

Suppose that  $a_{11}a_{22} > a_{12}a_{21}$ , then (C.4) have the solution:

$$\omega^2 = \frac{-(a_{11}^2 + a_{22}^2) + \sqrt{(a_{11}^2 + a_{22}^2)^2 - 4(a_{11}^2a_{22}^2 - a_{12}^2a_{21}^2)}}{2} \triangleq \eta. \quad (\text{C.5})$$

If we define

$$\tau_j = \frac{1}{\sqrt{\eta}} \left[ \arccos \frac{-\eta + a_{11}a_{22}}{a_{12}a_{21}} + 2j\pi \right], \quad j = 0, 1, 2, \dots, n, \quad (\text{C.6})$$

then Eq. (C.1) with  $\tau = \tau_j$  has a pair of purely imaginary roots  $\pm i\omega$ , and it is easy to see that  $\tau_0 < \tau_1 < \dots < \tau_n$ .

It can be found that (C.1) that

$$\left( \frac{d\lambda}{dt} \right)^{-1} = \frac{a_{11} + a_{22} - 2\lambda}{a_{12}a_{21}\lambda e^{-\lambda\tau}} - \frac{\tau}{\lambda} \quad (\text{C.7})$$

and thus

$$\text{Re} \left( \frac{d\lambda}{dt} \right)^{-1} \Big|_{\tau=\tau_j} = \frac{a_{11}^2 + a_{22}^2 + 2\omega^2}{a_{12}^2a_{21}^2} > 0. \quad (\text{C.8})$$

In other words, for the model (1) without spatial terms, the Hopf bifurcation will occur at  $\tau = \tau_j$ , and period solution will emerge when  $\tau > \tau_j$ .

## References

- [1] Abbott KC, Dwyer G. Food limitation and insect outbreaks: complex dynamics in plant-herbivore models. *J Anim Ecol* 2007;76:1004–14.
- [2] Abbott KC, Morris WF, Gross K. Simultaneous effects of food limitation and inducible resistance on herbivore populations dynamics. *Theor Popul Biol* 2008;73:63–78.
- [3] Andrewartha HG, Birch LC. The distribution and abundance of animals. Chicago, IL, USA: University of Chicago Press; 1954.
- [4] Bascompte J, Rodriguez MA. Self-disturbance as a source of spatiotemporal heterogeneity. *J Theor Biol* 1998;195:383–93.
- [5] Bjrnstad ON, Sait SM, Stenseth NC, Thompson DJ, Begon M. The impact of specialized enemies on the dimensionality of host dynamics. *Nature* 2001;409:1001–6.
- [6] Busenberg SN, Velasco-Hernandez JX. Habitat suitability and herbivore dynamics. *Biosystems* 1994;32:37–47.
- [7] Bylund H. Long-term interactions between the autumnal moth and mountain birch: the roles of resources, competitors, natural enemies and weather [Ph.D. thesis]. Swedish Univ. Agr. Sci., Uppsala; 1995, p. 1–116.
- [8] Cappuccino N, Price PW. Population dynamics: new approaches and synthesis. New York: Academic Press; 1995.
- [9] Coulson RN, Witter JA. Forest entomology: ecology and management. NY: Wiley; 1984.
- [10] Das K, Sarkar AK. Stability and oscillations of an autotroph-herbivore model with time delay. *Int J Sys Sci* 2001;32(5):585–90.
- [11] DeBach P. The role of weather and entomophagous species in the natural control of insect populations. *J Econ Entomol* 1958;51:474–84.
- [12] Dempster JP. The natural control of populations of butterflies and moths. *Biol Rev Cambridge Philos Soc* 1983;58:461–81.
- [13] Dwyer G, Dushoff J, Yee SH. The combined effects of pathogens and predators on insect outbreaks. *Oecologia* 2004;430:341–5.
- [14] Edelstein-Keshet L, Rausher MD. The effects of inducible plant defenses on herbivore populations. 1. mobile herbivores in continuous time. *Am Nat* 1989;133:787–810.
- [15] Elton CS. Periodic fluctuations in the number of animals: their causes and effects. *Br J Exp Biol* 1924;2:119–63.
- [16] Freedman HI, Rao VSH. The trade-off between mutual interaction and time lag in predator-prey system. *Bull Math Biol* 1983;43:991–1004.
- [17] Ginzburg LR, Taneyhill DE. Population cycles of forest lepidoptera: a maternal effect hypothesis. *J Anim Ecol* 1994;63:79–92.
- [18] Gourtley S, Kuang Y. A stage structured predator-prey model and its dependence on through-stage delay and death rate. *J Math Biol* 2004;49:188–200.
- [19] Gripenberg S. Spatial ecology of a specialist insect herbivore: the leaf-mining moth *Tischeria ekebladella* on the pedunculate oak *Quercus robur* [Ph.D. thesis]; 2007.
- [20] Guttal V, Jayaprakash C. Impact of noise on bistable ecological systems. *Ecol Model* 2007;201:420–8.
- [21] Hanski I, Hansson L, Henttonen H. Specialist predators, generalist predators, and the microtine rodent cycle. *J Anim Ecol* 1991;60:353–67.
- [22] Haukioja E. On the role of plant defenses in the fluctuation of herbivore populations. *Oikos* 1980;35:202–13.
- [23] Haukioja E, Neuvonen S. Insect population dynamics and induction of plant resistance: the testing of hypotheses. In: Barbosa P, Schultz JC, editors. *Insect outbreaks*. San Diego, CA: Academic Press; 1987.
- [24] Haukioja E, Pakarinen E, Niemela P, Iso-livari L. Crowding-triggered phenotypic responses alleviate consequences of crowding in *epirrita autumnata* [lep., geometridae]. *Oecologia* 1988;75:549–58.
- [25] Hawkins BA, Holyoak M. Transcontinental crashes of insect populations? *Am Nat* 1998;152:480–4.
- [26] Hunter AF. Traits that distinguish outbreaking and nonoutbreaking macrolepidoptera feeding on northern hardwood trees. *Oikos* 1991;60:275–82.
- [27] Huygens CH. *Horologioium oscilatorium*. Parisis, France; 1673.
- [28] Itamies J, Pulliainen E, Siekkinen J. Mobility of *epirrita autumnata* (lepidoptera, geometridae), as indicated by altitudinal light trap sampling in finnish forest lapland. *Oecologia Montana* 1995;4:27–30.
- [29] Kaplan I, Denno RF. Interspecific interactions in phytophagous insects revisited: a quantitative assessment of competition theory. *Ecol Lett* 2007;10:977–94.
- [30] Lawton JH, McNeill S. Between the devil and the deep blue sea: on the problem of being a herbivore. In: *Population dynamics*. Oxford UK: Blackwell Scientific; 1979.
- [31] Levy J, III JG, Scott M, Avioli L. Mineral homeostasis in neonates of streptozotocin-induced noninsulin-dependent diabetic rats and in their mothers during pregnancy and lactation. *Bone* 1987;8(1):1–6.
- [32] Lewis MA. Spatial coupling of plant and herbivore coupling: the contribution of herbivore dispersal to transient and persistent waves of damage. *Theor Popul Biol* 1994;45(3):277–312.
- [33] Liebhold A, Koenig WD, Bjrnstad ON. Spatial synchrony in population dynamics. *Ann Rev Ecol Evol Syst* 2004;35:467–90.
- [34] Liebhold AM, Elkinton JS, Muzika RM. What causes outbreaks of the gypsy moth in north america? *Popul Ecol* 2000;42:257–66.
- [35] Liebhold AM, Kamata N. Are population cycles and spatial synchrony a universal characteristic of forest insect populations? *Popul Ecol* 2000;42:205–9.
- [36] Liebhold AM, Kamata N, Jacobs T. Cyclicity and synchrony of historical outbreaks of the beech caterpillar. *quadricalcarifera punctatella* (motschulsky) in japan. *Res Popul Ecol* 1996;38:87–94.
- [37] Lundberg S, Jaremo J, Nilsson P. Herbivory, inducible defence and population oscillations: a preliminary theoretical analysis. *Oikos* 1994;71:537–40.
- [38] MacDonald N. Time lags in biological models. Lecture notes in biomathematics. Berlin: Springer-Verlag; 1985.
- [39] MacLean DA. Effects of spruce budworm outbreaks on the productivity and stability of balsam fir forests. *Forestry Chronicle* 1984;60:273–9.
- [40] Maron JL, Harrison S, Greaves M. Origin of an insect outbreak: escape in space or time from natural enemies? *Oecologia* 2001;126:595–602.
- [41] McCann K, Hastings A, Harrison S, Wilson WG. Population outbreaks in a discrete world. *Theor Popul Biol* 2000;57:97–108.
- [42] Miller WE, Epstein ME. Synchronous population fluctuations among moth species. *Environ Entomol* 1986;15:443–7.
- [43] Milne A. Theories of natural control of insect populations. *Cold Spring Harbor Symp Quant Biol* 1957;22:253–71.
- [44] Morris WF, Dwyer G. Population consequences of constitutive and inducible plant resistance: herbivore spatial spread. *Am Nat* 1997;149(6):1071–90.
- [45] Myers JH. Population cycles of western tent caterpillars: experimental introductions and synchrony of fluctuations. *Ecology* 1990;71:986–95.
- [46] Myers JH. Synchrony in outbreaks of forest lepidoptera: a possible example of the moran effect. *Ecology* 1998;79:1111–7.
- [47] Neuvonen S, Haukioja E. The effects of inducible resistance in host foliage on birch-feeding insects. In: Raupp M, Tallamy D, editors. *Phytochemical induction by herbivore*. New York: John Wiley and Sons; 1991. p. 277–91.
- [48] Neuvonen S, Ruohomki K, Bylund H, Kaitaniemi P. Insect herbivores and herbivory effects on mountain birch dynamics. In: Wielgolaski FE, editor, *Nordic mountain birch ecosystems, Man and biosphere series*, vol. 27; 2001. p. 207–222.
- [49] Nunney L. The effect of long time delays in predator-prey systems. *Theor Popul Biol* 1985;27:202–21.
- [50] Ohmart CP, Stewart LG, Thomas JR. Leaf consumption by insects in three eucalyptus forest types in southeastern australia and their role in short term nutrient cycling. *Oecologia* 1983;59:322–30.
- [51] Ortega-Cejas V, Fort J, Mendez V. The role of the delay time in the modeling of biological range expansions. *Ecology* 2004;85:258–64.
- [52] Ovaskainen O, Sato K, Bascompte J, Hanski I. Metapopulation models for extinction threshold in spatially correlated landscapes. *J Theor Biol* 2002;215:95–108.
- [53] Post E. Large-scale spatial gradients in herbivore population dynamics. *Ecology* 2005;86(9):2320–8.
- [54] Post E, Forchhammer MC. Synchronization of animal population dynamics by largescale climate. *Nature* 2002;420:168–71.
- [55] Raimondo S, Liebhold AM, Strazanac JS, Butler L. Population synchrony within and among lepidoptera species in relation to weather, phylogeny, and larval phenology. *Environ Entomol* 2004;29:96–105.
- [56] Root RB, Cappuccino N. Patterns in population change and the organization of the insect community associated with goldenrod. *Ecol Monogr* 1992;62:393–420.
- [57] Royama T. *Analytical population dynamics*. London, UK: Chapman and Hall; 1992.
- [58] Schowalter T, Hargrove W, Crossley D. Herbivory in forested ecosystems. *Ann Rev Entomol* 1986;31:177–96.

- [59] Shepherd RF, Bennett DD, Dale JW, Tunnock S, Dolph RE, Thier RW. Evidence of synchronized cycles in outbreak patterns of douglas-fir tussock moth, *orgyia pseudotsugata*. *Mem Entomol Soc Can* 1988;146:107–21.
- [60] Sherratt JA, Lambin X, Thomas CJ, Sherratt TN. Generation of periodic waves by landscape features in cyclic predator–prey systems. *Proc R Soc Biol Sci* 2002;269:327–34.
- [61] Sherratt JA, Lord GJ. Nonlinear dynamics and pattern bifurcations in a model for vegetation stripes in semi-arid environments. *Theor Popul Biol* 2007;71:1–11.
- [62] Southwood TRE, Comins HN. A synoptic population model. *J Anim Ecol* 1976;45:949–65.
- [63] Sun G-Q, Jin Z, Liu Q-X, Li L. Dynamical complexity of a spatial predator–prey model with migration. *Ecol Model* 2008;219:248–55.
- [64] Tenou O. The outbreaks of *opornia autumnata* (bkh.) and *operophthera* spp. (lep., geometridae) in the scandinavian mountain chain and northern finland 1862–1968. *Zool Bidr Uppsala Suppl* 1972;2:1–107.
- [65] Underwood N. The influence of plant and herbivore characteristics on the interaction between induced resistance and herbivore population dynamics. *Am Nat* 1999;153:282–94.
- [66] Underwood N. The influence of plant and herivore characteristics on the interaction between induced resistance and herbivore population dynamics. *Am Nat* 1999;153(3):282–94.
- [67] Underwood N, Anderson K, Inouye BD. Induced vs. constitutive resistance and the spatial distribution of herbivores among plants. *Ecology* 2005;86(3):594–602.
- [68] Underwood NC. The timing of induced resistance and induced susceptibility in the soybean-mexican bean beetle system. *Oecologia* 1998;114:376–81.
- [69] Veblen T, Hadley K, Nel E, Kitzberger T, Reid M, Villalba R. Disturbance regime and disturbance interactions in a rocky mountain subalpine forest. *J Ecol* 1994;82:125–35.
- [70] Wang R-H, Liu Q-X, Sun G-Q, Jin Z, van de Koppel J. Nonlinear dynamic and pattern bifurcations in a model for spatial patterns in young mussel beds. *JR Soc Interface* 2009;6(37):705–18.
- [71] White TCR. The abundance of invertebrate herbivores in relation to the availability of nitrogen in stressed food plants. *Oecologia* 1984;63:90–105.