

UCLA

UCLA Previously Published Works

Title

Salivary cortisol awakening levels are reduced in human subjects with intermittent explosive disorder compared with controls.

Permalink

<https://escholarship.org/uc/item/80k09131>

Authors

Meruelo, Alejandro

Timmins, Matthew

Irwin, Michael

et al.

Publication Date

2023-05-01

DOI

10.1016/j.psyneuen.2023.106070

Peer reviewed



Published in final edited form as:

Psychoneuroendocrinology. 2023 May ; 151: 106070. doi:10.1016/j.psyneuen.2023.106070.

SALIVARY CORTISOL AWAKENING LEVELS ARE REDUCED IN HUMAN SUBJECTS WITH INTERMITTENT EXPLOSIVE DISORDER COMPARED WITH CONTROLS

Alejandro D. Meruelo, M.D., Ph.D.⁽¹⁾, Matthew A. Timmins, Ph.D.⁽²⁾, Michael R. Irwin, M.D.⁽³⁾, Emil F. Coccaro, M.D.⁽²⁾

¹Department of Psychiatry, University of California, San Diego, San Diego, CA.

²Clinical Neuroscience and Psychotherapeutics Research Unit, Department of Psychiatry and Behavioral Health, The Ohio State University Wexner Medical Center, Columbus, OH.

³Department of Psychiatry & Biobehavioral Sciences and Medicine, Norman Cousins Center, and Semel Institute, UCLA School of Medicine, Los Angeles, CA.

Abstract

Background.—The role of the hypothalamic-pituitary-adrenal (HPA) axis in human aggressive behavior is poorly characterized, though some studies report that, unlike depression, circulating or salivary levels of cortisol are low compared with controls.

Methods.—In this study, we collected three salivary cortisol levels (two in the morning and one in the evening) on three separate days in 78 adult study participants with ($n = 28$) and without ($n = 52$) prominent histories of impulsive aggressive behavior. Plasma C-Reactive Protein (CRP) and Interleukin-6 (IL-6) were also collected in most study participants. Aggressive study participants meet DSM-5 criteria for Intermittent Explosive Disorder (IED) while non-aggressive participants either had a history of a psychiatric disorder or no such history (Controls).

Results.—Morning, but not evening, salivary cortisol levels were significantly lower in IED ($p < 0.05$), compared with control, study participants. In addition, salivary cortisol levels correlated with measures of trait anger (partial $r = -0.26$, $p < 0.05$) and aggression (partial $r = -0.25$, $p < 0.05$) but not with measures of impulsivity, psychopathy, depression, history of childhood maltreatment, or other tested variables that often differ in individuals with IED. Finally, plasma CRP levels correlated inversely with morning salivary cortisol levels (partial $r = -0.28$, $p < 0.05$);

Address Correspondence to: Emil F. Coccaro, M.D., Clinical Neuroscience and Psychotherapeutics Research Unit, Department of Psychiatry and Behavioral Neuroscience, The Ohio State University Wexner Medical Center, 1640 Upham Drive, Columbus, OH 43201, (773) 852-1338, emil.coccaro@osumc.edu.

AUTHORSHIP CONTRIBUTION STATEMENT

Meruelo: Writing

Timmins: Writing – review and editing.

Irwin: Plasma CRP/IL-6 assays, Writing – review and editing.

Coccaro: Conceptualization, Funding acquisition, Project management, Data collection, Statistical analysis, Writing – review and editing.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

plasma IL-6 levels showed a similar, though not statistically significant ($r_p = -0.20$, $p = 0.12$) relationship with morning salivary cortisol levels.

Conclusion.—The cortisol awakening response appears to be lower in individuals with IED compared with controls. In all study participants, morning salivary cortisol levels correlated inversely with trait anger, trait aggression, and plasma CRP, a marker of systemic inflammation. This suggests the present of a complex interaction between chronic-low level inflammation, the HPA axis, and IED that warrants further investigation.

Keywords

IED; Cortisol; C-Reactive Protein; Aggression; Impulsivity

1. INTRODUCTION

Intermittent Explosive Disorder (IED) (Coccaro and Lee, 2020) is a behavioral disorder with wide impact on both the afflicted individual and our communities that occurs, over the lifetime, in up to four percent in the U.S. population. IED is characterized by impulsive and aggressive verbal or physical outbursts that are disproportional to a preceding stressor or trigger (American Psychiatric Association, 2013) and that are associated with distress or impairment in social or work-related function. For example, untreated IED carries the risks of higher job loss, divorce, incarceration, and suicide (Coker et al., 2014; DeLisi et al., 2017; McLaughlin et al., 2012; Mundt et al., 2013; Scott et al., 2007; Smith et al., 2021).

IED, and impulsive aggression, is also characterized by a variety of neurobiological features including low serotonin (5-HT) function (Coccaro et al., 2009), reduced grey matter volume in cortico-limbic circuits (Coccaro et al., 2016), increased amygdala reactivity to social threat (Coccaro et al., 2007; McCloskey et al., 2016), compared with non-aggressive controls, all of which are influenced by the hypothalamic-pituitary-adrenal (HPA) axis. Specifically, brain 5-HT turnover (Heinz et al., 2002) and brain grey matter volume (Echouffo-Tcheugui et al., 2018) exhibit inverse relationships with endogenous cortisol. In addition, endogenous cortisol responses to emotional stimuli have been associated with an inverse relationship with ventro-medial pre-frontal and a direct relationship with amygdala, activity (Root et al., 2009), similar to what has been reported in fMRI studies in IED (Coccaro et al., 2007; McCloskey et al., 2016). These findings suggest a role for circulating cortisol in aggression but one that is still to be clarified. For example, the inverse relationship between cortisol and 5-HT turnover and grey matter volume suggests that cortisol should be high in those with IED while the positive relationship between cortisol and amygdala reactivity to social threat suggests that cortisol should also be high in those with IED. In individuals with aggressive behavior (Böhnke et al., 2010; Raine, 2002), however, cortisol appears to have an inverse relationship with aggression. Accordingly, it is of interest to explore the relationship between the HPA axis in those with IED compared with non-aggressive controls.

Levels of circulating cortisol typically follow a diurnal cycle (Van Cauter et al., 1996). In addition, cortisol is released in response to acute environmental stressors (Del Giudice et al., 2011). Cortisol levels typically reach maximum levels within the first 30 minutes of

waking (defined as the cortisol awakening response) and proceed to fall over the course of the day before reaching a low point in the evening (Van Cauter et al., 1996). A healthy HPA axis provides protective and adaptive responses including maintenance of energy production, memory retention, and our immune systems (McEwen and Seeman, 1999).

While high cortisol activity is often associated with depressive mood and depressive disorders (Jurueña et al., 2018), low cortisol activity has been linked to aggressive and violent behaviors (Böhnke et al., 2010; McBurnett et al., 2000; Raine, 2002). Lower morning cortisol activity, in the form of a reduction of the cortisol awakening response, has been observed in a number of clinical populations including those with Oppositional Defiant Disorder (Freitag et al., 2009), Conduct Disorder (Pajer et al., 2001), and those with psychopathic traits (Cima et al., 2008; Holi et al., 2006; O'Leary et al., 2006), disorders that include aggressive behavior but not necessarily impulsive aggressive behavior. In addition, a recent meta-analysis reported that nine of eleven studies conducted in adolescents found an inverse relationship between morning cortisol levels and "aggression" (Blankenstein et al., 2022). The measures used in these studies, primarily, was the "Disruptive or Aggressive Behavior" subscale from the Young Mania Rating Scale (Young et al., 1978). While IED is the prototypical disorder of impulsive aggression, to our knowledge, no work on the function of the HPA axis has been published in individuals with IED and few studies have prospectively used measures designed primarily to assess aggression.

In order to fill this gap, we measured morning and evening salivary cortisol levels in study participants with IED and in non-aggressive human control participants. Note that salivary cortisol reflects free cortisol, the active component most relevant to effects on brain (Hellhammer et al., 2009). Based on prior research (Blankenstein et al., 2022; Cima et al., 2008; Freitag et al., 2009; Holi et al., 2006; O'Leary et al., 2006; Pajer et al., 2001), we hypothesized that morning cortisol levels would be lower in study participants with IED compared to non-aggressive controls. In addition, we looked at the effect of basal inflammatory state on the cortisol awakening response given the inverse relationship between cortisol and inflammation (Silverman 2012), and reports of chronic low-level inflammation in individuals with aggression (Coccaro 2006; Marsland et al., 2008; Suarez, 2003, 2004) and/or IED (Coccaro et al., 2014). We hypothesized that there would be an inverse relationship between salivary cortisol levels and basal inflammatory state as manifest by circulating levels of C-Reactive Protein (CRP) and/or Interleukin-6 (IL-6).

2. MATERIALS AND METHODS

2.1 Participants.

Ninety physically healthy adult individuals of both sexes participated in this study from December 2015 through November 2019. Participants were recruited from a larger study of inflammatory markers and aggression involving a total of 135 adult individuals. Participants were recruited through public service announcements, newspaper, other media, and advertisements seeking out individuals who reported psychosocial difficulty related to impulsive aggression or to one or more psychiatric conditions, as well as healthy controls. All participants gave informed consent and signed the informed consent document approved

by The University of Chicago's Institutional Review Board. Study participants in this project were not the same as those in our previous studies (Coccaro, 2006; Coccaro et al., 2014).

2.2 Assessment

Diagnoses were made according to DSM-5 criteria (American Psychiatric Association 2013). Assessments were completed using information from: (a) the Structured Clinical Interview for DSM Diagnoses (SCID-I; First et al., 1997); (b) the Structured Interview for the Diagnosis of Personality Disorder (SIDP; Pfohl et al., 1997); (c) clinical interview by a research psychiatrist; and, (d) review of all other available clinical data. Research diagnostic interviews were conducted by individuals with a masters or doctorate degree in Clinical Psychology. All diagnostic raters went through a rigorous training program that included: lectures on DSM diagnoses and rating systems; videos of expert raters conducting SCID/SIDP interviews; and practice interviews/ratings until the raters were deemed reliable by the trainer. This process resulted in good to excellent inter-rater reliabilities (kappa of .84 (\pm) .05; range: .79 to .93) across the diagnosis of anxiety, mood, substance use, impulse control, and personality disorders. Final diagnoses were determined by team-based best-estimate consensus procedures involving research psychiatrists and clinical psychologists (Coccaro et al., 2012). While information for assigning syndromal (formally Axis I) diagnoses were collected through the use of the SCID-I, more than sufficient information was available to update these diagnoses from DSM-IV to those of DSM-5. DSM-5 diagnoses for personality disorders, based on the SIDP, are the same for DSM-IV. Participants with a diagnosis mutually exclusive to the diagnosis of IED were excluded from the study. Exclusion criteria included active substance use disorders, a lifetime history of bipolar disorder, schizophrenia (or other psychotic disorder), and those with an intellectual disability. Data for age, sex assigned at birth, and ethnicity were self-reported by the study participant; socio-economic status was estimated by the method of Hollingshead (Hollingshead, 1975).

Twenty-nine participants had no evidence of any past or present psychiatric diagnosis including personality disorder (Healthy Controls: HC); 29 met criteria for a lifetime diagnosis of a syndromal psychiatric and/or a personality disorder (Psychiatric Controls: PC), and 32 met criteria for Intermittent Explosive Disorder (IED). Of the 61 participants assessed with a psychiatric disorder, most (68.6%) reported history of formal psychiatric evaluation and/or treatment (51.4%) or of behavioral disturbance during which the participant, or others, thought they should have sought mental health services but did not (17.1%). Despite this history, no study participant was engaged in any psychologic, or psychiatric, treatment (i.e., medication-free) at the time of entry into, or during the duration of, the study.

2.3 Medical Clearance.

After signed informed consent, study participants were medically evaluated by medical history, physical examination, standard laboratory studies (e.g., hematology, chemistry, etc.) and ECG. In addition, this assessment confirmed that study participants had no signs of physical trauma, or skin rash, and had not received any vaccinations in the previous six months.

2.4 Collection, Processing, and Assays of Salivary Cortisol.

After signed informed consent, study participants were given a home salivary collection kit (Salimetrics, USA) containing nine tubes for collection of saliva for three days. Each tube included an outer tube, inner tube, absorption swab, and cap. Participants were instructed to collect saliva samples in the morning, immediately after waking up, 30 minutes after waking up, and in the evening before they went to sleep. Participants were also instructed not to take anything by mouth for 30 minutes before each sample, to write down the time and date for each sample, and to place the sample tube in their home refrigerator immediately after sample collection. Specific instructions included: a) removing the cap from the inner tube (without removing the inner tube from the outer tube, b) guiding the tube and swab to their mouth, c) chew on the absorption swab (without touching the absorption swab with their hands) for one minute at which point the swab is saturated with saliva and, d) spitting the absorption swab into the inner tube, then sealing the tube with the cap and placing it in a zip-locked bag in their home refrigerator. After the three samples for the day were collected, participants were instructed to take the samples from their refrigerator and place the tubes in the package provided to send the samples overnight (via FEDEX at room temperature) to our laboratory. Upon receipt, each tube was centrifuged for 15 minutes and the saliva supernatant was pipetted into a polypropylene tube and frozen at -80° Celsius until assay at the University of Chicago Clinical Research Center (CRC) core laboratory. Samples were assayed for unbound cortisol, in duplicate, using the Salimetrics HS Salivary Cortisol EIA Kit. This assay had a lower limit of sensitivity from 0.007 to 1.2 $\mu\text{g}/\text{dL}$, with a mean inter-assay CV between 3.9% and 7.1%, and mean inter-assay CV between 6.7% and 6.9%, for high and low concentrations, respectively. Twelve study participants (13.3%) did not provide complete samples for each of the three days and were excluded from analysis. The proportion excluded did not differ among the groups [HC: 3 of 29 (10.3%); PC: 4 of 30 (13.3%); IED: 5 of 31 (16.1%); Chi-Square = 0.43, $df = 2$, $p = 0.81$]. This left full three-day salivary cortisol data from 26 participants in each group. Excluded participants did not differ from included participants in mean age, distribution sex or ethnicity, or in mean Hollingshead Socio-Economic Status.

2.5 Collection, Processing, and Assays of Plasma Inflammatory Markers.

Finally, as part of this protocol, most (76%: 59 of 78) participants had three samples for plasma C-Reactive Protein (CRP) and Interleukin-6 (IL-6) collected during the overall time of collection of the salivary cortisol samples. These samples were collected by venipuncture, between 9 and 11 am, into EDTA tubes, placed on ice, centrifuged for plasma, and stored at -80° C until assayed. Inflammatory marker samples were assayed in the laboratory of one of the co-authors (MRI) and levels reported represent the mean of the duplicates. Plasma CRP was assayed using a high-sensitivity ELISA assay by R&D Systems (Minneapolis, MN) and plasma IL-6 was assayed using a multiplex assay by Meso Scale Discovery (Rockville, Maryland). The sensitivity of the assays for CRP and IL-6 were 0.20 mg/L and 0.21 pg/ml, respectively. Mean intra- and inter- assay CVs for CRP were 1.0% and 6.1%; mean intra- and inter- assay CVs for IL-6 were 1.0% and 4.0%.

2.6 Assessment of Aggression and Anger.

Aggression was assessed with the aggression scale of the Life History of Aggression (LHA; Coccaro et al., 1997). The LHA assesses history of overt aggressive behavior and is conducted as a semi-structured interview and has good psychometric properties ($\alpha = 0.86$). Anger was assessed using the Anger Expression Index (AEI) score from the Spielberger State-Trait Anger and Expression of Anger Inventory (STAXI-2; (Spielberger 1999). The AEI is a 32 item self-report assessment that considers anger expression ($\alpha = 0.89$) and anger control ($\alpha = 0.94$) in one variable with higher numbers signifying greater anger.

2.7 Assessment of Aggression Related Variables: Impulsivity and Psychopathy.

Impulsivity was assessed with the Barratt Impulsiveness Scale (BIS-11; Patton et al., 1995). The BIS-11 assesses the tendency to be impulsive and has good psychometric properties ($\alpha = 0.86$). Psychopathy as assessed with the Psychopathic Checklist-Screening Version (Hart et al., 2003) as previously described (Coccaro et al., 2014: $\alpha = 0.79$).

2.8 Assessment of State Depression, Perceived Stress, and History of Childhood Maltreatment.

State depression was assessed by the Beck Depression Inventory-II (BDI-II; Beck & Brown, 1996: $\alpha = 0.96$). Perceived stress over the past month was assessed by the Perceived Stress Scale (PSS; Cohen et al., 1983), a 10-item, Likert-scale questionnaire: $\alpha = 0.92$). History of childhood maltreatment was assessed by the Childhood Trauma Questionnaire (CTQ; Bernstein et al., 2003), a 28-item, Likert-scaled, questionnaire inquiring about physical, emotional, and sexual abuse as well as physical and emotional neglect: $\alpha = 0.87$). These variables were included because of their relevance to, and potential impact on, the HPA Axis (Counts, 2022; Nandam, 2019; Zorn, 2017).

2.9 Statistical Analysis and Data Reduction.

The primary outcome variables for categorical and dimensional analyses were the DSM-5 diagnosis of Intermittent Explosive Disorder, LHA Aggression, and STAXI-2 AEI Anger scores. Secondary outcome variables were BIS-11 Impulsivity and the PCL-SV Psychopathy scores. Other outcome variables included the BDI-II, PSS, and the CTQ. Statistical procedures included Chi-Square, t-test, Analyses of Covariance (ANCOVA), Repeated Analysis of Covariance (RM-ANCOVA), Pearson Correlation, Partial Correlation, and Multiple Regression. Analyses included age, sex assigned at birth, ethnicity, and socioeconomic status; body mass index was added as a covariate in analyses involving pro-inflammatory markers. Salivary cortisol levels (μdL), and CRP (mg/dL) / IL-6 (pg/ml) levels, were log-transformed because these variables were not normally distributed. A two-tailed alpha value of 0.05 was used to denote statistical significance. Statistical procedures were performed using SPSS 28.

3. RESULTS

3.1 Demographic Characteristics of Sample and Sub-Samples.

The 90 participants recruited for this study did not differ in demographic variables (age, sex assigned at birth, ethnicity, and socioeconomic status) from those remaining in the sample that participated in our larger study of inflammatory markers and aggression ($n = 135$; Supplemental Table 1a); there were also no differences in demographic variables between the 78 study participants in the final sample ($n = 78$) with complete salivary cortisol data and participants those excluded due to incomplete salivary cortisol data ($n = 12$; Supplementary Table 1b). Study participants with complete both salivary cortisol and inflammatory marker data ($n = 59$) also did not differ on demographic variables compared with those without inflammatory marker data ($n = 31$; Supplemental Table 1c).

3.2 Demographic, Psychometric, and Diagnostic Characteristics of the Final Sample.

Table 1 displays demographic, body mass index (BMI), and relevant psychometric variables for the final sample with full three-day salivary cortisol data. The groups did not differ in age, distribution of sex or ethnicity, or BMI, though IED participants had non-significantly lower SES scores compared with HC and PC controls (Table 1). As expected, the groups differed in the psychometric variables related to aggression, anger, impulsivity, and psychopathy, in which IED study participants displayed higher scores on each variable compared with HC and PC study participants. Variables related to depression, perceived stress, and history of childhood maltreatment were also elevated in IED study participants compared with both sets of controls. Despite this, PC and IED study participants did not meaningfully differ in current or lifetime DSM-5 disorders (Table 2).

3.3 Log Salivary Cortisol Levels in IED and Control Participants.

Repeated measures ANCOVA revealed that the two morning log salivary cortisol levels did not differ significantly by time ($F[1,71] = 0.85$, $p = 0.32$; Marginal Means + SEM: Time 1 = $1.36 + 0.02$; Time 2 = $1.41 + 0.03$) but did differ by HC/PC/IED diagnostic group ($F[1,71] = 3.11$, $p < 0.05$) with lower log morning salivary cortisol levels among participants with IED compared with participants in both control groups ($\eta^2 p = 0.081$), who did not differ in this respect. There was no effect of Age, Sex, Ethnicity, and SES, with morning log salivary cortisol levels, and no significant interactions among variables (all $p > 0.51$). For subsequent analyses, the two log morning cortisol levels were combined into one mean log morning cortisol level (figure 1, left). Log mean morning salivary cortisol levels did not differ as a function of Non-IED Disorders (e.g., Depressive Disorder, Anxiety Disorder, etc.). Finally, and in contrast, mean log evening (Time 3, at bedtime) salivary cortisol levels did not differ by group or as a function of any of the covariates (figure 1, right).

3.4 Log Salivary Morning Cortisol Levels and Aggression and Anger.

Partial correlation analyses revealed statistically significant inverse relationships between mean log salivary morning cortisol levels and LHA Aggression ($r = -0.25$, $p < 0.05$), and STAXI-2 Anger Expression Index ($r = -0.26$, $p < 0.05$), scores. The partial correlation using a Composite Anger/Aggression score was similar ($r = -0.24$, $p < 0.05$).

3.5 Log Salivary Morning Cortisol Levels and Impulsivity, Psychopathy, and Other Relevant Variables (Table 3).

Though correlated with LHA Aggression and/or STAXI Anger Expression Index scores, no statistically significant relationships were noted between mean log salivary morning cortisol levels and the other secondary behavioral outcome variables in this study. Adding these behavioral variables to the statistical model had no impact on the relationship between mean log salivary morning cortisol levels and Composite Anger/Aggression scores ($r_p = -0.29$, $p < 0.05$; i.e., with demographic variables and with BIS-11, PCL-SV, BDI-II, PSS, and CTQ scores as covariates); figure 2. Supplemental Table II lists all partial correlations between log salivary morning cortisol levels and all behavioral variables.

3.6 Mean Log Morning Salivary Cortisol Levels and Plasma Pro-Inflammatory Markers.

Partial correlation with demographic variables and body mass index also revealed a significant inverse relationship between mean log plasma CRP levels and mean log salivary morning cortisol levels ($r_p = -0.28$, $p < 0.05$). A similar inverse, but not statistically significant, correlation was also noted with mean log plasma IL-6 level and mean log salivary morning cortisol ($r_p = -0.20$, $p = 0.12$) levels. Multiple regression analysis with log plasma CRP levels and Composite Anger/Aggression scores as predictors of mean log salivary cortisol levels revealed a significant correlation with log plasma CRP level ($\beta = -0.31$, $p < 0.05$; figure 3) but a smaller, statistically non-significant, correlation with Composite Anger/Aggression scores ($\beta = -0.14$, $p = 0.31$).

4. DISCUSSION

We investigated the relationship between impulsive aggression and morning/evening salivary cortisol levels in males and females with Intermittent Explosive Disorder (IED) compared with both healthy and psychiatric controls. We also explored the presence of a relationship between salivary cortisol and markers of inflammation (CRP and IL-6) given the presence of an inverse relationship between cortisol and inflammation (Silverman 2012).

We found a reduction in morning, but not evening, cortisol levels in individuals with IED compared with all control subjects (who were no different from each other in salivary cortisol levels). Consistent with this finding we found statistically significant correlations between mean morning salivary cortisol level and trait anger expression and history of lifetime aggressive behavior. This relationship appeared to be specific to IED because similar analyses did not find significant differences in mean log morning cortisol level between those who had, or did not have, other Non-IED disorders.

The observation that other behavioral variables, such as impulsivity, depression, psychopathy, perceived stress, and history of childhood maltreatment did not correlate with mean morning salivary cortisol level suggests either a level of specificity for trait anger and aggression in this relationship or the presence of a Type I error. Supporting specificity for a relationship with trait anger and aggression are reports that show that anger induction reduces salivary cortisol levels in both healthy men (Herrero et al., 2010) and women

(Herrero et al., 2010; Kazén et al., 2012). To our knowledge, however, there are no reports regarding this dynamic in aggressive individuals.

The observation of lower morning cortisol levels in those with IED, and of an inverse relationship between morning cortisol levels and dimensional measures of trait anger and aggression, is consistent with several reports in the literature. Individuals with aggressive and violent behaviors (Böhnke, 2010; Raine, 2002) have been reported to exhibit lower cortisol activity as do individuals with other psychiatric disorders [e.g., Oppositional Defiant Disorder; (Freitag, 2009) and Conduct Disorder (Pajer, 2001)]. While studies of individuals with psychopathic traits (Cima et al., 2008; Holi et al., 2006; O’Leary et al., 2006) report the same, callous/unemotional psychopathic traits correlate highly with social deviant psychopathic traits, which are largely aggressive in nature (Coccaro et al., 2014). Thus, these findings do not necessarily contradict our observation of an inverse correlation between morning cortisol levels without a similar one for psychopathic traits. Finally, several studies of adolescents have reported inverse correlations between basal morning cortisol levels with secondary measures of “disruptive or aggressive behavior” (Blankenstein et al., 2022).

Lower morning cortisol activity is postulated to be a manifestation of blunted hypothalamic-pituitary axis (HPA) activity. Poor adaptation to chronic stress is thought to lead to flattening of diurnal rhythms of cortisol and subsequently attenuated typical morning rise of cortisol (Sephton et al., 2000). Adults who experienced early life difficulties have been posited to experience stress sensitivity that may reset the HPA axis to a blunted level of activity and responsivity (Miller et al., 2007). In addition, while the HPA axis of those with such stress sensitivity may be more responsive during childhood (e.g., in threatening family environments), such individuals may manifest attenuated HPA activity later during adulthood (Repetti et al., 2011). While we observed no dimensional relationship between morning cortisol levels and history of childhood maltreatment, or with perceived levels of current stress, study participants with IED were significantly higher than both sets of controls on these two variables.

The observation of an inverse relationship between plasma CRP and morning cortisol levels, pointing to a relationship between the HPA axis and basal inflammatory state is of note. In the homeostatic state, the HPA axis activates innate immunity and fosters the inhibition of proinflammatory mediators (Silverman 2012). Accordingly, this finding is consistent with the possibility that lower HPA activity is associated with higher levels of systemic inflammation. Further, we found that mean plasma CRP levels were a stronger correlate of morning salivary cortisol levels than the composite anger/aggression score. While this suggests that chronically elevated levels of inflammation may drive these relationships, this hypothesis requires further study. Despite this finding, it is important to note that only about half of study participants with IED had plasma CRP data and, while mean CRP levels in this group was higher compared with controls, this difference was not statistically significant. Accordingly, caution should be taken in the interpretation of these data.

Other explanations for these findings can be considered. For one, the acute induction of anger in healthy individuals is associated with a reduction in salivary cortisol levels that begins within 25 minutes of anger induction, persisting for at least a total of 45 minutes

(Kazén et al., 2012). Thus, it is possible that the frequent anger outbursts of those with IED is responsible for lower salivary cortisol levels in these individuals. Aggressive behavior is also related to serotonin (5-HT) and individuals with IED display evidence of reduced central serotonergic function (Coccaro et al., 2009; New et al., 2004; Siever et al., 1999). While 5-HT agents can acutely increase circulating cortisol levels (Coccaro and Kavoussi, 1994), the reduction of 5-HT through acute tryptophan depletion can reduce salivary cortisol levels (Sobczak et al., 2002). If so, low morning cortisol levels would be consistent with low 5-HT function, especially in those with IED (Coccaro et al., 2010).

Another mechanism underlying our findings may be through the relationship between anger and testosterone. This is because anger inducing events have been shown to be associated with increases in salivary testosterone (Peterson and Harmon-Jones, 2012) and testosterone and cortisol may be inversely related (Herrero et al., 2010). However, while there may be an inverse relationship between testosterone and 5-HT (Kindlundh et al., 2003), central levels of testosterone do not correlate with central levels of 5-HT metabolites in human (Virkkunen et al., 1994) and non-human primates (Higley et al., 1996), and high levels of aggression may be observed only in individuals with low central levels of 5-HT coupled with high central levels of testosterone (Higley, 1996; Virkkunen et al., 1994). That said, we did not measure testosterone in this study and, thus, these ideas regarding testosterone represent only speculation.

4.1 Strengths and Limitations.

This study has a number of strengths and limitations. A major strength was the use of male and female adults who were either healthy or struggled with impulsive aggressive behavior leading to distress and/or interpersonal difficulties. Our investigation of the LHA Aggression, STAXI-2 Anger Expression Index, BIS Impulsivity, BDI-II Depression, PCL-SV Psychopathy, PSS Perceived Stress, and CTQ Childhood Maltreatment scales for participants represents another major strength.

This study also has limitations including the exclusion of individuals with bipolar, and other psychotic, disorders that may also exhibit aggressiveness (Coccaro 2019) which limits the generalizability of these findings to a broader group of aggressive individuals. In addition, this study did not strictly follow recent recommendations for a standardized assessment of cortisol awakening responses (Stalder et al., 2016). This was because the present study was planned, funded, and implemented prior to the publication of those standards. As a result, this study: a) had only two samples for morning cortisol, and only one for evening cortisol, limiting the reliability of cortisol values and preventing the examination of an integrated Area Under the Curve variable, b) did not use an objective method to verify sampling accuracy (even though clear instructions to participants were given by study staff), and c) pre-study power analyses was not conducted. Unfortunately, a review of relevant publications since 2016, indicates that these standards have yet to be adopted by most investigators in the field (Stalder et al., 2022).

4.2 Conclusion.

In this case-control study, male and female adults with IED, compared to similar, but non-aggressive, individuals demonstrated a reduction in mean morning cortisol levels. The cortisol awakening response appears to be lower in individuals with IED compared with controls. In all study participants, morning salivary cortisol levels correlated inversely with trait anger, trait aggression, and plasma CRP, a marker of systemic inflammation. This suggests the present of a complex interaction between chronic-low level inflammation, the HPA axis, and IED that warrants further investigation.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

ACKNOWLEDGEMENTS

We would like to express our thanks to the participants for joining this study and the team members of the Clinical Neuroscience and Psychotherapeutics Research Unit, specifically, Ciara Zagaja, Sarina Farhardi, and Jocelyn Gomez, for their help in the coordination of this project.

CONFLICT OF INTEREST

This work was supported by a grant from the National Institutes of Health: RO1 MH 104673 (Dr. Coccaro). Dr. Coccaro reports being a member and consultant to the Scientific Advisory Boards of Azevan Pharmaceuticals, Inc., Avanir Pharmaceuticals, Inc., and Boehringer Ingelheim Pharmaceuticals, Inc. Drs. Meruelo, Timmins, and Irwin have nothing to declare.

REFERENCES

- American Psychiatric Association Task Force on DSM-5 (2013). Diagnostic and Statistical Manual of Mental Disorders: DSM-5 (5th ed.). American Psychiatric Association.
- Beck AT, Brown GK (1996). Beck Depression Inventory-II (BDI-II). The Psychological Corporation.
- Bernstein DP, Newcomb MD, Walker E, Pogge D, Ahluvalia T, Stokes J, Handlesman L, Medrano M, Desmond D, Zule W. (2003). Development and validation of a brief screening version of the Childhood Trauma Questionnaire. *Child Abuse Negl*, 27(2):169–190. [PubMed: 12615092]
- Blankenstein NE, Vandenbroucke ARE, de Vries R, Swaab H, Popma A, Jansen LMC (2022). Understanding aggression in adolescence by studying the neurobiological stress system: A systematic review. *Motivation Science*, 8(2):133–149.
- Böhnke R, Bertsch K, Kruk MR, Naumann E (2010). The Relationship between Basal and Acute HPA Axis Activity and Aggressive Behavior in Adults. *J Neural Trans*, 117(5): 629–637.
- Cima M, Smeets T, Jelicic M (2008). Self-reported trauma, cortisol levels, and aggression in psychopathic and non-psychopathic prison inmates. *Biol Psychiatry*, 78:75–86.
- Coccaro EF (2006). Association of C-reactive protein elevation with trait aggression and hostility in personality disordered subjects: a pilot study. *J Psychiatr Res*, 40(5):460–465. [PubMed: 15993896]
- Coccaro EF (2019). Psychiatric comorbidity in Intermittent Explosive Disorder. *J Psychiatr Res*, 118:38–43. [PubMed: 31487607]
- Coccaro EF, Berman ME, Kavoussi RJ (1997). Assessment of life history of aggression: development and psychometric characteristics. *Psychiatry Res*, 73(3):147–157. [PubMed: 9481806]
- Coccaro EF, Fitzgerald D, Lee R, McCloskey M, Phan KL. (2016). Frontolimbic Morphometric Abnormalities in Intermittent Explosive Disorder and Aggression. *Biol Psychiatry Cogn Neurosci Neuroimaging*, 1(1):32–38. [PubMed: 29560894]
- Coccaro EF, Kavoussi RJ (1994). Neuropsychopharmacologic challenge in biological psychiatry. *Clin Chemistry*, 40(2):319–327.

- Coccaro EF, Lee R, McCloskey MS (2014). Relationship between psychopathy, aggression, anger, impulsivity and Intermittent Explosive Disorder. *Aggressive Behavior*, 40:526–536. [PubMed: 24760575]
- Coccaro EF, Lee R, Kavoussi RJ (2010). Aggression, Suicidality, and Intermittent Explosive Disorder: Serotonergic Correlates in Personality Disorder and Healthy Control Subjects. *Neuropsychopharmacology*, 35(2):435–444. [PubMed: 19776731]
- Coccaro EF, Lee RJ (2020). Disordered Aggression and Violence in the United States. *J Clin Psychiatry*, 81(2).
- Coccaro EF, Lee R, Coussons-Read M (2014). Elevated plasma inflammatory markers in individuals with intermittent explosive disorder and correlation with aggression in humans. *JAMA Psychiatry*, 71(2):158–165. [PubMed: 24352431]
- Coccaro EF, McCloskey MS, Fitzgerald DA, Phan KL (2007). Amygdala and orbitofrontal reactivity to social threat in individuals with impulsive aggression. *Biol Psychiatry*, 62(2):168–178. [PubMed: 17210136]
- Coccaro EF, Nayer H, McCloskey MS (2012). Personality disorder-not otherwise specified evidence of validity and consideration for DSM-5. *Comprehensive Psychiatry*, 53(7):907–914. [PubMed: 22520088]
- Cohen S, Kamarck T, and Mermelstein R (1983). A global measure of perceived stress. *J Health Soc Behav*, 24:386–396.
- Coker KL, Smith PH, Westphal W, Zonana HV, McKee SA (2014). Crime and Psychiatric Disorders among Youth in the US Population: An Analysis of the National Comorbidity Survey-Adolescent Supplement. *JACAP*, 53(8):888–898.
- Counts CJ, Ginty AT, Larsen JM, Kampf TD, John-Henderson NA (2022). Childhood Trauma and Cortisol Reactivity: An Investigation of the Role of Task Appraisals. *Frontiers in Psychology* 13:803339. [PubMed: 35478771]
- Del Giudice M, Ellis BJ, Shirtcliff EA (2011). The Adaptive Calibration Model of Stress Responsivity. *Neuroscience and Biobehavioral Reviews*, 35(7): 1562–1592. [PubMed: 21145350]
- DeLisi M, Elbert M, Caropreso D, Tahja K, Heinrichs T, Drury A (2017). Criminally Explosive: Intermittent Explosive Disorder, Criminal Careers, and Psychopathology among Federal Correctional Clients. *International J Forensic Mental Health*, 16(4): 293–303.
- Echouffo-Tcheugui JB, Conner SC, Himali JJ, Maillard P, DeCarli CS, Beiser AS, Vasan RS, Seshadri S (2018). Circulating cortisol and cognitive and structural brain measures: The Framingham Heart Study. *Neurology*, 91(21): e1961–e1970. [PubMed: 30355700]
- First MB, Spitzer RL, Gibbon M, Williams JBW (1997). Structured Clinical Interview for DSM-IV Axis I Disorders (SCID). Psychiatric Institute, Biometrics Research.
- Freitag CM, Hänig S, Palmason H, Meyer J, Wüst S, Seitz C (2009). Cortisol Awakening Response in Healthy Children and Children with ADHD: Impact of Comorbid Disorders and Psychosocial Risk Factors. *Psychoneuroendocrinology*, 34(7):1019–1028. [PubMed: 19278790]
- Hart SD, Cox DN, Hare RD. (2003). Hare Psychopathy Checklist: Screening Version (PCL-SV). MHS.
- Heinz A, Jones DW, Bissette G, Hommer D, Ragan P, Knable M, Wellek S, Linnoila M, Weinberger DR (2002). Relationship between cortisol and serotonin metabolites and transporters in alcoholism [correction of alcoholism]. *Pharmacopsychiatry*, 35(4):127–134. [PubMed: 12163982]
- Hellhammer DH, Wüst S, Kudielka BM (2009). Salivary cortisol as a biomarker in stress research. *Psychoneuroendocrinology*, 34(2):167–171.
- Herrero N, Gadea M, Rodríguez-Alarcón G, Espert R, Salvador A (2010). What happens when we get angry? Hormonal, cardiovascular and asymmetrical brain responses. *Horm Behavior*, 57(3): 276–283.
- Higley JD, Mehlman PT, Poland RE, Taub DM, Vickers J, Suomi SJ, Linnoila M (1996). CSF testosterone and 5-HIAA correlate with different types of aggressive behaviors. *Biol Psychiatry*, 40(11):1067–1082. [PubMed: 8931909]
- Holi M, Auvinen-Lintunen L, Lindberg N, Tani P, Virkkunen M (2006). Inverse correlation between severity of psychopathic traits and serum cortisol levels in young adult violent male offenders. *Psychopathology*, 39:102–104. [PubMed: 16424682]

- Hollingshead AB (1975). Four factor index of social status. Unpublished manuscript.
- Juruena MF, Bocharova M, Agustini B, Young AH (2018). Atypical depression and non-atypical depression: Is HPA axis function a biomarker? A systematic review. *J Affect Disorders*, 233:45–67. [PubMed: 29150144]
- Kazén M, Kuenne T, Frankenberg H, Quirin M (2012). Inverse relation between cortisol and anger and their relation to performance and explicit memory. *Biol Psychology*, 91(1):28–35.
- Kindlundh AMS, Lindblom J, Bergström L, Nyberg F (2003). The anabolic-androgenic steroid nandrolone induces alterations in the density of serotonergic HHT1B and 5HT2 receptors in the male rat brain. *Neuroscience*, 119:113–1220. [PubMed: 12763073]
- Marsland AL, Prather AA, Petersen KL, Cohen S, Manuck SB (2008). Antagonistic characteristics are positively associated with inflammatory markers independently of trait negative emotionality. *Brain Behav Immun*, 22(5):753–761. [PubMed: 18226879]
- McBurnett K, Lahey BB, Rathouz PJ, Loeber R (2000). Low salivary cortisol and persistent aggression in boys referred for disruptive behavior. *Arch Gen Psychiatry*, 57(1):38–43. [PubMed: 10632231]
- McCloskey MS, Phan KL, Angstadt M, Fettich KC, Keedy S, Coccaro EF (2016). Amygdala hyperactivation to angry faces in intermittent explosive disorder. *J Psychiatr Res*, 79:34–41. [PubMed: 27145325]
- McEwen BS, Seeman T (1999). Protective and Damaging Effects of Mediators of Stress. Elaborating and Testing the Concepts of Allostasis and Allostatic Load. *Annals NY Acad Sciences*, 896:30–47.
- McLaughlin KA, Green JG, Hwang I, Sampson NA, Zaslavsky AM, Kessler RC (2012). Intermittent Explosive Disorder in the National Comorbidity Survey Replication Adolescent Supplement. *Arch Gen Psychiatry*, 69(11):1131–1139. [PubMed: 22752056]
- Miller GE, Chen E, Zhou ES (2007). If it goes up, must it come down? Chronic stress and the hypothalamic-pituitary-adrenocortical axis in humans. *Psychological Bulletin*, 133(1):25. [PubMed: 17201569]
- Mundt AP, Alvarado A, Fritsch R, Poblete C, Villagra C, Kastner S, Priebe S (2013). Prevalence Rates of Mental Disorders in Chilean Prisons. *PLoS One*, 8(7): e69109. [PubMed: 23894415]
- Nandam LS, Brazel M, Zhou M, Jhaveri DJ (2019). Cortisol and Major Depressive Disorder—Translating Findings From Humans to Animal Models and Back. *Front Psychiatry*, 10:974. [PubMed: 32038323]
- New AS, Trestman RF, Mitropoulou V, Goodman M, Koenigsberg HH, Silverman J, Siever LJ (2004). Low prolactin response to fenfluramine in impulsive aggression. *J Psychiatric Research*, 38(3):223–230.
- O’Leary MM, Loney BR, Eckel LA (2006). Gender differences in the association between psychopathic personality traits and cortisol response to induced stress.. *Psychoneuroendocrinology*, 32:183–191.
- Pajer K, Gardner W, Rubin RT, Perel J, Neal S (2001). Decreased Cortisol Levels in Adolescent Girls with Conduct Disorder. *Arch General Psychiatry*, 58(3):297–302.
- Patton J, Stanford M, Barratt E (1995). Factor structure of the Barratt impulsiveness scale. *J Clin Psychol*, 51(6):768–774. [PubMed: 8778124]
- Peterson CK, Harmon-Jones E (2012). Anger and testosterone: evidence that situationally-induced anger relates to situationally-induced testosterone. *Emotion*, 12(5):899–902. [PubMed: 21910539]
- Pfohl B, Blum N, Zimmerman M, & University of Iowa. Dept. of, Psychiatry (1997). Structured interview for DSM Personality Disorder: SIDP. American Psychiatric Press.
- Raine A (2002). Biosocial Studies of Antisocial and Violent Behavior in Children and Adults: A Review. *J Abnormal Child Psychology*, 30(4):311–326.
- Repetti RL, Robles TF, Reynolds B (2011). Allostatic Processes in the Family. *Development and Psychopathology*, 23(3):921–938. [PubMed: 21756442]
- Root JC, Tuescher O, Cunningham-Bussell A, Pan H, Epstein J, Altemus M, Cloitre M, Goldstein M, Silverman M, Furman D, Ledoux J, McEwen B, Stern E, Silbersweig D (2009). Frontolimbic function and cortisol reactivity in response to emotional stimuli. *Neuroreport*, 20(4):429–434. [PubMed: 19225430]

- Scott J, Cobb SW, Woods P, Matt GE, Meyer RA, Heaton RK, J Hampton Atkinson JR, Grant I (2007). Neurocognitive Effects of Methamphetamine: A Critical Review and Meta-Analysis. *Neuropsychology Review*, 17(3):275–297. [PubMed: 17694436]
- Sephton SE, Sapolsky RM, Kraemer HC, Spiegel D (2000). Diurnal Cortisol Rhythm as a Predictor of Breast Cancer Survival. *JNCI*, 92(12):994–1000. [PubMed: 10861311]
- Siever LJ, Buchsbaum MS, New AS, Spiegel-Cohen J, Wei T, Hazlett EA, Sevin E, Nunn M, & Mitropoulou V (1999). d,l-fenfluramine response in impulsive personality disorder assessed with [¹⁸F]fluorodeoxyglucose positron emission tomography. *Neuropsychopharmacology*, 20(5):413–423. [PubMed: 10192822]
- Silverman MN (2012). Glucocorticoid regulation of inflammation and its functional correlates: from HPA axis to glucocorticoid receptor dysfunction. *Ann. NY Acad. Sci.*, 1261:55–63. [PubMed: 22823394]
- Smith DM, Meruelo A, Campbell-Sills L, Sun X, Kessler RC, Ursano RJ, Jain S, Stein MB, Army STARRS Team. (2021). Pre-Enlistment Anger Attacks and Post-Enlistment Mental Disorders and Suicidality Among US Army Soldiers.. *JAMA Network Open* 4(9), e2126626. [PubMed: 34570207]
- Sobczak S, Honig A, Nicolson N, Reidel WJ (2002). Effects of Acute Tryptophan Depletion on Mood and Cortisol Release in First-degree Relatives of Type I and Type II Bipolar Patients and Healthy Matched Controls.. *Neuropsychopharmacology*, 27:834–842. [PubMed: 12431857]
- Spielberger CD (1999). STAXI-2 State Trait Anger Expression Inventory-2, Professional Manual. Psychological Assessment Resources, Inc.
- Stalder T, Kirschbaum C, Kudielka B,M, Adam EK, Pruessner JC, Wüst S, Dockray S, Smyth N, Evans P, Hellhammer DH, Miller R, Wetherell MA, Lupien SJ, Clow A (2016). Assessment of the cortisol awakening response: Expert consensus guidelines. *Psychoneuroendocrinology*, 63:414–432. [PubMed: 26563991]
- Stalder T, Lupien SJ, Kudielka BM, Adam EK, Pruessner JC, Wüst S, Dockray S, Smyth N, Evans P, Kirschbaum C, Miller R, Wetherell MA, Finke JB, Klucken T, Clow A (2022). Evaluation and update of the expert consensus guidelines for the assessment of the cortisol awakening response (CAR). *Psychoneuroendocrinology*, 146:105946. [PubMed: 36252387]
- Suarez EC (2003). Joint effect of hostility and severity of depressive symptoms on plasma interleukin-6 concentration. *Psychosom Med*, 65(4):523–527. [PubMed: 12883100]
- Suarez EC (2004). C-reactive protein is associated with psychological risk factors of cardiovascular disease in apparently healthy adults. *Psychosom Med*, 66(5):684–691. [PubMed: 15385692]
- Van Cauter E, Leproult R, Kupfer DJ (1996). Effects of Gender and Age on the Levels and Circadian Rhythmicity of Plasma Cortisol. *J Clin Endo Metabolism*, 81(7): 2468–2473.
- Virkkunen M, Rawlings R, Tokola R, Poland RE, Guidotti A, Nemeroff C, Bissette G, Kalogeras K, Karonen SL, Linnoila M (1994). CSF biochemistries, glucose metabolism, and diurnal activity rhythms in alcoholic, violent offenders, fire setters, and healthy volunteers. *Arch Gen Psychiatry*, 51(1):20–27. [PubMed: 7506515]
- Young B, Ziegler VE, Meyer DA (1978). A Rating Scale for Mania: Reliability, Validity and Sensitivity. *Br J Psychiatry*: 133:429–435. [PubMed: 728692]
- Zorn JV, Schür RR, Boks MP, Kahn RS, Joëls M, Vinkers CH (2017). Cortisol stress reactivity across psychiatric disorders: A systematic review and meta-analysis. *Psychoneuroendocrinology*, 77: 25–36. [PubMed: 28012291]

HIGHLIGHTS

HPA-axis dysfunction is evident in many behavioral disorders including those who display aggressive behavior.

Lower levels of morning cortisol may be linked with aggression, but this has not been studied in Intermittent Explosive Disorder (IED).

78 adults provided three saliva samples for morning/evening cortisol on three days and three morning CRP levels over an overlapping period.

Morning, but not evening, cortisol levels were significantly lower in IED and were inversely related to aggression and morning CRP levels.

This suggests a complex interaction between chronic low-level inflammation, the HPA axis, and IED that warrants further investigation.




Figure 1. Marginal means (+ SEM) of Log Morning (upon awakening and thirty minutes later), and Evening (prior to bedtime), Salivary Cortisol Levels in IED, Psychiatric (PC) and Healthy Control (HC) study participants. Asterisk indicates $p < 0.05$ from other groups.




Figure 2. Partial Correlation between Log Morning Salivary Cortisol levels and Composite Anger/Aggression scores in all subjects (covariates: age, sex, race, ses, and BIS-11, PCL-SV, BDI-II, PSS, CTQ scores). Closed circles designate study participants with IED, closed triangles designate psychiatric control, and open circles designate healthy control, study participants.




Figure 3. Partial regression plot for Log Salivary Morning Cortisol levels and Log CRP Plasma Levels in all subjects with these data (covariates: Age, Sex, Ethnicity, SES, BMI, and Composite Anger/Aggression score). Closed circles designate study participants with IED, closed triangles designate psychiatric control, and open circles designate healthy control, study participants.

TABLE I

Demographic and Psychometric Characteristics of Study Participants

	HC (N = 26)	PC (N = 26)	IED (N = 26)	p	Group Differences
<i>Demographic Variables</i>					
Age	31.2 + 9.1	29.9 + 8.9	33.7 + 9.4	= 0.310	HC = PC = IED
Sex (% Male)	26.9%	53.8%	46.2%	= 0.129	HC = PC = IED
Ethnicity (% White / African-American / Other)	31% / 46% / 23%	39% / 38% / 23%	31% / 50% / 19%	= 0.935	HC = PC = IED
SES Score	44.0 + 7.7	41.4 + 10.8	37.8 + 11.2	= 0.088	HC = PC > IED
<i>Psychometric Variables</i>					
Aggression (LHA)	3.7 + 3.3	5.4 + 3.0	18.5 + 3.7	< 0.001	HC = PC < IED
Anger Expression (STAXI-2)	-10.5 + 7.2	-6.0 + 9.2	13.7 + 7.8	< 0.001	HC = PC < IED
Impulsivity (BIS-11)	52.2 + 7.1	56.9 + 4.3	66.8 + 10.1	< 0.001	HC = PC < IED
Psychopathy (PCL-SV)	0.2 + 0.5	2.1 + 2.5	8.5 + 3.9	< 0.001	HC < PC < IED
Beck Depression Inventory	2.0 + 2.9	5.6 + 8.4	12.5 + 8.9	< 0.001	HC = PC < IED
Perceived Stress (PSS)	21.4 + 2.4	20.6 + 3.5	24.6 + 3.1	< 0.001	HC = PC < IED
Childhood Maltreatment (CTQ)	30.5 + 7.9	36.6 + 12.8	49.2 + 21.6	< 0.001	HC = PC < IED
<i>Other Variables</i>					
Body Mass Index (BMI)	26.4 + 4.0	25.2 + 2.9	26.1 + 3.8	= 0.464	HC = PC = IED

TABLE 2

DSM-5 Diagnoses Among Study Participants with a Psychiatric Disorder

	Psychiatric Controls (N = 26)	Intermittent Explosive Disorder (N = 26)	P	Group Differences*
Current Syndromal Disorders				
Any Depressive Disorder	1 (3.8%)	2 (7.7%)	0.557	PC = IED
Any Anxiety Disorder	6 (23.1%)	7 (26.9%)	0.749	PC = IED
Post-Traumatic Stress Disorder	2 (7.7%)	8 (30.8%)	0.075	PC = IED
Any Substance Use Disorder	0 (0.0%)	0 (0.0%)	0.999	PC = IED
Lifetime Syndromal Disorders				
Any Depressive Disorder	14 (53.8%)	11 (42.3%)	0.579	PC = IED
Any Anxiety Disorder	9 (34.6%)	10 (38.5%)	0.999	PC = IED
Post-Traumatic Stress Disorder	4 (15.4%)	13 (50.0%)	0.017	PC = IED
Any Substance Use Disorder	13 (50.0%)	11 (42.3%)	0.781	PC = IED
Personality Disorders				
Odd Cluster	0 (0.0%)	3 (11.3%)	0.235	PC = IED
Dramatic Cluster	2 (7.7%)	6 (19.2%)	0.419	PC = IED
Anxious Cluster	4 (15.4%)	10 (38.5%)	0.116	PC = IED
PD-Not Otherwise Specified	6 (23.1%)	11 (42.3%)	0.237	PC = IED

* $p < 0.05$ after correction for multiple comparisons (uncorrected $p < 0.0032$).