

UC Merced

UC Merced Previously Published Works

Title

Draft Genome Sequence and Description of Janthinobacterium sp. Strain CG3, a Psychrotolerant Antarctic Supraglacial Stream Bacterium.

Permalink

<https://escholarship.org/uc/item/80m675h1>

Journal

Microbiology Resource Announcements, 1(6)

ISSN

2169-8287

Authors

Smith, Heidi

Akiyama, Tatsuya

Foreman, Christine

et al.

Publication Date

2013-11-21

DOI

10.1128/genomeA.00960-13

Peer reviewed

Draft Genome Sequence and Description of *Janthinobacterium* sp. Strain CG3, a Psychrotolerant Antarctic Supraglacial Stream Bacterium

Heidi Smith,^{a,d} Tatsuya Akiyama,^{b,d} Christine Foreman,^{c,d} Michael Franklin,^{b,d} Tanja Woyke,^e Hazuki Teshima,^f Karen Davenport,^f Hajnalka Daligault,^f Tracy Erkkila,^f Lynne Goodwin,^f Wei Gu,^f Yan Xu,^f Patrick Chain^f

Department of Land Resources and Environmental Sciences, Montana State University, Bozeman, Montana, USA^a; Department of Microbiology, Montana State University, Bozeman, Montana, USA^b; Department of Chemical and Biological Engineering, Montana State University, Bozeman, Montana, USA^c; Center for Biofilm Engineering, Montana State University, Bozeman, Montana, USA^d; DOE Joint Genome Institute, Walnut Creek, California, USA^e; Genome Science Programs, Bioenergy and Biome Science, Los Alamos National Laboratory, Los Alamos, New Mexico, USA^f

Here we present the draft genome sequence of *Janthinobacterium* sp. strain CG3, a psychrotolerant non-violacein-producing bacterium that was isolated from the Cotton Glacier supraglacial stream. The genome sequence of this organism will provide insight as to the mechanisms necessary for bacteria to survive in UV-stressed icy environments.

Received 14 October 2013 Accepted 14 October 2013 Published 21 November 2013

Citation Smith H, Akiyama T, Foreman C, Franklin M, Woyke T, Teshima H, Davenport K, Daligault H, Erkkila T, Goodwin L, Gu W, Xu Y, Chain P. 2013. Draft genome sequence and description of *Janthinobacterium* sp. strain CG3, a psychrotolerant Antarctic supraglacial stream bacterium. *Genome Announc.* 1(6):e00960-13. doi:10.1128/genomeA.00960-13.

Copyright © 2013 Smith et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](https://creativecommons.org/licenses/by/3.0/).

Address correspondence to Christine Foreman, cforeman@montana.edu.

Psychrophiles often possess mechanisms to protect themselves from environmental conditions, including extremes of UV radiation, temperature, freeze-thawing events, and desiccation, which cause severe damage to nonadapted organisms (1). Reductions in stratospheric ozone have resulted in formation of the Antarctic ozone hole, causing increased levels of UV B radiation (UVB) to reach the Earth's surface, specifically affecting Antarctica (2). The highly damaging effects of short-wavelength radiation, UVB, and UV A radiation (UVA) to biological systems have been well documented (3). Short-wave radiation induces damage to a variety of macromolecules (e.g., DNA, RNA, proteins, and lipids) by indirectly increasing the formation of reactive oxygen species (4). The pigment violacein has been shown to possess strong antioxidant properties, protecting cellular lipid membranes from hydroxyl radicals by mitigating peroxidation (5).

Janthinobacterium sp. strain CG3 was isolated from a supraglacial fluvial system on the Cotton Glacier, Antarctica (77°07'S, 161°50'E). The bacterial strain was isolated on R2A agar medium incubated at 4°C in the dark for 12 days. The *Janthinobacterium* sp. strain CG3 isolate has characteristics of the genus *Janthinobacterium*, with strictly aerobic, Gram-negative, rod-shaped cells, but does not show evidence of violacein production (6). Additionally, *Janthinobacterium* sp. strain CG3 does not possess genetic evidence of the *viaA* operon, which is responsible for violacein production (7). As *Janthinobacterium* sp. strain CG3 lacks the genetic machinery necessary for violacein production, the genome sequence of this psychrotolerant organism will provide insight into strategies that have been evolved by organisms lacking pigments to survive in UV-stressed polar environments.

The draft genome sequence of *Janthinobacterium* sp. strain CG3 was generated at the DOE Joint Genome Institute (JGI) using Illumina data (8). Short-insert (insert length, 190 bp) and long-

insert (insert length, 10,800 bp) paired-end libraries generated 19,156,074 and 21,678,138 reads, respectively. A total of 6,125 Mbp of Illumina data were used to assemble the genome with Allpaths, Velvet, and Parallel Phrap. Initial assemblies generated by Allpaths version 39750 and Velvet version 1.1.05 (9) were computationally shredded into 10-kbp or 1.5-kbp overlapping fake reads, respectively. Overlapping fake reads from Allpaths and two Velvet assemblies were combined with the Illumina long-insert paired-end reads to generate the final assembly by Parallel Phrap version 4.24. Possible misassemblies were corrected with manual editing in Consed (10–12). Sequence gaps were closed by using repeat resolution software and sequencing bridging PCR fragments.

The draft genome sequence of *Janthinobacterium* sp. strain CG3 consists of 71 contigs in 7 scaffolds, with an estimated size of 6.4 Mbp. A total of 5,426 candidate protein-coding genes were predicted, with a total G+C content of 65.5%. The small-subunit rRNA gene sequences had 99% sequence identity to a *Janthinobacterium* sp. isolate from an Arctic glacier (GenBank accession number FM955878). Scaffolds 3, 4, and 7 were circularized with sequences of bridging PCR fragments, suggesting the presence of three plasmids in the *Janthinobacterium* sp. strain CG3 isolate.

Nucleotide sequence accession numbers. This whole-genome shotgun project has been deposited at DDBJ/EMBL/GenBank under accession number [APFF00000000](https://www.ncbi.nlm.nih.gov/nuccore/APFF00000000). The version described in this paper is version APFF01000000.

ACKNOWLEDGMENTS

Funding for this research came from the National Science Foundation (OPP-0838970 and 1141978). The work conducted by the U.S. Department of Energy Joint Genome Institute is supported by the Office of Science of the U.S. Department of Energy under contract no. DE-AC02-

05CH11231. H. Smith was supported by the NASA Earth and Space Science Fellowship (NESSF) program. Logistical support was provided by Raytheon Polar Services and Petroleum Helicopters Incorporated.

Any opinions, findings, and conclusions or recommendations expressed in this material are those of the authors and do not necessarily reflect the views of the National Science Foundation.

REFERENCES

1. Dieser M, Greenwood M, Foreman CM. 2010. Carotenoid pigmentation in Antarctic heterotrophic bacteria as a strategy to withstand environmental stresses. *Arct. Antarct. Alp. Res.* **40**:396–405.
2. Madronich S, McKenzie RL, Björn LO, Caldwell MM. 1998. Changes in biologically active ultraviolet radiation reaching the Earth's surface. *J. Photochem. Photobiol. B Biol.* **46**:5–19.
3. Buma AGJ, de Boer MK, Boelen P. 2001. Depth distribution of DNA damage in Antarctic marine phyto- and bacterioplankton exposed to summertime UV radiation. *J. Phycol.* **37**:200.
4. Lesser MP, Farrell JH, Walker CW. 2001. Oxidative stress, DNA damage and p53 expression in the larvae of Atlantic cod (*Gadus morhua*) exposed to ultraviolet (290–400 nm) radiation. *J. Exp. Biol.* **204**:157–164.
5. Konzen M, De Marco D, Cordova CAS, Vieira TO, Antônio RV, Creczynski-Pasa TB. 2006. Antioxidant properties of violacein: possible relation on its biological function. *Bioorg. Med. Chem.* **14**:8307–8313.
6. Sneath PHA. 1978. Classification of microorganisms. John Wiley & Sons, Chichester, United Kingdom.
7. Pantanella F, Berlutti F, Passariello C, Sarli S, Morea C, Schippa S. 2007. Violacein and biofilm production in *Janthinobacterium lividum*. *J. Appl. Microbiol.* **102**:992–999.
8. Bennett S. 2004. Solexa Ltd. *Pharmacogenomics* **5**:433–438.
9. Zerbino DR, Birney E. 2008. Velvet: algorithms for de novo short read assembly using de Bruijn graphs. *Genome Res.* **18**:821–829.
10. Ewing B, Hillier L, Wendl MC, Green P. 1998. Base-calling of automated sequencer traces using Phred. I. Accuracy assessment. *Genome Res.* **8**:175–185.
11. Ewing B, Green P. 1998. Base-calling of automated sequencer traces using Phred. II. Error probabilities. *Genome Res.* **8**:186–194.
12. Gordon D, Abajian C, Green P. 1998. Consed: a graphical tool for sequence finishing. *Genome Res.* **8**:195–202.