Lawrence Berkeley National Laboratory

Recent Work

Title

THE NUCLEAR DEFORMATION PARAMETERS AT HIGH EXCITATION ENERGIES

Permalink

<https://escholarship.org/uc/item/80s3q63x>

Author

Egido, J.L.

Publication Date

1986-04-01

Lawrence Berkeley Laboratory

UNIVERSITY OF CALIFORNIA

LAWRENCE
BERKELEY LABORATORY

JUN 18 1986

LIBRARY AND
DOCUMENTS SECTION

Submitted to Physics Letters B

THE NUCLEAR DEFORMATION PARAMETERS AT
HIGH EXCITATION ENERGIES

J.L. Egido, C. Dorso, J.O. Rasmussen,
and P. Ring

April 1986

TWO-WEEK LOAN COPY

This is a Library Circulating Copy
which may be borrowed for two weeks.

LBL-21461
c.2

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

The Nuclear Deformation Parameters at High Excitation Energies.[†]

J.L. Egido,* C. Dorso,** J.O. Rasmussen

Nuclear Science Division, Lawrence Berkeley Laboratory,
University of California, Berkeley, California 94720

and

P. Ring

Physik-Department der Technischen Universitaet Muenchen
Garching, West Germany

[†]This work was supported in part by a Fulbright/MEC grant and by the Director, Office of Energy Research, Division of Nuclear Physics of the Office of High Energy and Nuclear Physics, of the U.S. Department of Energy under Contract DE-AC03-76SF00098.

*Permanent address: Departamento de Fisica Teórica, Universidad Autonoma, Madrid, Spain.

**Permanent address: Facultad de Ciencias Exactas, Universidad de Buenos Aires, Argentina.

The Nuclear Deformation Parameters at High Excitation Energies.[†]

J.L. Egido,* C. Dorso,** J.O. Rasmussen

Nuclear Science Division, Lawrence Berkeley Laboratory,
University of California, Berkeley, California 94720

and

P. Ring

Physik-Department der Technischen Universitaet Muenchen,
Garching, West Germany

Abstract

The behavior of the deformation parameters with increasing excitation energy is analyzed in different theories. Realistic calculations are done for the nucleus ^{158}Er . The inclusion of fluctuations leads to large departures from the mean field values.

[†]This work was supported in part by a Fulbright/MEC grant and by the Director, Office of Energy Research, Division of Nuclear Physics of the Office of High Energy and Nuclear Physics, of the U.S. Department of Energy under Contract DE-AC03-76SF00098.

*Permanent address: Departamento de Física Teórica, Universidad Autónoma, Madrid, Spain.

**Permanent address: Facultad de Ciencias Exactas, Universidad de Buenos Aires, Argentina.

In recent years much experimental as well as theoretical effort has been devoted to understanding properties of nuclei under extreme conditions, such as high angular momentum and/or high excitation energy.¹ On the experimental side several new phenomena have been discovered. Some examples are the observed irregularities in the yrast band of a cold nucleus and the existence of Giant Resonances built on excited states². The quasi continuum is now the object of intensive study by several groups. The Self-consistent Cranking Model³ has been successfully used for the description of the yrast line of a cold heavy nucleus. For moderately higher excitation energy one still has to consider the pairing degree of freedom as well as the temperature dependence, which complicates the use of realistic forces in calculations of Hartree-Fock-Bogoliubov type. Consequently only the semiclassical method of Strutinsky generalized to finite temperature and superfluid nuclei has been used⁴ as an approximation. Recently the Finite Temperature Hartree-Fock-Bogoliubov (FTHFB) has been proposed⁵ and solved⁶ for separable forces with the Baranger-Kumar Hamiltonian and configuration space. A systematic study of the deformation parameters, as a function of temperature, has also been made⁶ in the FTHFB which as a mean field theory neglects contributions stemming from departures from the mean values.

In all the above calculations no attention has been paid to the fluctuations due to the finite temperature. However, simple models,^{7,8} as well as more realistic calculations,⁹ have shown that the effect of such fluctuations can be very important for quantities such as the pairing gap even at low temperatures. The motivation of this work was to extend such investigations to other degrees of freedom such as the deformation parameters a_0 and a_2

$$\begin{aligned} a_0 &= \beta \cos \gamma \\ a_2 &= \beta \sin \gamma \end{aligned} \quad , \quad (1)$$

which for a deformed nucleus are related to the experimentally observed quadrupole moments.

If one considers a_0 and a_2 as collective coordinates, it is possible to derive¹⁰ at finite temperature a classical Hamiltonian function in the Adiabatic Time-Dependent Hartree-Fock Theory given by

$$H(a_0, a_2, \dot{a}_0, \dot{a}_2) = \frac{1}{2} B_{00} \dot{a}_0^2 + \frac{1}{2} B_{22} \dot{a}_2^2 + B_{02} \dot{a}_0 \dot{a}_2 + V(a_0, a_2) \quad (2)$$

$V(a_0, a_2)$ is given by the temperature-dependent mean-field energy, and the mass parameters B_{ij} are given by

$$B_{\mu\nu} = \frac{1}{2} \sum_{k,l} \frac{q_{lk}^\mu q_{lk}^\nu}{|E_l - E_k|^3} |f_k - f_l| \quad (3)$$

where q^μ is the supermatrix,⁴

$$\begin{pmatrix} Q^{11} & Q^{20} \\ -Q^{02} & -Q^{11T} \end{pmatrix}$$

corresponding to the multipole operators $r^2 Y_{20}$ or $\frac{1}{\sqrt{2}}(r^2 Y_{22} + r^2 Y_{2-2})$. The quantities E_k are the quasiparticle energies,

$$\begin{pmatrix} E_k & 0 \\ 0 & -E_k \end{pmatrix}$$

and f_k the occupation factors of the Fermi-Dirac distribution at temperature T , in matrix form,

$$\begin{pmatrix} f_k & 0 \\ 0 & 1-f_k \end{pmatrix}.$$

The probability for the nucleus having the deformation a_0, a_2 is given^{11,9} by

$$P(a_0, a_2) \approx [B_{00}(a_0, a_2) B_{22}(a_0, a_2) - B_{20}^2(a_0, a_2)]^{1/2} \cdot \exp(-F(a_0, a_2)/T) \quad (4)$$

$F(a_0, a_2)$ is the free energy obtained from $V(a_0, a_2)$, varying a_0, a_2 independently. In principle, one should also consider the pairing degree of freedom as an independent variable, but then the calculations become very time consuming, so we approximate these calculations by solving the gap equations for each point of the plane a_0, a_2 to find the self-consistent gap for this deformation. The chemical potentials λ_Z, λ_N , were adjusted to get the right particle numbers. In our calculations for the deformed nucleus ^{168}Er we use for $V(a_0, a_2)$ the pairing plus quadrupole Hamiltonian of Baranger-Kumar¹², we also use their configuration space.

For each couple (a_0, a_2) we selfconsistently determine the gap parameter for protons and neutrons for different temperatures. In Fig. 1 contour plots of these quantities are depicted* in the β, γ plane for protons (upper part) and neutrons (lower part). For protons at $T=0.2$ MeV there are strong correlations in the whole β, γ plane ; at $T=0.4$ MeV they vanish only for β larger than 0.5. At $T=0.6$ MeV they disappear for $\beta > 0.3$ and for $T=0.8$ only very weak pairing correlations remain (maximum gap is 0.7 MeV for the spherical shape) for $\beta < 0.2$. For $T > 0.8$ MeV no significant pairing was found. For the neutron gap, see the lower part of the figure; a similar trend is observed, the only difference being that the quenching of the correlations is much faster. It is interesting the fact that the gap parameters show almost no dependence with the γ degree of freedom.

In calculating probabilities P , the mass dependence has been usually neglected⁷⁻⁸ under the implicit assumption that the mass parameters do not depend strongly on the parameters involved (in our case a_0 and a_2). Only recently, in Ref.[9] the probability (4) was calculated ,with and without mass dependence, taking the gap parameter as a coordinate. The results for the average gap value were, qualitatively, not very different in both approximations; only a small reduction in the first calculation with respect to the second was observed.

We again would like to do both kind of calculation for the deformations parameters. However, it is well known¹³ that the Cranking formula (3) has some inherent difficulties for the mass parameter in the region of a level crossing or pseudocrossing, since the denominators become very small. This difficulty increases at higher temperatures, where the gap has vanished and the particle-hole configurations give a finite contribution. In this letter it is not our intention to go beyond the approximation (3) to include further correlations which could wash out the sharpness of levels crossing. On the other hand, if one wants to have some information on the effect of the mass parameter on the probability P , some kind of modification has to be done in expression (3) to avoid the possible divergences in the level crossings. We adopt the pragmatic point of view of

*One has to distinguish between the coordinates a_0, a_2 and $\tilde{a}_0 = \chi \cdot \langle r^2 Y_{20} \rangle$, $\tilde{a}_2 = \chi \cdot \frac{1}{\sqrt{2}} \langle r^2 Y_{22} + Y_{2-2} \rangle$ where χ is the strength of the quadrupole-quadrupole interaction¹², they are equal only in the minimum. All results as well as the plots are calculated with $\tilde{a}_0$ and $\tilde{a}_2$ (in some cases the corresponding β, γ). We have omitted the tildes just for simpler notation.

introducing a minimum level separation ϵ , stemming from the neglected residual interaction, and replace $|E_k - E_l|$ by $|E_k - E_l| + \epsilon$ in (3). We have done calculations within the range $\epsilon = 0.2 - 3.0$ MeV and the results are qualitatively independent of ϵ . We shall later on come back on this point. In the final results we choose $\epsilon = 1$ MeV. In the following we denote by $P(a_0, a_2)$ the probability without mass dependence and by $P_M(a_0, a_2)$ the probability with mass dependence.

In Fig. 2a-f we show the probability distribution $P(\beta, \gamma)$ without mass dependence for the nucleus ^{168}Er at zero angular momentum for the temperatures $T = .2, .8, 1.0, 1.2, 1.4$ and 2.0 MeV. $P(\beta, \gamma)$ is normalized in such a way that the maximum value for a given temperature is unity. Each line in the contour plots differs by ± 1 from the adjacent lines. For $T = .2$ MeV we have a strongly peaked distribution around the axially symmetric minimum at $\beta = 0.23$. At $T = .8$ the distribution softens in all directions. At $T = 1.0$ the minimum is still on the prolate side, but on the oblate side at $\beta \sim .1$ there is an appreciable part of the probability density. At $T = 1.2$ the shifting towards the oblate side continues, while at $T = 1.4$ the maximum of the distribution is already at $\beta = 0$. At $T = 2.0$ MeV the distribution is essentially that of a spherical vibrator.

The full probability distribution $P_M(a_0, a_2)$, i.e., including the mass dependence, is shown in Fig. 3. For $T = 0.2$ MeV the exponential dependence in eq.(4) is too strong to be changed by the mass dependence on a_0, a_2 . For $T = 0.8$ MeV we already notice some difference with respect to Fig. 2, namely it softens faster toward the oblate shape. At temperature 1 MeV a second maximum appears at the oblate edge with equal probability to the prolate one. At $T = 1.2$ and 1.4 MeV the probability maximum appears for the oblate shape only. For $T = 2$ MeV the maximum extends around the spherical shape. The different behavior between $P(a_0, a_2)$ and $P_M(a_0, a_2)$ can be understood classically : for an axially symmetric spheroid the irrotational mass is given¹⁴ by $\frac{1}{5} (1 + \frac{1}{2} \frac{a^2}{c^2}) B$, where c is the semiaxis along the symmetry axis and a the transverse. Clearly since the oblate shape has a larger mass parameter the broad distribution which appears in Fig. 2 can be easily driven to the oblate shape.

In the upper part of Fig. 4 we show a_0^{sc} , the self-consistent value of a_0 , which corresponds to the value of a_0 in the free energy minimum in the plane a_0, a_2 for a given T . Since we are at zero angular momentum, the shape parameter γ is 0° and therefore $a_2^{sc} = 0$, ($a_2^{sc} = \beta$) for all temperatures. Also represented are the corresponding values of a_0 and a_2 but taking into account the temperature fluctuations, i.e.

$$\overline{a_K} = \langle a_K \rangle = \int a_K P(a_0, a_2) da_0 da_2, \quad K = 0, 2 \quad (5)$$

the scale of a_2^{sc} and $\overline{a_0}$ is on the left, and that of $\overline{a_2}$ is on the right. Corresponding shape parameters $\overline{a_{0M}}$ and $\overline{a_{2M}}$ are average values calculated taking into account the mass dependence of the probability, i.e. using $P_M(a_0, a_2)$ in eq. (5).

The integration in eq. 5 run over the area $0 \leq \beta < \infty$, $0 \leq \gamma \leq 60^\circ$ in the β, γ plane. This area includes all possible quadrupole deformations and each of them is counted only once. The average values $\overline{a_0}$ and $\overline{a_2}$ have to be considered with some care. A finite $\overline{a_2}$, for example, may arise from fluctuation away axial symmetry and not necessarily mean a most probable asymmetric shape. In order to understand the full picture, we have always to keep in mind the entire distributions as given in Figs. 2 and 3. At temperatures $0 < T < 0.5$ parameter a_2^{sc} (or β) slightly rises, mainly because of the reduction of pairing correlations with growing temperature. Between $T = 0.6$ and 1.1 MeV the temperature destroys the shell effects and we observe a diminution of the deformation. For $T > 1.1$ MeV the fluctuations are so large that the mean field approach breaks down, and the deformation drops dramatically from .16 at $T \sim 1.1$ MeV to zero at $T \sim 1.4$ MeV.

The average values $\overline{a_0}$ behave qualitatively in a similar manner to a_2^{sc} for temperatures $T \leq 1.1$ MeV, but for T larger than 1.1 MeV, $\overline{a_0}$ does not decrease but approaches the value $\sim .10$, as can be understood from the probability distribution $P(\beta, \gamma)$ in Fig. 2. The parameter a_2^{sc} , as mentioned above, is always zero because of axial symmetry; $\overline{a_2}$ rises with a rather steep slope from 0 to .045 at $T \sim 1.1$, then remains almost constant around this value. This behavior can also be understood from Fig. 2. Parameter $\overline{a_{2M}}$ is rather similar to $\overline{a_2}$ for all temperatures. Parameter $\overline{a_{0M}}$ looks like $\overline{a_0}$ for temperatures lower than 0.4 MeV but for higher temperatures

exhibits smaller values due to the fact that the maximum on the oblate side has a smaller β value. In both cases the sharp collapse of the deformation is smeared out by the temperature fluctuations. The neglect of the mass dependence overemphasizes this effect.

In Fig. 4b the quantities $\sigma(a_0)$ and $\sigma(a_2)$ are represented. The variance is defined by

$$\sigma^2(a_K) = \langle \bar{a}_K^2 \rangle - \langle \bar{a}_K \rangle^2, \quad K = 0, 2 \quad (6)$$

Again the left hand scale is for a_0 and the right hand scale for a_2 . Both of them behave in a similar way. They rise quickly from $T=0$ to $T \sim 1.1$ MeV and then stabilize, or slightly decrease, for higher temperatures. Finally to have some feeling about the dependence of the different quantities on the parameter ϵ introduced to avoid the possible singularities of the mass parameters, we show in Fig. 4c the average deformations $\bar{a}_{0M}$ and $\bar{a}_{2M}$ for some values of ϵ . The dashed lines correspond to $\epsilon = .5$, the dashed-dotted to $\epsilon = 1.0$ and the dotted to $\epsilon = 1.5$ MeV, the bold ones are for $\bar{a}_0$ and $\bar{a}_2$, i.e., when the mass dependence is neglected. From this figure one can conclude that the dependence on ϵ is rather smooth in a broad range. For very large values of ϵ the possible structure is washed out and we expect the limiting value of $\bar{a}_0$ and $\bar{a}_2$ respectively.

In summary, we have for the first time self-consistently taken into account the temperature fluctuation in the calculation of the deformation parameter for ^{168}Er in realistic calculations. The effects on a_0 are large for $T \geq 1.1$ MeV; on a_2 they become appreciable already for small temperatures.

ACKNOWLEDGEMENTS

One of the authors (J.L.E.) would like to acknowledge the Nuclear Science Division, especially R.M. Diamond and F.S. Stephens, for the hospitality extended to him during his stay in the Lawrence Berkeley Laboratory. W. Swiatecki is also thanked for a discussion

This work was supported in part by a Fulbright/MEC grant and by the Director, Office of Energy Research, Division of Nuclear Physics of the Office of High Energy and Nuclear Physics, of the U.S. Department of Energy under Contract DE-AC03-76SF00098.

References

1. B. Herskind, Nucl. Structure and Heavy Ion Dynamics, LXXVII Corso, Soc. Italiana de Fisica, Bologna, Italy, 1984, p. 68.
2. J.O. Newton, B.Herskind, R.M. Diamond, E.L. Dines, J.L. Draper, K.H. Lindenberg, C. Schuck, S. Shin, and F.S. Stephens, Phys. Rev. Lett. **46**, 1383 (1981).
3. J.L. Egidio and P. Ring. Nucl. Phys. A **388**, 19 (1982) and references therein.
4. P. Ring, L.M. Robledo, J.L. Egidio, and M. Faber, Nucl. Phys. A **419**, 261 (1984) and references therein.
5. A.L Goodman, Nucl. Phys. A **369**, 365 (1981);
K. Tanabe, K. Sugawara-Tanabe, and H.J. Mang, Nucl. Phys. A **357**, 20 (1981), Nucl. Phys. A **357**, 45 (1981).
6. J.L. Egidio, P. Ring, and H.J. Mang, Nucl. Phys. A **451**, 77 (1986).
7. L.G. Moretto, Phys. Lett. **44B**, 494 (1973).
8. A.L. Goodman, Phys. Rev. C **29**, 1887 (1984).
9. J.L. Egidio, P. Ring, S. Iwasaki, and H.J. Mang, Phys. Lett. B **154**, 1 (1985).
10. M. Baranger and M. Veneroni, Ann. Phys. (N.Y.) **114**, 123 (1978).
11. E.M. Lifshitz and L.P. Pitaevskii, Statistical Physics. (Pergamon, New York, 1980) Vol. 1
12. M. Baranger and K. Kumar, Nucl. Phys. A **110**, 490 (1968).
13. V.M. Strutinski, Z. Phys. A **280**, 99 (1977).
14. M. Lamb, Hydrodynamics, 6th ed. (Dover Publications, N. Y., 1945)

Fig. 1. Contour plots of the selfconsistent gap parameter for fixed (a_0, a_2) in the β, γ plane for different temperatures (in MeV) for the nucleus ^{158}Er . In the upper part the protons gap. The contour near the origin has the value : in (a) 1.7 MeV, in (b) 1.5 MeV, in (c) 1.35 MeV, and in (d) 0.7 MeV. The other contours decrease outwards with a stepsize of 0.150 MeV. In the lower part the same for neutrons. The contour near the origin has the value : in (a) 1.35 MeV, in (b) 1.2 MeV, in (c) 1.0 MeV. Further details as for the protons.

Fig. 2. The probability distribution $P(a_0, a_2)$, without mass dependence for different temperatures for the nucleus ^{158}Er . The outer contour corresponds to 0.1 it increases inwards with a stepsize of 0.1, the maximum value is 1.0.

Fig. 3. The probability distribution $P_M(a_0, a_2)$, with mass dependence for different temperatures for the nucleus ^{158}Er . The outer contour corresponds to 0.1, the stepsize of 0.1, the maximum value is 1.0. See text for further details

Fig. 4. In the upper part are shown the parameters a_0 and a_2 in different approximations (see text) versus the temperature. In the middle part are shown the quantities $\sigma(a_0)$, $\sigma(a_2)$. In both cases the left scale applies for a_0 , whereas the right hand one is for a_2 . In the lower part of the figure the parameters $\bar{a}_{0M}$ and $\bar{a}_{2M}$ are depicted for different values of the parameter ϵ .

^{158}Er

Fig. 1

XBL 864-10750

^{158}Er

XEL 364-10751

Fig. 2

^{158}Er

XBL 864-10752

Fig. 3

XBL 864-10753

Fig. 4

This report was done with support from the Department of Energy. Any conclusions or opinions expressed in this report represent solely those of the author(s) and not necessarily those of The Regents of the University of California, the Lawrence Berkeley Laboratory or the Department of Energy.

Reference to a company or product name does not imply approval or recommendation of the product by the University of California or the U.S. Department of Energy to the exclusion of others that may be suitable.

*LAWRENCE BERKELEY LABORATORY
TECHNICAL INFORMATION DEPARTMENT
UNIVERSITY OF CALIFORNIA
BERKELEY, CALIFORNIA 94720*