

UCSF

UC San Francisco Previously Published Works

Title

Comparing the effectiveness of a brief intervention to reduce unhealthy alcohol use among adult primary care patients with and without depression: A machine learning approach with augmented inverse probability weighting.

Permalink

<https://escholarship.org/uc/item/81q34524>

Authors

Papini, Santiago

Chi, Felicia

Schuler, Alejandro

et al.

Publication Date

2022-10-01

DOI

10.1016/j.drugalcdep.2022.109607

Peer reviewed



Published in final edited form as:

Drug Alcohol Depend. 2022 October 01; 239: 109607. doi:10.1016/j.drugalcdep.2022.109607.

Comparing the effectiveness of a brief intervention to reduce unhealthy alcohol use among adult primary care patients with and without depression: A machine learning approach with augmented inverse probability weighting

Santiago Papini^{a,*}, Felicia W. Chi^a, Alejandro Schuler^b, Derek D. Satre^{a,c}, Vincent X. Liu^a, Stacy A. Sterling^{a,c}

^aDivision of Research, Kaiser Permanente Northern California, 2000 Broadway, Oakland, CA 94612, USA

^bDivision of Biostatistics, UC Berkeley School of Public Health, 2121 Berkeley Way, Berkeley, CA 94704, USA

^cDepartment of Psychiatry and Behavioral Sciences, Weill Institute for Neurosciences, University of California, San Francisco, 675 18th Street, San Francisco, CA 94143, USA

Abstract

Background: The combination of unhealthy alcohol use and depression is associated with adverse outcomes including higher rates of alcohol use disorder and poorer depression course. Therefore, addressing alcohol use among individuals with depression may have a substantial public health impact. We compared the effectiveness of a brief intervention (BI) for unhealthy alcohol use among patients with and without depression.

Method: This observational study included 312,056 adult primary care patients at Kaiser Permanente Northern California who screened positive for unhealthy drinking between 2014 and 2017. Approximately half (48%) received a BI for alcohol use and 9% had depression. We examined 12-month changes in heavy drinking days in the previous three months, drinking days per week, drinks per drinking day, and drinks per week. Machine learning was used to estimate BI propensity, follow-up participation, and alcohol outcomes for an augmented inverse probability weighting (AIPW) estimator of the average treatment (BI) effect. This approach does not depend on the strong parametric assumptions of traditional logistic regression, making it more robust to model misspecification.

* Corresponding author.: santiago.x.papini@kp.org (S. Papini).

CRediT authorship contribution statement

Santiago Papini: Conceptualization, Methodology, Software, Formal analysis, Visualization, Writing - original draft, Writing - review & editing. **Felicia W. Chi:** Software, Data curation, Writing - original draft, Writing - review & editing. **Alejandro Schuler:** Methodology, Software, Supervision, Writing - original draft, Writing - review & editing. **Derek D. Satre:** Writing - review & editing. **Vincent X. Liu:** Resources, Writing - review & editing. **Stacy A. Sterling:** Conceptualization, Writing - review & editing, Supervision, Project administration, Funding acquisition.

Appendix A. Supporting information

Supplementary data associated with this article can be found in the online version at doi:10.1016/j.drugalcdep.2022.109607.

Results: BI had a significant effect on each alcohol use outcome in the non-depressed subgroup (-0.41 to -0.05 , all $ps < .003$), but not in the depressed subgroup (-0.33 to -0.01 , all $ps > .28$). However, differences between subgroups were nonsignificant (0.00 to 0.11 , all $ps > .44$).

Conclusion: On average, BI is an effective approach to reducing unhealthy drinking, but more research is necessary to understand its impact on patients with depression. AIPW with machine learning provides a robust method for comparing intervention effectiveness across subgroups.

Keywords

Alcohol brief intervention; Causal machine learning; Treatment heterogeneity; SBIRT; Depression; AIPW

1. Introduction

Unhealthy alcohol use is the most globally prevalent substance use problem (Degenhardt et al., 2018). Epidemiological research and meta-analyses provide evidence of a strong link between unhealthy drinking and depression: the presence of one is associated with an increased risk of the other (Boden and Fergusson, 2011; Grant et al., 2015; Lai et al., 2015; Sullivan et al., 2005). Compared to either problem on its own, the co-occurrence of unhealthy drinking and depression is associated with more severe alcohol use disorders (AUD), poorer course of depression, higher risk of suicide attempt, poorer global functioning, and lower quality of life and satisfaction (Brière et al., 2014; Levola et al., 2014; Sullivan et al., 2005). Unhealthy drinking patterns that do not meet criteria for an AUD are also highly prevalent among individuals with and without depression (Sullivan et al., 2005), who may be unaware that alcohol use is associated with adverse health consequences (National Institute on Alcohol Abuse and Alcoholism, 2005, updated 2007). Moreover, when patients with depression reduce unhealthy alcohol use, mental health symptoms often improve (Bahorik, 2016). Therefore, targeting unhealthy alcohol use can have public health impact in general, and on individuals with depression in particular.

In adult primary care, alcohol screening, brief intervention, and referral to treatment (SBIRT) is a population health approach that begins with routine alcohol screening to assess for unhealthy drinking, followed by brief intervention (BI) and/or referral to specialty treatment as needed (Babor et al., 2007). The most recent meta-analysis of randomized clinical trials (69 RCTs, 33,642 participants) found moderate-quality evidence in support of BIs for unhealthy alcohol use (Kaner et al., 2018). Despite these promising findings from RCTs, there is a paucity of effectiveness research of BI implemented in “real world” settings. An evaluation of SBIRT implementation at the United States Veterans Health Administration (30 facilities, 6210 participants with follow-up) did not find an association between BI and reduced rates of unhealthy drinking using multilevel logistic regression (Williams et al., 2014). Recently, we found small but significant effects of BI implemented in primary care of a large healthcare system (59 facilities, 312,056 participants) using marginal structural modeling, a method that accounts for potential biases arising from non-randomization of BI delivery and loss of follow-up by incorporating inverse probability weights (IPW) estimated with logistic regression (Chi et al., 2022). However, there is insufficient and inconsistent evidence on the effectiveness of alcohol BI in patients with

co-occurring depression (Boniface et al., 2017). For example, online (Montag et al., 2015) and face-to-face or telephone (Satre et al., 2013) BI led to significant improvement in drinking outcomes among patients with co-occurring depression. However, in studies with general practice patients (Grothues et al., 2008) and college students (Geisner et al., 2015), BI was less effective among participants with co-occurring depression.

Here our aim is to compare the effect of BI between depressed and non-depressed patients. We selected an augmented inverse probability weight (AIPW) approach with machine learning for several reasons. This approach, which can leverage data-adaptive machine learning algorithms to model complex nonlinear interactions, does not require parametric assumptions of traditional logistic regression that may not be realistic, such as the assumption of linear associations between covariates and outcomes (Kurz, 2022). AIPW is also doubly robust (i.e., reliable so long as either the treatment propensity or outcome model is correctly specified), which can reduce bias in estimates arising from misspecification of traditional logistic regression with or without IPW (Glynn and Quinn, 2010; Kurz, 2021, 2022).

2. Material and methods

2.1. Study design and cohort

Electronic health record (EHR) data were used to identify 440,882 patients who endorsed unhealthy drinking in Kaiser Permanente Northern California (KPNC) adult primary care between January 1, 2014, and December 31, 2017. For each patient, the index date was defined as the first positive screen for unhealthy drinking within the study window. The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) Diagram (Vandenbroucke et al., 2007) provided in Fig. 1 shows the exclusion criteria applied to define the cohort for this analysis, which consisted of 312,056 patients. Among them, 27,014 (8.7%) had an International Classification of Diseases (ICD) 9/10 diagnosis of depression. This study was approved by the Institutional Review Board at KPNC. Since this data-only study used existing protected health information in the EHR, waiver of informed consent was not required as data analysis is regulated by the Health Insurance Portability and Accountability Act (HIPAA) Privacy Rule.

2.2. Alcohol Screening, Brief Intervention, and Referral to Treatment (SBIRT)

KPNC conducts systematic SBIRT in adult primary care (Palzes et al., 2020). Medical assistants use National Institute on Alcohol Abuse and Alcoholism (NIAAA) screening tools embedded in the EHR to assess alcohol use. If a patient meets criteria for unhealthy drinking, defined as exceeding the NIAAA recommended daily limits (>3 drinks/day for women and men aged ≥ 66 , or >4 drinks/day for men aged 18–65) and/or weekly limits (>7 drinks/week for women and men aged ≥ 66 , or >14 drinks/week for men aged 18–65), physicians are trained to conduct a BI using Motivational Interviewing principles (Miller and Rollnick, 2012). Additionally, physicians can provide referral to specialty Addiction Medicine treatment programs within KPNC. The EHR codes used to determine delivery of BI are provided in the Supplementary Material.

2.3. Outcomes

Four 12-month outcomes were examined, defined as changes in: 1) heavy drinking days in the past 3 months, 2) drinking days per week, 3) drinks per drinking day, and 4) drinks per week. These outcomes are based on participant self-report. When a participant did not complete a 12-month screening, the previous screening was used (no less than 7 months post-index).

2.4. Baseline covariates

In a clinical trial, randomization is used to ensure that baseline differences between individuals that did and did not receive the intervention under evaluation occurred by chance. Effectiveness studies using observational data to estimate intervention effects must deal with potential confounds arising from nonrandomized intervention (Lu, 2009; Rosenbaum, 2002; Rubin, 1974). Non-random loss to follow-up may also introduce bias that can be mitigated by modeling the follow-up attrition process (Curtis et al., 2007). Additionally, both RCTs and observational studies can improve estimation of intervention effects by including covariates related to the outcome and must use appropriate methods to include patients with missing outcome data, since follow-up participation may be related to baseline characteristics, the type of intervention received, and their interactions (Rubin and Thomas, 2000). Accordingly, we selected a broad range of covariates hypothesized to be associated with intervention delivery, follow-up participation, and outcome.

Covariates were a total of 25 patient, provider, and facility characteristics. Patient characteristics were sex, age, race/ethnicity, insurance type categorized as either commercial, Medicaid, Medicare, or other/unknown, smoking status, physical activity, body mass index, Charlson comorbidity score (Charlson et al., 2008), presence of alcohol, drug, or other mental health disorders, neighborhood deprivation index from geocoded census data (Messer et al., 2006), and healthcare services utilization. Baseline alcohol consumption was defined as exceeding daily limit, weekly limit, or both daily and weekly limits (per NIAAA guidelines). Provider characteristics were age, sex, race/ethnicity, specialty (Internal Medicine, Family Practice, Other) and years of service. The year of screening, index facility, and department were also recorded.

2.5. Analyses

To estimate BI effects, we used augmented inverse probability weighting (AIPW) which incorporates estimates of intervention selection (i.e., propensity score), non-attrition (i.e., probability of having outcome data), and the outcome (Glynn and Quinn, 2010; Kurz, 2021). AIPW offers several advantages compared to traditional parametric regression-based analyses. By combining models for intervention, attrition, and outcome, AIPW can control multiple potential sources of bias. Moreover, the use of nonparametric machine learning models makes AIPW an efficient estimator for attaining the smallest theoretical sampling variance without imposing additional assumptions. Thus, this approach can obtain point estimates with less bias and smaller confidence intervals with better coverage. Both AIPW and machine learning are grounded in decades of theoretical and empirical research; advances in computing power along with availability of open-source machine learning software have facilitated the application of these advanced approaches in the biomedical

and behavioral sciences. The formulas used to estimate and compare BI effects across the depressed and non-depressed groups are provided in the Supplementary Material.

Each component of the AIPW models was estimated using stacked ensemble machine learning with the *h2o* package (H2O.ai, 2016) in the open-source programming software *R* (version 4.1.1) (R Core Team, 2013). Stacked ensembles (also referred to as super learners) are comprised of algorithms that estimate the outcomes of interest (i.e., the base learners) and an algorithm that integrates estimates from the base learners to make a final estimate (i.e., the meta-learner). This approach combines the strengths of diverse algorithms based on how accurately they estimate the outcome of interest. The added flexibility can protect against model misspecification because the algorithm(s) that works best (e.g., regression without interaction vs. data-adaptive tree-based methods with complex non-parametric interactions) may depend on the estimand (e.g., intervention outcome vs. intervention selection). Nested cross-estimation was used to protect against overfitting, such that all estimates used in the AIPW were based on models trained in separate data (Zivich and Breskin, 2021). Each meta-learner selected from a pool of up to 50 base learners comprised of generalized linear model with elastic net regularization, gradient boosting machines, distributed random forests, extremely randomized trees, and multi-layer artificial neural networks, each with different tuning parameters. The code used to run the analyses is included in the Supplementary Material, and a full description of the individual algorithms (including hyperparameters) is provided in the documentation for *h2o* (<https://docs.h2o.ai/h2o/1a-test-stable/h2o-docs/automl.html>).

3. Results

Fig. 2 summarizes the study results. AIPW estimates of the average intervention effect showed significantly greater reduction across the four drinking outcomes: 1) drinking days mean difference [95% CI], -0.05 [$-0.08, -0.02$], $p < 0.001$; 2) heavy drinking days, -0.41 [$-0.59, -0.22$], $p < 0.001$; 3) drinks per week -0.17 [$-0.28, -0.05$], $p = .004$; and 4) drinks per drinking day -0.06 [$-0.09, -0.03$], $p < .001$. The overall effects were consistent with what was estimated in the non-depressed subgroup (-0.41 to -0.05 , all $ps < .003$), whereas the depressed subgroup did not show significant effects (-0.33 to -0.01 , all $ps > .28$); however, comparisons between depressed and non-depressed subgroups were nonsignificant (0.00 to 0.11 , all $ps > .44$). AIPW estimates of the average intervention effect within each subgroup are provided in the Supplementary Material.

4. Discussion

Consistent with prior analyses that used marginal structural models with inverse probability weights estimated by logistic regression (Chi et al., 2022), our results showed modest but robust average BI effects across four drinking outcomes using an AIPW approach with machine learning, which requires fewer assumptions to address potential confounding in intervention delivery and follow-up participation and is more robust to model misspecifications. This highlights the utility of data-adaptive approaches to estimating intervention effectiveness as delivered to hundreds of thousands of patients in real world

settings, a key extension of the scientific support for BI that includes evidence from tightly controlled randomized trials in relatively smaller samples.

Extending this approach to compare the efficacy of BI between subgroups of patients with and without depression yielded a consistent, though nonsignificant, pattern of results suggesting that BI was slightly less effective for the depressed population. As captured by the confidence intervals in Fig. 2, the intervention effect in the depressed group was nonsignificant (i.e., the estimated effects overlapped with zero). A key source of the inflated variance in the depressed group is the relatively smaller sample size (27,014 depressed vs 285,042 non-depressed); this variability impacted inference on the difference in the intervention effect. Unobserved confounders may also have an impact on estimates, though their impact on our estimate of the difference in intervention effect would require that they have differential biasing effects across subgroups (Schuler and van der Laan, 2022). Together, these results suggest that further research is needed to improve our understanding of the impact of BI on alcohol use among patients with depression.

Further research may help clarify whether there are clinical explanations for the pattern of results we observed. Given that amotivation is a core feature of depression (Grahek et al., 2019; Smith, 2013), BI may not be sufficient for enhancing motivation to change alcohol use for patients with severe depression in general, or severe reductions in motivation in particular. There may be practice variation among primary care providers in how depression was considered when delivering BI, potentially leading to inconsistent effects. For example, it is possible that people with depression may benefit from a somewhat longer or more tailored brief intervention to reduce unhealthy alcohol use (Boniface et al., 2017; Satre et al., 2016). Additionally, some patients with depression may use alcohol to cope with negative emotional states (Magee and Connell, 2021); therefore, careful consideration of the interplay between alcohol use and depression (Boden and Fergusson, 2011) and more frequent monitoring of both outcomes (Hirschtritt et al., 2018) may be necessary to adequately address the complex needs of this subpopulation.

We note several limitations. Despite the advantages of AIPW with machine learning, any causal analysis must still rely on certain unverifiable assumptions. In our setting, the effects of interest can only be statistically identified if all confounders and common predictors of the outcome and loss to follow-up are observed and included in the analysis; there may be key unmeasured confounders that could impact results, such as depression severity, BI fidelity, and baseline motivation to change alcohol use. Additionally, we relied on patient self-report of alcohol use and there were no direct measurements of BI fidelity. Given the smaller size of the depressed subgroup, and the non-significance of statistical comparisons, cautious interpretation is warranted, and more research is necessary to better understand the efficacy of BI for patients with depression. Finally, we leveraged data from the EHR of a large integrated healthcare delivery system with membership that reflects the diversity of the U.S. population; further research is required to assess whether findings generalize to other healthcare systems or to uninsured populations.

5. Conclusions

We examined the impact of BI delivered in the primary care departments of a large, diverse, integrated healthcare system using AIPW with machine learning, a method that is flexible, efficient, robust, and interpretable, to compare intervention effectiveness across subgroups. On average, BI showed effectiveness in reducing unhealthy drinking, but more research is necessary to understand its impact on patients with depression, who showed a nonsignificant, though consistent, pattern of slightly weaker intervention effect.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We thank Agatha Hinman for editorial assistance with the manuscript.

Funding source

This study was supported by a grant from the National Institute on Alcohol Abuse and Alcoholism (R01AA025902, PI: Stacy A. Sterling). Dr. Papini's effort was supported by the Delivery Science Fellowship at Kaiser Permanente Northern California Division of Research and a grant from the National Institute of Mental Health (T32MH125792). Dr. Satre's effort was supported by a grant from the National Institute on Alcohol Abuse and Alcoholism (K24 AA025703). No other funding sources were declared. The funding sources had no role in the study design; in the collection, analysis, and interpretation of data; in the writing of the report; or in the decision to submit the article for publication.

References

- Babor TF, McRee BG, Kassebaum PA, Grimaldi PL, Ahmed K, Bray J, 2007. Screening, brief intervention, and referral to treatment (SBIRT): toward a public health approach to the management of substance abuse. *Subst. Abus.* 28 (3), 7–30. [PubMed: 18077300]
- Bahorik A, Leibowitz A, Sterling S, Weisner C, Travis A, Satre DD, 2016. The role of hazardous drinking reductions in predicting depression and anxiety symptom improvement among psychiatry patients: a longitudinal study. *J. Affect. Disord.* 206, 169–173. [PubMed: 27475887]
- Boden JM, Fergusson DM, 2011. Alcohol and depression. *Addiction* 106 (5), 906–914. [PubMed: 21382111]
- Boniface S, Malet-Lambert I, Coleman R, Deluca P, Donoghue K, Drummond C, Khadjesari Z, 2017. The effect of brief interventions for alcohol among people with comorbid mental health conditions: a systematic review of randomized trials and narrative synthesis. *Alcohol. Alcohol.* 53 (3), 282–293.
- Brière FN, Rohde P, Seeley JR, Klein D, Lewinsohn PM, 2014. Comorbidity between major depression and alcohol use disorder from adolescence to adulthood. *Compr. Psychiatry* 55 (3), 526–533. [PubMed: 24246605]
- Charlson ME, Charlson RE, Peterson JC, Marinopoulos SS, Briggs WM, Hollenberg JP, 2008. The Charlson comorbidity index is adapted to predict costs of chronic disease in primary care patients. *J. Clin. Epidemiol.* 61 (12), 1234–1240. [PubMed: 18619805]
- Chi FW, Parthasarathy S, Palzes VA, Kline-Simon AH, Metz VE, Weisner C, Satre DD, Campbell CI, Elson J, Ross TB, Lu Y, Sterling SA, 2022. Alcohol brief intervention, specialty treatment and drinking outcomes at 12 months: Results from a systematic alcohol screening and brief intervention initiative in adult primary care. *Drug Alcohol Depend.* 235, 109458. [PubMed: 35453082]
- Curtis LH, Hammill BG, Eisenstein EL, Kramer JM, Anstrom KJ, 2007. Using inverse probability-weighted estimators in comparative effectiveness analyses with observational databases. *Med. Care* S103–S107. [PubMed: 17909367]

- Degenhardt L, Charlson F, Ferrari A, Santomauro D, Erskine H, Mantilla-Herrera A, Whiteford H, Leung J, Naghavi M, Griswold M, 2018. The global burden of disease attributable to alcohol and drug use in 195 countries and territories, 1990–2016: a systematic analysis for the Global Burden of Disease Study 2016. *Lancet Psychiatry* 5 (12), 987–1012. [PubMed: 30392731]
- Geisner IM, Varvil-Weld L, Mittmann AJ, Mallett K, Turrisi R, 2015. Brief web-based intervention for college students with comorbid risky alcohol use and depressed mood: does it work and for whom? *Addict. Behav.* 42, 36–43. [PubMed: 25462652]
- Glynn AN, Quinn KM, 2010. An introduction to the augmented inverse propensity weighted estimator. *Political Anal.* 18 (1), 36–56.
- Grahek I, Shenhav A, Musslick S, Krebs RM, Koster EH, 2019. Motivation and cognitive control in depression. *Neurosci. Biobehav. Rev.* 102, 371–381. [PubMed: 31047891]
- Grant BF, Goldstein RB, Saha TD, Chou SP, Jung J, Zhang H, Pickering RP, Ruan WJ, Smith SM, Huang B, Hasin DS, 2015. Epidemiology of DSM-5 alcohol use disorder: results from the national epidemiologic survey on alcohol and related conditions III. *JAMA Psychiatry* 72 (8), 757–766. [PubMed: 26039070]
- Grothues JM, Bischof G, Reinhardt S, Meyer C, John U, Rumpf H-J, 2008. Effectiveness of brief alcohol interventions for general practice patients with problematic drinking behavior and comorbid anxiety or depressive disorders. *Drug Alcohol Depend.* 94 (1–3), 214–220. [PubMed: 18207336]
- H2O.ai, October 2016. R Interface for H2O, R package version 3.10.0.8. (<https://github.com/h2oai/h2o-3>) .
- Hirschtritt ME, Kline-Simon AH, Kroenke K, Sterling SA, 2018. Depression screening rates and symptom severity by alcohol use among primary care adult patients. *J. Am. Board Fam. Med.* 31 (5), 724–732. [PubMed: 30201668]
- Kaner EF, Beyer FR, Muirhead C, Campbell F, Pienaar ED, Bertholet N, Daeppen JB, Saunders JB, Burnand B, 2018. Effectiveness of brief alcohol interventions in primary care populations. *Cochrane Database Syst. Rev.* CD004148. [PubMed: 29476653]
- Kurz CF, 2021. Augmented inverse probability weighting and the double robustness property. *Med. Decis. Mak.* 42 (2), 156–157.
- Kurz CF, 2022. Augmented inverse probability weighting and the double robustness property. *Med. Decis. Mak.* 42 (2), 156–167.
- Lai HM, Cleary M, Sitharthan T, Hunt GE, 2015. Prevalence of comorbid substance use, anxiety and mood disorders in epidemiological surveys, 1990–2014: A systematic review and meta-analysis. *Drug Alcohol Depend.* 154, 1–13. [PubMed: 26072219]
- Levola J, Aalto M, Holopainen A, Cieza A, Pitkanen T, 2014. Health-related quality of life in alcohol dependence: a systematic literature review with a specific focus on the role of depression and other psychopathology. *Nord. J. Psychiatry* 68 (6), 369–384. [PubMed: 24228776]
- Lu CY, 2009. Observational studies: a review of study designs, challenges and strategies to reduce confounding. *Int J. Clin. Pract.* 63 (5), 691–697. [PubMed: 19392919]
- Magee KE, Connell AM, 2021. The role of substance use coping in linking depression and alcohol use from late adolescence through early adulthood. *Experimental and Clinical Psychopharmacology.*
- Messer LC, Laraia BA, Kaufman JS, Eyster J, Holzman C, Culhane J, Elo I, Burke JG, O'Campo P, 2006. The development of a standardized neighborhood deprivation index. *J. Urban Health* 83 (6), 1041–1062. [PubMed: 17031568]
- Miller WR, Rollnick S, 2012. *Motivational interviewing: Helping people change.* Guilford press.
- Montag AC, Brodine SK, Alcaraz JE, Clapp JD, Allison MA, Calac DJ, Hull AD, Gorman JR, Jones KL, Chambers CD, 2015. Effect of depression on risky drinking and response to a screening, brief intervention, and referral to treatment intervention. *Am. J. Public Health* 105 (8), 1572–1576. [PubMed: 26066915]
- National Institute on Alcohol Abuse and Alcoholism, 2005, updated 2007. *Helping patients who drink too much: a clinician's guide.*
- Palzes VA, Weisner C, Chi FW, Kline-Simon AH, Satre DD, Hirschtritt ME, Ghadiali M, Sterling S, 2020. The Kaiser Permanente Northern California Adult Alcohol Registry, an electronic health

records-based registry of patients with alcohol problems: Development and implementation. *JMIR Med Inf.* 8 (7), e19081.

- R Core Team, 2013. R: A language and environment for statistical computing.
- Rosenbaum PR, 2002. Overt bias in observational studies, *Observational Studies*. Springer, New York, pp. 71–104.
- Rubin DB, 1974. Estimating causal effects of treatments in randomized and nonrandomized studies. *J. Educ. Psychol.* 66 (5), 688–701.
- Rubin DB, Thomas N, 2000. Combining propensity score matching with additional adjustments for prognostic covariates. *J. Am. Stat. Assoc.* 95 (450), 573–585.
- Satre DD, Delucchi K, Lichtmacher J, Sterling SA, Weisner C, 2013. Motivational interviewing to reduce hazardous drinking and drug use among depression patients. *J. Subst. Abus. Treat.* 44 (3), 323–329.
- Satre DD, Leibowitz A, Sterling SA, Lu Y, Travis A, Weisner C, 2016. A randomized clinical trial of Motivational Interviewing to reduce alcohol and drug use among patients with depression. *J. Consult. Clin. Psychol.* 84 (7), 571. [PubMed: 26985728]
- Schuler A, van der Laan M, 2022. 2.3. Unidentified estimands. <https://alejandroschuler.github.io/mci/> . 2022).
- Smith B, 2013. Depression and motivation. *Phenomenol. Cogn. Sci.* 12 (4), 615–635.
- Sullivan LE, Fiellin DA, O'Connor PG, 2005. The prevalence and impact of alcohol problems in major depression: a systematic review. *Am. J. Med* 118 (4), 330–341. [PubMed: 15808128]
- Vandenbroucke JP, von Elm E, Altman DG, Gotzsche PC, Mulrow CD, Pocock SJ, Poole C, Schlesselman JJ, Egger M, initiative S, 2007. Strengthening the Reporting of Observational Studies in Epidemiology (STROBE): explanation and elaboration. *Ann. Intern. Med.* 147 (8), W163–W194. [PubMed: 17938389]
- Williams EC, Rubinsky AD, Chavez LJ, Lapham GT, Rittmueller SE, Achtmeyer CE, Bradley KA, 2014. An early evaluation of implementation of brief intervention for unhealthy alcohol use in the US V eterans H ealth A dministration. *Addiction* 109 (9), 1472–1481. [PubMed: 24773590]
- Zivich PN, Breskin A, 2021. Machine learning for causal inference: On the use of cross-fit estimators. *Epidemiology* 32 (3), 393–401. [PubMed: 33591058]




Fig. 1.
STROBE Diagram of the study cohort.




Fig. 2. Average BI effects for each 12-month alcohol use outcome. *Note.* Estimates are based on augmented inverse probability weighting models that used the complete sample of individuals that received BI ($n = 148,155$; 12,622 with depression) or did not receive BI ($n = 163,901$; 14,392 with depression).