

UCLA

UCLA Previously Published Works

Title

Exome-wide association study to identify rare variants influencing COVID-19 outcomes: Results from the Host Genetics Initiative.

Permalink

<https://escholarship.org/uc/item/8225n3bp>

Journal

PLoS Genetics, 18(11)

Authors

Butler-Laporte, Guillaume
Povysil, Gundula
Kosmicki, Jack
et al.

Publication Date

2022-11-01

DOI

10.1371/journal.pgen.1010367

Peer reviewed

RESEARCH ARTICLE

Exome-wide association study to identify rare variants influencing COVID-19 outcomes: Results from the Host Genetics Initiative

Guillaume Butler-Laporte^{1,2}, Gundula Povysil³, Jack A. Kosmicki⁴, Elizabeth T. Cirulli⁵, Theodore Drivas^{6,7,8}, Simone Furini⁹, Chadi Saad¹⁰, Axel Schmidt¹¹, Pawel Olszewski¹², Urszula Korotko^{12,13}, Mathieu Quinodoz^{14,15,16}, Elifnaz Çelik^{14,16}, Kousik Kundu^{17,18}, Klaudia Walter¹⁸, Junghyun Jung^{19,20}, Amy D. Stockwell²¹, Laura G. Sloofman²², Daniel M. Jordan²³, Ryan C. Thompson²⁴, Diane Del Valle²⁵, Nicole Simons²⁵, Esther Cheng²⁵, Robert Sebra²⁶, Eric E. Schadt²⁶, Seunghee Kim-Schulze²⁷, Sacha Gnjatich²⁵, Miriam Merad²⁸, Joseph D. Buxbaum²², Noam D. Beckmann²⁸, Alexander W. Charney²⁹, Bartłomiej Przychodzen³⁰, Timothy Chang³¹, Tess D. Pottinger³, Ning Shang³², Fabian Brand³³, Francesca Fava^{9,34,35}, Francesca Mari^{9,34,35}, Karolina Chwialkowska^{12,13}, Magdalena Niemira³⁶, Szymon Pula¹², J Kenneth Baillie^{37,38,39,40}, Alex Stuckey⁴¹, Antonio Salas^{42,43,44}, Xabier Bello^{42,43,44}, Jacobo Pardo-Seco^{42,43,44}, Alberto Gómez-Carballea^{42,43,44}, Irene Rivero-Calle^{43,44,45}, Federico Martínón-Torres^{43,44,45}, Andrea Ganna^{46,47}, Konrad J. Karczewski^{48,49}, Kumar Veerapen^{48,49}, Mathieu Bourgey^{50,51}, Guillaume Bourque^{50,51,52}, Robert JM Eveleigh^{50,51}, Vincenzo Forgetta², David Morrison², David Langlais^{51,52}, Mark Lathrop^{51,52}, Vincent Mooser⁵², Tomoko Nakanishi^{2,52,53,54}, Robert Frithiof⁵⁵, Michael Hultström^{1,2,55,56}, Miklos Lipcsey^{55,57}, Yanara Marincevic-Zuniga⁵⁸, Jessica Nordlund⁵⁸, Kelly M. Schiabor Barrett⁵, William Lee⁵, Alexandre Bolze⁵, Simon White⁵, Stephen Riffle⁵, Francisco Tanudjaja⁵, Efen Sandoval⁵, Iva Neveux⁵⁹, Shaun Dabe⁶⁰, Nicolas Casadei^{61,62}, Susanne Motamery^{63,64}, Manal Alaamery^{65,66}, Salam Massadeh^{65,66}, Nora Aljawini^{65,66}, Mansour S. Almutairi^{65,66}, Yaseen M. Arabi⁶⁷, Saleh A. Alqahtani^{68,69}, Fawz S. Al Harthi⁷⁰, Amal Almutairi⁷⁰, Fatima Alqubaishi⁷⁰, Sarah Alotaibi⁷⁰, Albandari Binowayn⁷⁰, Ebtehal A. Alsolm⁷⁰, Hadeel El Bardisy⁷⁰, Mohammad Fawzy⁷⁰, Fang Cai²¹, Nicole Soranzo¹⁸, Adam Butterworth¹⁸, COVID-19 Host Genetics Initiative[†], DeCOI Host Genetics Group[†], GEN-COVID Multicenter Study (Italy)[†], Mount Sinai Clinical Intelligence Center[†], GEN-COVID consortium (Spain)[†], GenOMICC Consortium[†], Japan COVID-19 Task Force[†], Regeneron Genetics Center[†], Daniel H. Geschwind³¹, Stephanie Arteaga³¹, Alexis Stephens⁷¹, Manish J. Butte^{71,72}, Paul C. Boutros⁷³, Takafumi N. Yamaguchi⁷³, Shu Tao⁷³, Stefan Eng⁷³, Timothy Sanders⁷⁴, Paul J. Tung⁷⁴, Michael E. Broudy⁷⁴, Yu Pan⁷⁴, Alfredo Gonzalez⁷³, Nikhil Chavan⁷⁴, Ruth Johnson⁷⁵, Bogdan Pasaniuc^{73,76,77}, Brian Yaspan²¹, Sandra Smieszek³⁰, Carlo Rivolta^{14,15,16}, Stephanie Bibert⁷⁸, Pierre-Yves Bochud⁷⁸, Maciej Dabrowski⁷⁹, Pawel Zawadzki^{79,80}, Mateusz Sypniewski⁷⁹, Elżbieta Kaja^{79,81}, Pajaree Chariyavilaskul⁸², Voraphoj Nilaratanakul⁸³, Nattiya Hirankarn⁸⁴, Vorasuk Shotelersuk⁸⁵, Monnat Pongpanich⁸⁶, Chureerat Phokaew⁸⁷, Wanna Chetruengchai⁸⁸, Katsushi Tokunaga⁸⁹, Masaya Sugiyama⁸⁹, Yosuke Kawai⁹⁰, Takanori Hasegawa⁹¹, Tatsuhiko Naito^{92,93}, Ho Namkoong⁹⁴, Ryuya Edahiro^{92,95}, Akinori Kimura⁹⁶, Seishi Ogawa^{97,98,99}, Takanori Kanai¹⁰⁰, Koichi Fukunaga¹⁰¹, Yukinori Okada^{92,93,102,103,104}, Seiya Imoto¹⁰⁵, Satoru Miyano⁹¹, Serghei Mangul¹⁹, Malak S. Abedalthagafi⁷⁰, Hugo Zeberg¹⁰⁶, Joseph J. Grzymalski⁵⁹, Nicole L. Washington⁵, Stephan Ossowski^{61,62}, Kerstin U. Ludwig^{11,107}, Eva C. Schulte^{108,109,110}, Olaf Riess^{61,62}, Marcin Moniuszko^{111,112}, Mirosław Kwasniewski^{12,13}, Hamdi Mbarek¹⁰, Said I. Ismail¹⁰, Anurag Verma^{7,8,113}, David B. Goldstein^{3,114}, Krzysztof Kiryluk^{3,32}, Alessandra Renieri^{9,34,35}, Manuel A. R. Ferreira⁴, J Brent Richards^{1,2,52,115,116,117*}



OPEN ACCESS

Citation: Butler-Laporte G, Povysil G, Kosmicki JA, Cirulli ET, Drivas T, Furini S, et al. (2022) Exome-wide association study to identify rare variants influencing COVID-19 outcomes: Results from the Host Genetics Initiative. *PLoS Genet* 18(11): e1010367. <https://doi.org/10.1371/journal.pgen.1010367>

Editor: Gregory M. Cooper, HudsonAlpha Institute for Biotechnology, UNITED STATES

Received: April 6, 2022

Accepted: July 29, 2022

Published: November 3, 2022

Copyright: © 2022 Butler-Laporte et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: Code availability Code guidance is available at <https://github.com/DrGBLWES.WGS>. Data availability The exome-wide burden test summary statistics are available in the manuscript’s supporting information files. The single variant association studies summary statistics are available openly on the GWAS Catalog [62] (study numbers GCST90132193, GCST90132194, and GCST90132195). Participant level data from each corresponding cohorts may be accessed according to the rules of each cohort’s

data sharing policies. Specifically, we refer readers to the following resources: - Biobanque Québécoise de la COVID-19 (BQC-19): <https://www.bqc19.ca/> - Sedish biobank: <https://swecovid.org/> - Columbia Biobank: <https://www.vagelos.columbia.edu/research/researchers/core-and-shared-facilities/new-instruments-and-facilities/columbia-university-biobank> - Geisinger Health Systems: <https://www.geisinger.edu/research> - Helix Exome+ and Healthy Nevada Project: <https://healthynev.org/> - Penn Medicine Biobank: <https://pmbb.med.upenn.edu/> - GEN-COVID Multicenter Study: <https://sites.google.com/dbm.unisi.it/gen-covid> - Qatar Genome Program: <https://www.qatargenome.org.qa/> - Deutsche COVID-19 OMICS Initiative (DeCOI): <https://decoi.eu/> - POLCOVID-Genomika: Medical University of Bialystok ethics board. - FHoGID: Commission cantonale d'éthique de la recherche sur l'être humain (CER-VD, <https://www.cer-vd.ch/>) - Interval: <https://www.intervalstudy.org.uk/> - Saudi Human Genome Program: <https://shgp.kacst.edu.sa/index.en.html> - Genentech: <https://www.gene.com/> - Mount Sinai Clinical Intelligence Center: <https://labs.icsahn.mssm.edu/minervalab/resources/data-ark/mscic-covid-19-biobank/> - Vanda COVID-19: <https://www.vandapharma.com/> - University of California Los Angeles: <https://www.uclahealth.org/precision-health/research> - Japan COVID-19 Taskforce: <https://www.covid19-taskforce.jp/en/home/> - Thai Biobank: Institutional Review Board of the Faculty of Medicine, Chulalongkorn University, Bangkok, Thailand (COA No. 691/2021) - MNM Bioscience Polish COVID WGS: <https://mnm-bioscience.com/> - UK Biobank: <https://www.ukbiobank.ac.uk/>.

Funding: Genome sequencing of Biobanque Québec COVID-19 was funded by the CanCOGeN HostSeq project, with contribution from Fonds de Recherche Québec Santé (FRQS), Génomique Québec, and the Public Health Agency of Canada. The Richards group is supported by the Canadian Institutes of Health Research (CIHR), the Lady Davis Institute of the Jewish General Hospital, the Canadian Foundation for Innovation, the NIH, Cancer Research UK, and FRQS. The Richards research group is supported by the Canadian Institutes of Health Research (CIHR: 365825; 409511, 100558, 169303), the McGill Interdisciplinary Initiative in Infection and Immunity (MI4), the Lady Davis Institute of the Jewish General Hospital, the Jewish General Hospital Foundation, the Canadian Foundation for Innovation, the NIH Foundation, Cancer Research UK, Genome Québec, the Public Health Agency of Canada, McGill University, Cancer Research UK [grant number C18281/A29019] and the Fonds de Recherche Québec Santé (FRQS). JBR is

1 Department of Epidemiology, Biostatistics and Occupational Health, McGill University, Montréal, Québec, Canada, **2** Lady Davis Institute, Jewish General Hospital, McGill University, Montréal, Québec, Canada, **3** Institute for Genomic Medicine, Columbia University, New York city, New York, United States of America, **4** Regeneron Genetics Center, Tarrytown, New York, United States of America, **5** Helix, San Mateo, California, United States of America, **6** Division of Human Genetics, Department of Pediatrics, Children's Hospital of Philadelphia, Philadelphia, Pennsylvania, United States of America, **7** Division of Translational Medicine and Human Genetics, Department of Medicine, Perelman School of Medicine, University of Pennsylvania, Philadelphia, Pennsylvania, United States of America, **8** Department of Genetics, Perelman School of Medicine, University of Pennsylvania, Philadelphia, Pennsylvania, United States of America, **9** Department of Medical Biotechnologies, Med Biotech Hub and Competence Center, University of Siena, Siena, Italy, **10** Qatar Genome Program, Qatar Foundation Research, Development and Innovation, Qatar Foundation, Doha, Qatar, **11** Institute of Human Genetics, School of Medicine and University Hospital Bonn, University of Bonn, Bonn, Germany, **12** IMAGENE.ME SA, Bialystok, Poland, **13** Centre for Bioinformatics and Data Analysis, Medical University of Bialystok, Bialystok, Poland, **14** Institute of Molecular and Clinical Ophthalmology Basel (IOB), Basel, Switzerland, **15** Department of Genetics and Genome Biology, University of Leicester, Leicester, United Kingdom, **16** Department of Ophthalmology, University Hospital Basel, Basel, Switzerland, **17** Department of Haematology, University of Cambridge, Cambridge, United Kingdom, **18** Department of Human Genetics, Wellcome Sanger Institute, Hinxton, United Kingdom, **19** Department of Clinical Pharmacy, School of Pharmacy, University of Southern California, Los Angeles, California, United States of America, **20** Department of Preventive Medicine, Keck School of Medicine, University of Southern California, Los Angeles, California, United States of America, **21** Genentech Inc, South San Francisco, California, United States of America, **22** Seaver Autism Center for Research and Treatment, Department of Psychiatry, Icahn School of Medicine at Mount Sinai, New York city, New York, United States of America, **23** Mount Sinai Clinical Intelligence Center, Charles Bronfman Institute for Personalized Medicine, Department of Genetics & Genomic Sciences, Icahn School of Medicine at Mount Sinai, New York city, New York, United States of America, **24** Icahn Institute of Data Science and Genomics Technology, New York city, New York, United States of America, **25** Icahn School of Medicine at Mount Sinai, New York city, New York, United States of America, **26** Department of Genetics and Genomic Sciences, Icahn School of Medicine at Mount Sinai, New York city, New York, United States of America, **27** Department of Oncological Science, Human Immune Monitoring Center, Precision Immunology Institute, Icahn School of Medicine at Mount Sinai, New York city, New York, United States of America, **28** Precision Immunology Institute, Tisch Cancer Institute, Icahn School of Medicine at Mount Sinai, New York city, New York, United States of America, **29** Mount Sinai Clinical Intelligence Center; Department of Genetics and Genomic Sciences, Icahn School of Medicine at Mount Sinai, New York city, New York, United States of America, **30** Vanda Pharmaceuticals, Washington, District of Columbia, United States of America, **31** Department of Neurology, David Geffen School of Medicine, University of California—Los Angeles, Los Angeles, California, United States of America, **32** Division of Nephrology, Department of Medicine, Vagelos College of Physicians & Surgeons, Columbia University, New York city, New York, United States of America, **33** Institute of Genomic Statistics and Bioinformatics, School of Medicine and University Hospital Bonn, University of Bonn, Bonn, Germany, **34** Genetica Medica, Azienda Ospedaliero-Universitaria Senese, Siena, Italy, **35** Medical Genetics, University of Siena, Siena, Italy, **36** Centre for Clinical Research, Medical University of Bialystok, Bialystok, Poland, **37** Roslin Institute, University of Edinburgh, Edinburgh, United Kingdom, **38** MRC Human Genetics Unit, Institute of Genetics and Molecular Medicine, University of Edinburgh, Western General Hospital, Edinburgh, United Kingdom, **39** Centre for Inflammation Research, The Queen's Medical Research Institute, University of Edinburgh, Edinburgh, United Kingdom, **40** Intensive Care Unit, Royal Infirmary of Edinburgh, Edinburgh, United Kingdom, **41** Genomics England, London, United Kingdom, **42** Unidade de Xenética, Instituto de Ciencias Forenses (INCFOR), Facultade de Medicina, Universidade de Santiago de Compostela, and GenPoB Research Group, Instituto de Investigaciones Sanitarias, Hospital Clínico Universitario de Santiago (SERGAS), Santiago de Compostela, Galicia, Spain, **43** Genetics, Vaccines and Infections Research Group (GENVIP), Instituto de Investigación Sanitaria de Santiago, Santiago de Compostela, Spain, **44** Centro de Investigación Biomédica en Red de Enfermedades Respiratorias (CIBER-ES), Madrid, Spain, **45** Translational Pediatrics and Infectious Diseases, Department of Pediatrics, Hospital Clínico Universitario de Santiago de Compostela, Santiago de Compostela, Spain, **46** Institute for Molecular Medicine Finland, Helsinki Institute of Life Science, University of Helsinki, Helsinki, Finland, **47** Massachusetts General Hospital, Harvard Medical School, Boston, Massachusetts, United States of America, **48** Stanley Center for Psychiatric Genetics, Broad Institute of Harvard and MIT, Cambridge, Massachusetts, United States of America, **49** Analytic and Translational Genetics Unit, Massachusetts General Hospital, Boston, Massachusetts, United States of America, **50** Canadian Centre for Computational Genomics, McGill University, Montréal, Québec, Canada, **51** McGill Genome Center, McGill University, Montréal, Québec, Canada, **52** Department of Human Genetics, McGill University, Montréal, Québec, Canada, **53** Kyoto-McGill International Collaborative School in Genomic Medicine, Graduate School of Medicine, Kyoto University, Kyoto, Japan, **54** Research Fellow, Japan Society for the Promotion of Science, Tokyo, Japan, **55** Anaesthesiology and Intensive Care Medicine, Department of Surgical Sciences, Uppsala University,

supported by a FRQS Mérite Clinical Research Scholarship. Support from Calcul Québec and Compute Canada is acknowledged. GBL is supported by FRQS and CIHR fellowships. The Columbia COVID-19 Biobank is supported by the Vagelos College of Physicians & Surgeons Office for Research, Precision Medicine Resource, and Biomedical Informatics Resource of the Columbia University Irving Institute for Clinical and Translational Research (CTSA). Columbia CTSA is funded by the National Center for Advancing Translational Sciences (UL1TR001873). The Columbia University COVID-19 Biobank was supported by Columbia University and the National Center for Advancing Translational Sciences (NCATS), National Institutes of Health (NIH), through grant no. UL1TR001873. DeCOI NGS was supported by funding from the German Research Foundation (DFG; INST 217/1011-1) and were performed at the DFG-funded NGS Competence Center Tübingen (INST 37/1049-1) – Project-ID 286/2020B01 – 428994620 and the DFG-funded West German Genome Center (INST 216/981-1) – Project Number 407493903. Funding for this work was further received from the Stiftung Universitätsmedizin Essen, Germany. The COMRI cohort is funded through in-house institutional funding of the Technical University of Munich, Munich, Germany. Individual grants are as follows: AS is supported by the BONFOR program of the Medical Faculty, University of Bonn (0-149.0134). KUL is supported by the Emmy-Noether programm of the German Research Foundation (DFG; LU 1944/3-1). RB was supported by the State of Saarland and the Dr. Rolf M. Schwiete Foundation. ECS is supported by the Munich Clinician Scientist Programm (MCSP) and the DFG (SCHU2419/2-1). JR received funding from DFG, RY159/3-1; DFG SFB1403; BMBF COVIM 01KX2021. MA and PS were supported by Netzwerk-Universitaetsmedizin-COVIM: (NaFoUniMedCovid19, FKZ: 01KX2021) and the BMFB (Idepico). For FhOGID: P-YB is supported by the Swiss National Science Foundation (31CA30_196036, 33IC30_179636 and 314730_192616), the Leenaards Foundation, the Santos-Suarez Foundation as well as grants allocated by Carigest. CR is supported by the Swiss National Science Foundation (31CA30_196036, 31003A_176097, and 310030_204285). The GEN-COVID Multicenter Study (Italy) was funded by the MIUR project “Dipartimenti di Eccellenza 2018-2020” to Department of Medical Biotechnologies University of Siena, Italy (Italian D.L. n.18 March 17, 2020), private donors for COVID-19 research, “Bando Ricerca COVID-19 Toscana” project to Azienda Ospedaliero-Universitaria Senese, charity fund 2020 from Intesa San Paolo dedicated to the

Uppsala, Sweden, **56** Integrative Physiology, Department of Medical Cell Biology, Uppsala University, Uppsala, Sweden, **57** Hedenstierna Laboratory, CIRRRUS, Anaesthesiology and Intensive Care Medicine, Department of Surgical Sciences, Uppsala University, Uppsala, Sweden, **58** Department of Medical Sciences, Science for Life Laboratory, Uppsala University, Uppsala, Sweden, **59** Center for Genomic Medicine, Desert Research Institute, Reno, Nevada United States of America, **60** Renown Health, Reno, Nevada, United States of America, **61** Institute of Medical Genetics and Applied Genomics, University of Tuebingen, Tuebingen, Germany, **62** NGS Competence Center Tuebingen, Institute of Medical Genetics and Applied Genomics, University of Tuebingen, Tuebingen, Germany, **63** West German Genome Center, site Cologne, University of Cologne, Cologne, Germany, **64** Cologne Center for Genomics, University of Cologne, Cologne, Germany, **65** Developmental Medicine Department, King Abdullah International Medical Research Center, King Saud Bin Abdulaziz University for Health Sciences, King Abdulaziz Medical City, Ministry of National Guard Health Affairs, Riyadh, Saudi Arabia, **66** Saudi Human Genome Project at King Abdulaziz City for Science and Technology, Riyadh, Saudi Arabia, **67** Ministry of the National Guard Health Affairs, King Abdullah International Medical Research Center and King Saud Bin Abdulaziz University for Health Sciences, Riyadh, Saudi Arabia, **68** Liver Transplant Unit, King Faisal Specialist Hospital and Research Centre, Riyadh, Saudi Arabia, **69** Division of Gastroenterology and Hepatology, Johns Hopkins University, Baltimore, Maryland, United States of America, **70** Genomics Research Department, Saudi Human Genome Project, King Fahad Medical City and King Abdulaziz City for Science and Technology, Riyadh, Saudi Arabia, **71** Department of Pediatrics, David Geffen School of Medicine, University of California—Los Angeles, Los Angeles, California, United States of America, **72** Department of Microbiology, Immunology, and Molecular Genetics (MIMG), David Geffen School of Medicine, University of California—Los Angeles, Los Angeles, California, United States of America, **73** Department of Human Genetics, David Geffen School of Medicine, University of California—Los Angeles, Los Angeles, California, United States of America, **74** Office of Health Informatics and Analytics, David Geffen School of Medicine, University of California—Los Angeles, Los Angeles, California, United States of America, **75** Department of Computer Science, McGill University, Montréal, Québec, Canada, **76** Department of Computational Medicine, David Geffen School of Medicine, University of California—Los Angeles, Los Angeles, California, United States of America, **77** Department of Pathology, David Geffen School of Medicine, University of California—Los Angeles, Los Angeles, California, United States of America, **78** Infectious Diseases Service, Department of Medicine, University Hospital and University of Lausanne, Lausanne, Switzerland, **79** MNM Bioscience Inc., Cambridge, Massachusetts, United States of America, **80** Faculty of Physics, Adam Mickiewicz University, Poznan, Poland, **81** Department of Medical Chemistry and Laboratory Medicine, Poznań University of Medical Sciences, Poznań, Poland, **82** Clinical Pharmacokinetics and Pharmacogenomics Research Unit, Department of Pharmacology, Faculty of Medicine, Chulalongkorn University, Bangkok, Thailand, **83** Healthcare-associated Infection Research Group STAR (Special Task Force for Activating Research) and Division of Infectious Diseases, Department of Medicine, Chulalongkorn University, Bangkok, Thailand, **84** Center of Excellence in Immunology and Immune-mediated Diseases, Department of Microbiology, Faculty of Medicine, Chulalongkorn University, Bangkok, Thailand, **85** Center of Excellence for Medical Genomics, Medical Genomics Cluster, and Department of Pediatrics, Faculty of Medicine, Chulalongkorn University, Bangkok, Thailand, **86** Department of Mathematics and Computer Science, Faculty of Science, Chulalongkorn University, Bangkok, Thailand, **87** Research Affairs, Faculty of Medicine, Chulalongkorn University, Bangkok, Thailand, **88** Center of Excellence for Medical Genomics, Medical Genomics Cluster, Faculty of Medicine, Chulalongkorn University, Bangkok, Thailand, **89** Genome Medical Science Project, Research Institute, National Center for Global Health and Medicine, Shinjuku-ku, Tokyo, Japan, **90** Genome Medical Science Project, National Center for Global Health and Medicine (NCGM), Tokyo, Japan, **91** M&D Data Science Center, Tokyo Medical and Dental University, Tokyo, Japan, **92** Department of Statistical Genetics, Osaka University Graduate School of Medicine, Suita, Japan, **93** Laboratory for Systems Genetics, RIKEN Center for Integrative Medical Sciences, Yokohama, Japan, **94** Department of Infectious Diseases, Keio University School of Medicine, Tokyo, Japan, **95** Department of Respiratory Medicine and Clinical Immunology, Osaka University Graduate School of Medicine, Suita, Japan, **96** Institute of Research, Tokyo Medical and Dental University, Tokyo, Japan, **97** Department of Pathology and Tumor Biology, Kyoto University, Kyoto, Japan, **98** Institute for the Advanced Study of Human Biology (WPI-ASHBi), Kyoto University, Kyoto, Japan, **99** Department of Medicine, Center for Hematology and Regenerative Medicine, Karolinska Institute, Stockholm, Sweden, **100** Division of Gastroenterology and Hepatology, Department of Medicine, Keio University School of Medicine, Tokyo, Japan, **101** Division of Pulmonary Medicine, Department of Medicine, Keio University School of Medicine, Tokyo, Japan, **102** Integrated Frontier Research for Medical Science Division, Institute for Open and Transdisciplinary Research Initiatives, Osaka University, Suita, Japan, **103** Laboratory of Statistical Immunology, Immunology Frontier Research Center (WPI-IFReC), Osaka University, Suita, Japan, **104** Center for Infectious Disease Education and Research (CiDER), Osaka University, Suita, Japan, **105** Division of Health Medical Intelligence, Human Genome Center, the Institute of Medical Science, the University of Tokyo, Tokyo, Japan, **106** Department of Neuroscience, Karolinska Institutet, Stockholm, Sweden, **107** West German Genome Center, site Bonn, University of Bonn, Bonn, Germany, **108** Institute of Psychiatric Phenomics and Genomics (IPPG), University Hospital, LMU Munich,

project N. B/2020/0119 “Identificazione delle basi genetiche determinanti la variabilità clinica della risposta a COVID-19 nella popolazione italiana”, the Italian Ministry of University and Research for funding within the “Bando FISR 2020” in COVID-19 and the Istituto Buddista Italiano Soka Gakkai for funding the project “PAT-COVID: Host genetics and pathogenetic mechanisms of COVID-19” (ID n. 2020-2016_RIC_3). The GEN-COVID (Spain) study received support from Instituto de Salud Carlos III (ISCIII): GePEM (PI16/01478/Cofinanciado FEDER; A.S.), DIAVIR (DTS19/00049/Cofinanciado FEDER; A.S.), Resvi-Omics (PI19/01039/Cofinanciado FEDER; A.S.), ReSVInext (PI16/01569/Cofinanciado FEDER; F.M.T.), Enterogen (PI19/01090/Cofinanciado FEDER; F.M.T.); Agencia Gallega para la Gestión del Conocimiento en Salud (ACIS): BI-BACVIR (PRIS-3; A.S.), and CovidPhy (SA 304 C; A.S.); Agencia Gallega de Innovación (GAIN): Grupos con Potencial de Crecimiento (IN607B 2020/08; A.S.), GEN-COVID (IN845D 2020/23; F.M.T.); Framework Partnership Agreement between the Consellería de Sanidad de la XUNTA de Galicia and GENVIP-IDIS - 2021-2024 (SERGAS-IDIS march 2021); and consorcio Centro de Investigación Biomédica en Red de Enfermedades Respiratorias (CB21/06/00103; F.M.T.). For Genentech: The COVACTA study was supported by F. Hoffmann-La Roche Ltd and, in part, by federal funds received from the U.S. Department of Health and Human Services, Office of the Assistant Secretary for Preparedness and Response, and Biomedical Advanced Research and Development Authority, under grant number HHSO100201800036C. For Helix+ and Healthy Nevada Project: Funding was provided to Desert Research Institute (DRI) by the Nevada Governor’s Office of Economic Development. Funding was provided to the Renown Institute for Health Innovation by Renown Health and the Renown Health Foundation. Thai Biobank (Host genetic factors in COVID-19 patients in relation to disease susceptibility, disease severity and pharmacogenomics) funding was obtained from the following sources: 1. Ratchadapiseksompotch Fund, Faculty of Medicine, Chulalongkorn University (RA(PO) 003/63 and 764002-HE01). 2. Grant for the Healthcare-associated Infection Research Group STAR (Special Task Force for Activating Research), Chulalongkorn University (STF 6100430002-1). 3. Grant for Development of New Faculty Staff, Ratchadapiseksomphot Endowment Fund (DNS 64_002_30_001_2). 4. The e-ASIA Joint Research Program (e-ASIA JRP) as administered by the National Science and Technology Development Agency. 5. Health Systems Research Institute, TSRI Fund

Munich, Germany, **109** Department of Psychiatry and Psychotherapy, University Hospital, LMU Munich, Munich, Germany, **110** Institute of Virology, Technical University Munich/Helmholtz Zentrum München, Munich, Germany, **111** Department of Regenerative Medicine and Immune Regulation, Medical University of Białystok, Białystok, Poland, **112** Department of Allergology and Internal Medicine, Medical University of Białystok, Białystok, Poland, **113** Corporal Michael Crescenz VA Medical Center, Philadelphia, Pennsylvania, United States of America, **114** Department of Genetics & Development, Columbia University, New York city, New York, United States of America, **115** Infectious Diseases and Immunity in Global Health Program, Research Institute of the McGill University Health Centre, Montréal, Québec, Canada, **116** Department of Twin Research, King’s College London, London, United Kingdom, **117** 5 Prime Sciences Inc, Montreal, Quebec, Canada

¶ Membership of DeCOI Host Genetics Group, COVID-19 Host Genetics Initiative, GEN-COVID Multicenter Study (Italy), Mount Sinai Clinical Intelligence Center, and GEN-COVID consortium (Spain) is provided in Supporting Information file [S2 Table](#).

* brent.richards@mcgill.ca

Abstract

Host genetics is a key determinant of COVID-19 outcomes. Previously, the COVID-19 Host Genetics Initiative genome-wide association study used common variants to identify multiple loci associated with COVID-19 outcomes. However, variants with the largest impact on COVID-19 outcomes are expected to be rare in the population. Hence, studying rare variants may provide additional insights into disease susceptibility and pathogenesis, thereby informing therapeutics development. Here, we combined whole-exome and whole-genome sequencing from 21 cohorts across 12 countries and performed rare variant exome-wide burden analyses for COVID-19 outcomes. In an analysis of 5,085 severe disease cases and 571,737 controls, we observed that carrying a rare deleterious variant in the SARS-CoV-2 sensor toll-like receptor *TLR7* (on chromosome X) was associated with a 5.3-fold increase in severe disease (95% CI: 2.75–10.05, $p = 5.41 \times 10^{-7}$). This association was consistent across sexes. These results further support *TLR7* as a genetic determinant of severe disease and suggest that larger studies on rare variants influencing COVID-19 outcomes could provide additional insights.

Author summary

COVID-19 clinical outcomes vary immensely, but a patient’s genetic make-up is an important determinant of how they will fare against the virus. While many genetic variants commonly found in the populations were previously found to be contributing to more severe disease by the COVID-19 Host Genetics Initiative, it isn’t clear if more rare variants found in less individuals could also play a role. This is important because genetic variants with the largest impact on COVID-19 severity are expected to be rarely found in the population, and these rare variants require different technologies to be studied (usually whole-exome or whole-genome sequencing). Here, we combined sequencing results from 21 cohorts across 12 countries to perform a rare variant association study. In an analysis comprising 5,085 participants with severe COVID-19 and 571,737 controls, we found that the gene for toll-like receptor 7 (*TLR7*) on chromosome X was an important determinant of severe COVID-19. Importantly, despite being found on a sex chromosome, this observation was consistent across both sexes.

(CU_FRB640001_01_30_10) and Thailand Research Fund (DPG6180001). Further, PC is supported by Ratchadapiseksompotch Fund, Faculty of Medicine, Chulalongkorn University, Bangkok, Thailand, Grant number RA(PO) 003/63. VN is supported by Ratchadapiseksompotch Fund, Faculty of Medicine, Chulalongkorn University, Bangkok, Thailand, Grant number RA(PO) 001/63. NH is supported by The e-ASIA Joint Research Program (e-ASIA JRP) as administered by the National Science and Technology Development Agency. VS is supported by Health Systems Research Institute (64-132) and the Ratchadapisek Sompoch Endowment Fund, Chulalongkorn University (764002-HE01), Bangkok, Thailand. Interval was funded by the NHS Blood and Transplant, the National Institute for Health Research, the UK Medical Research Council, and the British Heart Foundation Japan COVID-19 Task Force acknowledges the contribution of Japan Agency for Medical Research and Development (AMED) and Japan Science and Technology Agency (JST). The Japan NCGM-COVID-19 study was supported in part by Grants-in-Aid for Research from the National Center for Global Health and Medicine (20A2009) and the Agency for Medical Research and Development (AMED) (JP20fk0108416 and JP20fk0108104). MNM Diagnostics (Polish COVID WGS) partially supported by the Polish National Science Centre grant No. SZPITALE-JEDNOIMIENNE/2/2020 and by the Medical Research Agency grant No 2020/ABM/COVID19/0022. The Penn Medicine Biobank is supported by Perelman School of Medicine at University of Pennsylvania, a gift from the Smilow family, and the National Center for Advancing Translational Sciences of the National Institutes of Health under CTSA award number UL1TR001878. The POLCOVID-Genomika study was financially supported by the Polish Medical Research Agency (ABM) grant no. 2020/ABM/COVID19/0001. The Qatar Genome Program and Qatar Biobank are both Research, Development & Innovation entities within Qatar Foundation for Education, Science and Community Development. The Saudi COVID-19 acknowledges the Saudi Ministry of Health and King Abdulaziz City for Science and Technology (KACST). The Swedish Biobank received funding from the European Union's Horizon 2020 research and innovation program under grant agreement No 824110 (EASI-Genomics). Sequencing was performed by the National Genomics Infrastructure SNP&SEQ facility, which is supported by Science for Life Laboratory, the Swedish Research Council, and the Knut and Alice Wallenberg Foundation. UCLA acknowledges OCRC, Microsoft COVID Compute Funding, Illumina in-kind donation. We

Introduction

Despite successful vaccine programs, SARS-CoV-2 is still a major cause of mortality and widespread societal disruption [1,2]. While disease severity has correlated with well established epidemiological and clinical risk factors (e.g., advanced age, obesity, immunosuppression), these do not explain the wide range of COVID-19 presentations [3]. Hence, individuals without one of these known risk factors may have a genetic predisposition to severe COVID-19 [4]. These genetic determinants to severe disease can, in turn, inform about the pathophysiology underlying COVID-19 severity and accelerate therapeutics development [5,6].

Previous work on COVID-19 host genetics using genome-wide association studies (GWASs) revealed 23 statistically robust genetic loci associated with either COVID-19 severity or susceptibility [7–11]. Given that most GWASs use genetic data obtained from genome-wide genotyping followed by imputation to measure the association between a phenotype and genetic variation, their reliability and statistical power declines as a variant's frequency decreases, especially at allele frequencies of less than 1% [12]. Ascertainment of rare genetic variation can be improved with sequencing technology [13]. Rare variants are expected to be enriched for larger effect sizes, due to evolutionary pressure on highly deleterious variants, and may therefore provide unique insights into genetic predisposition to COVID-19 severity. Identifying such genes may highlight critical control points in the host response to SARS-CoV-2 infection.

Measuring the effect of rare genetic variants on a given phenotype (here COVID-19) is difficult. Specifically, while variants of large effect on COVID-19 are more likely to be rare, the converse is not true, and most rare variants are not expected to strongly impact COVID-19 severity [14]. Therefore, unless large sample sizes and careful statistical adjustments are used, most rare variant genetic associations studies risk being underpowered, and are at higher risk of false or inflated effect estimates if significant associations are found between COVID-19 and genetic loci. This is exemplified by the fact that several rare variant associations reported for COVID-19 have not been replicated in independent cohorts [15–17].

Here, we investigated the association of rare genetic variants on the risk of COVID-19 by combining gene burden test results from whole exome and whole genome sequencing. We build off recent work on exome-wide analyses [17] and include close to 5 times the number of severe cases, with a more genetically diverse cohort, to better study the effect of rare variants on COVID-19. To our knowledge, this is the first rare genetic variant burden test meta-analysis ever performed on a worldwide scale, including 21 cohorts, in 12 countries, including all main continental genetic ancestries.

Results

Study population and outcome

The final analysis included up to 28,159 individuals infected with SARS-CoV-2, and up to 597,165 controls from 21 cohorts in 12 countries (Fig 1). Most participants were of European genetic ancestry ($n = 576,389$), but the consortium also included participants of Admixed American ($n = 4,529$), African ($n = 25,465$), East Asian ($n = 4,716$), Middle Eastern ($n = 4,977$) and South Asian ancestries ($n = 9,943$). These resulted in a genetically diverse sample of participants (Fig 2). Participating cohorts enrolled patients based on local protocols, and both retrospective and prospective designs were used. Genetic sequencing was also performed locally, and cohorts were provided with a specific framework for quality control analyses, but each were allowed to deviate based on individual needs. Both exome ($n = 11$ cohorts) and genome

thank the UCLA COVID-19 Oversight Research Committee, Microsoft COVID Compute Funding, Illumina in-kind donation, and UCLA David Geffen School of Medicine - Eli and Edythe Broad Center of Regenerative Medicine and Stem Cell Research Award Program* for funding for this project under award #20-10 ("COVID-19 Host Genomics Registry at UCLA" PI:Pasaniuc). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing interests: I have read the journal's policy and the authors of this manuscript have the following competing interests: Biobanque Québécoise de la Covid-19: Brent Richards's institution has received investigator-initiated grant funding from Eli Lilly, GlaxoSmithKline and Biogen for projects unrelated to this research. He is the CEO of 5 Prime Sciences Inc (www.5primesciences.com). DeCOI: Oliver Witzke has received research grants for clinical studies, speaker's fees, honoraria and travel expenses from Amgen, Alexion, Astellas, Basilea, Biotest, Bristol-Myers Squibb, Correvio, Chiesi, Gilead, Hexal, Janssen, Dr. F. Köhler Chemie, MSD, Novartis, Roche, Pfizer, Sanofi, Takeda, TEVA and UCB. Kerstin U. Ludwig is co-founder and holds equity in the LAMPseq Diagnostics GmbH. UP serves as ad hoc advisor for Sanofi-Pasteur, BioNtech and Sobi and is member of the SAB of Leukocare. Christoph D. Spinner reports grants, personal fees from AstraZeneca, personal fees and non-financial support from BBraun Melsungen, grants, personal fees and non-financial support from Gilead Sciences, grants and personal fees from Janssen-Cilag, personal fees from Eli Lilly, personal fees from Formycon, personal fees from Roche, other from Apeiron, grants and personal fees from MSD, grants from Cepheid, personal fees from GSK, personal fees from Molecular partners, other from Eli Lilly, personal fees from SOBI during the conduct of the study; personal fees from AbbVie, personal fees from MSD, grants and personal fees from ViiV Healthcare, outside the submitted work. Jochen Schneider received grants and/or personal fees from Gilead Sciences, Janssen-Cilag, and from AbbVie outside the submitted work. Philipp Koehler reports grants or contracts from German Federal Ministry of Research and Education (BMBF) B-FAST (Bundesweites Forschungsnetz Angewandte Surveillance und Testung) and NAPKON (Nationales Pandemie Kohorten Netz, German National Pandemic Cohort Network) of the Network University Medicine (NUM) and the State of North Rhine-Westphalia; Consulting fees Ambu GmbH, Gilead Sciences, Mundipharma Research Limited, Noxxon N.V. and Pfizer Pharma; Honoraria for lectures from Akademie für Infektionsmedizin e.

sequencing ($n = 10$ cohorts) were included in the meta-analyses. The mean age of participants was 55.6 years, and 55.9% were females (**S3 Table**).

We studied three separate outcome phenotypes, as previously described by the COVID-19 Host Genetics Initiative (COVID-19 HGI)[8]. Briefly, the outcome cases were defined according to three standard COVID-19 HGI outcomes: A) severe disease: individuals with SARS-CoV-2 infection who died or required invasive respiratory support (extracorporeal membrane oxygenation, intubation with mechanical ventilation, high-flow oxygen support, or new bilevel or continuous positive airway pressure ventilation), B) hospitalisation: individuals with SARS-CoV-2 who died or required hospitalisation, and C) susceptibility to infection: any individual with SARS-CoV-2 infection. These are also referred to as A2, B2, and C2, respectively, in the COVID-19 HGI meta-analyses [8]. For all three phenotypes, controls were all individuals not classified as cases (including population controls with unknown COVID-19 status). The final meta-analyses included up to 5,085 cases and 571,737 controls for the severe disease outcome, 12,304 cases and 590,151 controls for the hospitalisation outcome, and 28,196 cases and 597,165 controls for the susceptibility outcome.

Single-variant analysis

We first performed an exome-wide association study using single variants with a MAF (minor allele frequency) higher than 0.1% and an allele count of 6 or more in at least one cohort, with the same additive model and covariates used in the COVID-19 HGI GWAS [8]. Analyses were performed separately by each cohort and each ancestry using Firth regression as applied in the Regenie software [18]. Firth regression is a penalized likelihood regression method that provides unbiased effect estimates even in highly unbalanced case-control analyses [19]. The summary statistics were then meta-analyzed with a fixed effect inverse-variance weighted model within each ancestry, and then with a DerSimonian-Laird random effect model across ancestries.

The previously described Neanderthal chromosome 3 locus associated with COVID-19 outcomes [8,20] was also found in all three phenotypes (**Figs 3 and S1**), with lead variants in the *CXCR6* gene for the severe COVID-19 phenotype (rs13059238), and in *FYCO1* in the hospitalisation phenotype (rs13069079), and for the *LIMD1* gene in the susceptibility phenotype (rs141045534). Reassuringly, each cohort provided summary statistics in the chromosome 3 locus, suggesting that the QC process was working as intended (allowing for sample sizes and number of cases) (**S2–S4 Figs**).

Three other loci were found for the hospitalization phenotypes. One at *SRRM1* (rs1479489847, OR: 4.17, 95% CI 2.60–6.70, $p = 3.25 \times 10^{-9}$), *IL6R* (rs911647797, OR: 6.19, 95% CI: 3.29–11.6, $p = 1.45 \times 10^{-8}$), and another at cytoskeleton *FRMD5* (rs1369031075, OR: 4.06, 95% CI: 2.70–6.11, $p = 1.75 \times 10^{-11}$). While these loci may hold biological plausibility (especially *IL6R*, given the use of IL-6 receptor inhibitors in the treatment of COVID-19), these associations were driven by two smaller cohorts (Genentech and Vanda, **S5 Fig**). However, the *SRRM1* locus is located between two stretches of T nucleotides, while both the *FRMD5* and the *IL-6* loci are within GC rich regions, making variant calling difficult. Hence, these findings will require validation, despite the biological plausibility.

Finally, all genetic inflation factors were below 1 (**S4 Table**). Summary statistics for genome-wide significant variants can be found in **S5 Table** and QQ-plots can be found in **S1 Fig**.

Burden test definition

Given the expected paucity of large-effect size rare deleterious variants, strategies have been devised to increase statistical power to test associations between rare variants and

V., Ambu GmbH, Astellas Pharma, BioRad Laboratories Inc., European Confederation of Medical Mycology, Gilead Sciences, GPR Academy Ruesselsheim, medupdate GmbH, MedMedia, MSD Sharp & Dohme GmbH, Pfizer Pharma GmbH, Scilink Comunicación Científica SC and University Hospital and LMU Munich; Participation on an Advisory Board from Ambu GmbH, Gilead Sciences, Mundipharma Resarch Limited and Pfizer Pharma; A pending patent currently reviewed at the German Patent and Trade Mark Office; Other non-financial interests from Elsevier, Wiley and Taylor & Francis online outside the submitted work. Oliver A. Cornely reports grants or contracts from Amplyx, Basilea, BMBF, Cidara, DZIF, EU-DG RTD (101037867), F2G, Gilead, Matinas, MedPace, MSD, Mundipharma, Octapharma, Pfizer, Scynexis; Consulting fees from Amplyx, Biocon, Biosys, Cidara, Da Volterra, Gilead, Matinas, MedPace, Menarini, Molecular Partners, MSG-ERC, Noxxon, Octapharma, PSI, Scynexis, Seres; Honoraria for lectures from Abbott, Al-Jazeera Pharmaceuticals, Astellas, Grupo Biotoscana/United Medical/Knight, Hikma, MedScape, MedUpdate, Merck/MSD, Mylan, Pfizer; Payment for expert testimony from Cidara; Participation on a Data Safety Monitoring Board or Advisory Board from Actelion, Allegra, Cidara, Entasis, IQVIA, Janssen, MedPace, Paratek, PSI, Shionogi; A patent at the German Patent and Trade Mark Office (DE 10 2021 113 007.7); Other interests from DGHO, DGI, EMM, ISHAM, MSG-ERC, Wiley, Genentech; Amy D Stockwell, Fang Cai, and Brian L Yaspan are, or were at the execution of the study, full time employees of Genentech with stock and stock options in Roche. Helix Exome+ and Healthy Nevada Project COVID-19 Phenotypes: Alexandre Bolze, Kelly M Schiabor Barrett, Simon White, Nicole L Washington, Francisco Tanudjaja, Stephen Riffle, Eren Sandoval, and Elizabeth T Cirulli are employees of Helix. Regeneron: Jack A Kosmicki and Manuel AR Ferreira are current employees and/or stockholders of Regeneron Genetics Center or Regeneron Pharmaceuticals. Vanda CALYPSO COVID-19: Bartłomiej Przychodzen and Sandra Smieszek are employees of Vanda Pharmaceuticals Inc.

biomedically-relevant outcomes. One such strategy is to use burden tests [21], where each variant is collapsed into larger sets of variants, and association is tested between groups of variants and an outcome. Here, we collapsed deleterious variants in each gene and devised the following burden test: for each gene, an individual received a score of 0 if they do not carry any deleterious variant, a score of 1 if they carry at least one non-homozygous deleterious variant, and a score of 2 if they carry at least 1 homozygous deleterious variant. As defined in previous studies on burden testing of rare variants [17,22] deleterious variants were chosen using three masks: 1) “M1” which uses only predicted loss of function variants, 2) “M3” which uses all variants in M1, as well as indels of moderate consequence as predicted by Ensembl [23], and missense variants classified as deleterious in 5 *in-silico* algorithms (see **Methods**), and 3) “M4”, which uses all variants in M1 and M3, and also adds all missense variants classified as deleterious in at least 1 of the *in-silico* algorithms.

The analyses were performed separately both for variants with MAF of less than 1%, and for variants of MAF less than 0.1%. We defined MAFs based on a combination of gnomAD [24] MAF annotations, and of cohort-specific common variant exclusion lists. These common variant exclusion lists included variants that achieved a MAF of >1% or >0.1% in at least one study population within the consortium. To reduce the effect of fluctuations due to sampling, a minor allele count (MAC) ≥ 6 in the corresponding study was required for inclusion in the common variant list. Such “blacklists” have been shown to increase statistical power by removing variants at lower risk of being highly deleterious, and it reduces the risk of having cohort-specific false-positive variants being retained on the overall analysis [25]. Hence for each MAF threshold, each cohort removed any variant with a MAF above the threshold in either gnomAD or the corresponding common variant exclusion list.

The resulting score (either 0, 1, or 2) for each mask was then regressed on each of our three phenotypes using logistic regression, controlling for age, age², sex, sex*age, sex*age², and 10 common variant (MAF > 1%) genetic principal components (the same covariates as for COVID-19 HGI GWASs [7,8]). Additionally, given that population genetic structure and its confounding effect on phenotypes is different at the rare variant level [26], we also used the first 20 genetic principal components from rare variants (MAF<1%) as covariates in all our analyses. Analyses were otherwise done using the same approach as for single-variant analyses.

Exome-wide burden test analyses results

Our meta-analysis included a total of 18,883 protein-coding genes, and all burden test genetic inflation factors, for all masks, were less than 1 (**S4 Table**), suggesting that our results were not biased by population stratification and that Firth regression adequately adjusted for unbalanced case-control counts. Using an exome-wide significance p-value threshold of 0.05/20,000 = 2.5×10^{-6} , we found 3 genes associated with one of the COVID-19 phenotypes in at least one mask in our meta-analyses (**Table 1 and S6–S12 Figs**). Of specific interest, we observed that carrying a predicted loss of function or *in-silico* highly deleterious missense variant (i.e., mask M3) in the toll-like receptor 7 (*TLR7*) gene was associated with a 5.3-fold increase (95% CI: 2.7–10.1, $p = 5.41 \times 10^{-7}$) in odds of severe COVID-19. *TLR7* is an important part of the innate viral immunity, encoding a protein that recognizes coronaviruses and other single-stranded RNA viruses, leading to upregulation of the type-1 and type-2 interferon pathway [27]. Results from the severe COVID-19 outcome analyses of *TLR7* with other masks also nearly reach our statistical significance threshold, with larger effects found in the M1 mask (OR: 13.6, 95% CI: 4.41–44.3, $p = 1.64 \times 10^{-5}$) and smaller effect in the M4 mask (OR: 3.12, 95% CI: 1.91–5.10, $p = 5.30 \times 10^{-6}$), though the latter was balanced by smaller standard errors due to the larger number of cases (3275 cases in M4 vs 1577 in M1), as expected. These findings




Fig 1. Maps of countries contributing data to the consortium. Sample sizes (cases and controls) for each phenotype were added and represented on the logarithmic scale by each circle. Relative contribution to each phenotype is represented by the three colors. Maps obtained from <https://www.naturalearthdata.com/>.

<https://doi.org/10.1371/journal.pgen.1010367.g001>

further support previous reports of *TLR7* errors of immunity underlying severe COVID-19 presentations [17,28–31].

In the meta-analyses, we also found that pLoFs (M1) in *MARK1* were associated with a 23.9-fold increase in the odds of severe COVID-19 (95% CI: 6.5–88.2, $p = 1.89 \times 10^{-6}$), and a 12.3-fold increase in the odds of hospitalisation due to COVID-19 (95% CI: 4.8–31.2,



Fig 2. Participant's genome projection on the first and second genetic principal components of the 1000G reference panel. AFR: African ancestry. AMR: admixed American ancestry. EAS: east Asian ancestry. EUR: European ancestry. MID: middle eastern ancestry. SAS: south Asian ancestry.

<https://doi.org/10.1371/journal.pgen.1010367.g002>

$p = 1.43 \times 10^{-7}$). While the number of *MARK1* pLoFs (M1) found in severe and hospitalized cases was small (MAC = 4 and MAC = 8, respectively), the signal was consistent in our three largest cohorts: UK Biobank, Penn Medicine, and Geisinger Health Services (S9–S12 Figs). *MARK1* is a member of the microtubule affinity-regulating kinase family, and is involved in multiple biological processes, chief among which is the promotion of microtubule dynamics [32]. *MARK1* has previously been shown to interact with the SARS-CoV-2 ORF9b protein [33], further supporting its potential role in COVID-19. Lastly, our meta-analyses also found marginal evidence for an association between severe COVID-19 and pLoFs (M1) in *RILPL1* (OR: 20.2, 95% CI: 5.8–70.7, $p = 2.42 \times 10^{-6}$), a gene that, like *MARK1*, is associated with microtubule formation and ciliopathy [34].

We then meta-analyzed p-values using the aggregated Cauchy association test [35] (ACAT). ACAT accounts for correlation between test statistics (as is expected here) by treating p-values as Cauchy random variables, and taking their weighted average, which also is Cauchy distributed. With ACAT, the association between *TLR7* and severe COVID-19 ($p = 1.58 \times 10^{-6}$), and between *MARK1* and hospitalisation ($p = 4.30 \times 10^{-7}$) remained exome-significant (Fig 4). Full summary statistics are available in S6 Table.




Fig 3. Single variant exome-wide association study Manhattan plot (MAF>0.1%). QQ-plot available in the [S1 Fig](#). Black dashed line demarcates the genome-wide significance threshold ($p < 5 \times 10^{-8}$).

<https://doi.org/10.1371/journal.pgen.1010367.g003>

Table 1. Exome-wide significant findings, as well as other *TLR7* results (for the severe phenotype only). Note that for Masks M1, all deleterious variants had a MAF<0.1%, and hence both burden tests (MAF<1% and 0.1%) gave the same results. Full results available in [S4 Table](#).

Gene	Mask	Phenotype	MAF	Beta	Standard Error	Odds Ratio	95% Confidence Interval	P-value	Heterogeneity p-value	N Cases 0 1 2 Burden Test	N Controls 0 1 2 Burden Test
Meta-Analysis Across Ancestries											
MARK1	M1	Severe COVID-19	<0.1%	3.17	0.67	23.9	6.5–88.2	1.89x10 ⁻⁶	0.883	1935 4 0	540031 92 0
MARK1	M1	Severe COVID-19	<1%	3.17	0.67	23.9	6.5–88.2	1.89x10 ⁻⁶	0.883	1935 4 0	540031 92 0
MARK1	M1	Hospitalisation	<0.1%	2.51	0.48	12.3	4.8–31.2	1.43x10 ⁻⁷	0.893	6132 8 0	547943 93 0
MARK1	M1	Hospitalisation	<1%	2.51	0.48	12.3	4.8–31.2	1.43x10 ⁻⁷	0.893	6132 8 0	547943 93 0
RILPL1	M1	Severe COVID-19	<0.1%	3.01	0.64	20.2	5.8–70.7	2.42x10 ⁻⁶	0.941	1745 4 0	558448 121 0
TLR7	M3	Severe COVID-19	<0.1%	1.66	0.33	5.25	2.75–10.05	5.41x10 ⁻⁷	0.755	3101 2 5	519047 83 47
TLR7	M3	Severe COVID-19	<1%	1.63	0.33	5.10	2.67–9.72	7.48x10 ⁻⁷	0.760	3275 2 5	519834 85 47
Other <i>TLR7</i> results for severe phenotype											
TLR7	M1	Severe COVID-19	<0.1%	2.61	0.60	13.6	4.14–44.4	1.64x10 ⁻⁵	0.820	1577 0 2	508987 13 11
TLR7	M1	Severe COVID-19	<1%	2.61	0.60	13.6	4.14–44.4	1.64x10 ⁻⁵	0.820	1577 0 2	508987 13 11
TLR7	M4	Severe COVID-19	<0.1%	1.14	0.25	3.12	1.91–5.10	5.30x10 ⁻⁶	0.854	3275 3 7	519616 210 139
TLR7	M4	Severe COVID-19	<1%	1.11	0.24	3.03	1.90–4.85	3.43x10 ⁻⁶	0.956	3273 5 8	521166 221 144

<https://doi.org/10.1371/journal.pgen.1010367.t001>

Finally, we note that for both *TLR7* and *MARK1*, the signal was driven by European ancestry participants. Further, while the larger biobanks contributed to these findings, smaller prospective cohorts also provided cases with rare variants at both genes, highlighting the importance of study design in rare variant association testing ([S6–S12 Figs](#)).

TLR7 sex stratified analyses

Given that *TLR7* is located on the X chromosome, we performed sex-stratified analyses of the severe disease phenotype to determine if the effect was also observed in females. These could only be done for the M3 and M4 masks due to very low number of M1 mask qualifying variants ([Fig 5](#)). In both we still see a clear effect among males with a 4.81-fold increase in the odds of severe COVID-19 in M3-variant carriers (95% CI: 2.41–9.59, 5 case carriers, 47 control carriers), and a 3.08-fold increase in M4-variant carriers (95% CI: 1.83–5.20, 7 case carriers, 143 control carriers). In females, we still observed a nominally significant signal in the M3 mask, with a 15.2-fold in odds of severe disease in M3-variant carriers (95% CI: 1.51–153.4). However, an M3-variant was observed in only one female with severe disease (heterozygous) in these analyses (compared to 76 heterozygous controls). In M4 variant, the analyses included 2 female heterozygous carriers (and 203 heterozygous controls), with a 4.86-fold in odds of severe disease (95% CI: 0.43–54.3).

Rare variants in interferon-related genes and at previously reported genome-wide significant loci

Despite a 7.7-fold increase in number of cases, and a 1,069-fold increase in number of controls, the previously reported associations of genes in the interferon pathway with COVID-19 outcomes [[15,16](#)] could not be replicated with either our exome-wide significance threshold ([S7 Table](#)) or a more liberal one of $p = 0.05/10 = 0.005$ (based on Bonferroni correction by the number of genes in the interferon pathway defined in a previous study [[15](#)]).

We also tested for rare variant associations between GWAS candidate genes from genome-wide significant loci in the COVID-19 HGI GWAS meta-analyses, but observed no exome-wide significant associations ([S8 Table](#)). However, at a more liberal Bonferroni threshold of $p = 0.05/46 = 0.001$ (correcting for the 46 genes in the COVID-19 HGI GWAS associated loci), we observed an increased burden of pLoF (M1) or missense variants (M3 mask) in *ABO* gene




Fig 4. Exome burden test ACAT p-value meta-analysis Manhattan plots and QQ plots. QQ-plot available in the [S6 Fig](#). Black dashed line demarcates the Bonferroni significance threshold ($p < 0.5/20,000$).

<https://doi.org/10.1371/journal.pgen.1010367.g004>




Fig 5. Sex-stratified *TLR7* analyses.

<https://doi.org/10.1371/journal.pgen.1010367.g005>

among those susceptible to SARS-CoV-2 infection (Table 2). For example, individuals carrying a pLoF (M1) with MAF<0.1% in *ABO* were at a 2.34-fold higher risk of having a positive SARS-CoV-2 infection (95% CI: 1.50–3.64, $p = 1.6 \times 10^{-4}$). The *ABO* results were driven mainly by European and African ancestry participants (S13 Fig). Note that deleterious variants in *ABO* often lead to blood groups A and B [36,37], which is consistent with the epidemiological association that non-type-O individuals are at higher risk of COVID-19[38]. However, more work is required to better understand the genetics of this locus as it relates to COVID-19 outcomes. Lastly, missense variants in *NSF* (mask M4, MAF<1%) were also associated with higher susceptibility to SARS-CoV-2 (OR: 1.48, 95% CI: 1.21–1.82, $p = 1.4 \times 10^{-4}$), but this association was not present in other masks (S9 Table).

Replication in GenOMICC

Data for the M1 mask for *TLR7* and *MARK1* in the severe COVID-19 phenotype was then replicated with the GenOMICC cohort [11], a prospective study enrolling critically ill individuals with COVID-19, with controls selected from the 100,000 genomes cohort[39]. Results are shown in Table 3. For *TLR7*, European ancestry individuals with a pLoF (M1) had a 4.70-fold increase in odds of severe disease (95% CI: 1.58 to 14.0, $p = 0.005$). In the sample of South Asian ancestry individuals, a pLoF (M1) was associated with a 1.90-fold increase in odds of severe disease, but the 95% confidence interval crossed the null (0.23 to 15.6, $p = 0.55$), which was likely due to a much smaller sample size than in the European ancestry subgroup (1,202 vs 10,645). Of interest, in both Europeans and South Asians, no pLoFs were observed in either of the control groups.

Table 2. Results of burden tests at genes identified from common variants GWAS in the COVID-19 HGI. Only genes with $p < 0.05/46$ are shown here. Full results available in S8 Table.

Gene	Mask	Phenotype	MAF	Beta	Standard Error	Odds Ratio	95% Confidence Interval	P-value	Heterogeneity p-value	N Cases 0 1 2 Burden Test	N Controls 0 1 2 Burden Test
NSF	M4	Susceptibility	<1%	0.395	0.104	1.484	1.21–1.82	1.44×10^{-4}	0.866	25752 127 2	585642 1907 5
ABO	M1	Susceptibility	<0.1%	0.851	0.226	2.341	1.50–3.65	1.68×10^{-4}	0.498	22778 27 0	572310 296 0
ABO	M1	Susceptibility	<1%	0.784	0.209	2.19	1.45–3.30	1.75×10^{-4}	0.826	23460 34 0	574608 364 0
ABO	M3	Susceptibility	<1%	0.729	0.195	2.073	1.41–3.04	1.89×10^{-4}	0.869	24455 42 0	575051 434 0
ABO	M1	Hospitalisation	<0.1%	1.33	0.395	3.78	1.74–8.20	7.56×10^{-4}	0.542	7859 12 0	561642 291 0
ABO	M1	Susceptibility	<0.1%	0.736	0.222	2.088	1.35–3.23	9.35×10^{-4}	0.512	23779 29 0	572799 320 0

<https://doi.org/10.1371/journal.pgen.1010367.t002>

Table 3. Replication of M1 mask, severe COVID-19, *MARK1* and *TLR7* results in the GenOMICC cohort. Note that the same variants were included in both the MAF<1% and MAF<0.1% replication, and the same results were obtained (shown here).

Gene	Ancestry	Beta	Standard Error	Odds Ratio	95% Confidence Interval	P-value	N Cases 0 1 2 Burden Test	N Controls 0 1 2 Burden Test
MARK1	EUR	0.195	1.42	1.21	0.075–19.7	0.891	5988 1 0	4655 1 0
MARK1	SAS	1.44	2.19	4.21	0.058–307.6	0.511	787 1 0	414 0 0
TLR7	EUR	1.55	0.555	4.70	1.58–14.0	0.005	5980 1 8	4566 0 0
TLR7	SAS	0.640	1.08	1.90	0.230–15.6	0.552	786 0 2	414 0 0

<https://doi.org/10.1371/journal.pgen.1010367.t003>

On the other hand, we could not replicate an effect from *MARK1*, which demonstrated an OR of 1.21 in European ancestry participants (95% CI 0.075 to 19.7, $p = 0.89$) and an OR of 4.21 in South Asian ancestry individuals (95% CI 0.058 to 307, $p = 0.51$).

Discussion

Whole genome and whole exome sequencing can provide unique insights into genetic determinants of COVID-19, by uncovering associations between rare genetic variants and COVID-19. Specifically, gene burden tests can be particularly helpful, because they test for coding variants, thereby pointing directly to a causal gene and often suggesting a direction of effect. However, such studies require careful control for population stratification and an adapted analysis method such as burden testing, in order to have enough statistical power to find those associations. In our study, we observed that individuals with rare deleterious variants at *TLR7* are at increased risk of severe COVID-19 (up to 13.1-fold increase in odds in those with pLoFs). Although this association was suggested by previous studies [28–30], our study provides the most definitive evidence for the role of *TLR7* in COVID-19 pathogenesis, with exome-wide significance for this gene in the discovery phase followed by strong replication in a large independent cohort. *TLR7* is a well-studied part of the antiviral immunity cascade and stimulates the interferon pathway after recognizing viral pathogen-associated molecular patterns. Given its location on the X chromosome, it has been hypothesized that it could partly explain the observed COVID-19 outcome differences between sexes [40–42], and to our knowledge, this is the first study to show that even in heterozygous females, this gene can potentially play a role in severe disease. Further, our results suggest that *TLR7* mediated genetic predisposition to severe COVID-19 may be a dominant or co-dominant trait, an observation that cannot be made in cohorts limited to male participants [28,30].

We also uncovered a potential role for cellular microtubule disruption in the pathogenesis of COVID-19 and the microtubule network is known to be exploited by other viruses during infections [43]. Indeed, the *MARK1* protein has been shown to interact with SARS-CoV-2 in previous *in-vitro* experiments [33]. Nevertheless, these findings at *MARK1* were not replicated in the GenOMICC cohort and will need to be tested in larger cohorts, especially given the small number of highly deleterious variants that we found in our consortium. Lastly, we found single variant associations at *IL6R*, *SRRM1*, and *FRMD5*. While *IL6R* is already a therapeutic target [44,45] for COVID-19, and *SRRM1* has been reported in a previous pre-print [46], these were found in smaller cohorts and will require replication.

To our knowledge, this is the first time a rare variant burden test meta-analysis has been attempted on such a large scale. Our framework allowed for easy and interpretable summary statistics results, while at the same time preventing participant de-identification or any breach of confidentiality that stems from sharing results of rare genetic variant analyses [47]. It also provides important insights into how these endeavours should be planned in the future. First, our burden test operated under the assumption that the effect of any of the deleterious variants

on the phenotype would be in the same direction and did not account for compound deleterious variant heterozygosity. This allowed for easier meta-analysis across cohorts, but may have decreased statistical power. Other methods may be needed in future analysis to soften this assumption, though some of these cannot be easily meta-analyzed across multiple cohorts directly from summary statistics (e.g., SKAT-O [48]). Similarly, methods that combine both rare and common variants might also provide additional insights into disease outcomes [31,49]. Second, our results highlight the importance at looking at different categories of variants through different masks to increase sensitivity and specificity of our burden tests. Third, while the largest biobanks contributed the most to the signal observed at *TLR7* and *MARK1*, many of our smaller prospective COVID-19 specific cohorts also contributed to the signal. This further highlights the importance of robust study design to improve statistical power, especially with rare variant associations. Lastly, work remains to be done to standardize sequencing and annotation pipelines to allow comparisons of results easily across studies and cohorts. Here, we provided a pipeline framework to every participating cohort, but there remains room for process harmonization. While the decentralized approach to genetic sequencing, quality control, and analyses allowed for more rapid generation of results, it may come at the cost of larger variance in our estimates. In the future, more sophisticated approaches may be required to increase statistical power of exome-wide rare variant association studies [50].

Our study had limitations. First, even if this is one of the world's largest consortia using sequencing technologies for the study of rare variants, we remain limited by a relatively small sample size. For example, in a recent analyses of UK Biobank exomes, many of the phenotypes for which multiple genes were found using burden tests had a much higher number of cases than in our analyses (e.g. blonde hair colour, with 48,595 cases) [22]. Further, rare variant signals were commonly found in regions enriched in common variants found in GWASs. The fact that *ABO* and *NSF* were the only genes from the COVID-19 HGI GWAS that were also identified in our burden test (albeit using a more liberal significance threshold), also suggests a lack of statistical power. Similarly, GenOMICC, a cohort of similar size, was also unable to find rare variant associations using burden tests [11]. However, their analysis methods were different from ours, making further comparisons difficult. Nevertheless, this provides clear guidance that smaller studies looking at the effect of rare variants across the genome are at considerable risk of finding both false positive and false negative associations. Second, many cohorts used population controls, which may have decreased statistical power given that some controls may have been misclassified. However, given that COVID-19 critical illness remains a rare phenomenon [51], our severe disease phenotype results are unlikely to be strongly affected by this. Finally, the use of population control is a long-established strategy in GWAS burden tests [7,8,11,22,52], and the statistical power gain from increasing our sample size is likely to have counter-balanced the misclassification bias.

In summary, we reproduced an exome-wide significant association with severe COVID-19 outcomes in carriers of rare deleterious variants at *TLR7*, for both sexes. Our results also suggest an association between the cellular microtubule network and severe disease, which requires further validation. More importantly, our results underline the fact that future genome-wide studies of rare variants will require considerably larger sample size, but our work provides a roadmap for such collaborative efforts.

Methods

Ethics statement

Each cohort had the following statement to make on ethics:

BQC-19. Each participant or their legal representative (if the participant was incapable to consent) provided informed consent to the biobank. If a participant regained capacity to give consent, informed consent was obtained again directly from the participant. The study was approved by the Jewish General Hospital and Centre Hospitalier de l'Université de Montréal institutional review boards.

Columbia Biobank: Recruitment and sequencing of participants from the Columbia COVID-19 Biobank were approved by the Columbia University Institutional Review Board (IRB) protocol AAAS7370 and the genetic analyses were approved under protocol AAAS7948. A subset of patients was included under a public health crisis IRB waiver of consent specifically for COVID-19 studies if patients were deceased, not able to consent, or if the study team was unable to contact them as per IRB protocol AAAS7370.

DeCOI. Informed consent was obtained from each participant or the legal representative. DeCOI received ethical approval by the Ethical Review Board (ERB) of the participating hospitals/centres (Technical University Munich, Munich, Germany; Medical Faculty Bonn, Bonn, Germany; Medical Board of the Saarland, Germany; University Duisburg-Essen, Germany; Medical Faculty Duesseldorf, Duesseldorf, Germany)

FHoGID. Each participant or their legal representative provided informed consent to the biobank. FHoGID received ethical approval by the Commission cantonale d'éthique de la recherche sur l'être humain.

GEN-COVID multicenter study: The patients were informed of this research and agreed to it through the informed consent process. The GEN-COVID is a multicentre academic observational study that was approved by the Internal Review Boards (IRB) of each participating centre (protocol code 16917, dated March 16, 2020 for GEN-COVID at the University Hospital of Siena).

Genentech. The protocol was reviewed by the institutional review board or ethics committee at each site. Written informed consent was obtained from all the patients or, if written consent could not be provided, the patient's legally authorized representative could provide oral consent with appropriate documentation by the investigator. Details on institutional review boards are provided in [S9 Table](#).

GenOMICC. GenOMICC was approved by the appropriate research ethics committees (Scotland, 15/SS/0110; England, Wales and Northern Ireland, 19/WM/0247). Informed consent was obtained for all participants.

Geisinger Health Systems: All subjects consented to participation and the analysis was approved by the Geisinger Institutional Review Board under project number 2006–0258.

Helix Exome+ and Healthy Nevada Project COVID-19 Phenotypes: informed consent was obtained for all participants. The Healthy Nevada Project study was reviewed and approved by the University of Nevada, Reno Institutional Review Board (IRB, project 956068–12)

Thai Biobank (). Informed consent was obtained for each participant via the biobank. The study was approved by the Institutional Review Board of the Faculty of Medicine, Chulalongkorn University, Bangkok, Thailand (COA No. 691/2021).

Japan COVID-19 Task Force: Each participant or their legal representative (if the participant was incapable to consent) provided informed consent to the biobank. Study was approved by the ethical committees of Keio University School of Medicine, Osaka University Graduate School of Medicine, and affiliated institutes.

Interval WGS. After reading study leaflets and participating in a discussion with donor carer staff, eligible donors were asked to complete the trial consent form before giving a blood donation. The National Research Ethics Service (United Kingdom) approved this study.

MNM Diagnostics (Polish Covid WGS). All participants, or their guardians/parents for the participants under 18), provided their informed consent before collecting their blood

samples. The study was approved by the Institutional Ethics Committee of the Central Clinical Hospital of the Ministry of Interior and Administration in Warsaw, Poland (decision nr: 41/2020 from 03.04.2020 and 125/2020 from 1.07.2020).

MSCIC. This research protocol was reviewed and approved by the Icahn School of Medicine at Mount Sinai Institutional Review Board (IRB) (STUDY-20-00341). During the height of the SARS-CoV-2 pandemic in New York City, all patients admitted to the Mount Sinai Health System were made aware of the research study by a notice included in their admission paperwork. The notice outlined details of the planned research, potential specimen collection and the opportunity to opt-out of research. Flyers announcing the study were also posted throughout the health system. Given the monumental hurdles of consenting sick and infectious patients in isolation rooms, the IRB allowed for specimen collection to occur prior to obtaining research consent at the time of clinical blood collection. Patients and/or their legally authorized representative provided consent to the research study, including genetic profiling for research and data sharing on an individual level. In a subset of individuals, who were unreachable following hospital discharge, we were unable to obtain written informed consent. In these cases, data cannot be shared further. All data used in these analyses were anonymized same as above.

Penn medicine. Recruitment of PMBB participants was approved under IRB protocol 813913 and supported by Perelman School of Medicine at University of Pennsylvania.

POLCOVID-Genomika. All study participants provided written informed consent and received detailed information on the study and associated risk before enrollment. The study was approved by the Bioethics Committee of the Medical University of Bialystok.

Qatar Genome Program. All QBB participants signed an Informed Consent Form prior to their participation; QBB study protocol ethical approval was obtained from the Hamad Medical Corporation Ethics Committee in 2011 and continued with QBB Institutional Review Board (IRB) from 2017 onwards and it is renewed on an annual basis

Saudi human genome program. Informed Consent was provided to each participant or their legal guardian (if the participant could not consent) by the corresponding institute. This study was approved the IRB of each participating hospitals, and the IRB at King Abdullah International Medical Research Centre, Ministry of National Guard–Health Affairs, Riyadh, Ministry of Health, and King Fahad Medical City.

Swedish Biobank. Informed consent was obtained for all study participants. The study was approved by the National Ethical Review Agency (Sweden) (No. 2020–01623).

UK Biobank. All subjects consented to participation. The UK Biobank was approved by the North West Multi-centre Research Ethics Committee (United Kingdom) (11/NW/0382). The work described herein was approved by the UK Biobank under application no. 26041.

University of California, Los Angeles biobank. Each participant or their legal representative (if the participant was incapable to consent) provided informed consent to the biobank. If a participant regained capacity to give consent, informed consent was obtained again directly from the participant. This study was considered human subjects research exempt because all genetic and electronic health records were de-identified. This study was approved by the UCLA Health Institutional Review Board.

Vanda COVID-19. All participants consented to WGS. The study was reviewed and approved by Advarra IRB; Pro00043096.

COVID-19 outcome phenotypes

For all analyses, we used three case-control definitions: A) Severe COVID-19, where cases were those who died, or required either mechanical ventilation (including extracorporeal

membrane oxygenation), high-flow oxygen supplementation, new continuous positive airway pressure ventilation, or new bilevel positive airway pressure ventilation, B) Hospitalized COVID-19, where cases were all those who died or were admitted with COVID-19, and C) Susceptibility to COVID-19, where cases are anyone who tested positive for COVID-19, self-reported an infection to SARS-CoV-2, or had a mention of COVID-19 in their medical record. For all three, controls were individuals who did not match case definitions, including population controls for which case status was unknown (given that most patients are neither admitted with COVID-19, nor develop severe disease [53]). These three analyses are also referred to as analyses A2, B2, and C2 by the COVID-19 Host Genetics Initiative [8], respectively.

Cohort inclusion criteria and genetic sequencing

Any cohort with access to genetic sequencing data and the associated patient level phenotypes were allowed in this study. Specifically, both whole-genome and whole-exome sequencing was allowed, and there were no limitations in the platform used. There were no minimal number of cases or controls necessary for inclusion. However, the first step of Regenie, which was used to perform all tests (see below), uses a polygenic risk score which implicitly requires that a certain sample size threshold be reached (which depends on the phenotype and the observed genetic variation). Hence, cohorts were included if they were able to perform this step. All cohorts obtained approval from their respective institutional review boards, and informed consent was obtained from all participants. More details on each cohort's study design and ethics approval can be found in the [S3 and S1 Tables](#).

Variant calling and quality control

Variant calling was performed locally by each cohort, with the pre-requisite that variants should not be joint-called separately between cases and controls. Quality control was also performed individually by each cohort according to individual needs. However, a general quality control framework was made available using the Hail software [54]. This included variant normalization and left alignment to a reference genome, removal of samples with call rate less than 97% or mean depth less than 20. Genotypes were set to unknown if they had genotype quality less than 20, depth less than 10, or poor allele balance (more than 0.1 for homozygous reference calls, less than 0.9 for homozygous alternative calls, and either below 0.25 or above 0.75 for heterozygous calls). Finally, variants were removed from if the mean genotype quality was less than 11, mean depth was less than 6, mean call rate less than or equal to 0.8, and Hardy-Weinberg equilibrium p-value less than or equal to 5×10^{-8} (10^{-16} for single variant association tests). Details on variant calling and quality control is described for each cohort in the [S3 Table](#).

Single variant association tests

We performed single variant association tests using a GWAS additive model framework, with the following covariates: age, age², sex, age*sex, age²*sex, 10 genetic principal components obtained from common genetic variants (MAF>1%). Each cohort performed their analyses separately for each genetic ancestry, but also restricted their variants to those with MAF>0.1% and MAC>6. Summary statistics were then meta-analyzed using a fixed effect model within each ancestry and using a DerSimonian-Laird random effect model across ancestries with the Metal package [55] and its random effect extension [56]. Lastly, given that multiple technologies were used for sequencing, and that whole-exome sequencing can provide variant calls of worse quality in its off-target regions [57], we used the UKB, GHS, and Penn Medicine whole-exome sequencing variants as our "reference panel" for whole-exome sequencing. Hence, only variants reported in at least one of these biobanks were used in the final single-variant analyses.

Variant exclusion list

For the burden tests, we also compiled a list of variants that had a MAF $> 1\%$ or $> 0.1\%$ in any of the participating cohorts. This list was used to filter out variants that were less likely to have a true deleterious effect on COVID-19, even if they were considered rare in other cohorts, or in reference panels [25]. We created two such variant exclusion lists: one to be used in our burden test with variants of MAF less than 1%, and the other for the analysis with MAF less than 0.1%. In any cohort, if a variant had a minor allele count of 6 or more, and a MAF of more than 1% (or 0.1%), this variant was added to our exclusion list. This list was then shared with all participating cohorts, and all variants contained were removed from our burden tests.

Gene burden tests

The following analyses generally followed the methods used by recent literature on large-scale whole-exome sequencing [22] and the COVID-19 HGI [8].

The burden tests were performed by pooling variants in three different variant sets (called masks), as described in recent UK Biobank whole-exome sequencing papers by Backman *et al.* [22] and Kosmicki *et al.* [17]: “M1” which included loss of functions as defined by high impact variants in the Ensembl database [23] (i.e. transcript ablation, splice acceptor variant, splice donor variant, stop gained, frameshift variant, stop lost, start lost, transcript amplification), “M3” which included all variants in M1 as well as moderate impact indels and any missense variants that was predicted to be deleterious based on all of the *in-silico* pathogenicity prediction scores used, and “M4” which included all variants in M3 as well as all missense variants that were predicted to be deleterious in at least one of the *in-silico* pathogenicity prediction scores used. For *in-silico* prediction, we used the following five tools: SIFT [58], LRT [59], MutationTaster [60], PolyPhen2 [61] with the HDIV database, and PolyPhen2 with the HVAR database. Protein coding variants were collapsed on canonical gene transcripts.

Once variants were collapsed into genes in each participant, for each mask, genes were given a score of 0 if the participant had no variants in the mask, a score of 1 if the participant had one or more heterozygous variant in this mask, and a score of 2 if the participant had one or more homozygous variant in this mask. These scores were used as regressors in logistic regression models for the three COVID-19 outcomes above. These regressions were also adjusted for age, age², sex, age*sex, age²*sex, 10 genetic principal components obtained from common genetic variants (MAF $>1\%$), and 20 genetic principal components obtained from rare genetic variants (MAF $<1\%$). The Regenie software [18] was used to perform all burden tests, and generate the scores above. Regenie uses Firth penalized likelihood to adjust for rare or unbalanced events, providing unbiased effect estimates.

All analyses were performed separately for each of six genetic ancestries (African, Admixed American, East Asian, European, Middle Eastern, and South Asian). Summary statistics were meta-analyzed as for the single variant analysis. Participant assignment to genetic ancestry was done locally by each cohort, more details on the methods can be found in the [S3 Table](#).

Lastly, we used ACAT [35] to meta-analyze p-values across masks, within each phenotype separately. ACAT is not affected by lack of independence between tests. These values were used to draw Manhattan and QQ plots in [Fig 2](#).

Supporting information

S1 Table. Acknowledgments per cohort.

(XLSX)

S2 Table. Members of DeCOI, GEN-COVID (Italy), GEN-COVID (Spain), Mount Sinai Biobank, COVID-19 Host Genetics Initiative.

(XLSX)

S3 Table. Cohorts demographics and sequencing details.

(XLSX)

S4 Table. Inflation factors.

(XLSX)

S5 Table. Single variant association tests significant results.

(XLSX)

S6 Table. Full Burden test results.

(XLSX)

S7 Table. IFN related genes burden test results.

(XLSX)

S8 Table. GWAS loci burden test results.

(XLSX)

S9 Table. Additional information on the Genentech cohort.

(XLSX)

S1 Fig. QQ plot and Manhattan plot for the exome-wide single variant association studies.

(DOCX)

S2 Fig. Chromosome 3 exome single variant association studies result by cohort for the severe disease phenotype. Note that all ancestries are shown together. Black dashed line represents the nominal statistical significance threshold ($p = 0.05$).

(DOCX)

S3 Fig. Chromosome 3 exome single variant association studies result by cohort for the severe hospitalized phenotype. Note that all ancestries are shown together. Black dashed line represents the nominal statistical significance threshold ($p = 0.05$)

(DOCX)

S4 Fig. Chromosome 3 exome single variant association studies result by cohort for the susceptibility phenotype. Note that all ancestries are shown together. Black dashed line represents the nominal statistical significance threshold ($p = 0.05$).

(DOCX)

S5 Fig. Single variant association study results at the three novel loci (OR and 95% CI).

(DOCX)

S6 Fig. QQ plot from exome burden test ACAT meta-analyses.

(DOCX)

S7 Fig. *TLR7* beta coefficients (on logistic scale) with 95% interval for the severe COVID-19 phenotype, $MAF < 0.1\%$. X-axes are cut at -10 and 10.

(DOCX)

S8 Fig. *TLR7* beta coefficients (on logistic scale) with 95% interval for the severe COVID-19 phenotype, $MAF < 1\%$. X-axes are cut at -10 and 10.

(DOCX)

S9 Fig. *MARK1* beta coefficients (on logistic scale) with 95% interval for the severe COVID-19 phenotype, MAF<0.1%. X-axes are cut at -10 and 10.
(DOCX)

S10 Fig. *MARK1* beta coefficients (on logistic scale) with 95% interval for the severe COVID-19 phenotype, MAF<1%. X-axes are cut at -10 and 10.
(DOCX)

S11 Fig. *MARK1* beta coefficients (on logistic scale) with 95% interval for the hospitalized COVID-19 phenotype, MAF<0.1%. X-axes are cut at -10 and 10.
(DOCX)

S12 Fig. *MARK1* beta coefficients (on logistic scale) with 95% interval for the hospitalized COVID-19 phenotype, MAF<1%. X-axes are cut at -10 and 10.
(DOCX)

S13 Fig. Ancestry stratified results for *TLR7*, *MARK1*, and *ABO*. X-axis cut at 50. Figures show odds ratios and 95% confidence intervals.
(DOCX)

Acknowledgments

We thank the patients who volunteered to all participating cohorts, and the researchers and clinicians who enrolled them into the respective studies. A full list of acknowledgments can be found in [S1](#) and [S2](#) Tables.

Author Contributions

Conceptualization: Guillaume Butler-Laporte, Jack A. Kosmicki, Elizabeth T. Cirulli, Theodore Drivas, Simone Furini, Chadi Saad, Axel Schmidt, Pawel Olszewski, Mathieu Quinodoz, Elifnaz Çelik, Kousik Kundu, Kumar Veerapen, Tomoko Nakanishi, Malak S. Abedalthagafi, Hugo Zeberg, Stephan Ossowski, Kerstin U. Ludwig, Eva C. Schulte, Hamdi Mbarek, David B. Goldstein, Krzysztof Kiryluk, Alessandra Renieri, Manuel A. R. Ferreira, J Brent Richards.

Data curation: Guillaume Butler-Laporte, Gundula Povysil, Jack A. Kosmicki, Elizabeth T. Cirulli, Theodore Drivas, Simone Furini, Chadi Saad, Axel Schmidt, Pawel Olszewski, Mathieu Quinodoz, Elifnaz Çelik, Kousik Kundu, Klaudia Walter, Junghyun Jung, Amy D. Stockwell, Laura G. Sloofman, Bartłomiej Przychodzen, Timothy Chang, Yosuke Kawai, Yukinori Okada, Kerstin U. Ludwig.

Formal analysis: Guillaume Butler-Laporte, Gundula Povysil, Jack A. Kosmicki, Elizabeth T. Cirulli, Theodore Drivas, Simone Furini, Chadi Saad, Axel Schmidt, Pawel Olszewski, Mathieu Quinodoz, Elifnaz Çelik, Kousik Kundu, Klaudia Walter, Junghyun Jung, Amy D. Stockwell, Laura G. Sloofman, Bartłomiej Przychodzen, Timothy Chang, Tess D. Pottinger, Alex Stuckey, Yosuke Kawai, Yukinori Okada, Alessandra Renieri.

Investigation: Guillaume Butler-Laporte, Gundula Povysil, Jack A. Kosmicki, Elizabeth T. Cirulli, Theodore Drivas, Simone Furini, Chadi Saad, Axel Schmidt, Pawel Olszewski, Urszula Korotko, Mathieu Quinodoz, Elifnaz Çelik, Kousik Kundu, Klaudia Walter, Junghyun Jung, Amy D. Stockwell, Laura G. Sloofman, Daniel M. Jordan, Ryan C. Thompson, Diane Del Valle, Nicole Simons, Esther Cheng, Robert Sebra, Eric E. Schadt, Seunghee Kim-Schulze, Sacha Gnjatic, Miriam Merad, Joseph D. Buxbaum, Noam D. Beckmann,

Alexander W. Charney, Bartłomiej Przychodzen, Timothy Chang, Tess D. Pottinger, Ning Shang, Fabian Brand, Francesca Fava, Francesca Mari, Karolina Chwialkowska, Magdalena Niemira, Szymon Pula, J Kenneth Baillie, Alex Stuckey, Antonio Salas, Xabier Bello, Jacobo Pardo-Seco, Alberto Gómez-Carballa, Irene Rivero-Calle, Federico Martínón-Torres, Andrea Ganna, Konrad J. Karczewski, Kumar Veerapen, Mathieu Bourgey, Guillaume Bourque, Robert JM Eveleigh, Vincenzo Forgetta, David Morrison, David Langlais, Mark Lathrop, Vincent Mooser, Tomoko Nakanishi, Robert Frithiof, Michael Hultström, Miklos Lipcsey, Yanara Marincevic-Zuniga, Jessica Nordlund, Kelly M. Schiabor Barrett, William Lee, Alexandre Bolze, Simon White, Stephen Riffle, Francisco Tanudjaja, Efen Sandoval, Iva Neveux, Shaun Dabe, Nicolas Casadei, Susanne Motameny, Manal Alaamery, Salam Massadeh, Nora Aljawini, Mansour S. Almutairi, Yaseen M. Arabi, Saleh A. Alqahtani, Fawz S. Al Harthi, Amal Almutairi, Fatima Alqubaishi, Sarah Alotaibi, Albandari Binowayn, Ebtehal A. Alsolm, Hadeel El Bardisy, Mohammad Fawzy, Fang Cai, Nicole Soranzo, Adam Butterworth, Daniel H. Geschwind, Stephanie Arteaga, Alexis Stephens, Manish J. Butte, Paul C. Boutros, Takafumi N. Yamaguchi, Shu Tao, Stefan Eng, Timothy Sanders, Paul J. Tung, Michael E. Broudy, Yu Pan, Alfredo Gonzalez, Nikhil Chavan, Ruth Johnson, Bogdan Pasaniuc, Brian Yaspan, Sandra Smieszek, Carlo Rivolta, Stephanie Bibert, Pierre-Yves Bochud, Maciej Dabrowski, Pawel Zawadzki, Mateusz Sypniewski, Elżbieta Kaja, Pajaree Chariyavilaskul, Voraphoj Nilaratanakul, Nattiya Hirankarn, Vorasuk Shotelersuk, Monnat Pongpanich, Chureerat Phokaew, Wanna Chetruengchai, Katsushi Tokunaga, Masaya Sugiyama, Yosuke Kawai, Takanori Hasegawa, Tatsuhiko Naito, Ho Namkoong, Ryuya Edahiro, Akinori Kimura, Seishi Ogawa, Takanori Kanai, Koichi Fukunaga, Yukinori Okada, Seiya Imoto, Satoru Miyano, Serghei Mangul, Malak S. Abedalthagafi, Hugo Zeberg, Joseph J. Grzymalski, Nicole L. Washington, Stephan Ossowski, Kerstin U. Ludwig, Eva C. Schulte, Olaf Riess, Marcin Moniuszko, Mirosław Kwasniewski, Hamdi Mbarek, Said I. Ismail, Anurag Verma, David B. Goldstein, Krzysztof Kiryluk, Alessandra Renieri, Manuel A. R. Ferreira.

Methodology: Gundula Povysil.

Project administration: Guillaume Butler-Laporte, Francesca Fava, Francesca Mari, Andrea Ganna, David B. Goldstein, Krzysztof Kiryluk, Alessandra Renieri, Manuel A. R. Ferreira, J Brent Richards.

Resources: Elizabeth T. Cirulli, Klaudia Walter, Alexander W. Charney, Francesca Mari, J Kenneth Baillie, Mathieu Bourgey, Guillaume Bourque, Robert JM Eveleigh, Vincenzo Forgetta, David Morrison, Mark Lathrop, Vincent Mooser, Robert Frithiof, Michael Hultström, Miklos Lipcsey, Bogdan Pasaniuc, Brian Yaspan, Sandra Smieszek, Carlo Rivolta, Stephanie Bibert, Pierre-Yves Bochud, Elżbieta Kaja, Yosuke Kawai, Yukinori Okada, Serghei Mangul, Malak S. Abedalthagafi, Hugo Zeberg, Stephan Ossowski, Kerstin U. Ludwig, Eva C. Schulte, Olaf Riess, Mirosław Kwasniewski, Hamdi Mbarek, Said I. Ismail, Anurag Verma, David B. Goldstein, Krzysztof Kiryluk, Alessandra Renieri, Manuel A. R. Ferreira, J Brent Richards.

Supervision: Elizabeth T. Cirulli, Klaudia Walter, Alexander W. Charney, Francesca Fava, Francesca Mari, J Kenneth Baillie, Vincent Mooser, Robert Frithiof, Michael Hultström, Miklos Lipcsey, Bogdan Pasaniuc, Brian Yaspan, Sandra Smieszek, Carlo Rivolta, Stephanie Bibert, Pierre-Yves Bochud, Elżbieta Kaja, Yosuke Kawai, Yukinori Okada, Serghei Mangul, Malak S. Abedalthagafi, Hugo Zeberg, Stephan Ossowski, Kerstin U. Ludwig, Eva C. Schulte, Olaf Riess, Mirosław Kwasniewski, Hamdi Mbarek, Said I. Ismail, Anurag Verma,

David B. Goldstein, Krzysztof Kiryluk, Alessandra Renieri, Manuel A. R. Ferreira, J Brent Richards.

Visualization: Guillaume Butler-Laporte.

Writing – original draft: Guillaume Butler-Laporte.

Writing – review & editing: Guillaume Butler-Laporte, Gundula Povysil, Jack A. Kosmicki, Elizabeth T. Cirulli, Theodore Drivas, Simone Furini, Chadi Saad, Axel Schmidt, Pawel Olszewski, Urszula Korotko, Mathieu Quinodoz, Elifnaz Çelik, Kousik Kundu, Klaudia Walter, Junghyun Jung, Amy D. Stockwell, Laura G. Sloofman, Daniel M. Jordan, Ryan C. Thompson, Diane Del Valle, Nicole Simons, Esther Cheng, Robert Sebra, Eric E. Schadt, Seunghee Kim-Schulze, Sacha Gnjatich, Miriam Merad, Joseph D. Buxbaum, Noam D. Beckmann, Alexander W. Charney, Bartłomiej Przychodzen, Timothy Chang, Tess D. Pottinger, Ning Shang, Fabian Brand, Francesca Fava, Francesca Mari, Karolina Chwialkowska, Magdalena Niemira, Szymon Pula, J Kenneth Baillie, Alex Stuckey, Antonio Salas, Xabier Bello, Jacobo Pardo-Seco, Alberto Gómez-Carballa, Irene Rivero-Calle, Federico Martínón-Torres, Andrea Ganna, Konrad J. Karczewski, Kumar Veerapen, Mathieu Bourgey, Guillaume Bourque, Robert JM Eveleigh, Vincenzo Forgetta, David Morrison, David Langlais, Mark Lathrop, Vincent Mooser, Tomoko Nakanishi, Robert Frithiof, Michael Hultström, Miklos Lipcsey, Yanara Marinovic-Zuniga, Jessica Nordlund, Kelly M. Schiabor Barrett, William Lee, Alexandre Bolze, Simon White, Stephen Riffle, Francisco Tanudjaja, Efren Sandoval, Iva Neveux, Shaun Dabe, Nicolas Casadei, Susanne Motameny, Manal Alaamery, Salam Massadeh, Nora Aljawini, Mansour S. Almutairi, Yaseen M. Arabi, Saleh A. Alqahtani, Fawz S. Al Harthi, Amal Almutairi, Fatima Alqubaishi, Sarah Alotaibi, Albandari Binowayn, Ebtehal A. Alsolm, Hadeel El Bardisy, Mohammad Fawzy, Fang Cai, Nicole Soranzo, Adam Butterworth, Daniel H. Geschwind, Stephanie Arteaga, Alexis Stephens, Manish J. Butte, Paul C. Boutros, Takafumi N. Yamaguchi, Shu Tao, Stefan Eng, Timothy Sanders, Paul J. Tung, Michael E. Broudy, Yu Pan, Alfredo Gonzalez, Nikhil Chavan, Ruth Johnson, Bogdan Pasaniuc, Brian Yaspan, Sandra Smieszek, Carlo Rivolta, Stephanie Bibert, Pierre-Yves Bochud, Maciej Dabrowski, Pawel Zawadzki, Mateusz Sypniewski, Elżbieta Kaja, Pajaree Chariyavilaskul, Voraphoj Nilaratanakul, Nattiya Hirankarn, Vorasuk Shotelersuk, Monnat Pongpanich, Chureerat Phokaew, Wanna Chetruengchai, Katsushi Tokunaga, Masaya Sugiyama, Yosuke Kawai, Takanori Hasegawa, Tatsuhiko Naito, Ho Namkoong, Ryuya Edahiro, Akinori Kimura, Seishi Ogawa, Takanori Kanai, Koichi Fukunaga, Yukinori Okada, Seiya Imoto, Satoru Miyano, Serghei Mangul, Malak S. Abedalthagafi, Hugo Zeberg, Joseph J. Grzymalski, Nicole L. Washington, Stephan Ossowski, Kerstin U. Ludwig, Eva C. Schulte, Olaf Riess, Marcin Moniuszko, Mirosław Kwasniewski, Hamdi Mbarek, Said I. Ismail, Anurag Verma, David B. Goldstein, Krzysztof Kiryluk, Alessandra Renieri, Manuel A. R. Ferreira, J Brent Richards.

References

1. Cai Y, Kwek S, Tang SSL, Saffari SE, Lum E, Yoon S, et al. Impact of the COVID-19 pandemic on a tertiary care public hospital in Singapore: Resources and economic costs. *J Hosp Infect.* 2021. <https://doi.org/10.1016/j.jhin.2021.12.007> PMID: 34902499
2. Mulholland RH, Wood R, Stagg HR, Fischbacher C, Villacampa J, Simpson CR, et al. Impact of COVID-19 on accident and emergency attendances and emergency and planned hospital admissions in Scotland: an interrupted time-series analysis. *J R Soc Med.* 2020; 113: 444–453. <https://doi.org/10.1177/0141076820962447> PMID: 33012218
3. Huang C, Wang Y, Li X, Ren L, Zhao J, Hu Y, et al. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet.* 2020; 395: 497–506. [https://doi.org/10.1016/S0140-6736\(20\)30183-5](https://doi.org/10.1016/S0140-6736(20)30183-5) PMID: 31986264

4. Nakanishi T, Pigazzini S, Degenhardt F, Cordioli M, Butler-Laporte G, Maya-Miles D, et al. Age-dependent impact of the major common genetic risk factor for COVID-19 on severity and mortality. *J Clin Invest.* 2021; 131: e152386. <https://doi.org/10.1172/JCI152386> PMID: 34597274
5. Zhou S, Butler-Laporte G, Nakanishi T, Morrison DR, Afilalo J, Afilalo M, et al. A Neanderthal OAS1 isoform protects individuals of European ancestry against COVID-19 susceptibility and severity. *Nat Med.* 2021; 27: 659–667. <https://doi.org/10.1038/s41591-021-01281-1> PMID: 33633408
6. Gaziano L, Giambartolomei C, Pereira AC, Gaulton A, Posner DC, Swanson SA, et al. Actionable drug-gable genome-wide Mendelian randomization identifies repurposing opportunities for COVID-19. *Nat Med.* 2021; 27: 668–676. <https://doi.org/10.1038/s41591-021-01310-z> PMID: 33837377
7. COVID-19 Host Genetics Initiative. A first update on mapping the human genetic architecture of COVID-19. *Nature.* 2022 Aug; 608(7921):E1–E10. Epub 2022 Aug 3. <https://doi.org/10.1038/s41586-022-04826-7>. PMID: 35922517; PMCID: PMC9352569.
8. COVID-19 Host Genetics Initiative, Niemi MEK, Karjalainen J, Liao RG, Neale BM, Daly M, et al. Mapping the human genetic architecture of COVID-19. *Nature.* 2021. <https://doi.org/10.1038/s41586-021-03767-x> PMID: 34237774
9. Ellinghaus D, Degenhardt F, Bujanda L, Buti M, Albillos A, Invernizzi P, et al. Genomewide Association Study of Severe Covid-19 with Respiratory Failure. *N Engl J Med.* 2020; 383: 1522–1534. <https://doi.org/10.1056/NEJMoa2020283> PMID: 32558485
10. Pairo-Castineira E, Clohisey S, Klaric L, Bretherick AD, Rawlik K, Pasko D, et al. Genetic mechanisms of critical illness in COVID-19. *Nature.* 2021; 591: 92–98. <https://doi.org/10.1038/s41586-020-03065-y> PMID: 33307546
11. Kousathanas A, Pairo-Castineira E, Rawlik K, Stuckey A, Odhams CA, Walker S, et al. Whole genome sequencing reveals host factors underlying critical Covid-19. *Nature.* 2022. <https://doi.org/10.1038/s41586-022-04576-6> PMID: 35255492
12. Tam V, Patel N, Turcotte M, Bossé Y, Paré G, Meyre D. Benefits and limitations of genome-wide association studies. *Nat Rev Genet.* 2019; 20: 467–484. <https://doi.org/10.1038/s41576-019-0127-1> PMID: 31068683
13. Taliun D, Harris DN, Kessler MD, Carlson J, Szpiech ZA, Torres R, et al. Sequencing of 53,831 diverse genomes from the NHLBI TOPMed Program. *Nature.* 2021; 590: 290–299. <https://doi.org/10.1038/s41586-021-03205-y> PMID: 33568819
14. Ganna A, Satterstrom FK, Zekavat SM, Das I, Kurki MI, Churchhouse C, et al. Quantifying the Impact of Rare and Ultra-rare Coding Variation across the Phenotypic Spectrum. *Am J Hum Genet.* 2018; 102: 1204–1211. <https://doi.org/10.1016/j.ajhg.2018.05.002> PMID: 29861106
15. Qian Z, Paul B, Zhiyong L, Jérémie LP, Marcela M-V, Jie C, et al. Inborn errors of type I IFN immunity in patients with life-threatening COVID-19. *Science (80-).* 2020; 370: eabd4570. <https://doi.org/10.1126/science.abd4570> PMID: 32972995
16. Povysil G, Butler-Laporte G, Shang N, Wang C, Khan A, Alaamery M, et al. Rare loss-of-function variants in type I IFN immunity genes are not associated with severe COVID-19. *J Clin Invest.* 2021; 131. <https://doi.org/10.1172/JCI147834> PMID: 34043590
17. Kosmicki JA, Horowitz JE, Banerjee N, Lanche R, Marcketta A, Maxwell E, et al. Pan-ancestry exome-wide association analyses of COVID-19 outcomes in 586,157 individuals. *Am J Hum Genet.* 2021; 108: 1350–1355. <https://doi.org/10.1016/j.ajhg.2021.05.017> PMID: 34115965
18. Mbatchou J, Barnard L, Backman J, Marcketta A, Kosmicki JA, Ziyatdinov A, et al. Computationally efficient whole-genome regression for quantitative and binary traits. *Nat Genet.* 2021; 53: 1097–1103. <https://doi.org/10.1038/s41588-021-00870-7> PMID: 34017140
19. Wang X. Firth logistic regression for rare variant association tests. *Frontiers in Genetics.* 2014. Available: <https://www.frontiersin.org/article/10.3389/fgene.2014.00187> PMID: 24995013
20. Zeberg H, Pääbo S. The major genetic risk factor for severe COVID-19 is inherited from Neanderthals. *Nature.* 2020; 587: 610–612. <https://doi.org/10.1038/s41586-020-2818-3> PMID: 32998156
21. Cirulli ET. The Increasing Importance of Gene-Based Analyses. *PLOS Genet.* 2016; 12: e1005852. Available: <https://doi.org/10.1371/journal.pgen.1005852> PMID: 27055023
22. Backman JD, Li AH, Marcketta A, Sun D, Mbatchou J, Kessler MD, et al. Exome sequencing and analysis of 454,787 UK Biobank participants. *Nature.* 2021; 599: 628–634. <https://doi.org/10.1038/s41586-021-04103-z> PMID: 34662886
23. Howe KL, Achuthan P, Allen J, Allen J, Alvarez-Jarreta J, Amode MR, et al. Ensembl 2021. *Nucleic Acids Res.* 2021; 49: D884–D891. <https://doi.org/10.1093/nar/gkaa942> PMID: 33137190
24. Karczewski KJ, Francioli LC, Tiao G, Cummings BB, Alföldi J, Wang Q, et al. The mutational constraint spectrum quantified from variation in 141,456 humans. *Nature.* 2020; 581: 434–443. <https://doi.org/10.1038/s41586-020-2308-7> PMID: 32461654

25. Maffucci P, Bigio B, Rapaport F, Cobat A, Borghesi A, Lopez M, et al. Blacklisting variants common in private cohorts but not in public databases optimizes human exome analysis. *Proc Natl Acad Sci*. 2019; 116: 950 LP–959. <https://doi.org/10.1073/pnas.1808403116> PMID: 30591557
26. Mathieson I, McVean G. Differential confounding of rare and common variants in spatially structured populations. *Nat Genet*. 2012; 44: 243–246. <https://doi.org/10.1038/ng.1074> PMID: 22306651
27. Petes C, Odoardi N, Gee K. The Toll for Trafficking: Toll-Like Receptor 7 Delivery to the Endosome. *Frontiers in Immunology*. 2017. p. 1075. Available: <https://www.frontiersin.org/article/10.3389/fimmu.2017.01075> PMID: 28928743
28. van der Made CI, Simons A, Schuurs-Hoeijmakers J, van den Heuvel G, Mantere T, Kersten S, et al. Presence of Genetic Variants Among Young Men With Severe COVID-19. *JAMA*. 2020; 324: 663–673. <https://doi.org/10.1001/jama.2020.13719> PMID: 32706371
29. Fallerini C, Daga S, Mantovani S, Benetti E, Picchiotti N, Francisci D, et al. Association of Toll-like receptor 7 variants with life-threatening COVID-19 disease in males: findings from a nested case-control study. van de Veerdonk FL, van der Meer JWM, editors. *Elife*. 2021; 10: e67569. <https://doi.org/10.7554/eLife.67569> PMID: 33650967
30. Asano T, Boisson B, Onodi F, Matuozzo D, Moncada-Velez M, Maglorius Renkilaraj MRL, et al. X-linked recessive TLR7 deficiency in ~1% of men under 60 years old with life-threatening COVID-19. *Sci Immunol*. 2021; 6. <https://doi.org/10.1126/sciimmunol.abl4348> PMID: 34413140
31. Mantovani S, Daga S, Fallerini C, Baldassarri M, Benetti E, Picchiotti N, et al. Rare variants in Toll-like receptor 7 results in functional impairment and downregulation of cytokine-mediated signaling in COVID-19 patients. *Genes Immun*. 2022; 23: 51–56. <https://doi.org/10.1038/s41435-021-00157-1> PMID: 34952932
32. Drewes G, Ebneth A, Preuss U, Mandelkow EM, Mandelkow E. MARK, a novel family of protein kinases that phosphorylate microtubule-associated proteins and trigger microtubule disruption. *Cell*. 1997; 89: 297–308. [https://doi.org/10.1016/s0092-8674\(00\)80208-1](https://doi.org/10.1016/s0092-8674(00)80208-1) PMID: 9108484
33. Gordon DE, Jang GM, Bouhaddou M, Xu J, Obernier K, White KM, et al. A SARS-CoV-2 protein interaction map reveals targets for drug repurposing. *Nature*. 2020; 583: 459–468. <https://doi.org/10.1038/s41586-020-2286-9> PMID: 32353859
34. Schaub JR, Stearns T. The Rilp-like proteins Rilp1 and Rilp2 regulate ciliary membrane content. *Mol Biol Cell*. 2013; 24: 453–464. <https://doi.org/10.1091/mbc.E12-08-0598> PMID: 23264467
35. Liu Y, Chen S, Li Z, Morrison AC, Boerwinkle E, Lin X. ACAT: A Fast and Powerful p Value Combination Method for Rare-Variant Analysis in Sequencing Studies. *Am J Hum Genet*. 2019; 104: 410–421. <https://doi.org/10.1016/j.ajhg.2019.01.002> PMID: 30849328
36. Ying Y, Hong X, Xu X, Chen S, He J, Zhu F, et al. Molecular Basis of ABO Variants Including Identification of 16 Novel ABO Subgroup Alleles in Chinese Han Population. *Transfus Med Hemotherapy*. 2020; 47: 160–166. <https://doi.org/10.1159/000501862> PMID: 32355476
37. Hult AK, Yazer MH, Jørgensen R, Hellberg Å, Hustinx H, Peyrard T, et al. Weak A phenotypes associated with novel ABO alleles carrying the A2-related 1061C deletion and various missense substitutions. *Transfusion*. 2010; 50: 1471–1486. <https://doi.org/10.1111/j.1537-2995.2010.02670.x> PMID: 20456702
38. Zietz M, Zucker J, Tatonetti NP. Associations between blood type and COVID-19 infection, intubation, and death. *Nat Commun*. 2020; 11: 5761. <https://doi.org/10.1038/s41467-020-19623-x> PMID: 33188185
39. 100,000 Genomes Pilot on Rare-Disease Diagnosis in Health Care—Preliminary Report. *N Engl J Med*. 2021; 385: 1868–1880. <https://doi.org/10.1056/NEJMoa2035790> PMID: 34758253
40. Spiering AE, de Vries TJ. Why Females Do Better: The X Chromosomal TLR7 Gene-Dose Effect in COVID-19. *Frontiers in Immunology*. 2021. Available: <https://www.frontiersin.org/article/10.3389/fimmu.2021.756262> PMID: 34858409
41. Arnold CG, Libby A, Vest A, Hopkinson A, Monte AA. Immune mechanisms associated with sex-based differences in severe COVID-19 clinical outcomes. *Biol Sex Differ*. 2022; 13: 7. <https://doi.org/10.1186/s13293-022-00417-3> PMID: 35246245
42. Solanich X, Vargas-Parra G, van der Made CI, Simons A, Schuurs-Hoeijmakers J, Antolí A, et al. Genetic Screening for TLR7 Variants in Young and Previously Healthy Men With Severe COVID-19. *Front Immunol*. 2021; 12: 719115. <https://doi.org/10.3389/fimmu.2021.719115> PMID: 34367187
43. Simpson C, Yamauchi Y. Microtubules in Influenza Virus Entry and Egress. *Viruses*. 2020. <https://doi.org/10.3390/v12010117> PMID: 31963544
44. RECOVERY Collaborative Group. Tocilizumab in patients admitted to hospital with COVID-19 (RECOVERY): a randomised, controlled, open-label, platform trial. *Lancet (London, England)*. 2021; 397: 1637–1645. [https://doi.org/10.1016/S0140-6736\(21\)00676-0](https://doi.org/10.1016/S0140-6736(21)00676-0) PMID: 33933206

45. Interleukin-6 Receptor Antagonists in Critically Ill Patients with Covid-19. *N Engl J Med.* 2021; 384: 1491–1502. <https://doi.org/10.1056/NEJMoa2100433> PMID: 33631065
46. Roberts GHL, Park DS, Coignet M V, McCurdy SR, Knight SC, Partha R, et al. AncestryDNA COVID-19 Host Genetic Study Identifies Three Novel Loci. *medRxiv.* 2020; 2020.10.06.20205864. <https://doi.org/10.1101/2020.10.06.20205864>
47. Sankararaman S, Obozinski G, Jordan MI, Halperin E. Genomic privacy and limits of individual detection in a pool. *Nat Genet.* 2009; 41: 965–967. <https://doi.org/10.1038/ng.436> PMID: 19701190
48. Lee S, Emond MJ, Bamshad MJ, Barnes KC, Rieder MJ, Nickerson DA, et al. Optimal Unified Approach for Rare-Variant Association Testing with Application to Small-Sample Case-Control Whole-Exome Sequencing Studies. *Am J Hum Genet.* 2012; 91: 224–237. <https://doi.org/10.1016/j.ajhg.2012.06.007> PMID: 22863193
49. Wainschtein P, Jain D, Zheng Z, Aslibekyan S, Becker D, Bi W, et al. Assessing the contribution of rare variants to complex trait heritability from whole-genome sequence data. *Nat Genet.* 2022. <https://doi.org/10.1038/s41588-021-00997-7> PMID: 35256806
50. Warnat-Herresthal S, Schultze H, Shastry KL, Manamohan S, Mukherjee S, Garg V, et al. Swarm Learning for decentralized and confidential clinical machine learning. *Nature.* 2021; 594: 265–270. <https://doi.org/10.1038/s41586-021-03583-3> PMID: 34040261
51. Williamson EJ, Walker AJ, Bhaskaran K, Bacon S, Bates C, Morton CE, et al. Factors associated with COVID-19-related death using OpenSAFELY. *Nature.* 2020; 584: 430–436. <https://doi.org/10.1038/s41586-020-2521-4> PMID: 32640463
52. Cirulli ET, White S, Read RW, Elhanan G, Metcalf WJ, Tanudjaja F, et al. Genome-wide rare variant analysis for thousands of phenotypes in over 70,000 exomes from two cohorts. *Nat Commun.* 2020; 11: 542. <https://doi.org/10.1038/s41467-020-14288-y> PMID: 31992710
53. Mutambudzi M, Niedzwiedz C, Macdonald EB, Leyland A, Mair F, Anderson J, et al. Occupation and risk of severe COVID-19: prospective cohort study of 120 075 UK Biobank participants. *Occup Environ Med.* 2021; 78: 307 LP– 314. <https://doi.org/10.1136/oemed-2020-106731> PMID: 33298533
54. Hail Team. Hail 0.2. 2021. Available: <https://github.com/hail-is/hail>
55. Willer CJ, Li Y, Abecasis GR. METAL: fast and efficient meta-analysis of genomewide association scans. *Bioinformatics.* 2010/07/08. 2010; 26: 2190–2191. <https://doi.org/10.1093/bioinformatics/btq340> PMID: 20616382
56. Hemani G. random-metal. In: GitHub repository [Internet]. 2017 [cited 15 Mar 2022]. Available: <https://github.com/explodecomputer/random-metal>
57. Guo Y, Long J, He J, Li C-I, Cai Q, Shu X-O, et al. Exome sequencing generates high quality data in non-target regions. *BMC Genomics.* 2012; 13: 194. <https://doi.org/10.1186/1471-2164-13-194> PMID: 22607156
58. Vaser R, Adusumalli S, Leng SN, Sikic M, Ng PC. SIFT missense predictions for genomes. *Nat Protoc.* 2016; 11: 1–9. <https://doi.org/10.1038/nprot.2015.123> PMID: 26633127
59. Chun S, Fay JC. Identification of deleterious mutations within three human genomes. *Genome Res.* 2009; 19: 1553–1561. <https://doi.org/10.1101/gr.092619.109> PMID: 19602639
60. Schwarz JM, Rödelsperger C, Schuelke M, Seelow D. MutationTaster evaluates disease-causing potential of sequence alterations. *Nat Methods.* 2010; 7: 575–576. <https://doi.org/10.1038/nmeth0810-575> PMID: 20676075
61. Adzhubei I, Jordan DM, Sunyaev SR. Predicting Functional Effect of Human Missense Mutations Using PolyPhen-2. *Curr Protoc Hum Genet.* 2013; 76: 7.20.1–7.20.41. <https://doi.org/10.1002/0471142905.hg0720s76> PMID: 23315928
62. Buniello A, MacArthur JAL, Cerezo M, Harris LW, Hayhurst J, Malangone C, et al. The NHGRI-EBI GWAS Catalog of published genome-wide association studies, targeted arrays and summary statistics 2019. *Nucleic Acids Res.* 2019, Vol. 47 (Database issue): D1005–D1012.