UC San Diego

UC San Diego Previously Published Works

Title

Intermediate baseline appearance experiments and three-neutrino mixing schemes

Permalink

<https://escholarship.org/uc/item/8282c2pc>

Journal

Physics Letters, Section B: Nuclear, Elementary Particle and High-Energy Physics, 413(3-4)

ISSN

0370-2693

Authors

Cardall, CY
Fuller, GM
Cline, DB

Publication Date

1997-11-13

DOI

10.1016/S0370-2693(97)01057-5

Peer reviewed

ELSEVIER

13 November 1997

PHYSICS LETTERS B

Physics Letters B 413 (1997) 246–252

Intermediate baseline appearance experiments and three-neutrino mixing schemes

Christian Y. Cardall^{a,1}, George M. Fuller^a, David B. Cline^b

^a Department of Physics, University of California, San Diego, La Jolla, CA 92093-0319, USA

^b Physics & Astronomy Department, Box 951547, University of California Los Angeles, Los Angeles, CA 90095-1547, USA

Received 23 June 1997

Editor: W. Haxton

Abstract

Three-neutrino mixing schemes suggested by Cardall & Fuller and Acker & Pakvasa are compared and contrasted. Both of these schemes seek to solve the solar and atmospheric neutrino problems *and* to account for the possible neutrino oscillation signal in the LSND experiment. These neutrino oscillation schemes have different atmospheric and solar neutrino signatures that will be discriminated by Super-Kamiokande and SNO. They will also have different signatures in proposed long-baseline accelerator and reactor experiments. In particular, both of these schemes would give dramatic (and dramatically different) signals in an “intermediate baseline” experiment, such as the proposed ICARUS detector in the Jura mountains 17 km from CERN. © 1997 Elsevier Science B.V.

PACS: 14.60.Pq; 26.65.+t; 95.85.Ry

1. Three-mixing schemes

At the present time there are three hints of neutrino mixing: the solar (e.g. [1]) and atmospheric (e.g. [2]) neutrino problems, and the signal in the LSND experiment [3]. Each of these can be solved by or interpreted as neutrino flavor oscillations. If an independent neutrino mass difference is associated with each of these three hints, four neutrino flavors are necessary. Since it is known that only three neutrino flavors participate in the weak interaction [4], the fourth neutrino flavor must be “sterile” [an SU(2) singlet]. Accounting for all three hints with

only three active neutrino flavors would require that at least two of the phenomena must “share” one of the two available independent mass differences.

The possibility of a three-neutrino mixing scheme along these lines was noted by Cardall & Fuller [5]. Their scheme uses the smaller neutrino mass difference to account for the solar neutrino problem, and the larger mass difference to account for the atmospheric neutrino problem and the LSND signal [6]. They proposed the following mass/mixing parameters:

$$\delta m_{21}^2 \approx 7 \times 10^{-6} \text{ eV}^2, \quad (1)$$

$$\delta m_{31}^2 \approx \delta m_{32}^2 \approx 0.3 \text{ eV}^2, \quad (2)$$

$$U_{\text{CF, sma}} \approx \begin{pmatrix} 0.994 & 0.044 & 0.100 \\ -0.108 & 0.530 & 0.841 \\ -0.015 & -0.847 & 0.532 \end{pmatrix}. \quad (3)$$

¹ Current address: Department of Physics and Astronomy, State University of New York at Stony Brook, Stony Brook NY 11794-3800.

Here $\delta m_{ji}^2 = m_j^2 - m_i^2$, where $m_{i,j}^2$ are the i th and j th squared neutrino mass eigenvalues, and we take $m_j^2 > m_i^2$. The unitary matrix $U_{\text{CF},\text{sma}}$ (where “CF” stands for “Cardall & Fuller” and “sma” stands for “small mixing angle”) connects the neutrino flavor and mass eigenstates: $\nu_\alpha = \sum_i U_{\alpha i} \nu_i$, where $\alpha = e, \mu$, or τ and $i = 1, 2, 3$. These mass/mixing parameters were designed to employ the small mixing angle MSW solution to the solar neutrino problem (see below). However, as can readily be seen from the analysis of Ref. [7], the general framework of the Cardall & Fuller scheme can also accommodate the solar neutrino large mixing angle MSW solution:

$$\delta m_{21}^2 \approx 2 \times 10^{-5} \text{ eV}^2, \quad (4)$$

$$\delta m_{31}^2 \approx \delta m_{32}^2 \approx 0.3 \text{ eV}^2, \quad (5)$$

$$U_{\text{CF},\text{lma}} \approx \begin{pmatrix} 0.873 & 0.478 & 0.100 \\ -0.330 & 0.428 & 0.841 \\ 0.359 & -0.767 & 0.532 \end{pmatrix}, \quad (6)$$

and the solar neutrino “just-so” vacuum oscillation solution:

$$\delta m_{21}^2 \approx 6 \times 10^{-11} \text{ eV}^2, \quad (7)$$

$$\delta m_{31}^2 \approx \delta m_{32}^2 \approx 0.3 \text{ eV}^2, \quad (8)$$

$$U_{\text{CF},\text{vac}} \approx \begin{pmatrix} 0.807 & 0.582 & 0.100 \\ -0.381 & 0.384 & 0.841 \\ 0.451 & -0.717 & 0.532 \end{pmatrix}. \quad (9)$$

Another variation on the Cardall & Fuller scenario has very recently been proposed in Ref. [8].

A rather different scheme has been suggested recently by Acker & Pakvasa [9], in which the smaller neutrino mass difference accounts for both the solar and atmospheric neutrino problems, and the larger mass difference accounts for the LSND signal. They propose the following mass/mixing parameters:

$$\delta m_{21}^2 \approx 10^{-2} \text{ eV}^2, \quad (10)$$

$$\delta m_{31}^2 \approx \delta m_{32}^2 \approx 1\text{--}2 \text{ eV}^2, \quad (11)$$

$$U_{\text{AP}} \approx \begin{pmatrix} 0.700 & 0.700 & 0.140 \\ -0.714 & 0.689 & 0.124 \\ -0.010 & -0.187 & 0.982 \end{pmatrix}. \quad (12)$$

Relatively small alterations of this mixing matrix could also allow this scheme to accommodate slightly

larger or smaller mass differences to account for a significant fraction of the LSND signal range.

2. Consequences for solar neutrino experiments

The solar neutrino problem arises from the observation that the measured solar ν_e fluxes in experiments with chlorine [10], water [11], and gallium [12] detectors are only a fraction of the fluxes expected from models of the sun. Furthermore, the fact that the detectors employing the above substances have different thresholds and that they observe different ν_e flux deficits points to an energy dependence of the ν_e suppression. This energy dependence picks out values of δm^2 that may provide a neutrino mixing resolution to the puzzle (e.g. [7,13]): $\delta m^2 \approx 10^{-5} \text{ eV}^2$ for an MSW effect (matter-enhanced neutrino flavor conversion) solution, with both small and large mixing angle solutions as possibilities; or $\delta m^2 \approx 10^{-10} \text{ eV}^2$ for a “just-so” vacuum oscillation solution, with large mixing angle.

While the Cardall & Fuller schemes employ these standard solar neutrino solutions, the Acker & Pakvasa scheme employs an “energy independent” solar neutrino solution with $\delta m^2 \approx 10^{-2}$ [14,15]. Their motivation is to use the same mass difference as that suggested by the claimed zenith angle dependence of the Kamiokande multi-GeV atmospheric neutrino data (see below). Such an energy independent solution is strongly disfavored if all solar neutrino experiments and the latest standard solar models are used [16], but is marginally possible if the ^8B neutrino flux is significantly (factor of ~ 0.6) smaller than that predicted by the standard solar model. The possibility of a ^8B neutrino flux smaller than that of the SSM has received some motivation from a recent measurement [17] of the reaction $\gamma + ^8\text{B} \rightarrow ^7\text{Be} + \text{p}$, the inverse of a reaction important to the determination of that flux. However, issues surrounding the ability to extract the forward cross section from measurements of the inverse reaction have been controversial [18]. Another possibility for allowing a larger neutrino mass difference than those in the “standard” solar neutrino solutions is to ignore either the chlorine or gallium experiments [15,16].

The current evidence for neutrino oscillations associated with the sun rests simply on an observed

neutrino flux deficit as compared with solar models. Two next-generation solar neutrino experiments, SNO [19] and Super-Kamiokande [20], will allow tests that can provide more conclusive evidence of neutrino mixing. These tests include time variable solar neutrino signals (seasonal for vacuum oscillations [21,22], day-night for the MSW solutions [23,24]) and distortions of the recoil electrons from neutrino interactions in the detector [22,24,25] (relevant for both the MSW solutions and the “just so” vacuum oscillation solution).² These effects will be visible for the Cardall & Fuller schemes since they adopt the standard neutrino oscillation solutions. However, these effects will be absent in the “energy independent” solar neutrino solution of Acker & Pakvasa. This is a major observational difference between the Cardall & Fuller and Acker & Pakvasa mixing schemes.

3. Consequences for atmospheric neutrino experiments

Fluxes of ν_e and ν_μ arise from the interactions of cosmic rays with the earth’s atmosphere. In order to minimize uncertainties associated with neutrino cross sections and a lack of knowledge of the absolute magnitude of the neutrino fluxes, the “ratio of ratios” R is a commonly reported observable:

$$R = \frac{(\nu_\mu/\nu_e)_{\text{data}}}{(\nu_\mu/\nu_e)_{\text{Monte Carlo}}} \quad (13)$$

Studies involving large water detectors (Kamiokande [26] and IMB [27]) have reported values of $R \sim 0.6$. Experiments using iron calorimeter detectors have had mixed results: Soudan 2 [28] has observed a deficit in R comparable to that seen in the water detectors, while NUSEX [29] and Fréjus [30] have not seen evidence for values of R significantly different from unity.

An interesting aspect of the atmospheric neutrino puzzle is the zenith-angle dependence of R reported by the Kamiokande group for multi-GeV neutrinos

[31]. According to their analysis, R decreases from zero zenith angle (neutrinos from the atmosphere immediately above the detector) to maximum zenith angle (neutrinos from the atmosphere on the other side of the earth). This dependence suggests that neutrinos from overhead do not travel far enough to oscillate significantly, while those coming from across the earth do travel sufficiently far to oscillate. Such a scenario restricts neutrino mass differences to the range 10^{-3} – 10^{-1} eV².

A neutrino mass difference of about 10^{-2} eV² has therefore come to be associated with the atmospheric neutrino problem. This is the mass difference employed in the Acker & Pakvasa neutrino oscillation scheme. The mass difference of ~ 0.3 eV² in the Cardall & Fuller scheme requires one to discount the zenith angle dependence of the Kamiokande multi-GeV data. This position is not entirely implausible, as the statistical significance of this effect has been questioned [32], and the IMB group saw no zenith angle dependence in their data [33].

Super-Kamiokande will be able to settle the question of zenith angle dependence definitively. Preliminary analyses already confirm a deficit in R averaged over all directions, but nothing definitive can yet be said about the zenith angle dependence [34]. This is perhaps interesting in view of the fact that Super-Kamiokande already has more events than were obtained with the previous Kamiokande detector.

Another important point is that in the Acker & Pakvasa solution the atmospheric neutrino oscillations are primarily $\nu_\mu \leftrightarrow \nu_e$ to allow a simultaneous solution to the solar neutrino problem, while in the Cardall & Fuller scheme the atmospheric neutrino oscillation channel is primarily $\nu_\mu \leftrightarrow \nu_\tau$ to avoid conflict with ν_e disappearance experiments. Due to uncertainties in the ν_e and ν_μ absolute fluxes, it is not currently possible to determine the oscillation channel of atmospheric neutrinos. However, higher precision cosmic ray measurements may help reduce the ambiguity.

4. Consequences for accelerator and reactor experiments

While solar and atmospheric neutrinos have provided tantalizing suggestions of neutrino mixing, it is

² SNO, with its use of heavy water, will allow the very important measurement of the ratio of charged current/neutral current events for neutrino interactions with deuterium.

highly desirable to directly observe neutrino oscillations in controlled terrestrial experiments. We point out that the neutrino mass difference associated with solar neutrinos is so small that all three of the Cardall & Fuller solutions, corresponding to different solar neutrino solutions, will have the same effects in the terrestrial experiments discussed here.

4.1. Short baseline experiments

The LSND experiment, which has reported an excess of $\bar{\nu}_e$ in a $\bar{\nu}_\mu$ beam from pion decays at rest, continues to run. KARMEN [35], a similar experiment, has recently received an upgrade that should eventually be able to confirm this LSND signal. While the LSND signal is compatible with large neutrino mass differences, negative results from CCFR [36], CHORUS [37], and NOMAD [38] restrict the viable range of neutrino mass difference to $\delta m^2 \leq 10 \text{ eV}^2$. In addition, LSND has recently completed a study of excess ν_e events in a ν_μ beam from pion decays in flight [39], a channel whose backgrounds and systematics are different from those of the $\bar{\nu}_e$ beam from pion decays at rest. This excess can also be accounted for by neutrino oscillations, with mixing parameters that overlap those suggested by the decay at rest data. Finally, COSMOS [40], the proposed short baseline counterpart to the MINOS experiment, should just be able to see $\nu_\mu \leftrightarrow \nu_\tau$ oscillations in both the Acker & Pakvasa and Cardall & Fuller schemes.

4.2. Long baseline experiments

Long baseline experiments are designed to explore the region of neutrino mixing parameter space suggested by the atmospheric neutrino problem, neutrino mass differences of $\sim 10^{-2} \text{ eV}^2$ in particular. Long baseline accelerator experiments such as MINOS (Fermilab to the Soudan mine) [41], KEK to Super-Kamiokande [42], and CERN to Gran Sasso [43] involve a ν_μ beam and should be able to distinguish between $\nu_\mu \leftrightarrow \nu_\tau$ and $\nu_\mu \leftrightarrow \nu_e$ oscillations. Fig. 1 shows the oscillation (flavor conversion) probabilities for these channels implied by the Cardall & Fuller and Acker & Pakvasa schemes for a baseline of 735 km, which coincidentally is the approximate baseline length for both the CERN to Gran

Fig. 1. Probabilities for (a) ν_e and (b) ν_τ appearance in a ν_μ beam, as a function of neutrino energy, for a baseline of 735 km (CERN to Gran Sasso and Fermilab to the Soudan mine). “CF” and “AP” stand for the Cardall & Fuller and Acker & Pakvasa oscillation schemes respectively.

Sasso and MINOS experiments. The differences are evident. The Acker & Pakvasa scheme has large amplitude $\nu_\mu \leftrightarrow \nu_e$ mixing, with oscillations in energy space that could probably be observed experimentally. The Cardall and Fuller scheme has large amplitude $\nu_\mu \leftrightarrow \nu_\tau$ mixing, with the oscillations in energy space being so rapid that in practice an average would probably be measured.

There are also forthcoming long baseline reactor experiments, CHOOZ [44] and Palo Verde [45]. These are $\bar{\nu}_e$ disappearance experiments which will be sensitive to neutrino mass differences as low as $\sim 10^{-3} \text{ eV}^2$, but which will not provide increased sensitivity to the oscillation amplitude. These experiments are therefore sensitive to a $\nu_\mu \leftrightarrow \nu_e$ solution to the atmospheric neutrino problem (which has a very large effective mixing angle). They would see neutrino oscillations predicted by the Acker & Pakvasa

Fig. 2. Probabilities for (a) ν_e and (b) ν_τ appearance in a ν_μ beam, as a function of neutrino energy, for a baseline of 17 km (CERN to the Jura mountains). “CF” and “AP” stand for the Cardall & Fuller and Acker & Pakvasa oscillation schemes respectively.

scheme, but *not* those predicted by the Cardall & Fuller solution.

4.3. The Jura mountain neutrino detector

The ICARUS Collaboration, which is producing neutrino detectors for the CERN to Gran Sasso long

baseline experiment, has recently proposed placing one of their detector modules behind the Jura mountains 17 km from CERN [46]. This “intermediate baseline” experiment is of great interest in light of the three-neutrino mixing schemes discussed here, since the neutrino flight distance divided by the neutrino oscillation length is such that neutrino oscillations should be directly observable in energy space, as illustrated in Fig. 2 [the estimated energy resolution for ICARUS is (5–15)%]. This experiment was proposed with the Acker & Pakvasa scheme in mind, but Fig. 2 shows that the Cardall & Fuller scheme predicts an even more spectacular signal in the ν_τ appearance channel. We emphasize that an intermediate baseline experiment such as this is important to distinguish between the Acker & Pakvasa and Cardall & Fuller schemes, and to directly observe two features of those schemes which are presently only inferred from phenomenological considerations: (1) a neutrino mass difference in the range $0.1 \leq \delta m^2 \leq 1 \text{ eV}^2$, and (2) significant $\nu_\mu \leftrightarrow \nu_\tau$ mixing, which is (at least in part) indirectly responsible for the $\nu_\mu \leftrightarrow \nu_e$ signal observed at LSND.

5. Conclusion

There exist at least two three-neutrino mixing schemes designed to satisfy three hints of neutrino mixing (the solar and atmospheric neutrino problems, and the signal at LSND). Some of their signatures in future experiments are given in Table 1. Accommodating all three of these hints would normally require four neutrino flavors; use of a three generation framework requires ignoring some aspect of the data. Cardall & Fuller have chosen to ignore

Table 1
Some experimental signatures of the Cardall & Fuller and Acker & Pakvasa schemes

	Experimental signature	Cardall & Fuller	Acker & Pakvasa
Solar neutrinos	Energy dependence	Yes	No
	Spectral distortions	Yes	No
	Time dependence	Yes	No
Atmospheric neutrinos	Zenith angle variation	No	Yes
Long baseline reactor	$\bar{\nu}_e$ disappearance	No	Yes
Long baseline accelerator	ν_e appearance	No	Yes (unaveraged)
	ν_τ appearance	Yes (averaged)	No
Intermediate baseline accelerator	ν_e appearance	No	No
	ν_τ appearance	Yes (unaveraged)	Yes (unaveraged)

the zenith angle dependence of the atmospheric neutrino data, while Acker & Pakvasa have chosen to ignore the energy dependence of the solar neutrino data. The former choice has recently been identified in a thorough analysis as the choice of “minimum sacrifice” [47], and the validity of the above assumptions regarding atmospheric and solar neutrinos will eventually be tested by Super-Kamiokande and SNO. Nevertheless, direct experimental verification of the predictions of these schemes in a controlled terrestrial experiment is highly desirable. In particular, an “intermediate baseline” experiment such as the proposed ICARUS detector to be placed behind the Jura mountains 17 km from CERN is important to be able to see, in energy space, unaveraged $\nu_\mu \leftrightarrow \nu_\tau$ oscillations driven by a neutrino mass difference of $0.1 \leq \delta m^2 \leq 1 \text{ eV}^2$. Further impetus for the careful exploration of this range of neutrino mass difference also derives from the fact that such neutrino mass/mixing parameters may make possible successful r -process nucleosynthesis in supernovae [48].

6. Note added

Another “intermediate baseline” baseline experiment – BooNE (Booster Neutrino Experiment) – is being proposed [49]. BooNE would use a neutrino beam derived from the 8 GeV proton booster at FNAL, and in its final phase should be able to measure a neutrino mass difference in the range $0.1 \leq \delta m^2 \leq 1 \text{ eV}^2$.

Acknowledgements

We thank M. Goodman and W. Haxton for helpful communications. This work was supported by grant NSF PHY95-03384 at UCSD.

References

- [1] J.N. Bahcall, *Astrophys. J.* 467 (1996) 475; W. Haxton, *Ann. Rev. Astron. Astrophys.* 33 (1995) 459.
- [2] T.K. Gaisser, F. Halzen, T. Stanev, *Phys. Rep.* 258 (1995) 173.
- [3] C. Athanassopoulos et al. (LSND Collaboration), *Phys. Rev. Lett.* 75 (1995) 2650; 77 (1996) 3082.
- [4] D. Buskulic et al. (ALEPH Collaboration), *Z. Phys. C* 60 (1993) 71; B. Adeva et al. (L3 Collaboration), *Phys. Lett. B* 237 (1990) 136.
- [5] C.Y. Cardall, G.M. Fuller, *Phys. Rev. D* 53 (1996) 4421.
- [6] That recognition that, in: a three-neutrino mixing scheme, $\nu_e \rightarrow \nu_\mu$ flavor conversion could occur at separate scales of δm^2 relevant to solar neutrinos and accelerator experiments was expressed by K.S. Babu, J.C. Pati, F. Wilcek, *Phys. Lett. B* 359 (1995) 351. However, that work did not attempt to include a solution to the atmospheric neutrino anomaly.
- [7] G.L. Fogli, E. Lisi, D. Montanino, *Phys. Rev. D* 54 (1996) 2048.
- [8] E. Ma, P. Roy, preprint TIFR/TH/97-12, UCRHEP-T186, hep-ph/9706309 (1997).
- [9] A. Acker, S. Pakvasa, *Phys. Lett. B* 397 (1997) 209.
- [10] B.T. Cleveland et al., *Nucl. Phys. B (Proc. Suppl.)* 31 (1995) 47; R. Davis, *Prog. Part. Nucl. Phys.* 32 (1994) 13.
- [11] Y. Fukuda et al. (Kamiokande Collaboration), *Phys. Rev. Lett.* 77 (1996) 1683.
- [12] W. Hampel et al. (GALLEX Collaboration), *Phys. Lett. B* 388 (1996) 384; J.N. Abdurashitov et al. (SAGE Collaboration), *Phys. Rev. Lett.* 77 (1996) 4708.
- [13] N. Hata, P. Langacker, preprint IASSNS-AST 97/29, UPR-751T, hep-ph/9705339 (1997); J.N. Bahcall, P.I. Krastev, *Phys. Rev. D* 53 (1996) 4211.
- [14] A. Acker, J.G. Learned, S. Pakvasa, T.J. Weiler, *Phys. Lett. B* 298 (1993) 149; G. Conforto et al., *Astropart. Phys.* 5 (1996) 147.
- [15] P.F. Harrison, D.H. Perkins, W.G. Scott, *Phys. Lett. B* 349 (1995) 137.
- [16] P.I. Krastev, S.T. Petcov, preprint IASSNS-AST 96/58, SISSA 166/96/EP, hep-ph/9612243 (1996).
- [17] T. Motobayashi et al., *Phys. Rev. Lett.* 73 (1994) 2680.
- [18] W.C. Haxton, *Nucl. Phys. B (Proc. Suppl.)* 48 (1996) 317; R. Shyam, I.J. Thompson, A.K. Dutt-Mazumder, *Phys. Lett. B* 371 (1996) 1; J. v. Schwarzenberg et al., *Phys. Rev. C* 53 (1996) R2598; T. Kikuchi et al., *Phys. Lett. B* 391 (1997) 261.
- [19] G.T. Ewan et al. (SNO Collaboration), preprint SNO-87-12 (1987).
- [20] Y. Totsuka, *Nucl. Phys. B (Proc. Suppl.)* 48 (1996) 547.
- [21] G.L. Fogli, E. Lisi, D. Montanino, preprint BARI-TH/276-97, hep-ph/9706228, 1997.
- [22] P.I. Krastev, S.T. Petcov, *Nucl. Phys. B* 449 (1995) 605; P.I. Krastev, S.T. Petcov, *Phys. Lett. B* 299 (1993) 99.
- [23] J.N. Bahcall, P.I. Krastev, preprint IASSNS-AST 97/31, hep-ph/9706239, 1997; P.I. Krastev, talk presented at the DPF '96 meeting, preprint hep-ph/9610339, 1996.
- [24] M. Maris, S.T. Petcov, preprint SISSA 17/97/EP, hep-ph/9705392, 1997.
- [25] J.N. Bahcall, P.I. Krastev, E. Lisi, *Phys. Rev. C* 55 (1997) 494.

- [26] K.S. Hirata et al. (Kamiokande Collaboration), *Phys. Lett. B* 280 (1992) 146.
- [27] D. Casper et al. (IMB Collaboration), *Phys. Rev. Lett.* 66 (1991) 2561.
- [28] W.W.M. Allison et al. (Soudan Collaboration), *Phys. Lett. B* 391 (1997) 491.
- [29] M. Aglietta et al. (NUSEX Collaboration), *Europhys. Lett.* 8 (1989) 611.
- [30] Ch. Berger et al. (Fréjus Collaboration), *Phys. Lett. B* 245 (1990) 305.
- [31] Y. Fukuda et al. (Kamiokande Collaboration), *Phys. Lett. B* 335 (1994) 237.
- [32] D. Saltzberg, *Phys. Lett. B* 355 (1994) 237;
G.L. Fogli, E. Lisi, *Phys. Rev. D* 52 (1995) 2775.
- [33] R. Clark et al. (IMB Collaboration), *preprint* (1997).
- [34] T. Haines, talk presented at the INPAC meeting, 2 May 1997, Berkeley, California, USA.
- [35] J. Kleinfeller for the KARMEN Collaboration, *Nucl. Phys. B (Proc. Suppl.)* 51 (1996) 249.
- [36] K.S. McFarland et al. (CCFR Collaboration), *Phys. Rev. Lett.* 75 (1995) 3993.
- [37] D. Saltzberg et al. (CHORUS Collaboration), in: M. Baldo (Ed.), the Proceedings of the 7th International Conference on Neutrino Telescopes, Venice, Italy, 1996, Ceolin, University of Padova, Italy, 1996, p. 91.
- [38] A. Rubbia et al. (NOMAD Collaboration), in: M. Baldo (Ed.), the Proceedings of the 7th International Conference on Neutrino Telescopes, Venice, Italy, 1996, Ceolin, University of Padova, Italy, 1996, p. 101.
- [39] C. Athanassopoulos et al. (LSND Collaboration), *preprint* LA-UR-97-1998, 1997.
- [40] R. Sidwell et al. (COSMOS Collaboration), *preprint* NuMI-S-94, 1995.
- [41] D. Ayres et al. (MINOS Collaboration), *preprint* NuMI-L-62, 1995.
- [42] K. Nishikawa et al., Proposal for a Long Baseline Neutrino Oscillation Experiment, using KEK-PS and Super-Kamiokande, 1995.
- [43] F. Cavanna (for the ICARUS Collaboration), *preprint* CERN-PPE/95-133, 1995.
- [44] H. de Kerret et al. (CHOOZ Collaboration), Proposal to Search for Neutrino Vacuum Oscillations to $\Delta m^2 = 10^{-3} \text{ eV}^2$ Using a 1 km Baseline Reactor Neutrino Experiment, 1993.
- [45] F. Boehm et al. (Palo Verde Collaboration), Review of Progress, The Palo Verde Experiment, 1996.
- [46] A. Rubbia et al. (ICARUS Collaboration), *preprint* CERN/SPSLC 96-58, SPSLC/P 304, 1996.
- [47] G.L. Fogli, E. Lisi, D. Montanino, G. Scioscia, *preprint* BARI-TH/273-97, hep-ph/9706230, 1997.
- [48] G.M. Fuller, Y.-Z. Qian, in preparation, 1997.
- [49] J. Conrad, private communication, 1997.