Lawrence Berkeley National Laboratory

Recent Work

Title

193 nm photodissociation of thiophene Probed using synchrotron radiation

Permalink

<https://escholarship.org/uc/item/8353d76c>

Journal

Journal of Physical Chemistry A, 103(42)

Author

Qi, Fei

Publication Date

1999-06-01

LBLN-43495
Preprint

ERNEST ORLANDO LAWRENCE BERKELEY NATIONAL LABORATORY

193 nm Photodissociation of Thiophene Probed Using Synchrotron Radiation

Fei Qi, Osman Sorkhabi, Abbas H. Rizvi,
and Arthur G. Suits

Chemical Sciences Division

June 1999

Submitted to
*Journal of Physical
Chemistry A*

REFERENCE COPY |
Does Not |
Circulate |
Bldg. 50 Library - Ref.
Lawrence Berkeley National Laboratory

LBLN-43495

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

193 nm Photodissociation of Thiophene Probed Using Synchrotron Radiation

Fei Qi, Osman Sorkhabi, Abbas H. Rizvi,
and Arthur G. Suits

Chemical Sciences Division
Ernest Orlando Lawrence Berkeley National Laboratory
University of California
Berkeley, California 94720

June 1999

This work was supported by the Director, Office of Science, Office of Basic Energy Sciences, Chemical Sciences Division, of the U.S. Department of Energy under Contract No. DE-AC03-76SF00098. The Advanced Light Source is supported by the Director, Office of Science, Office of Basic Energy Sciences, Materials Sciences Division, of the U.S. Department of Energy under the same contract.

193 nm Photodissociation of Thiophene Probed Using Synchrotron Radiation

Fei Qi, Osman Sorkhabi, Abbas H. Rizvi, and Arthur G. Suits*

Chemical Sciences Division

Ernest Orlando Lawrence Berkeley National Laboratory

Berkeley, CA 94720

Abstract

The photodissociation dynamics of thiophene, $c\text{-C}_4\text{H}_4\text{S}$, have been studied at 193nm using tunable synchrotron undulator radiation as a universal product probe. Five primary dissociation channels have been observed, and the translational energy release distributions and photoionization efficiency spectra have been recorded for all products. The evidence suggests that dissociation occurs on the ground state surface following internal conversion and thus have direct relevance to high-temperature pyrolysis of thiophene.

*Corresponding author: agsuits@lbl.gov

I. INTRODUCTION

Although detailed photochemical studies of simple molecules have provided considerable insight into the dynamics of elementary chemical reactions, analogous studies for more complex molecules have been less ardently pursued, largely owing to experimental challenges attending such investigations. The availability of a universal, selective vacuum ultraviolet (VUV) probe on the Chemical Dynamics Beamline¹ has recently made possible new approaches to the study of the photodissociation dynamics of large molecules. In this paper, we extend our recent investigations of the photochemistry of complex cyclic molecules² to include thiophene (Th), a five-membered sulfur-containing heterocyclic aromatic molecule.

Thiophene is one of the principal sulfur carrying compounds in fossil fuels, and the study of its photodissociation can provide insights into the properties of sulfur-containing molecules and radicals of importance in fossil fuel combustion.³ The ultraviolet (UV) photochemistry of thiophene was studied in the gas phase at 213.9, 228.8 and 253.7 nm by Wiebe and Heicklen in the late 1960s using end-product analysis.⁴ The primary photofragments inferred from the experiments included C_2H_2 , $CH_2=C=CH_2$, $CH_3C\equiv CH$, CS_2 , $CH_2=CH-C\equiv CH$, and C_2H_2S . Infrared laser multiphoton excitation/dissociation of thiophene was studied by Nayak et al.⁵ They proposed a mechanism which involved breakage of the C-S bond in thiophene to form an unstable 1,5-diradical which further decomposed via different channels. Krishnamachari and Venkitachalam observed a transient absorption spectrum in the region 377 to 417 nm following flash photolysis of thiophene.⁶ They tentatively assigned this absorption to C_4H_3 . The first direct examination of primary photoproducts of thiophene at 193 nm was carried out by Myers using photofragment translational spectroscopy, and reported in his thesis work but not otherwise published.⁷ Six primary dissociation channels were identified as discussed below. Ng and co-workers examined the isomeric structures of the primary photoproducts formed in the 193 nm photodissociation of thiophene using photodissociation-photoionization detection of the hydrocarbons and 2 + 1 resonance-enhanced multiphoton ionization (REMPI) detection of the sulfur atom. Only two dissociative channels, $C_2H_2S + C_2H_2$ and $C_4H_4 + S$, were observed in their studies and they found that the S atoms were produced predominantly ($\geq 96\%$) in the 3P_J states.⁸

Thiophene is the sulfur containing analog of furan,² the subject of a recent photodissociation study from our laboratory. A comparison of the behavior of these systems promises insight into the

role of issues such as energetics and spin-orbit mediated processes in the photodissociation dynamics of closely related systems. Our study of the photodissociation of furan at 193 employed photofragment translational spectroscopy with tunable VUV probe provided by synchrotron radiation, just as in the present thiophene experiments. Three primary channels were observed from furan: $\text{HCO} + \text{C}_3\text{H}_3$, $\text{CO} + \text{C}_3\text{H}_4$ and $\text{H}_2\text{CCO} + \text{C}_2\text{H}_2$. The evidence suggested that the two closed shell channels occur on the ground state potential energy surface (PES) following internal conversion, while the radical channel likely takes place on an excited PES. All channels exhibited a barrier for dissociation with the acetylene + ketene channel having the largest value at about 25 kcal/mol. Angular distribution measurements showed anisotropy only for the radical channel. These findings were found to be consistent with a rapid excited state dissociation for the radical channel and slow dissociation for the other two pathways.

In this study, we have measured the time-of-flight (TOF) and photoionization efficiency spectra (PIE) of photofragments from the photolysis of thiophene using photofragment translational spectroscopy with synchrotron undulator radiation on the Chemical Dynamics Beamline at the Advanced Light Source. Five primary dissociation channels have been identified, including three closed shell channels and two radical channels, as summarized below.

Channels analogous to 2, 3 and 5 were observed in the dissociation of furan at 193 nm.² Two additional channels, 1 and 4, were observed in the present study. Fig. 1 shows the energy diagram of the observed photodissociation pathways for thiophene at 193 nm.

II. EXPERIMENTAL

The experiments were performed on beamline 9.0.2.1 of the Advanced Light Source using a rotatable source molecular beam apparatus described in detail elsewhere.^{9,1} Helium was bubbled through a thiophene sample held at room temperature. The total pressure was about 800 Torr, thus

yielding a 10% mixture of Th in He. The mixture was fed through a pulsed valve and expanded from a nozzle heated to $\sim 80^{\circ}\text{C}$ into the main chamber. The resulting molecular beam was collimated with two skimmers and its velocity and speed ratio were found to be 1280 ± 10 m/s and 7 ± 1 , respectively. Thiophene (99%) was obtained from Aldrich and used without further purification.

The molecular beam was intersected at 90° with an ArF excimer laser beam. The laser beam was focused to a spot of size 2×4 mm and aligned perpendicular to the plane containing the molecular beam and detector axes, on the axis of rotation of the molecular beam source. Photofragments entering the triply differentially pumped detector region were photoionized 15.2 cm downstream from the interaction region using tunable synchrotron radiation. The characteristics of the light source have been discussed in detail elsewhere,¹ but include a flux of 10^{16} photons/sec (quasi-continuous), an energy bandwidth of 2.2%, and a cross section in the probe region of 0.2×0.1 mm. The photoionized products were mass selected by a quadrupole mass filter and the ions were counted with a Daly ion counter. Time-of-flight spectra of the neutral products were measured with a multichannel scaler (MCS) whose bin width was fixed at 1 μsec for the measurements reported here. Timing sequences for the laser, pulsed valve, and the MCS were controlled by a digital delay generator. Eight quartz plates fixed at Brewster's angle were used for the polarization measurements to give $87 \pm 5\%$ polarized light. A half wave plate (Karl Lambrecht) was used to rotate the angle of polarization with respect to the detector axis. Care was taken to ensure that the TOF spectra shown here are free of multiphoton effects (except where indicated.)

The tunability of the VUV light source allows measurements of photoionization efficiency (PIE) spectra and for the selective ionization of products with very low background counts. A series of TOF spectra were recorded at a fixed angle for different photoionization energies, then they were normalized for the probe photon flux and integrated to obtain the PIE spectra. A gas filter filled with about 25 Torr of Ar was used to eliminate higher harmonics of the undulator radiation.¹⁰ A MgF_2 optical filter also was used to eliminate small contamination of the probe light by higher energy photons when probing below 10.5 eV.

III. RESULTS AND DISCUSSION

TOF spectra of photofragments from photodissociation of Th at 193 nm have been measured at several angles from 20° to 50°. Signal at $m/e = 58, 52, 51, 45, 44, 40, 39, 33, 32$, and 26 were detected. These correspond to C_2H_2S , C_4H_4 , C_4H_3 , HCS, CS, C_3H_4 , C_3H_3 , HS, S and C_2H_2 respectively. In all the TOF spectra presented here, the open circles represent the experimental data, the dash lines and the dash-dot lines are single channel contributions to the forward convolution fit, and the solid lines are the overall fit to the data. A forward convolution fit to data was used to get center of mass translational distributions, $P(E)$.^{11,12} Although we typically show TOF spectra for only one angle for each product, multiple angles were recorded and fitted for each channel shown. Fig. 1 shows the energy diagram including the five observed primary dissociation channels. Table I lists the energetics of the isomers of the various species observed in this study.

Fig. 2 shows PIE spectra of the fragment C_4H_4 (a), C_3H_3 (b), C_2H_2S (c), C_3H_4 (d), C_4H_3 (e), and CS (f). The onset in each PIE curve is roughly determined by linear extrapolation. Onsets obtained in this study are also listed in Table I. We will present the experimental results and data fits according to the approximate relative signal intensity of each dissociation channel.

A. The $C_4H_4 + S$ channel

TOF spectra of the C_4H_4 and S fragments measured at different angles are shown in Fig. 3. The photoionization energy was 11.0 and 10.5 eV, respectively. The contribution of dissociative ionization of larger fragments (C_2H_2S , HCS, CS and HS) to sulfur atom may be excluded since a MgF_2 window was used to cut off high-energy scattered radiation. However, these contributions were observed using electron impact ionization in Myers' study. The TOF spectra of mass 52 (C_4H_4) were fitted using the translational energy distribution shown where indicated in Fig. 4. This $P(E)$ fits all measured angles. The $P(E)$ is peaked at 4 kcal/mol with an average translational energy of 4.9 kcal/mol. This $P(E)$ extends out to about 17 kcal/mol.

Depending on the structure of mass 52 (C_4H_4), three possible dissociation channels are allowed energetically.⁸ They are:

The analogous channel, $C_4H_4 + O$, was not observed for furan eventhough above three dissociation channels were energetically allowed.² It is likely that the weaker C-S bond contributes to the observation of these channels for Th.

The PIE spectrum for C_4H_4 is shown in Fig. 2a. It is clear that the C_4H_4 fragment is produced predominantly as vinyl acetylene ($H_2C=CH-C\equiv CH$). Our assignment is based upon the following considerations. The onset is 9.40 eV labeled by an arrow and is very close to the ionization potential (IP) of vinyl acetylene (9.58 eV).¹³ IP for 1,2,3-butatriene ($CH_2=C=C=CH_2$) and methylenecyclopropene ($C=C-C=CH_2$) are 0.25 eV and 1.25 eV lower than the observed onset, respectively. The latter is thus ruled out as a major contributor to the m/e 52 product. The formation of 1,2,3-butatriene (process **1b**) is not excluded by our results, but energetic considerations suggest that **1a** is likely to dominate. Finally, the PIE curve drops down slowly at the region of 9.5 to 9.0 eV. The C_4H_4 fragment likely carries some internal energy, causing the observed threshold to shift toward low energies. These conclusions are in good agreement with previous studies.⁸

This channel likely occurs on the ground potential energy surface. Angular distribution measurements for this channel yielded an isotropic product angular distribution (i.e. $\beta = 0$), implying that the dissociation process takes place on a time scale longer than several molecular rotation periods. The $P(E)$ for this channel indicates that 16% of the available energy goes into product translational energy. The C_4H_4 fragment therefore has a very 'hot' internal energy distribution. This is consistent with red-shift of the onset shown in the PIE curve of the C_4H_4 fragment.

Since the first excited S (1D) state lies 26.52 kcal/mol above the ground (3P) state,¹⁴ S (1D) can be also formed together with vinyl acetylene in process **1a**. Some evidence for excited S (1D) was observed in this study: that is, S atom product appearing below the IP for S(3P). However, this product was found at translational energies only accessible with multiphoton excitation, leading us to conclude that it is not a product of single photon dissociation of thiophene.

B. The $C_2H_2S + C_2H_2$ channel

TOF spectra of the C_2H_2S and C_2H_2 fragments were measured at 20°, 35° and 50° respectively. Photon energies of 10.0 eV and 12.5 eV were used for the mass 58 (C_2H_2S) and 26

(C₂H₂) respectively. Fig. 5 shows the TOF spectra for the fragments: C₂H₂S and C₂H₂. The translational energy distribution used to fit the TOF spectra for the C₂H₂S fragment is shown in Fig. 4. Using this translational energy distribution, the fast part shown in the TOF spectra at mass 26 (C₂H₂) can clearly be momentum matched. The slow part of the C₂H₂ product needs additional contributions to complete the fit. These contributions are from the dissociative ionization of C₄H₄ and C₂H₂S fragments caused by the small amount of higher energy photons. The translational energy distribution (Fig.4) is peaked at 16.2 kcal/mol and extends out to 44.0 kcal/mol, with an average energy release of 19.4 kcal/mol. This result indicates that the C₂H₂ loss channel has a substantial exit barrier.

The energetics for the C₂H₂S + C₂H₂ channel for the different C₂H₂S isomers are:⁸

The C₂H₂S fragment is likely thioketene (H₂C=C=S), the more stable of the three isomers according to the following rationale. (1) The onset shown in Fig. 2(c) is at 8.72 eV. This onset is close to the IP (8.77 eV) of thioketene (H₂C=C=S).¹³ It is reasonably lower than IP of the thioketene because the C₂H₂S fragment has significant internal energy. (2) The peak of the translational energy distribution occurs at 44.0 kcal/mol, and is larger than the available energies of the processes 2b and 2c. Hence, thioketene is clearly the dominant, if not the exclusive C₂H₂S product isomer formed from thiophene dissociation at 193 nm. The analogous channel was observed in furan photodissociation at 193 nm.² In the furan study, identity of the C₂H₂O fragment was determined to be ketene (H₂C=C=O). Our result is also consistent with previous studies.⁸ Preliminary analysis indicates that channel 2a occurs on the ground state potential energy surface following internal conversion. Laser polarization measurements for this channel showed an isotropic angular distribution.

The secondary dissociation processes:

are essentially thermoneutral. Although the process 2d and 2e are energetically allowed, the yield will be negligible since it can only occur for the primary products formed with zero translational energy

C. The CS + C₃H₄ channel

This is the weakest of the closed-shell channels. Fig. 6 shows TOF spectra of the CS and C₃H₄ fragments obtained at the indicated angles. Photon energies of 12.5 eV and 10.5 eV were used for measuring the TOF spectra of mass 44 and 40, respectively. In measuring the TOF spectrum of mass 40, the MgF₂ window was used to cut off high-energy scattered radiation, so we need not consider the contribution of larger fragment dissociative ionization. However, a contribution was seen from the stronger radical product at m/e 39 (C₃H₃) owing to a small amount of mass leakage in the quadrupole. This is seen clearly as the slow component in the bimodal TOF spectrum for m/e 40 at 20°. The TOF spectra of the CS fragment were also fitted using two components represented by the CS + C₃H₄ channel (a dashed line) and the HCS + C₃H₃ channel (a dash-dot line). Fig. 4 shows the P(E) used to fit this channel. The P(E) shows momentum-matching between the fast part of mass 40 (C₃H₄) and mass 44 (CS). The P(E) in Fig. 4 has peaks at 10.0 kcal/mol and extends out to approximately 35 kcal/mol with an average translational energy release of 14.3 kcal/mol. This suggests an exit barrier of at least 10 kcal/mol for this channel.

The C₃H₄ fragment also has different isomers. The dissociations including three energetically allowed structures are listed as follows:

It is difficult to distinguish which structure is formed in this process judging solely from the reaction energy, since the energetics for 3a and 3b are very close. According to the onset (9.60 eV) of the C₃H₄ fragment shown in Fig. 2(d), a closer match to allene (H₂C=C=CH₂, 9.69 eV), than propyne (H₃C-C≡CH, 10.37) or cyclopropene (9.67 eV). However, this may still be propyne, with the ionization onset red-shifted by the internal energy. The probability for ring formation is likely to be low, although the IP of cyclopropene is also close to our experimental value. A similar

channel was observed at 193 nm photodissociation from furan. However, in the furan case, this was assigned to propyne, based largely on the results of theoretical studies.¹⁵ Similar arguments may well hold for thiophene. The PIE curve (shown in Fig. 2(f)) of the CS fragment was also measured. The onset is around 10.95 eV (IP(CS): 11.33 eV). A tail was observed at range of 11.2 – 10.5 eV. This is likely owing to vibrational excitation of the CS fragment.

D. The HCS + C₃H₃ channel

Two free radical channels were observed besides the three closed-shell channels mentioned above. The first, HCS + C₃H₃, was seen prominently in furan photodissociation at 193 nm.² The TOF spectra of mass 39 (C₃H₃) and mass 45 (HCS) measured at different angles are shown in Fig. 7, probed with 9.5 eV and 10.0 eV photon energies, respectively. A MgF₂ window was used in order to avoid the dissociative ionization of the C₃H₄ fragment to produce the C₃H₃ ion. The P(E) in Fig. 4 was used to fit the HCS + C₃H₃ channel. This P(E) decreases out to about 14.0 kcal/mol with an average translational energy release of 2.9 kcal/mol. The fast part of the TOF spectra of the C₃H₃ fragment have a small contribution on the leading edge that was neither momentum-matched to the HCS, nor energetically accessible for a one-photon dissociation. It likely represents a small multiphoton contribution. The C₃H₃ fragment was determined to be propargyl radical (HC≡C-C•H₂), by comparing ionization potentials of the C₃H₃ isomers and the onset shown in PIE curve (Fig. 2(b)). The formation of cyclopropenyl radical is not allowed energetically. HSC is less stable by 39.6 kcal/mol compared with HCS.¹⁶ The process for forming HSC from thiophene at 193 nm excitation is thus not energetically allowed. We conclude that the mass 45 product is the HCS free radical.

In the 193 nm photodissociation of furan, angular distribution measurements showed anisotropy for the analogous HCO + C₃H₃ channel, indicating that it likely occurs directly on an electronically excited potential energy surface (PES).² In contrast, laser polarization measurements for this channel in thiophene show an isotropic angular distribution.

E. The HS + C₄H₃ channel

Of all five primary dissociation pathways, the HS + C₄H₃ channel is the weakest one, perhaps due in part to the instability of the C₄H₃ radical. The TOF spectra of mass 33 (HS) and

mass 51 (C_4H_3) at different angles are shown in Fig. 8. The probe energy was 12.0 eV and 11.0 eV for ionization of the fragments HS and C_4H_3 , respectively. The translational energy distribution shown in Fig. 4 was used to fit the HS + C_4H_3 channel. The $P(E)$ peaks near zero and extends out to 7.0 kcal/mol with an average energy release of 1 kcal/mol. The m/e 51 signal includes a contribution from the stronger m/e 52 signal. The m/e 33 fragment also has the contribution of the mass leakage from the much more intense m/e 32 signal.

An *ab initio* study of the structures and properties of the free radical C_4H_3 was completed by Ha and Gey.¹⁷ Two low-lying isomers 2-dehydro-buta-1-ene-3-yne (iso- C_4H_3) and 1-dehydro-buta-1-ene-3-yne (n- C_4H_3) were found to be the most stable.¹⁷⁻¹⁹ Miller et al. noted that iso- C_4H_3 is thermodynamically more stable (9.6 kcal/mol) than n- C_4H_3 and thus most of the C_4H_3 should be in the form of the iso isomer even at combustion temperatures.²⁰ It seems likely that the C_4H_3 fragment from photodissociation of thiophene survives also as the iso- C_4H_3 isomer ($HC\equiv C-C\cdot=CH_2$). The PIE curve of the C_4H_3 fragment is shown in Fig. 2e with an onset of about 9.2 eV. No experimental results for the IP of this fragment are available for comparison. Theoretical calculations will be useful for determination of the IP of C_4H_3 isomers. Myers reported one additional channel, giving rise to m/e 82, that he ascribed predominantly to dimer photodissociation. The absence of signal for m/e 82 in our experiments provides additional support for this assignment.

F. Energy Partitioning

From the translational energy distributions of the different channels, the average translational energy release for each channel can be determined. Table II lists the obtained maximum translation energies, the average translational energies, the average internal energies, and their ratios for each channels. It is clear that the internal energy partitioning of the photodissociation products from thiophene at 193 nm is always larger than the translational energy release, with the latter ranging from 5% to 24% of the total energy available.

G. Dissociation mechanism

Several aspects of the dissociation dynamics in thiophene contrast sharply with the behavior observed for furan. Firstly, the different energetics open the channels to $C_4H_4 + S$ and $C_4H_3 + HS$, for which analogous processes were not seen in furan. More significantly, the $C_3H_3 + HCO$ radical

channel which dominates the UV photodissociation of furan is found to be only a minor channel in Th. Secondly, the anisotropy in the photofragment angular distribution for this channel in furan is absent in Th. These observations suggest that the competition between direct dissociation on the excited state surface, and internal conversion to the ground state, which was inferred for furan, does not take place in Th. Instead, it appears likely that internal conversion to the ground state is rapid, and leads to statistical decomposition of the hot, ground state molecules. The approximate product yields that we observe are entirely consistent with what one would expect for statistical decomposition to all the various product channels, given the energetics shown in Fig. 1.

The dissociation mechanism of Th has not been investigated in detail. Infrared laser multiphoton dissociation⁵ and UV photodissociation⁸ studies suggested that breakage of the C-S bond in Th forms an unstable 1,5-diradical ($\bullet\text{CHCHCHCHS}\bullet$) as the initial step. For the acetylene loss channel, Myers argued that a bond would form between the sulfur atom and C(3), while C(4) and C(5) form the acetylene triple bond and break their bonds with the sulfur atom and C(3). Mass 58 was considered to be thiirene structure in Myers' thesis. However, Hsu et al.⁸ and our PIE measurements both indicate that mass 58 is thioketene structure. This suggests that the mechanism proposed by Myers would not explain the acetylene loss channel. Hsu and coworkers also suggested a mechanism that breakage of the C(2)-C(3) and C(4)-C(5) bonds, together with cyclization between C(2) and C(5), would give $\text{CH}\equiv\text{CH}$ and $c\text{-CH=CHS}$. The formation of thioketene can be realized if the internal energy content of $c\text{-CH=CHS}$ is higher than the rearrangement barrier between $c\text{-CH=CHS}$ and $\text{CH}_2=\text{C}=\text{S}$. The C-S bond strength is weaker than the C=C bond for thiophene. It seems unlikely that two double bonds break and the weaker C-S bond persists. We suspect that scission of the weak C-S bond forms the 1,5-diradical as the initial step, followed by 2,3-H atom migration; the H-atom transfer should be fast because more than 60 kcal/mol average internal energy is available for this channel.

I. The acetylene loss channel

For the sulfur atom elimination channel, Myers thought that the closest match for mass 52 would be 3-methylene cyclopropene. One would expect a barrier somewhere along the reaction coordinate for Myers' mechanism since H atom transfer to yield the CH₂ group attached to the ring must occur. In addition, the probability of forming the ring would be small. However, we have shown that mass 58 corresponds to vinyl acetylene, which is in agreement with a previous study.⁸ The 1,5-diradical with extensive internal energy could readily undergo bond rearrangement and H atom migration to form vinyl acetylene and sulfur atom.

II. The sulfur atom elimination channel:

The other three channels would be similar to the two channels discussed above, with formation of the 1,5-diradical as the initial step. The H atom migration plays an important function in unimolecular decomposition of some other aromatic compounds as well.^{15,21-24}

III. The CS loss channel:

IV. The HCS loss channel:

V. The HS loss channel:

Thiophene photodissociation was found to be a very complex process. The discussion of dissociation mechanisms above is schematic, and based on qualitative arguments and 'chemical intuition' according to the identified products. The full elucidation of the detailed mechanism will require theoretical calculations.

IV. CONCLUSION

We have presented results for the primary photodissociation dynamics of thiophene, c-C₄H₄S 193nm using tunable synchrotron undulator radiation as a universal product probe. Three closed-shell and two radical channels were identified as primary dissociation processes. Mass 58, 52, 51, 45, 44, 40, 39, 33, 32, and 26 were identified to be thioketene, vinyl acetylene, iso-C₄H₃, HCS radical, CS, allene, propargyl radical, HS, S and acetylene, respectively. The evidence suggests that dissociation occurs on the ground state potential surface following internal conversion; the results thus have direct relevance to the high-temperature pyrolysis of thiophene.

Acknowledgments

This work was supported by the Director, Office of Science, Office of Basic Energy Sciences, Chemical Sciences Division of the US Department of Energy under contract No. DE-AC03-76SF00098. The Advanced Light Source is supported by the Director, Office of Science, Office of Basic Energy Sciences, Materials Sciences Division of the US Department of Energy under the same contract.

Table I The Energetics for Isomers of the observed fragments

Mass detected ^a	neutral species ^b	$\Delta H_{f,298}$ (kcal/mol) ^c	Literature IP (eV) ^c	This work (eV)
84 (C ₄ H ₄ S)	C ₄ H ₄ S (thiophene)	27.5	8.87	
52 (C ₄ H ₄)	CH ₂ =CH-C≡CH	73	9.58	9.4
	CH ₂ =C=C=CH ₂	83	9.15
	C=C-C=CH ₂	101	8.15	
51 (C ₄ H ₃)	CH≡C-C•=CH ₂	116 ^d		~9.2
	CH≡C-CH=C•H	126 ^d		
40 (C ₃ H ₄)	CH ₃ -C≡CH	44.2 ^e	10.36 ^e	9.6
	CH ₂ =C=CH ₂	45.5 ^e	9.69 ^e
	66.2 ^e	9.67 ^e	
39 (C ₃ H ₃)	CH≡C-C•H ₂	81.1 ^f	8.67 ^g	8.6
	105 ^h	6.6	
58 (C ₂ H ₂ S)	CH ₂ =C=S	39	8.77	8.7
	c-CH=CHS	~87 ⁱ		
	•CH=CHS•	~101 ⁱ		
45 (HCS)	HC•S	71.7 ^j	7.4 ⁱ	
	HSC•	111.3 ^k		
44 (CS)	CS	67 ^l	11.33 ^e	
26 (C ₂ H ₂)	HC≡CH	54.4	11.40	
33 (HS)	HS	33.3 ^l	10.42 ^m	
32(S)	S	66.25 (³ P) ^e	10.36 ^e	

^a ions observed which are attributable to VUV photoionization of photofragments for Th at 193 nm.

^b Possible isomers for photoproducts of Th formed at 193 nm.

^c Heat of formation at 298 K. Values are from ref ¹³ unless.

^d References 17 and 18. ^e Reference 25. ^f Reference 26. ^g Reference 27. ^h Reference 28. ⁱ References 8 and 29. ^j Reference 30. ^k References 31 and. ^l Reference 32. ^m Reference 33

Table II Energy partition of different dissociation channels of thiophene at 193 nm excitation. All energies shown here are in kcal/mol

Dissociation Channel (193 nm excitation)	Total available energy E_{tot}	Maximum E_{trans}	Averaged E_{trans}	Averged E_{int}	$E_{\text{trans}}/E_{\text{int}}$
$\text{H}_2\text{C}=\text{CH}-\text{C}\equiv\text{CH} + \text{S}(^3\text{P})$	36	17	4.9	31.1	14/86
$\text{H}_2\text{C}=\text{C}=\text{S} + \text{HC}\equiv\text{CH}$	82	44	19.4	62.6	24/76
$\text{H}_3\text{C}-\text{C}\equiv\text{CH} + \text{CS}$	64	35	14.3	49.7	22/78
$\text{HC}\cdot\text{S} + \text{HC}\equiv\text{C}-\text{C}\cdot\text{H}_2$	23	14	2.9	20.1	13/87
$\text{HS}\cdot + \text{HC}\equiv\text{C}-\text{C}\cdot=\text{CH}_2$	26	7	1.0	25.0	4/96

REFERENCES

- (1) Heimann, P. A.; Koike, M.; Hsu, C. W.; Blank, D.; Yang, X. M.; Suits, A. G.; Lee, Y. T.; Evans, M.; Ng, C. Y.; Flaim, C.; Padmore, H. A. *Rev. Sci. Instrum.* **1997**, *68*, 1945-1951.
- (2) Sorkhabi, O.; Qi, F.; Rizvi, A.; Suits, A. G. *J. Chem. Phys.* **1999**, *In press*.
- (3) Weissberger, A.; Taylor, E. C.; Gronowitz, S. Thiophene and Its Derivatives. In *The Chemistry of Heterocyclic Compounds*, 1992; Vol. 44.
- (4) Wiebe, H. A.; Heicklen, J. *Can. J. Chem.* **1969**, *47*, 2965.
- (5) Nayak, A. K.; Sarkar, S. K.; Karve, R. S.; Parthasarathy, V.; Rama Rao, K. V. S.; Mittal, J. P.; Krishnamachari, S. L. N. G.; Venkitachalam, T. V. *App. Phys. B* **1989**, *B48*, 437.
- (6) Krishnamachari, S. L. N. G.; Venkitachalam, T. V. *Chem. Phys. Lett.* **1978**, *55*, 116-18.
- (7) Myers, J. D. . Ph.D. Thesis, University of California, 1993.
- (8) Hsu, C. W.; Liao, C. L.; Ma, Z. X.; Ng, C. Y. *J. Phys. Chem.* **1995**, *99*, 1760-1767.
- (9) Yang, X.; Lin, J.; Lee, Y. T.; Blank, D. A.; Suits, A. G.; Wodtke, A. M. *Rev. Sci. Instrum.* **1997**, *68*, 3317-3326.
- (10) Suits, A. G.; Heimann, P.; Yang, X. M.; Evans, M.; Hsu, C. W.; Lu, K. T.; Lee, Y. T.; Kung, A. H. *Rev. Sci. Instrum.* **1995**, *66*, 4841-4844.
- (11) Wodtke, A. M. . Ph.D., University of California at Berkeley, 1986.
- (12) Zhao, X. . Ph.D., University of California at Berkeley, 1988.
- (13) Lias, S. G.; Bartmess, J. E.; Liebman, J. F.; Holmes, J. L.; Levin, R. D.; Mallard, W. *G. J. Phys. Chem. Ref. Data* **1988**, *17*, Suppl. 1.
- (14) Moore, C. E. *Atomic Energy Levels*; National Bureau of Standards: Washington, D.C., 1949; Vol. I.
- (15) Liu, R. F.; Zhou, X. F.; Zhai, L. *Journal of Computational Chemistry* **1998**, *19*, 240-249.
- (16) Kaiser, R. I.; Sun, W.; Suits, A. G. *J. Chem. Phys.* **1997**, *106*, 5288-5291.
- (17) Ha, T. K.; Gey, E. *Journal of Molecular Structure: Theochem* **1994**, *112*, 197-202.
- (18) Kaiser, R. I.; Stranges, D.; Lee, Y. T.; Suits, A. G. *J. Chem. Phys.* **1996**, *105*, 8721-8733.
- (19) Walch, S. P. *J. Chem. Phys.* **1995**, *103*, 8544.
- (20) Miller, J. A.; Mellius, C. F. *Combust. Flame* **1992**, *91*, 21.

- (21) Zhai, L.; Zhou, X.; Liu, R. *J. Phys. Chem. A* **1999**, *103*, 3917.
- (22) Lifshitz, A.; Bidani, M.; Bidani, S. *J. Phys. Chem.* **1986**, *90*, 5373.
- (23) Lifshitz, A.; Tambura, C.; Shashua, R. *J. Phys. Chem. A* **1997**, *101*, 1018.
- (24) Lifshitz, A.; Tambura, C.; Shashua, R. *J. Phys. Chem. A* **1998**, *102*, 10655.
- (25) Lide, D. R. *Handbook of Chemistry and Physics*; CRC: , 1995; Vol. 75 th Edition.
- (26) Tsang, W. *Heats of Formation of Organic Free Radicals by Kinetic Methods*; Blackie Academic and Professional: London, 1996.
- (27) Kaiser, R. I.; Lee, Y. T.; Suits, A. G. *J. Chem. Phys.* **1996**, *105*, 8705-8720.
- (28) Minsek, D. W.; Chen, P. *J. Phys. Chem.* **1990**, *94*, 8399.
- (29) Verkoczy, B.; Sherwood, A. G.; Safarik, I.; Strausz, O. P. *Can. J. Chem.* **1983**, *61*, 2268.
- (30) Ruscic, B.; Berkowitz, J. *J. Chem. Phys.* **1993**, *98*, 2568.
- (31) Kaiser, R. I.; Ochsenfeld, C.; Head-Gordon, M.; Lee, Y. T. *Science* **1998**, *279*, 1181.
- (32) Chase, M. W., Jr. *J. Phys. Chem. Ref. Data* **1998**, *9*, 1-1951.
- (33) Hsu, C. W.; Baldwin, D. P.; Liao, C. L.; Ng, C. Y. *J. Chem. Phys.* **1994**, *100*, 8047-8054.

FIGURE CAPTIONS

1. Energy diagram showing all observed product channels from 193nm dissociation of thiophene, along with barriers estimated from the results.
2. Low-resolution photoionization efficiency spectra for the indicated photoproducts, along with approximate thresholds.
3. Time-of-flight (TOF) spectra for the C_4H_4+S channel at 20° .
4. Translational energy distributions ($P(E)$'s) for the indicated product channel.
5. TOF spectra for m/e 26 and 58 (acetylene and thioketene) at 20° .
6. TOF spectra for m/e 40 and 44 (C_3H_4 and CS) at the indicated laboratory angles.
7. TOF spectra for m/e 45 and 39 (HCS and C_3H_3) at 20° .
8. TOF spectra for m/e 33 and 51 (HS and C_4H_3) at the indicated laboratory angles.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8

ERNEST ORLANDO LAWRENCE BERKELEY NATIONAL LABORATORY
ONE CYCLOTRON ROAD : BERKELEY, CALIFORNIA 94720

2