

UCLA

UCLA Previously Published Works

Title

Expanding CRISPR/Cas9 Genome Editing Capacity in Zebrafish Using SaCas9

Permalink

<https://escholarship.org/uc/item/83s2x8j3>

Journal

G3: Genes, Genomes, Genetics, 6(8)

ISSN

2160-1836

Authors

Feng, Yan
Chen, Cheng
Han, Yuxiang
et al.

Publication Date

2016-08-01

DOI

10.1534/g3.116.031914

Peer reviewed

Expanding CRISPR/Cas9 Genome Editing Capacity in Zebrafish Using SaCas9

Yan Feng,* Cheng Chen,* Yuxiang Han,* Zelin Chen,*[†] Xiaochan Lu,* Fang Liang,* Song Li,* Wei Qin,*¹ and Shuo Lin*^{†,1}

*Laboratory of Chemical Genomics, School of Chemical Biology and Biotechnology, Peking University Shenzhen Graduate School, 518055, China, and [†]Department of Molecular, Cell and Developmental Biology, University of California, Los Angeles, California 90095

ABSTRACT The type II CRISPR/Cas9 system has been used widely for genome editing in zebrafish. However, the requirement for the 5'-NGG-3' protospacer-adjacent motif (PAM) of Cas9 from *Streptococcus pyogenes* (SpCas9) limits its targeting sequences. Here, we report that a Cas9 ortholog from *Staphylococcus aureus* (SaCas9), and its KKH variant, successfully induced targeted mutagenesis with high frequency in zebrafish. Confirming previous findings, the SpCas9 variant, VQR, can also induce targeted mutations in zebrafish. Bioinformatics analysis of these new Cas targets suggests that the number of available target sites in the zebrafish genome can be greatly expanded. Collectively, the expanded target repertoire of Cas9 in zebrafish should further facilitate the utility of this organism for genetic studies of vertebrate biology.

KEYWORDS

CRISPR/Cas9
SaCas9
KKH SaCas9
variant
gene editing
zebrafish

Targeted genome engineering offers new genetic tools for understanding biological processes, and for developing therapeutics of human diseases. Following the development of Zinc Finger (ZFs) and transcription activator-like effector (TALEs) technology (Doyon *et al.* 2008; Huang *et al.* 2011; Meng *et al.* 2008), the clustered regularly interspaced short palindromic repeat/CRISPR-associated (CRISPR/Cas) system has emerged as the current gene editing tool of choice. CRISPR/Cas system has the advantages of ease of handling, low cost, and universal applicability in different cell types and organisms. CRISPR/Cas can be classified into six types based on the presence of "signature genes" (Makarova *et al.* 2011, 2015; Shmakov *et al.* 2015; Wright *et al.* 2016). Among them, Cas9 from *Streptococcus pyogenes* (SpCas9), which belongs to the type II CRISPR/Cas system, has been demonstrated to be effective in inducing targeted DNA

double strand breaks (DSBs) in a variety of organisms (Chang *et al.* 2013; Cong *et al.* 2013; Dickinson *et al.* 2013; Friedland *et al.* 2013; Gratz *et al.* 2013; Hwang *et al.* 2013; Jinek *et al.* 2013; Mali *et al.* 2013; Qin *et al.* 2015; Shalem *et al.* 2014; Wang *et al.* 2014; Yang *et al.* 2013a). SpCas9 nuclease DNA sequence specificity relies on a guide RNA with a protospacer-adjacent motif (PAM) sequence at the 3' end of a 20-bp target sequence. The most widely used SpCas9 recognizes a short 5'-NGG-3' PAM. Since PAM sequences are different in different CRISPR/Cas systems, alternative PAMs would provide more flexibility for targeting strategies such as precise knock-in mutations.

Recently, Cas9 orthologs with distinct DNA binding specificity and PAM recognition, including *Neisseria meningitidis* (NmCas9), *Streptococcus thermophilus* 1 (St1Cas9), and *Staphylococcus aureus* (SaCas9) have been applied for genome editing in human cells (Hou *et al.* 2013; Karvelis *et al.* 2013; Ran *et al.* 2015). Among them, Cas9 from *S. aureus* (SaCas9) is smaller, and has a longer PAM of 5'-NNGRRT-3' sequence. These features allow easier delivery by viral expression vectors, and higher sequence specificity, which would be more desirable for therapeutic applications. Recently, a SaCas9 variant (KKH SaCas9) with partially relaxed 5'-NNNRRT-3' PAM specificities has been demonstrated to show robust genome editing activities in human cells, which further increases the SaCas9 targeting range (Kleistner *et al.* 2015a).

Here, we demonstrate that SaCas9, with its KKH SaCas9 variant, can edit the zebrafish genome with high targeting efficiency. This increases the frequency of available target sites, and expands the utility of CRISPR/Cas9 in zebrafish by targeting those previously inaccessible Cas9 sites in the genome.

Copyright © 2016 Feng *et al.*

doi: 10.1534/g3.116.031914

Manuscript received April 29, 2016; accepted for publication June 6, 2016; published Early Online June 16, 2016.

This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Supplemental material is available online at www.g3journal.org/lookup/suppl/doi:10.1534/g3.116.031914/-/DC1

¹Corresponding authors: School of Chemical Biology and Biotechnology, Peking University Shenzhen Graduate School, Lishui Road, Xili Town, Nanshan District, 518055 Shenzhen, China. E-mail: qinwei@pkusz.edu.cn; and Department of Molecular, Cell and Developmental Biology, University of California, Los Angeles, 621 Charles E. Young Drive South, Los Angeles, CA 90095-1606. E-mail: shuolin@ucla.edu

(Bassett *et al.* 2013). Although SaCas9 can target all 5'-NNGRR-3' PAMs, it cleaves the sequence of 5'-NNGRR-3' most efficiently (Ran *et al.* 2015). We therefore selected this type of PAM sequence from the zebrafish genome for initial testing. Since pigmentation defect is a convenient phenotype indication for mutation in the tyrosinase (*tyr*) gene, which encodes an enzyme that converts tyrosine into melanin (Camp and Lardelli 2001), we designed a gRNA targeting *tyr*. After coinjection of *tyr* gRNA and SaCas9 mRNA into one-cell-stage zebrafish embryos, pigmentation reduction was observed in 42 embryos ($n = 73$), some of which totally lacked pigmentation (Figure 1B). Injected embryos developed normally, no obvious toxicity was observed. We randomly selected six embryos, extracted genomic DNA from pools of embryos and assessed the mutagenesis rates using the T7 endonuclease I (*T7EI*) assay and further confirmed indels at the expected sites by sequencing (Figure 1D).

To test if SaCas9/gRNA can be used generally in zebrafish, we constructed eight additional gRNAs, one targeting transgene EGFP and others targeting endogenous genes. When EGFP SaCas9/gRNA was injected into *tg (mylpfa:EGFP)* zebrafish eggs, loss of EGFP expression was observed in muscle cells (Figure 1C). For other specific sites, indels were detected in all injected embryos with a frequency of 13–76%. Again, the indels were sequencing confirmed at the expected genomic loci (Figure 1E and Figure S1). Taken together, these results demonstrate that SaCas9 can effectively edit genome in zebrafish.

Gene editing in zebrafish using KKH SaCas9

Native SaCas9 recognizes a longer 5'-NNGRR-3' PAM, which occurs on average every 32 bps of random DNA. Obtaining variants with relaxed or partially relaxed specificity to expand the repertoire of gene editing tools in zebrafish is desirable. Toward this goal, we investigated two recently described SaCas9 and SpCas9 variants with different specificities (Figure 2A) (Kleinstiver *et al.* 2015a, 2015b). One SaCas9 variant is called KKH SaCas9, which can target 5'-NNNRRT-3' PAM in human genome (Kleinstiver *et al.* 2015a). We evaluated the activity of KKH SaCas9 by targeting four endogenous genes in zebrafish, and observed indels in all of them (Figure 2B and Figure S2). Similarly, we evaluated activity of the SpCas9 VQR variant with an NGA PAM, which has been shown previously to induce indels in zebrafish and *Caenorhabditis elegans* (Bell *et al.* 2016; Kleinstiver *et al.* 2015b), and obtained expected indels at the two predicted sites (Figure 2B and Figure S2). Collectively, these results demonstrate that both KKH SaCas9 and VQR SpCas9 variants are functional in zebrafish. Since 5'-NNNRRT-3' represents a relaxed PAM, it is conceivable that the off-target rate might be increased. In zebrafish, this potential problem could be addressed by analyzing the linkage between the targeted mutation, and the specific phenotype in large numbers of individual fish through breeding multiple generations.

The SpCas9 system is an efficient tool for genome editing in many organisms including bacteria, yeast, human cells, *C. elegans*, zebrafish, and mice (Hsu *et al.* 2014; Sternberg and Doudna 2015). Due to the restriction of having to use the 5'-NGG-3' PAM required by SpCas9, it is challenging to find target sites with high efficiency in a specific genome locus. Improving the capacity to target very specific loci is highly desirable for DNA homology-directed gene repair, which is most efficient when a DSB is placed within 10–20 bp of a desired alteration (Elliott *et al.* 1998; Findlay *et al.* 2014; Yang *et al.* 2013b). Here, we demonstrate that using SaCas9 and its variant enable genome editing in a significantly wider range in zebrafish. Since Cas9-based gene editing is most commonly applied to induce or repair mutations in coding exons of a gene, we produced a BED file to annotate six types of target sites

(NGG, NGA, NNGRRN, NNGRRT, NNNRRN, and NNNRRT) on the UCSC gene browser.

One of the key issues in using CRISPR/Cas9 system successfully is to assure high activity. First, similar to the zebrafish-codon-optimized SpCas9, which increases gRNA targeting efficiency from ~4% to ~39% (Liu *et al.* 2014), we recommend constructing a zebrafish codon-optimized SaCas9 for future study. Second, it is also possible that direct SaCas9 protein injections could further increase efficiency (Burger *et al.* 2016; Gagnon *et al.* 2014). Finally, as modifying the SpCas9/gRNA structure by extending the duplex length has been shown to improve knockout efficiency (Dang *et al.* 2015), we speculate that the same strategy could be applied to SaCas9/gRNA structure in zebrafish.

ACKNOWLEDGMENTS

We thank Bo Zhang (Peking University) for providing the pT7-nls-zCas9-nls plasmid. We also thank Zenghou Tang for zebrafish husbandry. This work was supported by funding from the Science and Technology Program of Shenzhen (CXZZ20140903101756864, JCYJ20150924110425180, and JCYJ201510301707 55264) and by the China Postdoctoral Science Foundation (to W.Q.). Z. C. was supported by the International Postdoctoral Exchange Fellowship Program of China Postdoctoral Council (No. 20130010).

LITERATURE CITED

- Bassett, A. R., C. Tibbit, C. P. Ponting, and J. L. Liu, 2013 Highly efficient targeted mutagenesis of *Drosophila* with the CRISPR/Cas9 system. *Cell Reports* 4: 220–228.
- Bell, R.T., B.X. Fu, and A.Z. Fire, 2016 Cas9 variants expand the target repertoire in *Caenorhabditis elegans*. *Genetics* 202: 381–388.
- Burger, A., H. Lindsay, A. Felker, C. Hess, C. Anders *et al.*, 2016 Maximizing mutagenesis with solubilized CRISPR-Cas9 ribonucleoprotein complexes. *Development* 143: 2025–2037.
- Camp, E., and M. Lardelli, 2001 Tyrosinase gene expression in zebrafish embryos. *Dev. Genes Evol.* 211: 150–153.
- Chang, N., C. Sun, L. Gao, D. Zhu, X. Xu *et al.*, 2013 Genome editing with RNA-guided Cas9 nuclease in zebrafish embryos. *Cell Res.* 23: 465–472.
- Cong, L., F. A. Ran, D. Cox, S. Lin, R. Barretto *et al.*, 2013 Multiplex genome engineering using CRISPR/Cas systems. *Science* 339: 819–823.
- Dang, Y., G. Jia, J. Choi, H. Ma, E. Anaya *et al.*, 2015 Optimizing sgRNA structure to improve CRISPR-Cas9 knockout efficiency. *Genome Biol.* 16: 280.
- Dickinson, D. J., J. D. Ward, D. J. Reiner, and B. Goldstein, 2013 Engineering the *Caenorhabditis elegans* genome using Cas9-triggered homologous recombination. *Nat. Methods* 10: 1028–1034.
- Doyon, Y., J. M. McCammon, J. C. Miller, F. Faraji, C. Ngo *et al.*, 2008 Heritable targeted gene disruption in zebrafish using designed zinc-finger nucleases. *Nat. Biotechnol.* 26: 702–708.
- Elliott, B., C. Richardson, J. Winderbaum, J. A. Nickoloff, and M. Jasin, 1998 Gene conversion tracts from double-strand break repair in mammalian cells. *Mol. Cell Biol.* 18: 93–101.
- Findlay, G. M., E. A. Boyle, R. J. Hause, J. C. Klein, and J. Shendure, 2014 Saturation editing of genomic regions by multiplex homology-directed repair. *Nature* 513: 120–123.
- Friedland, A. E., Y. B. Tzur, K. M. Esvelt, M. P. Colaiacovo, G. M. Church *et al.*, 2013 Heritable genome editing in *C. elegans* via a CRISPR-Cas9 system. *Nat. Methods* 10: 741–743.
- Gagnon, J. A., E. Valen, S. B. Thyme, P. Huang, L. Akhmetova *et al.*, 2014 Efficient mutagenesis by Cas9 protein-mediated oligonucleotide insertion and large-scale assessment of single-guide RNAs. *PLoS One* 9: e98186.
- Gratz, S. J., A. M. Cummings, J. N. Nguyen, D. C. Hamm, L. K. Donohue *et al.*, 2013 Genome engineering of *Drosophila* with the CRISPR RNA-guided Cas9 nuclease. *Genetics* 194: 1029–1035.

- Hou, Z., Y. Zhang, N. E. Propson, S. E. Howden, L. F. Chu *et al.*, 2013 Efficient genome engineering in human pluripotent stem cells using Cas9 from *Neisseria meningitidis*. *Proc. Natl. Acad. Sci. USA* 110: 15644–15649.
- Hsu, P. D., E. S. Lander, and F. Zhang, 2014 Development and applications of CRISPR-Cas9 for genome engineering. *Cell* 157: 1262–1278.
- Huang, P., A. Xiao, M. Zhou, Z. Zhu, S. Lin *et al.*, 2011 Heritable gene targeting in zebrafish using customized TALENs. *Nat. Biotechnol.* 29: 699–700.
- Hwang, W. Y., Y. Fu, D. Reyon, M. L. Maeder, S. Q. Tsai *et al.*, 2013 Efficient genome editing in zebrafish using a CRISPR-Cas system. *Nat. Biotechnol.* 31: 227–229.
- Jinek, M., A. East, A. Cheng, S. Lin, E. Ma *et al.*, 2013 RNA-programmed genome editing in human cells. *eLife* 2: e00471.
- Karvelis, T., G. Gasiunas, A. Miksys, R. Barrangou, P. Horvath *et al.*, 2013 crRNA and tracrRNA guide Cas9-mediated DNA interference in *Streptococcus thermophilus*. *RNA Biol.* 10: 841–851.
- Kimmel, C. B., W. W. Ballard, S. R. Kimmel, B. Ullmann, and T. F. Schilling, 1995 Stages of embryonic development of the zebrafish. *Dev. Dyn.* 203: 253–310.
- Kleinstiver, B. P., M. S. Prew, S. Q. Tsai, N. T. Nguyen, V. V. Topkar *et al.*, 2015a Broadening the targeting range of *Staphylococcus aureus* CRISPR-Cas9 by modifying PAM recognition. *Nat. Biotechnol.* 33: 1293–1298.
- Kleinstiver, B. P., M. S. Prew, S. Q. Tsai, V. V. Topkar, N. T. Nguyen *et al.*, 2015b Engineered CRISPR-Cas9 nucleases with altered PAM specificities. *Nature* 523: 481–485.
- Liu, D., Z. Wang, A. Xiao, Y. Zhang, W. Li, Y. Zu, S. Yao, S. Lin, and B. Zhang, 2014 Efficient gene targeting in zebrafish mediated by a zebrafish-codon-optimized cas9 and evaluation of off-targeting effect. *J. Genet. Genomics* 41: 43–46.
- Makarova, K. S., L. Aravind, Y. I. Wolf, and E. V. Koonin, 2011 Unification of Cas protein families and a simple scenario for the origin and evolution of CRISPR-Cas systems. *Biol. Direct* 6: 38.
- Makarova, K. S., Y. I. Wolf, O. S. Alkhnbashi, F. Costa, S. A. Shah *et al.*, 2015 An updated evolutionary classification of CRISPR-Cas systems. *Nat. Rev. Microbiol.* 13: 722–736.
- Mali, P., L. Yang, K. M. Esvelt, J. Aach, M. Guell *et al.*, 2013 RNA-guided human genome engineering via Cas9. *Science* 339: 823–826.
- Meng, X., M. B. Noyes, L. J. Zhu, N. D. Lawson, and S. A. Wolfe, 2008 Targeted gene inactivation in zebrafish using engineered zinc-finger nucleases. *Nat. Biotechnol.* 26: 695–701.
- Qin, W., F. Liang, Y. Feng, H. Bai, R. Yan *et al.*, 2015 Expansion of CRISPR/Cas9 genome targeting sites in zebrafish by Csy4-based RNA processing. *Cell Res.* 25: 1074–1077.
- Ran, F. A., L. Cong, W. X. Yan, D. A. Scott, J. S. Gootenberg *et al.*, 2015 *In vivo* genome editing using *Staphylococcus aureus* Cas9. *Nature* 520: 186–191.
- Shalem, O., N. E. Sanjana, E. Hartenian, X. Shi, D. A. Scott *et al.*, 2014 Genome-scale CRISPR-Cas9 knockout screening in human cells. *Science* 343: 84–87.
- Shmakov, S., O. O. Abudayyeh, K. S. Makarova, Y. I. Wolf, J. S. Gootenberg *et al.*, 2015 Discovery and functional characterization of diverse class 2 CRISPR-Cas systems. *Mol. Cell* 60: 385–397.
- Sternberg, S. H., and J. A. Doudna, 2015 Expanding the biologist's toolkit with CRISPR-Cas9. *Mol. Cell* 58: 568–574.
- Wang, T., J. J. Wei, D. M. Sabatini, and E. S. Lander, 2014 Genetic screens in human cells using the CRISPR-Cas9 system. *Science* 343: 80–84.
- Wright, A. V., J. K. Nunez, and J. A. Doudna, 2016 Biology and applications of CRISPR systems: harnessing nature's toolbox for genome engineering. *Cell* 164: 29–44.
- Yang, H., H. Wang, C. S. Shivalila, A. W. Cheng, L. Shi *et al.*, 2013a One-step generation of mice carrying reporter and conditional alleles by CRISPR/Cas-mediated genome engineering. *Cell* 154: 1370–1379.
- Yang, L., M. Guell, S. Byrne, J. L. Yang, A. De Los Angeles *et al.*, 2013b Optimization of scarless human stem cell genome editing. *Nucleic Acids Res.* 41: 9049–9061.

Communicating editor: A. S. McCallion