



Powerful images. Clear answers.



Manage Patient's concerns about
Atypical Femur Fracture*



Vertebral Fracture Assessment –
a critical part of a complete
fracture risk assessment



Advanced Body Composition®
Assessment – the power to
see what's inside

Contact your Hologic rep today at BSHSalesSupportUS@hologic.com

PAID ADVERTISEMENT

*Incomplete Atypical Femur Fractures imaged with a Hologic densitometer, courtesy of Prof. Cheung, University of Toronto

ADS-02018 Rev 003 (10/19) Hologic Inc. ©2019 All rights reserved. Hologic, Advanced Body Composition, The Science of Sure and associated logos are trademarks and/or registered trademarks of Hologic, Inc., and/or its subsidiaries in the United States and/or other countries. This information is intended for medical professionals in the U.S. and other markets and is not intended as a product solicitation or promotion where such activities are prohibited. Because Hologic materials are distributed through websites, eBroadcasts and tradeshows, it is not always possible to control where such materials appear. For specific information on what products are available for sale in a particular country, please contact your local Hologic representative.

www.hologic.com | dxaperformance.com | 1.800.442.9892

American Society for Bone and Mineral Research- Orthopaedic Research Society Joint Task Force Report on Cell-Based Therapies

Regis J O'Keefe,¹ Rocky S Tuan,² Nancy E Lane,³ Frank Barry,⁴ Bruce A Bunnell,⁵ Céline Colnot,⁶ Matthew T Drake,⁷ Hicham Drissi,⁸ Lisa A Fortier,⁹ Robert E Guldberg,¹⁰ David G Little,¹¹ Mary F Marshall,¹² Jeremy J Mao,¹³ Norimasa Nakamura,¹⁴ Pamela G Robey,¹⁵ Vicki Rosen,¹⁶ David W Rowe,¹⁷ and Edward M. Schwarz¹⁸

¹Department of Orthopaedic Surgery, School of Medicine, Washington University, St. Louis, MO, USA

²The Chinese University of Hong Kong, Institute for Tissue Engineering and Regenerative Medicine, Hong Kong

³Department of Medicine, University of California, Davis, CA, USA

⁴Regenerative Medicine Institute, National University of Ireland Galway, Galway, Ireland

⁵Department of Pharmacology, School of Medicine, Tulane University, New Orleans, LA, USA

⁶INSERM UMR 1163, Imagine Institute, Paris, France

⁷Department of Endocrinology, Mayo Clinic, Rochester, MN, USA

⁸Department of Orthopaedics, Emory Healthcare, Emory University, Tucker, GA, USA

⁹College of Veterinary Medicine, Cornell University, Ithaca, NY, USA

¹⁰Phil and Penny Knight Campus for Accelerating Scientific Impact, University of Oregon, Eugene, OR, USA

¹¹Orthopaedic Research and Biotechnology, Kids Research Institute, Westmead, Australia

¹²Center for Biomedical Ethics and Humanities, University of Virginia, Charlottesville, VA, USA

¹³Division of Orthodontics, College of Dental Medicine, Columbia University, New York, NY, USA

¹⁴Institute for Medical Science in Sports, Osaka Health Science University, Osaka, Japan

¹⁵Craniofacial and Skeletal Diseases Branch, National Institute of Dental and Craniofacial Research, National Institutes of Health, U.S. Department of Health and Human Services, Bethesda, MD, USA

¹⁶Department of Developmental Biology, Harvard School of Dental Medicine, Harvard University, Boston, MA, USA

¹⁷Department of Reconstructive Sciences, University of Connecticut Health Center, Farmington, CT, USA

¹⁸Department of Orthopaedics, University of Rochester, Rochester, NY, USA

ABSTRACT

Cell-based therapies, defined here as the delivery of cells *in vivo* to treat disease, have recently gained increasing public attention as a potentially promising approach to restore structure and function to musculoskeletal tissues. Although cell-based therapy has the potential to improve the treatment of disorders of the musculoskeletal system, there is also the possibility of misuse and misrepresentation of the efficacy of such treatments. The medical literature contains anecdotal reports and research studies, along with web-based marketing and patient testimonials supporting cell-based therapy. Both the American Society for Bone and Mineral Research (ASBMR) and the Orthopaedic Research Society (ORS) are committed to ensuring that the potential of cell-based therapies is realized through rigorous, reproducible, and clinically meaningful scientific discovery. The two organizations convened a multidisciplinary and international Task Force composed of physicians, surgeons, and scientists who are recognized experts in the development and use of cell-based therapies. The Task Force was charged with defining the state-of-the-art in cell-based therapies and identifying the gaps in knowledge and methodologies that should guide the research agenda. The efforts of this Task Force are designed to provide researchers and clinicians with a better understanding of the current state of the science and research needed to advance the study and use of cell-based therapies for skeletal tissues. The design and implementation of rigorous, thorough protocols will be critical to leveraging these innovative treatments and optimizing clinical and functional patient outcomes. In addition to providing specific recommendations and ethical considerations for preclinical and clinical investigations, this report concludes with an outline to

Received in original form February 15, 2019; revised form May 28, 2019; accepted June 13, 2019.

Address correspondence to: Regis J O'Keefe, MD, PhD, Department of Orthopaedic Surgery, School of Medicine, Washington University, Campus Box 8233, 660 Euclid Avenue, St. Louis, MO 63110, USA. E-mail: okeefe@wudosis.wustl.edu

Additional Supporting Information may be found in the online version of this article.

Journal of Bone and Mineral Research, Vol. XX, No. X, Month 2019, pp XXXX-XXXXDOI: 10.1002/jbmr.3839© 2019 American Society for Bone and Mineral Research

Journal of Bone and Mineral Research, Vol. 00, No. 00, Month 2019, pp 1–15.

DOI: 10.1002/jbmr.3839

© 2019 American Society for Bone and Mineral Research

Executive Summary

Despite great advances in restorative surgeries involving prosthetic devices, there has been limited progress in the development of biological technologies for musculoskeletal repair. Cell-based therapies, defined here as the delivery of cells in vivo to treat disease, have recently gained increasing public attention as a potentially promising approach to restore structure and function to musculoskeletal tissues. Although cell-based therapy has the potential to improve the treatment of genetic, degenerative, inflammatory, and traumatic disorders of the musculoskeletal system, there is also the possibility of misuse and misrepresentation of the efficacy of such treatments. The medical literature contains anecdotal reports and research studies, along with web-based marketing and patient testimonials supporting cell-based therapy.

Both the American Society for Bone and Mineral Research (ASBMR) and the Orthopaedic Research Society (ORS) are committed to ensuring that the potential of cell-based therapies is realized through rigorous, reproducible, and clinically meaningful scientific discovery. In response, the two organizations convened a multidisciplinary and international Task Force composed of physicians, surgeons, and scientists who are recognized experts in the development and use of cell-based therapies. The Task Force charge was to define the state-of-the-art in cell-based therapies and to identify the gaps in knowledge and methodologies that should guide the research agenda.

Task Force charges

The objective of this Task Force report is to provide guidance to investigators, clinicians, and the general public about the potential and challenges of cell-based therapies for both soft and mineralized skeletal structures. See Table 1 for the complete Task Force charges that informed development of this report and the recommendations included herein. Candidate sources of stem/progenitor cells are reviewed, and optimized experimental protocols for assessing their progenitor and healing properties in animal models are presented. The Task Force examined the diverse range of affected tissues to assess the current state of cell therapy and developed recommendations for investigators and observers regarding evidence of potential clinical efficacy of cell based therapy

Task Force review process

The ASBMR-ORS Task Force reviewed nearly 400 manuscripts in which cell-based therapies were used in animal models to promote tissue repair in musculoskeletal tissues. Outcomes were classified into five categories: (i) radiographic; (ii) histologic, including tissue organization and biochemical or molecular composition of resulting tissue; (iii) donor cell tracking; (iv) function of resulting tissue; and (v) non-target or systemic effects (Table 2). The overall judgment of how these outcomes were evaluated included:

1. Radiographic and histologic assessment: The methods of X-ray, micro-computed tomography (μ CT), and magnetic resonance imaging were among the most widely used for outcome measures. Also widely used were histological methods that assessed the efficacy of cell-based therapy. However, histological methods primarily relied on non-quantifiable approaches (eg, hematoxylin and eosin staining) and often did not employ more rigorous analytical histomorphometric approaches. The use of more detailed analysis of the repair tissue using specialized staining, immunohistochemistry, or RNA expression assays, such as in situ hybridization, was less frequent and the utilization varied between tissues. When specific matrix components were assessed, only a limited number of protein markers were used, most commonly collagen types and aggrecan.
2. Functional criteria: Surprisingly, relatively few studies carefully considered functional criteria, such as the measurement of mechanical properties and pain. This may be partly due to the generally short duration of the studies and the challenge of defining function in different animal models. In addition, few manuscripts addressed non-target or systemic effects of treatment. Because transplanted cells, eg, mesenchymal stem/progenitor cells (MSCs) may have biological effects, such as influencing host immune-inflammatory responses,^(1,2) it is essential that perturbations of immune function, and other non-target functions, be considered in future studies.

Table 1. ASBMR-ORS Task Force Charges

Task Force charge
1. Make recommendations for provisional case definitions of cell-based therapies, including cell sources and target tissues, so that subsequent studies will report using common language and avoid ambiguity due to the complexity of the cell preparatory steps.
2. Address the specific requirements of tissue type, anatomical site and location, underlying disease state, host (gender and/or age), and local environmental status.
3. Carefully review the current available information to assess what is known and what is not known regarding different cell-based therapies, cellular sources, and protocols for addressing specific target tissues. Specifically, the literature will be reviewed and characterized according to whether the evidence is based on in vitro, in vivo (ie, animal models), or clinical reports.
4. Review the available noninvasive diagnostic (eg, biomarkers) and imaging techniques for characterizing the outcome of cell-based therapies.
5. Identify the key questions that the scientific community should address, and recommend a research agenda to elucidate the best approaches for cell-based therapy.
6. Establish criteria for assessing potential biological and clinical efficacy, and develop guidelines appropriate for the claimed use of each cell-based therapy.

Table 2. Percentage of Studies That Measured Various Outcome Criteria

Outcome criteria	Bone	Cartilage	Disc	Meniscus	Tendon/ligament
Radiographic	64	40	70	20	26
Histological	81	78	92	76	82
Origin of cellular components	45	20	68	56	36
Tissue organization	41	58	74	84	85
Biochemical/molecular	21	56	50	32	62
Functional	26	42	18	36	13
Off-target tissue effects	11	0	2	44	15

All values are percentages.

3. Cell fate: Evaluation of cell fate is critical for assessing the efficacy of cell-based therapies. Despite its importance, few studies performed cell tracing or described the localization or persistence of donor cells in the host tissue. In studies where donor cell fate was followed, cells were tracked using a variety of techniques, including those using detectable transgene reporters encoding fluorescent proteins (ie, green fluorescent protein [GFP], red fluorescent protein [RFP]) or enzyme (LacZ) for direct detection of donor cells or species-specific markers/antigens in cases of xenogeneic transplantation.

Overview of Task Force findings and recommendations

Task Force recommendations are summarized in Table 3. Guidance for conducting preclinical and clinical studies of cell-based therapies is offered as a part of these recommendations (see Key Question 5, below). The Appendices A through F provide more details regarding publications related to cell based therapies in various musculoskeletal tissues and also detail supplementary information on how to determine the cell autonomous and non-autonomous effects of a donor population being used for bone regeneration.

Conclusions

The efforts of this Task Force are designed to provide researchers and clinicians with a better understanding of the current state of the science and research needed to advance the study and use of cell-based therapies for skeletal tissues. The design and implementation of rigorous, thorough protocols will be critical to leveraging these innovative treatments and optimizing clinical and functional patient outcomes. In addition to providing specific recommendations and ethical considerations for preclinical and clinical investigations, this report concludes with an outline to address knowledge gaps in how to determine the cell autonomous and non-autonomous effects of a donor population used for bone regeneration. Currently, there is no proof of efficacy of these treatments given the lack of rigorous clinical studies and randomized clinical trials, and that these therapies should thus be considered experimental.

Introduction

Bone, articular/hyaline cartilage, intervertebral disc, meniscus, tendon, and ligaments are all musculoskeletal tissues that function to provide mechanical support and permit locomotion. The support comes from highly organized extracellular matrices, including collagens, proteoglycans, and glycoproteins, intermixed with carbonate-rich apatite (in the case of bone and hypertrophic cartilage), that modulate the mechanical properties of the tissues. Among the musculoskeletal tissues, there is wide

variability in terms of mechanical properties and function, the level of tissue oxygenation, cell turnover, and regenerative capacity. Loss of structure and function of musculoskeletal tissues is the principal cause of physical disability. Furthermore, this loss poses severe challenges to quality of life and presents heavy disease burden, particularly for the aged population.

Despite the great advances in restorative surgeries involving prosthetic devices, there has been limited progress in the development of biological technologies for musculoskeletal repair. Cell-based therapies have recently gained increasing public attention as a potentially promising approach to restore structure and function to musculoskeletal tissues. In fact, the technique of autologous chondrocyte implantation (ACI), developed two decades ago, which involves surgical implantation of a patient's own cartilage cells and chondrocytes to repair focal articular cartilage defects in the joint, is the earliest effective clinical cell therapy procedure that remains continuously practiced.⁽³⁾ A recent entry is stem/progenitor cell-based therapy, which has captured the public's attention because of the possibility of supplying sufficient numbers of cells that can differentiate into skeletal cells and provide tissue-appropriate signals necessary for tissue regeneration.

Although cell-based therapy has potential to improve the treatment of genetic, degenerative, inflammatory, and traumatic disorders of the musculoskeletal system, there is also the possibility of misuse and misrepresentation of the efficacy of such treatments. The medical literature contains research studies and anecdotal reports and research studies, along with web-based marketing and patient testimonials supporting cell-based therapy.

- Both the ASBMR and ORS are committed to ensuring that the potential of cell-based therapies is realized through thorough, rigorous, reproducible, and clinically meaningful scientific discovery. In response, the two organizations convened a multidisciplinary and international Task Force composed of physicians, surgeons, and scientists who are recognized experts in the development and use of cell-based therapies. The task force leaders and members were selected by each society for their expertise. Individual subgroup members searched the PubMed database for studies published in English between years 1995 and 2019 related to the subject area, the studies were reviewed by the subgroup, and studies were included in the report if they provided information relevant to the study topic, and utilized good experimental design and technique. In completion of Appendices B through F (Bone, Cartilage, Disc, Meniscus, and Tendon and Ligament cell-based therapy articles), the Task Force performed a detailed search with inclusion of articles published through 2015.

Each task force was asked to review all published studies in English published between years 1995 and 2019. This report reflects the findings and recommendations of the Task Force in

Table 3. Summary of Findings and Recommendations From the ASBMR-ORS Task Force

Key question	Primary findings and recommendations
1. What is the current knowledge of limitations of cell-based therapies according to cellular source?	<ul style="list-style-type: none">• Pluripotent stem/progenitor cells present unique challenges in terms of differentiation efficacy (compared to multipotent stem/progenitor cells) and in terms of safety. Further standardization and research into safety methods are needed before these models can be better leveraged for clinical use.• Connective tissue-specific stem/progenitor cells require better justification and nomenclature in order to differentiate the source of the progenitor population used for a particular cell therapy.• Markers for human and murine BMSCs are currently too nonspecific and nonexclusive.• The extent to which non-skeletal-derived MSCs contribute to the production of functional skeletal tissues remains to be determined.
2. What is the current knowledge of the therapeutic utility of cell-derived products based on cellular source?	<ul style="list-style-type: none">• The efficacy of platelet-rich plasma is not yet established, but evidence exists supporting its use in bone regeneration, cartilage repair, osteoarthritis treatment, and tendon/ligament and meniscal repair.• Animal models using cell-derived conditioned medium preparations have demonstrated a benefit in healing skeletal tissues, but human studies are lacking and the efficacy of conditioned medium preparations remains unclear.• EVs derived from cells may mediate intracellular communication and thereby affect repair and disease processes.
3. What is the current knowledge of tissue-specific animal models of cell-based therapies?	<ul style="list-style-type: none">• Mice are commonly used to study cell-based therapies for osteoporosis, osteogenesis imperfecta, fracture healing, and ectopic bone formation. Findings usually require replication in larger animals before being studied in clinical protocols.• For cartilage models, larger animals—including horses, dogs, goats, and sheep—are preferable to rodents and rabbits, although rabbit models are frequently used.• Rabbits and rats are commonly used in intervertebral disc disease and injury models, but larger animals such as sheep and dogs are preferable due to better generalizability to humans.• Sheep, pigs, and primates are useful in models of meniscus healing because they possess knee joint anatomy similar to humans, but these models still have certain limitations (eg, differences in biped versus quadruped meniscus and cartilage contact mechanics).• Rabbit and rodent models are convenient to study stem/progenitor cell therapies for acute tendon injury and repair. However, larger animal models (ie, horse, pig, dog, and sheep) may better meet FDA guidelines for demonstrating the efficacy of cell-based therapies and delivery devices in humans. The suitability of an animal model for tendon/ligament repair will depend on the tissue being examined and the objectives of the study.
4. What are the Task Force–recommended criteria for interpreting a cell-based regenerative experiment?	<ul style="list-style-type: none">• Success should be measured by the ability of cell-based therapies to regenerate or repair degenerated or injured tissue, decrease pain, and restore structure, mechanical properties, and function.• Interventions must be compared to vehicle control using validated outcome measures that include both functional and pain assessments.• Studies should consider cell autonomous and non-cell autonomous mechanisms of influence of cell-based therapies.
5. What are the Task Force recommendations for preclinical and clinical studies of cell-based therapies?	<p>Recommendations for preclinical studies</p> <ul style="list-style-type: none">• The Task Force found no preferred, standardized animal model for preclinical studies of cell-based therapies. Both large-animal and small-animal models have deficiencies and either singly and in combination fail to reproduce the biomechanics or biology seen in humans.• The Task Force recommends animal models be chosen based on size and anatomical considerations as well as protocol design and objectives, including cost, technical challenges, use of both autologous and allogeneic cells, potential complications related to immune

(Continues)

rejection, and the degree to which the model mimics human anatomy and disease.

- The optimal preclinical approach would be to initiate studies in small animals that focus on cellular, molecular, functional outcomes, mechanical properties, and genetic characterization, and, if these models provide proof of principle, perform follow-up trials in larger, more clinically relevant animals if indicated.
- Immune reactions in animals should be considered when assessing reparative potential in humans.
- A combination of evidence from in vitro and large and small animal in vivo studies may be needed to obtain FDA clearance.
- Interpretation of the role of donor stem/progenitor cells in tissue repair is critical.
- More research is needed investigating cell-based therapies in various mesenchymal tissues as well as noninvasive assessments of tissue composition, structure, and function.
- Centralized data resources, such as the NIH-supported National Swine Resource and Research Center, can play a valuable role in advancing this line of research. Recommendations for clinical studies
- The study methodology must be of the highest quality, including the use of appropriate design, blinding, techniques to prevent bias, validated quantitative outcome measures, and appropriate statistical techniques.
- Development and use of noninvasive measures of human tissue composition, structure, and mechanical function is essential.
- Patient-reported outcome measures are sensitive and valuable tools to assess functional improvement, tissue structure, pain, and quality of life. Research and clinical ethical considerations
- All animal studies should be conducted with strict adherence to ethical guidelines and with approval from an Institutional Animal Care and Use Committee Review Board.
- Although cell-based therapies may seem appealing due to their novelty and innovativeness, patients should be clearly informed that little data exists in either larger-animal preclinical studies or randomized clinical trials to support the use of cell-based therapies. However, cell-based therapies thus far appear to be generally safe and well-tolerated.
- Patients also should be made aware that use of these therapies is often “off label” and unlikely to be reimbursed by medical insurance.
- For patients to truly give informed consent, a neutral or second-opinion physician should be consulted to explain the benefits and potential risks for patients regarding receiving or not receiving the treatment.
- Given the lack of rigorous evidence, the Task Force cannot currently recommend local or systemic stem/progenitor cell therapy for skeletal tissue repair and regeneration and encourage clinical trials for treatment protocols to receive FDA approval.

BMSC = bone marrow stromal cell; MSC = mesenchymal stem/progenitor cell; EV = extracellular vesicle; FDA = U.S. Food and Drug Administration; NIH = National Institutes of Health.

response to its charge of defining the state of the art in cell-based therapies and identifying the gaps in knowledge and methodologies that should guide the research agenda.

Key Question 1: What Is the Current Knowledge of Limitations of Cell-Based Therapies According to Cellular Source?

- Pluripotent stem/progenitor cells present unique challenges in terms of differentiation efficacy (compared to multipotent

stem/progenitor cells) and in terms of safety. Therapeutic use of these cells in clinical practice is not recommended until further standardization and safety validation.

- Connective tissue-specific stem/progenitor cells require additional justification and nomenclature in order to differentiate the source of the progenitor population used for a particular cell therapy.
- Markers for human and murine bone marrow stromal cells (BMSCs) are currently too nonspecific and nonexclusive.
- It is unclear the extent to which non-skeletal-tissue-derived MSCs are useful in producing functional skeletal tissues.

Stem/progenitor cells

Based on the fact that virtually all tissues undergo renewal, albeit at highly variable rates, enduring tissue regeneration depends on the presence of a subset of tissue-specific stem/progenitor cells within a given population that can fuel tissue renewal. Hence, considerable effort has been focused on identifying cell sources that have the ability to maintain tissue homeostasis, including musculoskeletal tissues.⁽⁴⁾ Stem/progenitor cells in mammals can be divided into two broad categories: pluripotent stem/progenitor cells and postnatal stem/progenitor cells from various sources (see <https://stemcells.nih.gov/info/basics/1.htm> for more information).

Pluripotent stem cells (embryonic stem cells) and induced pluripotent stem cells

Embryonic stem cells (ESCs) are pluripotent stem cells derived experimentally by extraction from the inner cell mass of an early-stage embryo, the blastocyst, whereas postnatal stem/progenitor cells are found in different organs and tissues. Pluripotency refers to the potential of a stem cell to differentiate into cells of all three germ layers—endoderm, mesoderm, and ectoderm. In addition to ESCs, induced pluripotent stem cells (iPSCs) can be generated from adult somatic cells by reprogramming with essential pluripotency transcription factors by a variety of methods and are nearly identical to ESCs.

Because of their highly uncommitted state, pluripotent ESCs and iPSCs present more challenges in terms of being induced to differentiate into a specific musculoskeletal cell type compared to multipotent stem/progenitor cells, such as postnatal stem/progenitor cells. Osteogenic differentiation provides an example. There have been a number of reports on the differentiation of human ESCs (hESCs) and iPSCs into osteogenic cells through the formation of embryoid bodies, spontaneous differentiation, indirect co-culture with osteogenic cells, treatment with conditioned medium generated by osteogenic cells, or use of various schemes for direct osteogenic differentiation.⁽⁵⁾ However, for the most part, the results have relied on *in vitro* differentiation assays that may not reflect capabilities *in vivo*, and results from limited studies involving *in vivo* transplantation are not conclusive.^(6–9) Furthermore, most studies use ectopic transplantation sites rather than sites within an injured skeletal tissue that would better mimic a clinical scenario. The lack of differentiation into a functional tissue is likely because these cells have not undergone a developmental process that commits them to a particular lineage.⁽¹⁰⁾ Thus, there is the issue of differentiation efficacy. Another concern is safety, such as (i) the potential for tumor formation, and (ii) the potential for immunological rejection when allogeneic ESCs or iPSCs are introduced *in vivo*.⁽¹¹⁾ More standardization and safer methods are clearly needed to bring pluripotent stem cells into clinical use.

Connective tissue-specific stem/progenitor cells

BMSCs are non-hematopoietic adherent cells first identified and characterized by Friedenstein⁽¹²⁾ A subset of BMSCs are skeletal progenitor cells that differentiate into cartilage, bone, hematopoiesis-supportive stroma, and marrow adipocytes based on rigorous clonal and differentiation assays performed *in vivo* (bone, fat) and *in vitro* (cartilage). However, the original concept of a tissue-specific stem/progenitor cell for bone, was later altered to include other mesodermal derivatives such as muscle, tendon, and ligament, and BMSC terminology was altered so that these cells are considered synonymous with

MSCs. A better justification and nomenclature of stem/progenitor cells from other connective tissue sources need to be developed to clearly differentiate the source of a progenitor population used for a particular cell therapy.

A number of markers of BMSCs have been identified (such as CD73, CD90, CD105, and CD146, along with lack of expression of CD11b, CD14, CD19, CD34, CD45, and HLA-DR).⁽¹³⁾ However, the markers used are neither specific nor exclusive, because they are also expressed in many adherent fibroblastic cell populations. Although such “markers” have been utilized to identify cells derived from periosteum, synovium, dental pulp, and periodontal ligament cells (tissues associated with the skeleton) that are very similar to BMSCs and are capable of differentiating into cartilage, bone, and fat *in vitro*,⁽¹⁴⁾ it should be noted that these various cell types are not identical in their *in vivo* differentiation capacity.^(15–18) Furthermore, the standard *in vitro* assays used to claim “tri-lineage” differentiation are often not predictive of *in vivo* differentiation capacity.⁽¹⁹⁾ Whether and how these differences may have arisen from their different native tissue microenvironment needs to be examined.

In the last several years, lineage tracing studies have identified a self-renewing multipotent cell population called the skeletal stem cell (SCC) present in bone tissue of humans (PDPN+, CD146–, CD73+, and CD164+) and mice (Tie2–, integrin AlphaV+, Thy–, 6C3–, CD105–, and CD200+) that has differentiation restricted to the osteoblast, chondrocyte, and stromal cell lineage, and not adipocytes. In both mice and humans, these cells, when implanted into the renal capsule form ectopic bone and cartilage, and the cells support marrow formation. The SCC population is expanded in culture by BMP-2 and *in vivo* by bone fracture.^(20–22) Another study identified a rare population of Gremlin 1+ (Grem1+) cells in the metaphysis and tissues adjacent to the growth plate in mice. Similar to the SCC population, Grem1+ cells are BMP-2 responsive, self-renewing, and formed bone, cartilage, and stromal tissues, with limited adipocyte differentiation. During development and aging, the Grem1+ population generates articular and growth plate cartilage, and is found in periosteum, osteoblasts, and osteocytes. Like SCC, the Grem1+ cells expand in fractures. Moreover, Grem1+ cells transplanted into fractures engraft, self-renew, and form osteoblasts, and Grem1+ cells can be harvested and re-expanded. The elimination of Grem1+ cells in developing mice results in reduced bone mass.⁽²³⁾

Thus, there is a rare self-renewing SCC population in bone that is separate from the bone sinusoids that does not strictly meet the definition of an MSC, because it lacks capacity for adipocyte differentiation. The SCC is necessary for development of bone and cartilage, maintenance of the adult skeleton, and bone repair. These approaches show the utility of cell lineage tracing in defining progenitor cell populations *in vivo*. However, the manner in which these rare cell populations are applicable for a cell-based therapy approach remains to be determined. Similarly, it is unclear whether an analogous tissue lineage-restricted stem cell is present in other musculoskeletal tissues, such as tendon, ligament, or disc.

Nonskeletal tissue-derived MSCs

MSC-like cells obtained from nonskeletal sources (eg, muscle, cord blood, Wharton’s jelly, dermal, adipose, and amniotic fluid MSCs) have also been promoted as another source of progenitors for cell therapy of skeletal tissues.^(24,25) However, many of the cells used in these studies were pretreated with chondrogenic/osteogenic

factors, and whether these nonskeletal cells produce functional bone and associated tissues, or whether they induce local cells to undergo a repair process *in vivo*,^(26,27) is not known.

Perivascular stromal cells from various tissues have stem cell characteristics and differentiate into osteoblasts, chondrocytes, and adipocytes. In bone marrow, cell lineage studies show that LepR+ perivascular cells located around both arterioles and sinusoids account for nearly all of the fibroblastic colony-forming unit (CFU-F) in bone marrow. Lineage tracing shows that LepR+ perivascular cells are the major source of bone and adipocytes in bone marrow in adult mice. Moreover, they are involved in the regeneration of bone marrow following radiation and participate in fracture healing.⁽²⁸⁾ Perivascular cells isolated from fat, muscle, pancreas, skin, lung, brain, eye, gut, bone marrow, and umbilical cord are NG2+, CD146+, PDGF-R beta+, and alpha SMA+.⁽²⁹⁾ These cells are multipotent and differentiate into osteoblasts, chondrocytes, and adipocytes. In addition, a Gli1+ stromal cell population present in the adventitial layer of arteries has stem cell characteristics. Gli1+ cells differentiate into osteoblasts and are responsible for the vascular calcifications that occur in atherosclerosis and in chronic kidney disease.⁽³⁰⁾ Thus, perivascular stromal cells, as well as adventitial cells, are a key source of MSCs in nonskeletal tissue-derived tissues. These cells have been used in animal models in cell-based therapy approaches.^(31–35)

Differentiated skeletal cells

Fully committed cells associated with the skeleton, such as osteoblasts, chondrocytes, and tenocytes, have limited ability to self-renew. However, they may be considered therapeutically useful in situations where there is low tissue turnover. ACI, which uses culture-expanded autologous chondrocytes harvested from the less weight-bearing articular cartilage of the joint, has shown efficacy in the repair of focal cartilage defects.⁽³⁾ It is, however, generally acknowledged that the cultured chondrocytes undergo dedifferentiation and/or hypertrophy *in vitro*, thus compromising the quality of the regenerate cartilage, which is often more fibrous in nature or sometimes undergoes overgrowth and/or calcification. Also, currently ACI procedures are recommended only in young patients and exclude older adults (≥ 45 years old),^(36–38) because this age group has a higher susceptibility to degenerative joint diseases.

Key Question 2: What Is the Current Knowledge About the Therapeutic Utility of Cell-Derived Products Based on Cellular Sources?

- The efficacy of platelet-rich plasma is not yet established, but evidence exists supporting its use in bone regeneration, cartilage repair, osteoarthritis treatment, and tendon/ligament and meniscal repair.
- Animal models using cell-derived conditioned medium preparations have shown a benefit in healing skeletal tissues, but human studies are lacking and the efficacy of conditioned medium preparations remains unclear.
- Extracellular vesicles derived from cells may mediate intracellular communication and thereby affect repair and disease processes.

There are many examples of cell therapy in which positive outcomes were observed without evidence of the donor cells within the repair field. This is particularly true in experiments in

which the treating cells were administered systemically. It is now well established that when administered systemically, skeletal progenitor cells do not efficiently or stably home to a target tissue.^(39,40) Consequently, some successful outcomes are attributed to factors transiently produced by the donor cells that influence a successful host reparative response.^(39,41) This cellular process is classified as a donor cell–nonautonomous effect in that the long-term presence of the donor cells is not required because they did not participate directly in the repair process. If these factors could be identified, they could be used directly without the need for supplying the producing cells. Some of the candidate factors that have been investigated include platelet-rich plasma, conditioned medium, and extracellular vesicles (EVs).⁽⁴²⁾

Platelet-rich plasma

Derived from megakaryocytes, platelets contain a long list of (approximately 300) growth factors, such as platelet-derived growth factor, transforming growth factor beta, and vascular endothelial growth factor, and others such as cytokines, coagulation factors, fibrinolytic factors and proteins, proteases, antiproteases, and lipids. Upon platelet activation, eg, as a result of tissue injury, these factors are released for acute repair but are unlikely to have sustained activity or localization. Consequently, there are efforts to develop preparations, such as platelet-rich plasma (PRP) gels or PRP combined with other biological materials such as demineralized bone matrix or cells, to extend the biological activity of PRP.⁽⁴³⁾ PRP has been used in a number of clinical settings for bone regeneration⁽⁴⁴⁾ cartilage repair,⁽⁴⁵⁾ treatment of osteoarthritis,⁽⁴⁶⁾ and tendon/ligament and meniscal repair.⁽⁴⁷⁾ However, the biological action of PRP, eg, in regulating differentiation of multipotent MSCs,⁽⁴⁸⁾ remains unclear. In addition, the efficacy of PRP in the repair of hard tissues has yet to be established, in view of the wide variability in methods and measures of the studies that have been conducted.

Conditioned medium

Cells from connective tissues (including skeletal tissues) secrete a broad array of growth factors, cytokines, and other biologically active factors. It has been suggested that conditioned medium can exert a beneficial effect on healing of skeletal tissues based on paracrine, immunomodulatory, and immunosuppressive effects that encourage local cells to begin the repair process. A large number of animal studies⁽⁴⁹⁾ and one human study⁽⁵⁰⁾ have used this approach to treat injured skeletal tissues; however, efficacy of this type of therapy remains unestablished.

EVs

Perhaps the first description of EVs (eg, exosomes, ectosomes, microvesicles, microparticles) in the literature comes from the finding of matrix vesicles in cartilage.⁽⁵¹⁾ EVs, first termed microparticles, have since been detected in many bodily fluids. It is now thought that in culture, virtually all cell types release some sort of EV, formed by different processes and of varying size. Results derived from the use of conditioned medium may be due to the presence of EVs in addition to other factors. EVs are proposed to mediate intercellular communications and have been implicated in repair and disease processes by virtue of the cargo that they carry, which may include proteins, biofactors, and RNAs, including specific miRNAs, among others.^(52,53)

Key Question 3: What Is the Current Knowledge About Tissue-Specific Animal Models of Cell-Based Therapies?

- Mice are commonly used to study cell-based therapies for osteoporosis, osteogenesis imperfecta, fracture healing, and ectopic bone formation. Findings usually require replication in larger, more clinically relevant animals before being studied in clinical protocols.
- For cartilage models, larger animals—including dogs, goats, and sheep—are preferable to rodents and rabbits, although rabbit models are frequently used. Horse models have unique benefits but are costly and have large housing-space demands.
- Rabbits and rats are commonly used in intervertebral disc disease and injury models, but larger animals such as sheep and dogs are preferable because of better generalizability to humans.
- Sheep, goats, pigs, and primates are useful in models of meniscus healing because they possess knee joint anatomy similar to humans, but these models still have certain limitations (eg, differences in biped versus quadruped meniscus and cartilage contact mechanics).
- Rabbit and rodent models are convenient to study stem/progenitor cell therapies for acute tendon injury and repair. However, larger animal models (ie, horse, pig, dog, and sheep) may better meet US Food and Drug Administration (FDA) guidelines for demonstrating the efficacy of cell-based therapies and delivery devices in humans. The suitability of an animal model for tendon/ligament repair will depend on the tissue being examined and the objectives of the study.

Bone

Mice are commonly used to study cell-based therapies (eg, BMSCs, adipose-derived stromal cells) in models of osteoporosis, osteogenesis imperfecta, fracture healing, and ectopic bone formation (subcutaneous or intramuscular). Mice allow genetic approaches and have a high tolerance to xenograft engraftment. Humanized mice permit modeling of human cell activity and function. The reviewers recommend that the animal model yield convincing evidence with regard to bone architecture and mechanical properties. Impactful studies should clearly define the changes associated with cell therapy and the mechanism of cell involvement in the regenerative process.

A disadvantage of mice is that confirmation of findings typically requires subsequent experimentation in a larger animal model which more closely mimics the structure and mechanical features of human tissues prior to consideration for use in clinical trials. The rat is also an inexpensive small-animal model, and because it is relatively larger, it may have some advantages for mechanical testing. Gene-editing technologies are now being applied to the rat to provide genetic models that previously were only available in the mouse. Successful outcome should then be tested in larger animal models.

The models of bone disease/injury identified in the review included calvarial defects (46 studies), fracture (seven studies), heterotopic ossification (intramuscular, three studies), heterotopic ossification (subcutaneous, 38 studies), long-bone cortical defect (drill, 13 studies), long-bone segment (acute repair, 28 studies), long-bone segment (chronic repair, three studies),

osteoporosis (nine studies), or other (heterotopic, two studies) (Appendix B).

Cartilage

Rabbits and rats were the most commonly used animals among the 49 preclinical studies that examined cell-based therapies for the regeneration of articular cartilage (Table 4). Rabbit models were the most frequently used. Rabbits are widely available, are relatively low-cost, have simple handling requirements, and have a robust base of literature for comparison.⁽⁵⁶⁾ However, there are several disadvantages inherent in rabbit studies. Pure chondral lesions are difficult to create in the thin cartilage of this species. Also, researchers need to consider the potential for a natural healing response given the smaller size of cartilage defects (<3 mm)⁽³⁾ and the difference in mechanical loading in the rabbit knee relative to humans. Although rodents are widely used to screen new biomaterials and cell-laden constructs, the use of rodents for articular implants is less practical because of their very thin articular cartilage.^(57,58) Furthermore, unlike larger animals, rodent growth plates remain open during adulthood. The epiphysis is therefore more highly vascularized, a feature that may contribute to more robust intrinsic cartilage repair.

Considering clinical relevance, a larger animal model is preferred to approximate the area and thickness of articular cartilage in the human.⁽⁵⁹⁾ Full-thickness cartilage defects can be created without damage to the subchondral plate in dogs.⁽⁶⁰⁾ However, the dog is rarely used for cartilage repair⁽⁶¹⁾ because its status as a companion animal makes it a less attractive option. Goats and sheep are frequently used in cartilage repair models because the knee joints are large enough to create lesions as large as those treated in patients. The larger defect size and thicker cartilage layer permit biochemical assays of cartilage repair tissue as well as biomechanical testing.⁽⁶²⁾

Domestic pigs, minipigs, goats, and sheep, although more expensive, are attractive because of the thicker articular cartilage that more closely mimics the human joint.⁽⁶³⁾ The main advantages of the horse model are the large joint size and thick articular cartilage layer with easy arthroscopic joint access, as well as the actual clinical need in equine veterinary care.^(64,65) Horses can be monitored with respect to the clinical pain response to cartilage repair. A second-look arthroscopy with biopsy is usually the cornerstone of equine studies, allowing assessment of repair progression. However, specialized facilities and care are usually required, and thus this model is often used for late-stage development and pivotal studies.

In our review of 49 preclinical studies of cell-based therapy to promote cartilage repair, two used horses, six used sheep, 15 used rabbits, 10 used rats, six used goats, eight used pigs, one used dogs, and two used mice (Appendix C). Most of the models used an osteochondral defect model (42 studies). Five other studies used a partial-thickness cartilage-defect model, whereas two others used a model of global osteoarthritis from either injury (anterior cruciate ligament [ACL] and medial meniscectomy) or from chemically-induced osteoarthritis.

A total of 14 human studies were identified that examined cell-based therapy for articular cartilage repair. Because ACL therapy is already an FDA-approved procedure, studies using articular chondrocytes as a cell source were not included. All of the studies included full-thickness osteochondral defects. Bone marrow aspirate was the most common source of cells (10 studies) followed by peripheral blood-derived progenitors. One study each used either nasal chondrocytes or synovial progenitor cells.

Table 4. Individual Characteristics for Tissue Repair

Tissue	Indications for treatment	Structural repair goals
Bone	<ul style="list-style-type: none"> Augmentation of standard therapies including: <ul style="list-style-type: none"> Non-union fractures. Delayed-union fractures. Joint or spine fusions or to augment bone formation in osteonecrosis or osteoporosis. May be used in combination with matrix products to augment the healing of bone defects. 	<ul style="list-style-type: none"> Mechanical stability across the fracture/defect site. Union of the fracture/defect at the anatomical site via presence of a cortical bridge or outer cortical shell. Properly aligned bone after healing. Bone mineral density is appropriate for the area.
Cartilage	<ul style="list-style-type: none"> Individuals who are experiencing chronic joint pain. Augmentation of standard therapies, including repair of damaged or degenerated cartilage tissue. May be used to augment osteoarthritis therapy. 	<ul style="list-style-type: none"> Mechanical stability of the joint. Normal and painless range of motion.
Disc	<ul style="list-style-type: none"> Individuals who are experiencing chronic low back pain (greater than 6 months) not responding to conservative therapy, excluding those with spondylolisthesis. 	<ul style="list-style-type: none"> Sufficient strength to support painless weight bearing. Provide structural stability between vertebral bodies and protect spinal canal and nerve roots.
Meniscus	<p>Augmentation of standard therapies including:</p> <ul style="list-style-type: none"> Enhancing or enabling repair of tears with poor potential healing capacity, including those with marginal vascularity, meniscus revision surgeries, meniscal tissue with intrinsic degenerative changes, or meniscal tears in older patients. Replacement of seriously damaged menisci with engineered meniscal constructs populated with endogenous and/or transplanted stem/progenitor cells. Repair/regeneration of horizontal cleavage tears found in degenerative meniscus tissue. 	<ul style="list-style-type: none"> Restoration of full knee range of motion. Absence of joint line tenderness. Absence of pain reproduction with provocative maneuvers such as rotation. Absence of joint effusion. Restoration of the mechanical function of the meniscus in sharing load transmission across the tibiofemoral joint. Enhanced joint stability.
Tendon/ligament	<ul style="list-style-type: none"> Improving the repair and reconstruction of injured tendons and ligaments. Modulating inflammation associated with chronic tendinopathies.⁽⁵⁴⁾ Facilitating tendon-graft or ligament-graft integration in reconstructive surgeries. Promoting graftless ligament repair by inducing the native ligament tissue to regenerate.⁽⁵⁵⁾ Development of artificial tendon and ligament for grafting. 	<ul style="list-style-type: none"> Full restoration of passive and active range of motion. Stability of the joint. Absence of inflammation. Return to daily activities and sports. Restoration of force transmission from muscle to bone or tendon, or the stabilization of a diarthrodial joint or ligament. Improved range of motion and joint stability.

In 6 studies, mesenchymal progenitor cells from bone marrow (four studies), synovium (one study), and nasal cartilage (one study) were expanded in cell cultures prior to delivery into the joint.

Each of the papers was scored according to whether the study assessed the following features in the context of cartilage repair: radiographic criteria, histology, origin of the cell-tissue organization, biochemical criteria, functional outcomes, fate of the transplanted cells, and non-target tissue effects. Each criterion was scored as 0 (not examined); 1 (superficial analysis); or 2 (detailed analysis). A range of scores was identified, but consistently show higher scores in the preclinical models. All of the human studies had level of evidence 3 or 4, and analysis was limited because of the morbidity associated with tissue procurement in humans.

Intervertebral disc

Although there are murine genetic models of intervertebral disc disease, they often lack normal disc formation or develop severe structural alterations during development and thus have not been extensively used to study the role of cell-based therapies for disc regeneration.⁽⁶⁶⁾ The small size of mouse and rat discs

compared to human discs limits their use in the assessment of disc repair methodologies, because nutrient diffusion is not fully evaluated. In addition, the morphology of the rodent disc differs somewhat from humans. Although rabbits and guinea pigs are larger, their discs are still several-fold smaller than human discs, and the tissues do not experience the same degree of loading. In comparison, larger, quadruped animals such as sheep have intravertebral discs that are closer in size and loading to those of humans.

Cell-based therapy for disc repair has been reported in at least 50 studies. Animals utilized include sheep (one study), dogs (five studies), rabbits (19 studies), minipigs (four studies), rats (10 studies), and mice (seven studies) (Appendix D). Models of disease occurring through either natural means (ie, aging as in human disc degeneration) or induced by removing various amounts of nucleus pulposus (NP) tissue (nucleotomy; the majority of animal studies), removal of herniated tissue (at the time of discectomy, two human studies), stab injuries to the annulus fibrosus with or without removal of NP tissue (two studies), or removal of a whole disc (three studies). Most studies tested treatment in the acute injury setting, but in three studies, cell therapy was used at 4, 6, or 12 weeks postinjury in dogs.^(67–69) Chondrodystrophic canines

have been used, but whether these animals had developed premature degenerative disc disease at the time of implant evaluation was not specified.

Meniscus

Rabbits (43%) and rodents (16%) have been frequently used in studies that evaluate the underlying biological and molecular mechanisms of meniscus healing. Rabbits have also been used for preliminary analyses of new stem-cell or progenitor-cell therapy and tissue engineering approaches for the treatment of meniscal injury. Rabbits are less expensive than larger animal models and allow for the use of the larger group sample sizes that are necessary for more comprehensive outcome measures including histology, biochemical and molecular analyses of the repair tissue, and biomechanical testing.

Sheep (8%), pigs (8%), and primates (4%) have been used as large animal models to study meniscus healing. These animals possess knee joints similar in size to the human knee, enabling testing of implants that could be used in humans. However, unlike humans, these animals are quadrupeds with obvious differences in meniscus and cartilage contact mechanics when compared to humans. There is also an inability to control postoperative weight-bearing in most large animal models.

For meniscus, the majority of models used were local acute injury (19 studies). Chronic injury studies (four studies) were also examined. Most meniscus models evaluate tissue formation in a punch defect, although linear tear models have also been used (Appendix E).

Tendon/ligament

Smaller animal models, including rabbits^(70–81) and rodents (rats and mice),^(12,82–97) have been used to evaluate the effectiveness of stem-cell/progenitor-cell therapies in acute tendon injury and repair. Early-stage studies and anatomical considerations might necessitate the use of smaller animals, such as in the case of rotator cuff repair, where the rat is a commonly used model because it resembles the human shoulder anatomy.⁽⁹⁸⁾ Furthermore, if the objective is to track the fate of implanted cells and to mechanistically evaluate their contributions to the repair tissue, rodent models (rats and especially mice) have the distinct advantage of the availability of reporter gene models (eg, GFP, RFP, mTmG, nTnG) and genetic models of conditional gene deletion.

Large animal models, including equine,^(99–102) ovine,^(100,103) swine,^(104,105) and canine,^(106,107) might better meet the FDA's preference for demonstrating efficacy of cellular therapies and delivery devices prior to human approval. The suitability of an animal model for tendon/ligament repair depends on the tissue being examined and the objectives of the study. For example, the canine model is commonly used to study flexor tendon repair because of the ability to simulate zone II injuries, reproduce clinical protocols of multistrand suture repair, and implement physical therapy.^(106,107) Similar arguments can be made in favor of using large animals (eg, pigs) for evaluating cellular therapies in ACL repair and reconstruction.⁽¹⁰⁴⁾

For tendon/ligament, various acute injury and repair models have been examined, including flexor tendon,^(72,73,99–101,105–107) rotator cuff tendon,^(83–86) Achilles tendon,^(12,75,81,89,91,95,103) patellar tendon,^(70,71,74,76–80,87,88,94,96,97) and ACL.^(92,93,104) Furthermore, a number of equine studies have focused on veterinary clinical applications of autologous stem/progenitor cells to treat chronic overuse injury of the superficial digital flexor tendon in racehorses using ultrasound-guided intratendinous injections^(99–102) (Appendix F).

Key Question 4: What Are the Task Force–Recommended Criteria for Interpreting a Cell-Based Regenerative Experiment?

- Success should be measured by the ability of cell-based therapies to regenerate or repair degenerated or injured tissue and to restore functioning.
- Interventions must be compared to vehicle control using validated outcome measures that include functional and pain assessments.
- Studies should consider cell autonomous and non-cell autonomous mechanisms of influence of cell-based therapies.

Understanding the role of donor cells in the repair process will provide specific information on the mechanism of action and the basis for therapeutic product development. The measure of success of cell-based therapies is the ability of the intervention to result in regeneration or repair of degenerated or injured tissue with restoration of structure, mechanical properties, and function. A key aspect is comparison to a vehicle control using validated outcome measurements that include functional assessment and evaluation of pain.

There are two major mechanisms through which cell-based therapies may influence the regenerative or reparative process, and both are equally relevant. These mechanisms are (i) cell autonomous and (ii) non-cell autonomous. In cell-autonomous therapies, the delivered cells have a therapeutic effect in part through being incorporated in the regenerating tissue and participating directly in the repair process. In non-cell autonomous therapies, the cells secrete factors, such as growth factors, cytokines, extracellular vesicles, or other regulatory signals, that influence the behavior of the host cell population in a manner that leads to tissue repair or regeneration. This can occur by their acting directly on host tissue-specific progenitor cells or can be indirect by influencing other processes such as vascularization or regulation of the immune response. It should be recognized that cell-based therapies which demonstrate a cell autonomous mechanism may additionally also act in a non-cell autonomous manner at the same time.

Key Question 5: What Are the Task Force Recommendations for Preclinical and Clinical Studies of Cell-Based Therapies?

Recommendations for preclinical studies

- The Task Force found no preferred, standardized animal model for preclinical studies of cell-based therapies.
- The Task Force recommends animal models be chosen based on size and anatomical considerations as well as protocol design and objectives, including cost, technical challenges, use of both autologous and allogeneic cells, potential complications related to immune rejection, and the degree to which the model mimics human anatomy and disease.
- The optimal preclinical approach would be to initiate studies in small animals that focus on cellular, molecular, functional outcomes, mechanical properties, and genetic characterization, and, if these models provide proof of principle, perform follow-up trials in larger, more clinically relevant animals if indicated.
- Immune reactions in animals should be considered when assessing the potential for use in humans.

- A combination of evidence from in vitro and large and small animal in vivo studies may be needed to obtain FDA clearance.
- Interpretation of the role of donor stem/progenitor cells in tissue repair is critical.
- More research is needed investigating cell-based therapies in various mesenchymal tissues as well as noninvasive assessments of tissue composition, structure, and function.
- Large-animal models are essential but are limited by expenses and technical demand. Centralized data resources, such as the NIH-supported National Swine Resource and Research Center (<http://www.nsrc.missouri.edu/index.asp>), can play a valuable role in advancing this line of research.

Recommendations for clinical studies

- Study methodology must be of the highest quality, including the use of appropriate design, blinding, techniques to prevent bias, validated outcome measures, and appropriate statistical techniques.
- Development and use of noninvasive measures of human tissue composition, structure, and mechanical function is essential.
- Patient-reported outcome measures (PROMs) are sensitive and valuable tools to assess functional improvement, pain, and quality of life.

Research and clinical ethical considerations

- All animal studies should be conducted with strict adherence to ethical guidelines and with approval from the appropriate institutional animal care and use committee.
- Although these treatments may seem appealing because of their novelty and innovation, patients should be clearly informed that little data exists in either larger-animal preclinical studies or randomized clinical trials to support the use of cell-based therapies. However, cell-based therapies thus far appear to be safe and well-tolerated.
- Patients also should be made aware that use of these therapies is often “off label” and unlikely to be reimbursed by medical insurance.
- For patients to truly give informed consent, a neutral or second-opinion physician should be consulted to explain the benefits and potential risks of the patient receiving or not receiving the treatment.
- Given the lack of rigorous evidence, the Task Force cannot currently recommend local or systemic stem-cell/progenitor-cell therapy for skeletal tissue repair and regeneration.

Preclinical animal models

Overall, the reviewers found no preferred, standard animal model for a preclinical study. Thus, the Task Force recommends that, in addition to size and anatomical considerations, the choice of the animal model should thoughtfully consider other aspects of the study objectives and design, including the cost, technical challenges, the potential to use both autologous and allogeneic cells, and the degree to which the model mimics human disease.

An optimal experimental approach to evaluating stem/progenitor cells for enhancing musculoskeletal tissue repair/regeneration would be to initiate studies in small animals that focus on cellular, molecular, functional, and mechanical outcome measures, and allow the examination of genetic factors influencing regeneration. Once these models provide proof of principle for the utility of a specific cell population in augmentation of repair, additional

investigation would be completed in larger animals that more closely model the anatomical size and weight-bearing characteristics of human skeletal tissues. Subsequent successful outcomes in large-animal models, with inclusion of appropriate safety and efficacy profiles, would identify prime candidates for human clinical trials.

In studies where the reparative potential of human cells is assessed, immune reaction in the animal model is a challenge. Human cells are sometimes assessed in less-optimal models, including immunodeficient nude rat models or larger animals with drug-mediated immunosuppression, conditions that may influence the repair process. Still, with regard to immunological considerations of the efficacy of allogeneic progenitor cell therapy, mouse models of selective depletion of a subpopulation of T cells⁽¹⁰⁸⁾ or pig models of genetically manipulated major histocompatibility complex (ie, swine leukocyte antigen)⁽¹⁰⁹⁾ can be particularly useful.

The FDA will require clear and unequivocal evidence for the cellular basis of stem/progenitor-based therapies for musculoskeletal tissue repair. A combination of models will provide support for cell-based therapeutics, including in vitro three-dimensional models of tissue regeneration/tissue chip models,⁽¹¹⁰⁾ small-animal models; and large-animal models. Fundamental requirements for adequate interpretation of cell-based therapies include the ability to: (i) track the donor cell population; (ii) determine cellular fate and differentiation; and (iii) define the role of the cell population in restoring a stable and mechanically functional tissue. (Additional information about determining cell fate can be found in Appendix A).

Interpretation of the role of donor stem/progenitor cells in the repair process is critical. Persistence of donor-derived cells during the repair should preferably be observed, although non-cell autonomous mediation of therapeutic effects must also be considered. Donor cells may undergo terminal differentiation and remain in the tissue as mature cells or may have a critical early and more transient role in driving the repair process. Cell-based therapy should be investigated in relevant models across the various mesenchymal tissues. Because mesenchymal tissues (ie, bone, cartilage, tendon, ligament, and disc) each require unique mechanical properties, the model should permit assessment of whether an appropriately functioning tissue forms. There is a need for the development of additional in vitro and in vivo models, as well as computational approaches. In particular, noninvasive assessments of tissue composition, structure, and function need further development as specific cell-based therapies are extended to human trials.

Large-animal models, although essential, are expensive and technically demanding. Large-animal studies are currently limited to autologous-based experiments, in which markers distinguishing host from donor are limited. The NIH-supported National Swine Resource and Research Center, a large-animal resource, has been developed. The center can provide pig lines with GFP reporters and also has pigs with genetic backgrounds similar to the mouse NSG.NOG strain. Successful results from studies using human progenitor cells in the mouse could transition to the immunocompromised pig as a route to FDA-approved clinical trials of a cell-based therapy.

Human clinical trials

Appropriate assessment of cell-based therapies in human studies will require appropriate study design, including the use of appropriate controls, blinding and elimination of investigator bias, appropriate validated outcome measures, and rigorous statistical analysis. Because of the difficulty and/or morbidity associated with

harvesting tissues from human subjects, the development and use of sensitive, noninvasive measures of tissue composition, structure, and mechanical function is essential.

In humans, sensitive, validated PROMs will provide a powerful tool to assess functional improvement. PROMs utilize patient-based assessments of their own functional status and perception of pain. Many validated outcome measures are now commonly used to assess the efficacy of various medical and surgical therapies. For upper extremities, functional outcome measures include grip strength and assessment of arm, shoulder, and hand function (ie, Disabilities of the Arm, Shoulder, and Hand [DASH] score) based on questionnaires. Lower-extremity fracture functional assessments could include the Timed Up and Go Test or the Lower Extremity Functional Scale. The Harris Hip Score and the 36-item Short Form Survey (SF-36) can measure hip and lower-extremity PROMs of function. Finally, functional criteria to assess the success of a spinal fusion could include the Oswestry Disability Index or the SF-12 or SF-36. Through a Common Fund initiative, the NIH developed a comprehensive PROM system called PROMIS that uses computer-adapted testing and thus has increased efficiency and sensitivity. This is being increasingly adopted as a PROM for patients with musculoskeletal disease.

The primary clinical goals of cell therapies are to treat symptoms, especially pain and instability, and to allow for return of normal function of the targeted tissue. Success can be defined as: (i) improvements in pain and/or physical function; (ii) a durable response (years); and (iii) a return of structural support or joint mobility that permits movement. Structural repair is a secondary outcome, implying that the tissue repair or implanted tissue has value to the degree which it permits painless function. The individual characteristics and properties necessary for regeneration are specific to the various musculoskeletal tissues (Table 4).

For therapeutic product development, strong preclinical evidence in model organisms on healing and function is an absolute requirement for the initiation of randomized clinical trials, which involve objective clinical outcome and follow-up measurements. Rigorous, scientifically based understanding of the mechanism of action is ultimately required to justify the therapeutic application of cell-based therapy. It is noteworthy that adult stem/progenitor cell therapy, which is often being practiced in an unregulated, non-standardized manner, has recently resulted in several highly publicized studies. To ensure safety and enhance efficacy, adult stem-cell therapy should be practiced only in a regulated and standardized manner. In addition, clinical trials submitted for publication should adhere to the Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) or Consolidated Standards of Reporting Trials (CONSORT) guidelines¹.

Ethical considerations

Animal studies

Animal studies should be conducted with strict adherence to ethical principles. The NIH provides guidelines (<http://grants.nih.gov/grants/olaw/links.htm>) and experiments should be approved by institutional animal use committees.

¹ Clinical data utilizing these recommendations is not currently available and assessing efficacy is not possible at this time. Refer to the following FDA websites: <https://www.fda.gov/BiologicsBloodVaccines/GuidanceComplianceRegulatoryInformation/Guidances/default.htm>; <https://www.fda.gov/MedicalDevices/DeviceRegulationandGuidance/GuidanceDocuments/default.htm>; and <https://www.fda.gov/CombinationProducts/default.htm>.

and experiments should be approved by institutional animal use committees.

Cell-based therapy in humans

Currently, there is widespread use of cell-based therapeutics, often “off-label,” in the clinical setting. The use of “stem/progenitor cells” to improve tissue healing has a distinct appeal to patients, because it suggests cutting-edge science with the potential to regenerate lost or damaged tissues. When combined with the reality that existing treatment options are often suboptimal for many soft-tissue injuries and that many patients seeking treatment are frequently relatively young and desirous of returning to an active lifestyle, an environment has been created in which patients are vulnerable and will seek such stem/progenitor cell treatments. However, although great potential exists for the use of cell-based approaches to improve tissue healing and regeneration, the reality is that current data for the use of cell-based therapies for augmentation of tissue healing is highly variable.^(111,112)

Given this significant limitation, it is imperative that patients are informed that there exists little high-level animal model or randomized clinical trial data to support the use of stem/progenitor cell-based approaches for tissue healing. It should also be communicated to patients that use of these treatment approaches are most often off-label, and that the treating facility often has a financial incentive to perform a stem/progenitor cell procedure which is unlikely to be covered by standard medical insurance. At the same time, however, autologous therapies appear to be generally safe and well-tolerated, although their effectiveness is often largely based on testimonial evidence. Given this confluence of issues, it is critical that a neutral or second-opinion physician is consulted, such that truly informed consent may be provided. In the absence of these ethical considerations, the Task Force cannot support the current application of stem/progenitor cell therapy, whether administered locally or systemically, until the treatment protocols have obtained FDA approval.

Acknowledgments

The Task Force members would like to thank Fei Wang, Ph.D. from the National Institute of Arthritis and Musculoskeletal and Skin Diseases (NIAMS) for serving as an Advisory Liaison. We also express our appreciation to the late Paolo Bianco, M.D. and Christopher R. Jacobs, Ph.D., whose contributions to this work were of great significance.

The Task Force would also like to acknowledge the dedicated staff of the ASBMR, particularly Mr. Doug Fesler and Ms. Lynn Mirigian, without whose enthusiastic participation and administrative skills this Task Force report would not have been possible.

References

1. Aggarwal S, Pittenger MF. Human mesenchymal stem cells modulate allogeneic immune cell responses. *Blood*. 2005;105(4):1815–22.
2. Le Blanc K, Frassoni F, Ball L, et al. Mesenchymal stem cells for treatment of steroid-resistant, severe, acute graft-versus-host disease: a phase II study. *Lancet*. 2008;371(9624):1579–86.
3. Brittberg M, Nilsson A, Lindahl A, Ohlsson C, Peterson L. Rabbit articular cartilage defects treated with autologous cultured chondrocytes. *Clin Orthop Relat Res*. 1996;326:270–83.

4. Robey PG. Cell sources for bone regeneration: the good, the bad, and the ugly (but promising). *Tissue Eng Part B Rev.* 2011;17(6):423–30.
5. Bastami F, Nazeman P, Moslemi H, Rezai Rad M, Sharifi K, Khojasteh A. Induced pluripotent stem cells as a new getaway for bone tissue engineering: a systematic review. *Cell Prolif.* 2017;50(2):e12321.
6. Kuhn LT, Liu Y, Boyd NL, et al. Developmental-like bone regeneration by human embryonic stem cell-derived mesenchymal cells. *Tissue Eng Part A.* 2014;20(1–2):365–77.
7. Kuznetsov SA, Cherman N, Robey PG. In vivo bone formation by progeny of human embryonic stem cells. *Stem Cells Dev.* 2011;20(2):269–87.
8. Marolt D, Campos IM, Bhumiratana S, et al. Engineering bone tissue from human embryonic stem cells. *Proc Natl Acad Sci U S A.* 2012;109(22):8705–9.
9. Phillips MD, Kuznetsov SA, Cherman N, et al. Directed differentiation of human induced pluripotent stem cells toward bone and cartilage: in vitro versus in vivo assays. *Stem Cells Transl Med.* 2014;3(7):867–78.
10. Diederichs S, Shine KM, Tuan RS. The promise and challenges of stem cell-based therapies for skeletal diseases: stem cell applications in skeletal medicine: potential, cell sources and characteristics, and challenges of clinical translation. *Bioessays.* 2013;35(3):220–30.
11. Simonson OE, Domogatskaya A, Volchkov P, Rodin S. The safety of human pluripotent stem cells in clinical treatment. *Ann Med.* 2015;47(5):370–80.
12. Owen M, Friedenstein AJ. *Ciba Found. Symp.* 1998;136:42–60.
13. Lv FJ, Tuan RS, Cheung KM, Leung VY. Concise review: the surface markers and identity of human mesenchymal stem cells. *Stem Cells.* 2014;32(6):1408–19.
14. da Silva Meirelles L, Chagastelles PC, Nardi NB. Mesenchymal stem cells reside in virtually all post-natal organs and tissues. *J Cell Sci.* 2006;119(Pt 11):2204–13.
15. Colnot C, Zhang X, Knothe Tate ML. Current insights on the regenerative potential of the periosteum: molecular, cellular, and endogenous engineering approaches. *J Orthop Res.* 2012;30(12):1869–78.
16. Djouad F, Bony C, Häupl T, et al. Transcriptional profiles discriminate bone marrow-derived and synovium-derived mesenchymal stem cells. *Arthritis Res Ther.* 2005;7(6):R1304–15.
17. Gronthos S, Mankani M, Brahimi J, Robey PG, Shi S. Postnatal human dental pulp stem cells (DPSCs) in vitro and in vivo. *Proc Natl Acad Sci U S A.* 2000;97(25):13625–30.
18. Seo BM, Miura M, Gronthos S, et al. Investigation of multipotent postnatal stem cells from human periodontal ligament. *Lancet.* 2004;364(9429):149–55.
19. Bianco P, Cao X, Frenette PS, et al. The meaning, the sense and the significance: translating the science of mesenchymal stem cells into medicine. *Nat Med.* 2013;19(1):35–42.
20. Chan CK, Seo EY, Chen JY, et al. Identification and specification of the mouse skeletal stem cell. *Cell.* 2015;160(1–2):285–98.
21. Chan CKF, Gulati GS, Sinha R, et al. Identification of the human skeletal stem cell. *Cell.* 2018;175(1):43–56.e21.
22. Gulati GS, Murphy MP, Marecic O, et al. Isolation and functional assessment of mouse skeletal stem cell lineage. *Nat Protoc.* 2018;13(6):1294–309.
23. Worthley DL, Churchill M, Compton JT, et al. Gremlin 1 identifies a skeletal stem cell with bone, cartilage, and reticular stromal potential. *Cell.* 2015;160(1–2):269–84.
24. Levi B, Longaker MT. Concise review: adipose-derived stromal cells for skeletal regenerative medicine. *Stem Cells.* 2011;29(4):576–82.
25. Tsang WP, Shu Y, Kwok PL, et al. CD146+ human umbilical cord perivascular cells maintain stemness under hypoxia and as a cell source for skeletal regeneration. *PLoS One.* 2013;8(10):e76153.
26. Wang KX, Xu LL, Rui YF, et al. The effects of secretion factors from umbilical cord derived mesenchymal stem cells on osteogenic differentiation of mesenchymal stem cells. *PLoS One.* 2015;10(3):e0120593.
27. Wang W, He N, Feng C, et al. Human adipose-derived mesenchymal progenitor cells engraft into rabbit articular cartilage. *Int J Mol Sci.* 2015;16(6):12076–91.
28. Zhou BO, Yue R, Murphy MM, Peyer JG, Morrison SJ. Leptin-receptor-expressing mesenchymal stromal cells represent the main source of bone formed by adult bone marrow. *Cell Stem Cell.* 2014;15(2):154–68.
29. Crisan M, Yap S, Casteilla L, et al. A perivascular origin for mesenchymal stem cells in multiple human organs. *Cell Stem Cell.* 2008;3(3):301–13.
30. Kramann R, Goettsch C, Wongboonsin J, et al. Adventitial MSC-like cells are progenitors of vascular smooth muscle cells and drive vascular calcification in chronic kidney disease. *Cell Stem Cell.* 2016;19(5):628–42.
31. Askarinam A, James AW, Zara JN, et al. Human perivascular stem cells show enhanced osteogenesis and vasculogenesis with Nel-like molecule 1 protein. *Tissue Eng Part A.* 2013;19(11–12):1386–97.
32. Bouacida A, Rosset P, Trichet V, et al. Pericyte-like progenitors show high immaturity and engraftment potential as compared with mesenchymal stem cells. *PLoS One.* 2012;7(11):e48648.
33. Chung CG, James AW, Asatrian G, et al. Human perivascular stem cell-based bone graft substitute induces rat spinal fusion. *Stem Cells Transl Med.* 2014;3(10):1231–41.
34. Meyers CA, Xu J, Zhang L, et al. Skeletogenic capacity of human perivascular stem cells obtained via magnetic activated cell sorting. *Tissue Eng Part A.* Epub 2019 Aug 16. <https://doi.org/10.1089/ten.TEA.2019.0031>.
35. Tawonsawatruk T, West CC, Murray IR, Soo C, Péault B, Simpson AHRW. Adipose derived pericytes rescue fractures from a failure of healing—non-union. *Sci Rep.* 2016;6:22779.
36. Kon E, Filardo G, Condello V, et al. Second-generation autologous chondrocyte implantation: results in patients older than 40 years. *Am J Sports Med.* 2011;39(8):1668–75.
37. Niemeyer P, Köstler W, Salzmann GM, Lenz P, Kreuz PC, Südkamp NP. Autologous chondrocyte implantation for treatment of focal cartilage defects in patients age 40 years and older: a matched-pair analysis with 2-year follow-up. *Am J Sports Med.* 2010;38(12):2410–6.
38. Rosenberger RE, Gomoll AH, Bryant T, Minas T. Repair of large chondral defects of the knee with autologous chondrocyte implantation in patients 45 years or older. *Am J Sports Med.* 2008;36(12):2336–44.
39. Nitzsche F, Müller C, Lukomska B, Jolkkonen J, Deten A, Boltze J. Concise review: MSC adhesion cascade—insights into homing and transendothelial migration. *Stem Cells.* 2017;35(6):1446–60.
40. Granero-Molto F, Weis JA, Miga MI, et al. Regenerative effects of transplanted mesenchymal stem cells in fracture healing. *Stem Cells.* 2009;27(8):1887–98.
41. Marigo I, Dazzi F. The immunomodulatory properties of mesenchymal stem cells. *Semin Immunopathol.* 2011;33(6):593–602.
42. Chu CR, Rodeo S, Bhutani N, et al. Optimizing clinical use of biologics in orthopaedic surgery: consensus recommendations from the 2018 AAOS/NIH U-13 Conference. *J Am Acad Orthop Surg.* 2019;15;27(2):e50–63.
43. Rodriguez IA, Growney Kalaf EA, Bowlin GL, Sell SA. Platelet-rich plasma in bone regeneration: engineering the delivery for improved clinical efficacy. *Biomed Res Int.* 2014;2014:392398.
44. de Queiroz Fernandes J, de Lima VN, Bonardi JP, Filho OM, Queiroz SBF. Bone regeneration with recombinant human bone morphogenetic protein 2: a systematic review. *J Maxillofac Oral Surg.* 2018;17(1):13–8.
45. Xie X, Zhang C, Tuan RS. Biology of platelet-rich plasma and its clinical application in cartilage repair. *Arthritis Res Ther.* 2014;16(1):204.
46. Lane NE, Corr M. Osteoarthritis in 2016: anti-NGF treatments for pain - two steps forward, one step back? *Nat Rev Rheumatol.* 2017;13(2):76–8.
47. Braun HJ, Wasterlain AS, Dragoo JL. The use of PRP in ligament and meniscal healing. *Sports Med Arthrosc Rev.* 2013;21(4):206–12.
48. Liou JJ, Rothrauff BB, Alexander PG, Tuan RS. Effect of platelet-rich plasma on chondrogenic differentiation of adipose- and bone

- marrow-derived Mesenchymal stem cells. *Tissue Eng Part A*. 2018; 24(19–20):1432–43.
49. Dai M, Zhang Y, Yu M, Tian W. Therapeutic applications of conditioned medium from adipose tissue. *Cell Prolif*. 2016;49(5):561–7.
 50. Katagiri W, Osugi M, Kawai T, Hibi H. First-in-human study and clinical case reports of the alveolar bone regeneration with the secretome from human mesenchymal stem cells. *Head Face Med*. 2016; 12:5.
 51. Bourne GH. The biochemistry and physiology of bone. 2nd ed. New York: Academic Press; 1971.
 52. Cobelli NJ, Leong DJ, Sun HB. Exosomes: biology, therapeutic potential, and emerging role in musculoskeletal repair and regeneration. *Ann N Y Acad Sci*. 2017;1410(1):57–67.
 53. El Andaloussi S, Mäger I, Breakefield XO, Wood MJ. Extracellular vesicles: biology and emerging therapeutic opportunities. *Nat Rev Drug Discov*. 2013;12(5):347–57.
 54. Gaspar D, Spanoudes K, Holladay C, Pandit A, Zeugolis D. Progress in cell-based therapies for tendon repair. *Adv Drug Deliv Rev*. 2015;84: 240–56.
 55. Fu SC, Cheuk YC, Yung SH, Rolf CG, Chan KM. Systematic review of biological modulation of healing in anterior cruciate ligament reconstruction. *Orthop J Sports Med*. 2014;2(3):2325967114526687.
 56. Hurtig MB, Buschmann MD, Fortier LA, et al. Preclinical studies for cartilage repair: recommendations from the International Cartilage Repair Society. *Cartilage*. 2011;2(2):137–52.
 57. Ghadially FN, Thomas I, Oryschak AF, Lalonde JM. Long-term results of superficial defects in articular cartilage: a scanning electron-microscope study. *J Pathol*. 1977;121(4):213–7.
 58. Firth EC, Klarenbeek A. Transphyseal vessel involvement in repair of metaphyseal retained cartilage. *Aust Vet J*. 1995;72(12):452–5.
 59. Aigner T, Minas T, Hsu HP, Nehrer S, Sledge CB, Spector M. Histopathology atlas of animal model systems - overview of guiding principles. *Osteoarthritis Cartilage*. 2010;18(Suppl 3):S2–6.
 60. Breinan HA, Minas T, Hsu HP, Nehrer S, Sledge CB, Spector M. Effect of cultured autologous chondrocytes on repair of chondral defects in a canine model. *J Bone Joint Surg Am*. 1997;79(10):1439–51.
 61. Cook SD, Patron LP, Salkeld SL, Rueger DC. Repair of articular cartilage defects with osteogenic protein-1 (BMP-7) in dogs. *J Bone Joint Surg Am*. 2003;85-A(Suppl 3):116–23.
 62. Hoemann CD, Sun J, Chrzanowski V, Buschmann MD. A multivalent assay to detect glycosaminoglycan, protein, collagen, RNA, and DNA content in milligram samples of cartilage or hydrogel-based repair cartilage. *Anal Biochem*. 2002;300(1):1–10.
 63. Ando W, Fujie H, Moriguchi Y, et al. Detection of abnormalities in the superficial zone of cartilage repaired using a tissue engineered construct derived from synovial stem cells. *Eur Cell Mater*. 2012;24:292–307.
 64. Frisbie DD, Cross MW, McIlwraith CW. A comparative study of articular cartilage thickness in the stifle of animal species used in human pre-clinical studies compared to articular cartilage thickness in the human knee. *Vet Comp Orthop Traumatol*. 2006;19(3):142–6.
 65. Frisbie DD, Trotter GW, Powers BE, et al. Arthroscopic subchondral bone plate microfracture technique augments healing of large chondral defects in the radial carpal bone and medial femoral condyle of horses. *Vet Surg*. 1999;28(4):242–55.
 66. Marfia G, Campanella R, Navone SE, et al. Potential use of human adipose mesenchymal stromal cells for intervertebral disc regeneration: a preliminary study on biglycan-deficient murine model of chronic disc degeneration. *Arthritis Res Ther*. 2014;16(5):457.
 67. Hiyama A, Mochida J, Iwashina T, et al. Transplantation of mesenchymal stem cells in a canine disc degeneration model. *J Orthop Res*. 2008;26(5):589–600.
 68. Ganey T, Hutton WC, Moseley T, Hedrick M, Meisel HJ. Intervertebral disc repair using adipose tissue-derived stem and regenerative cells: experiments in a canine model. *Spine (Phila Pa 1976)*. 2009; 34(21):2297–304.
 69. Ganey T, Libera J, Moos V, et al. Disc chondrocyte transplantation in a canine model: a treatment for degenerated or damaged intervertebral disc. *Spine (Phila Pa 1976)*. 2003;28(23):2609–20.
 70. Awad HA, Boivin GP, Dressler MR, Smith FN, Young RG, Butler DL. Repair of patellar tendon injuries using a cell-collagen composite. *J Orthop Res*. 2003;21(3):420–31.
 71. Awad HA, Butler DL, Boivin GP, et al. Autologous mesenchymal stem cell-mediated repair of tendon. *Tissue Eng*. 1999;5(3):267–77.
 72. Behfar M, Javanmardi S, Sarrafzadeh-Rezaei F. Comparative study on functional effects of allotransplantation of bone marrow stromal cells and adipose derived stromal vascular fraction on tendon repair: a biomechanical study in rabbits. *Cell J*. 2014;16(3):263–70.
 73. Behfar M, Sarrafzadeh-Rezaei F, Hobbenaghi R, Delirez N, Dalir-Naghadeh B. Enhanced mechanical properties of rabbit flexor tendons in response to intratendinous injection of adipose derived stromal vascular fraction. *Curr Stem Cell Res Ther*. 2012;7(3):173–8.
 74. Dressler MR, Butler DL, Boivin GP. Effects of age on the repair ability of mesenchymal stem cells in rabbit tendon. *J Orthop Res*. 2005;23(2):287–93.
 75. Juncosa-Melvin N, Boivin GP, Galloway MT, Gooch C, West JR, Butler DL. Effects of cell-to-collagen ratio in stem cell-seeded constructs for Achilles tendon repair. *Tissue Eng*. 2006;12(4):681–9.
 76. Juncosa-Melvin N, Boivin GP, Galloway MT, et al. Effects of cell-to-collagen ratio in mesenchymal stem cell-seeded implants on tendon repair biomechanics and histology. *Tissue Eng*. 2005;11(3–4):448–57.
 77. Juncosa-Melvin N, Boivin GP, Gooch C, et al. The effect of autologous mesenchymal stem cells on the biomechanics and histology of gel-collagen sponge constructs used for rabbit patellar tendon repair. *Tissue Eng*. 2006;12(2):369–79.
 78. Juncosa-Melvin N, Matlin KS, Holdcraft RW, Nirmalanandhan VS, Butler DL. Mechanical stimulation increases collagen type I and collagen type III gene expression of stem cell-collagen sponge constructs for patellar tendon repair. *Tissue Eng*. 2007;13(6):1219–26.
 79. Juncosa-Melvin N, Shearn JT, Boivin GP, et al. Effects of mechanical stimulation on the biomechanics and histology of stem cell-collagen sponge constructs for rabbit patellar tendon repair. *Tissue Eng*. 2006;12(8):2291–300.
 80. Nirmalanandhan VS, Juncosa-Melvin N, Shearn JT, et al. Combined effects of scaffold stiffening and mechanical preconditioning cycles on construct biomechanics, gene expression, and tendon repair biomechanics. *Tissue Eng Part A*. 2009;15(8):2103–11.
 81. Young RG, Butler DL, Weber W, Caplan AI, Gordon SL, Fink DJ. Use of mesenchymal stem cells in a collagen matrix for Achilles tendon repair. *J Orthop Res*. 1998;16(4):406–13.
 82. Chen X, Yin Z, Chen JL, et al. Scleraxis-overexpressed human embryonic stem cell-derived mesenchymal stem cells for tendon tissue engineering with knitted silk-collagen scaffold. *Tissue Eng Part A*. 2014;20(11–12):1583–92.
 83. Gulotta LV, Kovacevic D, Ehteshami JR, Dagher E, Packer JD, Rodeo SA. Application of bone marrow-derived mesenchymal stem cells in a rotator cuff repair model. *Am J Sports Med*. 2009;37(11):2126–33.
 84. Gulotta LV, Kovacevic D, Montgomery S, Ehteshami JR, Packer JD, Rodeo SA. Stem cells genetically modified with the developmental gene MT1-MMP improve regeneration of the supraspinatus tendon-to-bone insertion site. *Am J Sports Med*. 2010;38(7):1429–37.
 85. Gulotta LV, Kovacevic D, Packer JD, Deng XH, Rodeo SA. Bone marrow-derived mesenchymal stem cells transduced with scleraxis improve rotator cuff healing in a rat model. *Am J Sports Med*. 2011; 39(6):1282–9.
 86. Gulotta LV, Kovacevic D, Packer JD, Ehteshami JR, Rodeo SA. Adenoviral-mediated gene transfer of human bone morphogenetic protein-13 does not improve rotator cuff healing in a rat model. *Am J Sports Med*. 2011;39(1):180–7.
 87. Hankemeier S, Hurschler C, Zeichen J, et al. Bone marrow stromal cells in a liquid fibrin matrix improve the healing process of patellar tendon window defects. *Tissue Eng Part A*. 2009;15(5):1019–30.
 88. Hankemeier S, van Griensven M, Ezechieli M, et al. Tissue engineering of tendons and ligaments by human bone marrow stromal cells in a liquid fibrin matrix in immunodeficient rats: results of a histologic study. *Arch Orthop Trauma Surg*. 2007;127(9):815–21.
 89. Huang TF, Yew TL, Chiang ER, et al. Mesenchymal stem cells from a hypoxic culture improve and engraft Achilles tendon repair. *Am J Sports Med*. 2013;41(5):1117–25.

90. Lee CH, Yew TL, Chiang ER, et al. Harnessing endogenous stem/progenitor cells for tendon regeneration. *J Clin Invest*. 2015;125(7):2690–701.
91. Lee JY, Zhou Z, Taub PJ, et al. BMP-12 treatment of adult mesenchymal stem cells in vitro augments tendon-like tissue formation and defect repair in vivo. *PLoS One*. 2011;6(3):e17531.
92. Lui PP, Wong OT, Lee YW. Application of tendon-derived stem cell sheet for the promotion of graft healing in anterior cruciate ligament reconstruction. *Am J Sports Med*. 2014;42(3):681–9.
93. Mifune Y, Matsumoto T, Takayama K, et al. Tendon graft revitalization using adult anterior cruciate ligament (ACL)-derived CD34+ cell sheets for ACL reconstruction. *Biomaterials*. 2013;34(22):5476–87.
94. Ni M, Lui PP, Rui YF, et al. Tendon-derived stem cells (TDSCs) promote tendon repair in a rat patellar tendon window defect model. *J Orthop Res*. 2012;30(4):613–9.
95. Okamoto N, Kushida T, Oe K, Umeda M, Ikehara S, Iida H. Treating Achilles tendon rupture in rats with bone-marrow-cell transplantation therapy. *J Bone Joint Surg Am*. 2010;92(17):2776–84.
96. Tan C, Lui PP, Lee YW, Wong YM. Scx-transduced tendon-derived stem cells (tdscs) promoted better tendon repair compared to mock-transduced cells in a rat patellar tendon window injury model. *PLoS One*. 2014;9(5):e97453.
97. Xu W, Wang Y, Liu E, et al. Human iPSC-derived neural crest stem cells promote tendon repair in a rat patellar tendon window defect model. *Tissue Eng Part A*. 2013;19(21–22):2439–51.
98. Soslowsky LJ, Carpenter JE, DeBano CM, Banerji I, Moalli MR. Development and use of an animal model for investigations on rotator cuff disease. *J Shoulder Elbow Surg*. 1996;5(5):383–92.
99. Godwin EE, Young NJ, Dudhia J, Beamish IC, Smith RK. Implantation of bone marrow-derived mesenchymal stem cells demonstrates improved outcome in horses with overstrain injury of the superficial digital flexor tendon. *Equine Vet J*. 2012;44(1):25–32.
100. Martinello T, Bronzini I, Perazzi A, et al. Effects of in vivo applications of peripheral blood-derived mesenchymal stromal cells (PB-MSCs) and platelet-rich plasma (PRP) on experimentally injured deep digital flexor tendons of sheep. *J Orthop Res*. 2013;31(2):306–14.
101. Pacini S, Spinabella S, Trombi L, et al. Suspension of bone marrow-derived undifferentiated mesenchymal stromal cells for repair of superficial digital flexor tendon in race horses. *Tissue Eng*. 2007;13(12):2949–55.
102. Smith RK. Mesenchymal stem cell therapy for equine tendinopathy. *Disabil Rehabil*. 2008;30(20–22):1752–8.
103. Lacitignola L, Staffieri F, Rossi G, Francioso E, Crovace A. Survival of bone marrow mesenchymal stem cells labelled with red fluorescent protein in an ovine model of collagenase-induced tendinitis. *Vet Comp Orthop Traumatol*. 2014;27(3):204–9.
104. Fan H, Liu H, Toh SL, Goh JC. Anterior cruciate ligament regeneration using mesenchymal stem cells and silk scaffold in large animal model. *Biomaterials*. 2009;30(28):4967–77.
105. Liu W, Chen B, Deng D, Xu F, Cui L, Cao Y. Repair of tendon defect with dermal fibroblast engineered tendon in a porcine model. *Tissue Eng*. 2006;12(4):775–88.
106. Gelberman RH, Shen H, Korpakis I, et al. Effect of adipose-derived stromal cells and BMP12 on intrasynovial tendon repair: a biomechanical, biochemical, and proteomics study. *J Orthop Res*. 2016;34(4):630–40.
107. Zhao C, Shimura H, Reisdorf RL, et al. Effects of lubricant and autologous bone marrow stromal cell augmentation on immobilized flexor tendon repairs. *J Orthop Res*. 2016;34(1):154–60.
108. Haudebourg T, Poirier N, Vanhove B. Depleting T-cell subpopulations in organ transplantation. *Transpl Int*. 2009;22(5):509–18.
109. Lunney JK, Ho CS, Wysocki M, Smith DM. Molecular genetics of the swine major histocompatibility complex, the SLA complex. *Dev Comp Immunol*. 2009;33(3):362–74.
110. Low LA, Tagle DA. Tissue chips - innovative tools for drug development and disease modeling. *Lab Chip*. 2017;17(18):3026–36.
111. Borakati A, Mafi R, Mafi P, Khan WS. A systematic review and meta-analysis of clinical trials of mesenchymal stem cell therapy for cartilage repair. *Curr Stem Cell Res Ther*. 2018;13(3):215–25.
112. McIntyre JA, Jones IA, Han B, Vangsness CT Jr. Intra-articular mesenchymal stem cell therapy for the human joint: a systematic review. *Am J Sports Med*. 2018;46(14):3550–63.