

UC Davis

UC Davis Previously Published Works

Title

Metabolic diversity and co-occurrence of multiple *Ferrovum* species at an acid mine drainage site

Permalink

<https://escholarship.org/uc/item/85d9m09d>

Journal

BMC Microbiology, 20(1)

ISSN

1471-2180

Authors

Grettenberger, Christen L

Havig, Jeff R

Hamilton, Trinity L

Publication Date

2020-12-01

DOI

10.1186/s12866-020-01768-w

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

RESEARCH ARTICLE

Open Access



Metabolic diversity and co-occurrence of multiple *Ferrovum* species at an acid mine drainage site

Christen L. Grettenberger¹, Jeff R. Havig² and Trinity L. Hamilton^{3,4*}

Abstract

Background: *Ferrovum* spp. are abundant in acid mine drainage sites globally where they play an important role in biogeochemical cycling. All known taxa in this genus are Fe(II) oxidizers. Thus, co-occurring members of the genus could be competitors within the same environment. However, we found multiple, co-occurring *Ferrovum* spp. in Cabin Branch, an acid mine drainage site in the Daniel Boone National Forest, KY.

Results: Here we describe the distribution of *Ferrovum* spp. within the Cabin Branch communities and metagenome assembled genomes (MAGs) of two new *Ferrovum* spp. In contrast to previous studies, we recovered multiple 16S rRNA gene sequence variants suggesting the commonly used 97% cutoff may not be appropriate to differentiate *Ferrovum* spp. We also retrieved two nearly-complete *Ferrovum* spp. genomes from metagenomic data. The genomes of these taxa differ in several key ways relating to nutrient cycling, motility, and chemotaxis.

Conclusions: Previously reported *Ferrovum* genomes are also diverse with respect to these categories suggesting that the genus *Ferrovum* contains substantial metabolic diversity. This diversity likely explains how the members of this genus successfully co-occur in Cabin Branch and why *Ferrovum* spp. are abundant across geochemical gradients.

Keywords: *Ferrovum*, Carbon fixation, Iron oxidation, Metagenome, Acidophilic, Biofilm, Microcosm, Co-occurrence, Nitrogen, Denoised sequence variants

Background

Iron-oxidizing bacteria are common in mine-impacted water and acid mine drainage (AMD) environments which are typically characterized by low pH and high concentrations of dissolved metals [1–3]. Betaproteobacteria of the genus *Ferrovum* play important roles in biogeochemical cycling in AMD environments including carbon fixation and rapid oxidation of iron [4–9] and could be of value in bioremediation [10]. However, *Ferrovum* spp. are challenging and labor-intensive to culture and isolate because

they often co-occur with heterotrophic *Acidiphilium* spp. or other Fe(II) oxidizers [7, 11, 12]. Indeed, *Ferrovum myxofaciens* strain P3 is the only *Ferrovum* spp. in culture to date, highlighting the value and need for characterization of this enigmatic genus via molecular techniques.

Ferrovum spp. occur in AMD environments with diverse geochemistry. *Ferrovum* spp. are abundant at sites with pH ranging from less than 3 to 7 [4–6, 13, 14] and with iron concentrations ranging from 2 μ M [5] to 71 mM [7]. High-throughput sequencing and ‘omics techniques have aided in characterizing the metabolic potential of *Ferrovum* via non-culture-based techniques [15–18] across these large gradients of Fe and pH. These studies indicate that *Ferrovum* is a diverse genus composed of six clades (Groups I - VI) of closely related species and strains [18]. Still, only four *Ferrovum* genomes are publicly

* Correspondence: trinityh@umn.edu

³Department of Plant and Microbial Biology, University of Minnesota, 218 Cargill Building, St. Paul, MN 55108, USA

⁴The BioTechnology Institute, University of Minnesota, St. Paul, MN 55108, USA

Full list of author information is available at the end of the article



© The Author(s). 2020 **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

available and these are from *Ferrovum* Groups I and IV. No genomes have been reported for Groups, II, III, or V. The published genomes and culture studies suggest that *Ferrovum* spp. are autotrophs that pair Fe(II) oxidation with carbon fixation [7, 15, 16, 18, 19]. *Ferrovum* spp. play important roles in nutrient cycling and some members of the genus also fix nitrogen [7, 19]. However, sequenced *Ferrovum* genomes have variable genomes especially with genes associated with motility, chemotaxis, biofilm formation, and nitrogen metabolism [15]. Furthermore, *Ferrovum* dominated communities exhibit morphological differences, including forming large streamers [5, 7, 14] or low extracellular polymeric substances [4] though it can be difficult to identify EPS in low pH environments [16]. These differences may help explain how *Ferrovum* is able to occupy a large range of pH and ferrous iron concentrations in AMD environments. However, the potential genetic and metabolic diversity within the genus is under-represented because there are no cultured representatives or genomes available for several *Ferrovum* clades.

In our previous studies of Cabin Branch, we reported the predominance of a single operational taxonomic unit (OTU; 97% similarity) in our 16S rRNA amplicon data most closely related to *F. myxofaciens* at this AMD site in the Daniel Boone National Forest in southern Kentucky. This taxon was abundant in diverse morphotypes including filament, floc, and 'brain' or 'spongy' mat biofilms over a pH range of 2.1 to 2.4 and Fe(II) concentrations of 448 to 5200 $\mu\text{mol/L}$ [5]. Based on these data, we employed a cultivation-independent approach to characterize the genetic diversity and functional potential of this taxon using 16S rRNA amplicons and by examining the gene content of metagenome assembled genomes (MAGs) from the site.

Results

Geochemistry

All aqueous geochemistry data are reported in Table 1. pH values were acidic, ranging from 2.90 to 2.97 from the source to Rose Pool (Fig. 1). Temperatures were highest at the source (14.6 °C) and decreased down the outflow channel to 8.1 °C at the Rose Pool, consistent with warm groundwater cooling after emerging due to relatively cool ambient temperatures (~ 0 °C at the time of sampling). Conductivity increased from 940 $\mu\text{S/cm}$ at the emergence to 2063 $\mu\text{S/cm}$ at Rose Pool. Dissolved oxygen increased from a low of 77.5 $\mu\text{mol/L}$ at the emergence sampling site to a high of 401 $\mu\text{mol/L}$ (close to saturation) at Rose Pool. Anion concentrations in Cabin Branch were dominated by sulfate (9.51 to 10.08 mmol/L), with lower concentration of chloride (166 to 220 $\mu\text{mol/L}$). Cation concentrations were highest at the emergence and nearby first outflow sites, and lowest at the Rose Pool, including calcium (1.20 to 0.39 mmol/L), magnesium (0.69 to 0.20

Table 1 Geochemical and physical features for Cabin Branch samples

Location:	Emergence		LLC		Rose Pool	
Parameter (Units)	Value	S.D.	Value	S.D.	Value	S.D.
pH	2.90		2.92		2.97	
mV	235		232		224	
Temperature (°C)	14.6		13.7		8.1	
Conductivity ($\mu\text{S/cm}$)	940		1853		2063	
D.O. ($\mu\text{mol/L}$)	77.5		266		401	
Fe ²⁺ ($\mu\text{mol/L}$)	403.2		882.0		11.16	
NH ₄ (T) ($\mu\text{mol/L}$)	35.7		50.0		10.7	
DIC (mmol/L)	1.67		0.48		0.30	
$\delta^{13}\text{CVPDB}$ (‰)	-15.96	0.08	-13.03	0.08	-11.79	0.08
DOC ($\mu\text{mol/L}$)	40.51	0.90	44.19	3.27	36.82	6.42
$\delta^{13}\text{CVPDB}$ (‰)	-22.50	1.53	-23.09	0.84	-23.95	0.29
Cl ($\mu\text{mol/L}$)	175.58	1.10	220.04	61.99	166.13	0.44
SO ₄ (mmol/L)	9.96	0.16	9.51	0.14	10.08	0.06
P ($\mu\text{mol/L}$)	9.55	0.49	4.49	0.35	bdl	
Na (mmol/L)	0.12	0.00	0.12	0.03	0.03	0.00
K (mmol/L)	0.17	0.00	0.18	0.04	0.03	0.00
Ca (mmol/L)	1.121	0.028	1.198	0.256	0.390	0.007
Mg (mmol/L)	0.666	0.012	0.686	0.156	0.201	0.001
Al ($\mu\text{mol/L}$)	1755	125	1201	0.26	187.4	1.2
Si ($\mu\text{mol/L}$)	1528	35	1454	66	nd	
Mn ($\mu\text{mol/L}$)	76.84	0.63	43.19	0.13	40.23	0.28
Fe _{Total} ($\mu\text{mol/L}$)	4684	22	4154	50	249.6	1.8

Geochemical and physical measurements: meters/probes (pH, mV, Temperature,

Conductivity, Dissolved Oxygen (D.O.)), spectrophotometry (Fe(II), NH₄(T)), ion chromatography (Cl, SO₄), IR-MS (dissolved inorganic carbon (DIC), dissolved organic carbon (DOC), ICP-OES (P, Na, K, Ca, Mg, Si, italicized values for Al and Fe_{Total}), and ICP-MS (Al, Mn, Fe). LLC = limestone-lined channel, S.D. = standard deviation, bdl = below detection limits,

mmol/L), potassium (0.18 to 0.03 mmol/L), and sodium (0.12 to 0.03 mmol/L).

Dissolved inorganic carbon (DIC, predominantly present as dissolved CO₂ due to the low pH) concentration was highest at the emergence (1.67 mmol/L) and decreased down the outflow channel to 0.30 mmol/L at the Rose Pool. DIC $\delta^{13}\text{C}$ values were more negative at the source ($\delta^{13}\text{C} = -15.96$ ‰) and became more positive down the outflow channel with the most positive values at the Rose Pool ($\delta^{13}\text{C} = -11.79$ ‰), consistent with preferential loss of ¹²CO₂ through volatilization and uptake by autotrophs. Dissolved organic carbon concentration was consistently low at all sites (36.8 to 44.2 $\mu\text{mol/L}$) and had similar $\delta^{13}\text{C}$ values (-22.50 to -23.95 ‰).

Potential nutrients measured included NH₄(T), P, Mn, Fe(II), and Fe_(total). NH₄(T) concentration was highest at the first outflow site (50.0 $\mu\text{mol/L}$) and lowest at Rose Pool (10.7 $\mu\text{mol/L}$). P concentration was highest at the

emergence (9.55 $\mu\text{mol/L}$) and decreased to below detection limits at the Rose Pool. Mn concentration decreased from the highest value at the emergence (76.84 $\mu\text{mol/L}$) to the lowest at the Rose Pool (40.23 $\mu\text{mol/L}$). Total iron (Fe_{total}) exhibited a similar trend as Mn and P, with the highest concentration at the emergence (4684 $\mu\text{mol/L}$) and the lowest at the Rose Pool in the outflow (249.6 $\mu\text{mol/L}$), though Fe(II) had concentrations significantly lower than Fe_{total} with the highest concentration at the outflow site downstream of the emergence (882 $\mu\text{mol/L}$) and the lowest at the Rose Pool (11.2 $\mu\text{mol/L}$).

Inorganic carbon assimilation

Autotrophic incorporation of added $^{13}\text{CO}_2$ was quantified in mesocosm experiments using treatments carried out under either light or dark (wrapped in aluminum foil) to characterize photoautotrophic and chemoautotrophic uptake, respectively. At the emergence, both green and white biomass was observed and autotrophic incorporation of $^{13}\text{CO}_2$ was quantified in both biofilm types. Because we assumed the majority of uptake in the green biomass would be due to photoautotrophy and we could not control for oxygen production in this treatment, we performed one set of microcosms on the bulk biofilm in the light and then separated the biofilm types (green or white) for microcosms performed in the dark. In the light treatments of full biofilms (green and white biomass) C-uptake rates averaged 32.547 (± 7.949) and 21.691 (± 6.36) $\mu\text{g C uptake/g C}_{\text{biomass}}/\text{hr.}$ after one- and two-hour incubations, respectively. Dark treatments of the green biomass returned averages of 1.632 (± 0.454) $\mu\text{g C uptake/g C}_{\text{biomass}}/\text{hr.}$ after 1 hour and 0.746 (± 0.174) $\mu\text{g C uptake/g C}_{\text{biomass}}/\text{hr.}$ after 2 hours, while dark treatments of the white biomass returned averages of 0.305 (± 0.403) $\mu\text{g C uptake/g C}_{\text{biomass}}/\text{hr.}$ after 1 hour and 0.662 (± 0.240) $\mu\text{g C uptake/g C}_{\text{biomass}}/\text{hr.}$ after 2 hours. Biofilms with both green and white biomass together from the limestone-lined channel immediately downstream of the emergence returned average C-uptake rates of 35.947 (± 5.516) and 15.085 (± 0.106) $\mu\text{g C uptake/g C}_{\text{biomass}}/\text{hr.}$ for light treatments of one and two hours respectively and 2.586 (± 0.148) and 1.245 (± 0.035) $\mu\text{g C uptake/g C}_{\text{biomass}}/\text{hr.}$ for dark treatments after one and two hours. Rose Pool sediment incubations returned average C-uptake rates of 1.372 (± 0.061) $\mu\text{g C uptake/g C}_{\text{biomass}}/\text{hr.}$ for light treatments of 2 hours and 0.066 (± 0.132) $\mu\text{g C uptake/g C}_{\text{biomass}}/\text{hr.}$ for dark treatments of 2 hours. All C-uptake results are reported in Supplementary Table S1.

16S rRNA

Thirty two thousand one hundred twenty four quality-controlled sequences were retrieved from the


emergence, 41,988 from the limestone-lined channel, and 37,546 from the Rose Pool. These sequences were affiliated with 82, 276, and 286 denoised sequence variants (DSVs) respectively. Cabin Branch contains nine DSVs closely related to *Ferroplasma myxofaciens* (Fig. 2). Four DSVs are present in the emergence sample, five in the limestone lined channel, and eight in Rose Pool (Figs. 1 and 2). DSVs affiliated with *Ferroplasma* are most abundant in the limestone lined channel where they compose 32.8% of the community and least abundant in the Rose Pool where they compose 5.0% of the community. *Ferroplasma* DSVs compose 29.6% of the emergence community. The abundance of each DSV varies between samples. DSVs 4, 10, 12, and 81 are the most abundant *Ferroplasma* DSVs in the Emergence community, 4, 10, 12, 34, and 59 in the limestone lined channel, and DSVs 56 and 81 in the Rose Pool (Fig. 3). The DSVs are phylogenetically placed within *Ferroplasma* groups I, II, III and between Groups IV and V (Fig. 2).

Metagenomics

74,021,852 to 101,886,314 metagenomic reads were retrieved per sample with an average GC content of 55% (Supplementary Table S2). Fourteen bins were retrieved that were phylogenetically placed within the genus *Ferroplasma*. The resulting MAGs ranged from low- to high-quality drafts based on quality and contamination, but lack 16S rRNA gene sequences [20]. Five of these bins were retrieved from individually assembled metagenomes and 9 from the combined assembly. Based on average nucleotide identity (ANI), each of the *Ferroplasma* bins from the individual assemblies was also present in the combined assembly and one bin was represented in the emergence, terrace, and combined assemblies (Supplementary Table S3). For MAGs that were present in more than one assembly, the highest quality bin from that MAG was used for further analysis (Table 2). Nine unique *Ferroplasma* MAGs were present in the dataset. These MAGs shared 68.5 to 88.5% ANI with each other and between 68.6 and 99.4% ANI with published *Ferroplasma* genomes (Supplementary Table S3). Based on a concatenated tree of ribosomal protein genes, eight of the nine MAGs occupy the phylogenetic space between the Group I *Ferroplasma* sp. JA12 and sp. PN-J185 and the Group IV *Ferroplasma* sp. Z-31 and the type strain (Fig. 4). Two MAGs, MAG-4 and MAG-7 are > 85% complete and have limited contamination. Therefore, we have described them in more detail below.

MAG-4

MAG-4 is represented by a single bin which was retrieved from the combined assembly and is the most complete MAG recovered in this study. It is 95.7% complete with 0.4% contamination. MAG-4 contains



2250 gene coding sequences, 1466 of which were annotated by GhostKoala (Table 2). MAG-4 shares 69–74% ANI with the other MAGs and the published *Ferrovum* genomes (Supplementary Table S3). The MAG is phylogenetically positioned between the Group I *Ferrovum* spp. JA12 and PN-J185 and the Group VI type strain and sp. Z-31 (Fig. 4)

Energy metabolism

MAG-4 contains three homologs of the high molecular weight, Cyc-2-like protein previously found in *Ferrovum* spp. (Figs. 5 and 6). This protein is thought to facilitate the oxidation of Fe(II) to Fe(III) [16]. The MAG also contains homologs of the genes necessary for oxidative phosphorylation using a B/A type NADH dehydrogenase (*nuoA-nuoN*) and succinate dehydrogenase (*sdhA-D*), a E/B/A type cytochrome *c* reductase, *cbb₃* and *bd* type terminal oxidases, and an F-type ATPase (Fig. 6).

Carbohydrate metabolism

MAG-4 encodes the enzymes necessary to fix carbon via the Calvin-Benson cycle except the enzyme needed to form ribose 5-phosphate from sedoheptulose-7 phosphate. The MAG

also contains the genes for a nearly complete TCA cycle. It contains a nearly complete suite of genes for glycolysis.

Nutrient acquisition

MAG-4 encodes the structural proteins necessary for nitrogen fixation (*NifHDK*) and denitrification (*NasAB*, *NirBD*), but lacks the genes necessary to import inorganic nitrogen into the cell (Fig. 6). The MAG contains many of the genes necessary for assimilatory sulfate reduction but is missing PAPSS which reduces ammonium persulfate (APS) to 3-phosphoadenosine-5-phosphosulfate (PAPS). It also contains homologs of *DsrA* and *DsrB* which can be used to reduce sulfite to sulfide or to oxidize sulfide to sulfite (Fig. 6). The MAG also contains the genes necessary to import inorganic phosphate and use it in biosynthetic pathways (*PST*, *ppx*, *ppk*).

Motility

MAG-4 contains all the genes necessary to construct flagella except *motXY* which encode proteins are involved in flagellar rotation and *fliJT*. The MAG also contains the genes necessary for chemotaxis (*cheABCD*, *cheR*, and *cheVWZ*)





Table 2 Metagenome assembled genomes. When a taxon was found in multiple assemblies, the MAG with the highest completeness and lowest contamination was used. The MAGs used for further analysis are shown in black text whereas those not used are shown in grey text. The MAGs used in the comparative analysis are highlighted by in grey

Name	Bin ID	Completeness	Contamination	Heterogeneity	Protein Coding Sequences	Annotated Sequences
MAG 1	Combined_89	20.69	0	0		
MAG 2	Combined_19	44.72	0.42	66.67		
MAG 3	Rose Pool 43	68.75	4.13	18.18		
	Combined_70	58.28	5.17	40		
MAG 4	Combined_34	95.76	0.5	0	2250	1466
MAG 5	Combined_31	43.89	0	0		
	Rose Pool 56	36.51	0	0		
MAG 6	Combined_120	42.16	1.72			
MAG 7	Limestone Lined Terr. 25	86.7	0.63	100	1764	1158
	Emergence 33	72.59	0	0		
	Combined_101	70.86	0	0		
MAG 8	Combined_72	86.37	33.74	80.26		
	Emergence 23	83.17	54.31	81.58		
MAG 9	Combined_5	30.17	0	0		



MAG-7

MAG-7 is represented by bins retrieved from the emergence, limestone lined channel, and co-assembled metagenomes (Table 2). The highest quality bin is from the limestone lined terrace and is 86.7% complete with 0.63% contamination. The other two bins are 72.6 and 70.9% complete with 0% contamination. Because the most complete bin is only 86.7% complete, when genes were absent from that bin, we searched the other two less complete bins

Energy metabolism

MAG-7 contains one homolog to the high-molecular weight Cyc2 type protein found in other *Ferroplasma* taxa

(Figs. 5 and 6). The Cyc2 sequence is phylogenetically placed as a sister group to those from the type strain and *Ferroplasma* sp. Z31 (Fig. 4). MAG-7 also contains homologs of the genes necessary for oxidative phosphorylation with a B/A type NADH dehydrogenase (*nuoA-nuoN*) and succinate dehydrogenase (*sdhA-D*), an E/B/A type cytochrome *c* reductase, *bo*, *bd*, and *cbb₃* type terminal oxidases, and an F-type ATP-ase.

Nutrient acquisition

MAG-7 encodes the genes necessary for reducing nitrate to nitrite (*nasAB*), but not those to further reduce the nitrate to ammonia (*nirBD*; Fig. 6). The MAG also contains the genes necessary to import inorganic phosphate and use it in biosynthetic pathways (*PST*, *ppx*, *ppk*). MAG-7 does not encode protein necessary for nitrogen fixation and lacks the genes necessary to transform urea to ammonia.

Carbohydrate metabolism

MAG-7 contains most of the genes necessary for carbon fixation via the Calvin-Benson cycle but like MAG-4 does not encode the enzyme needed to form ribose 5-phosphate from sedoheptulose-7 phosphate. The MAG encodes the proteins necessary for a complete TCA cycle. It contains a nearly complete suite of genes for glycolysis.


Motility

MAG-7 contains all the genes necessary to synthesize a functional flagellum except *motX* and *motY* which are also missing from MAG-4. The MAG also contains genes involved in bacterial chemotaxis including *mcp*, *aer*, *tar*, *cheABCD*, *cheR*, and *cheVY*, but it does not appear to contain *cheW* and *cheZ* which are present in MAG-4.

Discussion

Co-occurrence of *Ferroplasma* spp. in cabin branch communities


All known *Ferroplasma* spp. are obligate, Fe (II) oxidizing autotrophs that use O₂ as a terminal electron acceptor [7, 16, 17]. Therefore, co-occurring *Ferroplasma* spp. would presumably compete for common resources if they exploited identical resources, possibly including Fe (II) and oxygen, essential nutrients (e.g., N and P), and trace elements. At Cabin Branch, Fe (II) is present in mmol/L amounts constantly supplied by mine-impacted sources and is unlikely to be limiting. Dissolved oxygen is also present although the concentration is lower at the emergence and increases downstream. Ammonia and phosphate were detected at all sites. Alternately, gene content differences could explain co-existence of *Ferroplasma* populations at Cabin Branch. Co-occurring strains




of the same species have been found in a variety of environments including cyanobacterial blooms [21], and solar saltern/crystallization ponds [22, 23]. In some communities, successful co-occurrence results from genomic variation or differential gene expression between the strains of the co-occurring species. For example, six co-occurring strains of the Epsilonproteobacteria *Lebetimonas* acquired new functional genes via lateral gene transfer in sea mounts in the Mariana Arc [24]. Similarly, strains of *Salinabacter ruber* contain hypervariable regions in their genome which are associated with differences in surface properties [22]. Co-occurring strains of *S. ruber* also express different metabolite pools [23]. These differences likely allow co-occurring strains to avoid competing within the environment [22, 24]. Like these communities, the *Ferrovum* spp. at Cabin Branch

may be able to co-exist due to variations in their genomes or in gene expression [7, 16, 17].

Ferrovum spp. compose over 25% of the bacterial community in the emergence and limestone lined channel communities and 5% of the Rose Pool community, based on the 16S rRNA amplicon libraries. The relative abundance of *Ferrovum* spp. appears to increase with the rate of non-photosynthetic inorganic carbon assimilation (Fig. 7). These data suggest that these populations are important primary producers in the environment because *Ferrovum* spp. are the only abundant chemoautotroph and therefore, likely the main contributor to dark carbon fixation. Our previous work on samples collected in 2015 reported a single *Ferrovum* OTU in Cabin Branch. However, that analysis relied on a definition of 97% identity in 16S rRNA sequences [5] which is the





common cutoff for delineating species based on 16S rRNA gene sequence identity level, but likely underestimates bacterial diversity [25]. Currently, methods using unique sequences are becoming more common and may be more robust [26, 27]. Using a DSV approach on samples collected in 2017 as part of this research, we recovered nine unique *Ferrovum* taxa: four *Ferrovum* DSVs co-occur in the emergence community, five in the limestone lined channel, and eight in the Rose Pool sediments (Figs. 2 and 3). Genome-based studies using average nucleotide identity (ANI) and multilocus phylogenetic analysis are more robust than 16S rRNA-based analyses and are the best practice for assigning species affiliations in genomes [28]. Metagenomic data from Cabin Branch support the presence of multiple *Ferrovum* spp. at each site, suggesting that the diversity of *Ferrovum* spp. at AMD sites globally may be underestimated because studies have traditionally used an OTU approach [4, 5].

To co-occur successfully, the strains may partition the niche to use different resources as is seen in closely-related taxa or intraspecific competitors in eukaryotic systems [29–31]. If phylogenetic distance correlates with metabolic diversity, we might expect co-occurring strains of *Ferrovum* would be distantly related. Indeed, those that co-occur in the Cabin Branch communities span the Group I, II, and III *Ferrovum* clades and the limestone lined channel and Rose Pool also include DSVs that are not placed within a clade (Fig. 1). However, some co-occurring *Ferrovum* spp. are closely related to one another. Multiple DSVs from the same


Ferrovum group are found in the same microbial community. For example, two Group II DSVs are found in all three communities sampled and the Rose Pool community contains three closely related DSVs phylogenetically placed basally to the Group V *Ferrovum* (Fig. 2). Therefore, phylogenetic distance may not determine whether taxa can coexist. However, published *Ferrovum* taxa vary in their gene content with respect to genes associated with motility and nitrogen cycling among others, likely due to lateral gene transfer [15, 16, 18]. Therefore, we can examine how genome content may allow co-occurring taxa to partition resources and avoid competition.

Relating two new *Ferrovum* spp. (MAG-4 and MAG-7) to published species

Here, we compared our *Ferrovum* MAGs to two published *Ferrovum* species from *Ferrovum* Groups I (*F. myxofaciens* sp. P3G) and IV (*Ferrovum* sp. JA12) to explore if genomic variability could contribute to the co-occurrence of these taxa. Our two *Ferrovum* MAGs and the Group I and Group IV genomes share numerous features, but differ in key ways relating to energy metabolism, nitrogen cycling, and motility. Because the MAGs are incomplete, we only consider a feature to be missing if all of genes that code for key enzymes related to that feature are absent from the MAG.

Energy metabolism

All cultured *Ferrovum* spp. can grow autotrophically via iron oxidation [7, 15, 16, 18, 19] and we predict that the



two MAGs presented here share that ability. Each genome contains high molecular weight, Cyc2-like proteins found in other *Ferrovum* taxa [16]. In the acidophilic Fe oxidizers *Acidithiobacillus ferrooxidans* and a *Leptospirillum* spp. Cyc2 is an Fe oxidase [32, 33]. Cyc2-like proteins are also highly expressed in neutrophilic Fe(II) oxidizers [33–35]. Therefore, this protein likely plays a role in Fe(II) oxidation in *Ferrovum*. In *Acidithiobacillus ferrooxidans* and *Gallionella ferruginea*, Fe(II) oxidation is paired with carbon fixation. The MAGs encode a nearly full suite of genes for carbon fixation via the CBB pathway but differ in the types of terminal oxidases they use for oxidative phosphorylation (Fig. 5). MAG-7, *F. myxofaciens* sp. P3G, and *Ferrovum* sp. JA12 all contain cytochrome *bo*₃ type oxidases, but MAG-4 does not contain this type of cytochrome. Similarly, MAGs 4 and 7

and *Ferrovum* sp. JA12 contain a cytochrome *bd*-type quinol oxidase which is not present in *F. myxofaciens* sp. P3G (Fig. 5). Terminal oxidases differ in their affinity for oxygen and are differentially expressed depending on the concentration of oxygen [34]. For example, *E. coli* express cytochrome *bo*₃ under oxic conditions, and cytochrome *bd*, which has a higher affinity for O₂, under oxygen limiting conditions [34]. MAG-4 lacks genes for cytochrome *bo*₃ which may indicate that this taxon does not colonize well-oxygenated environments that may be more suitable for MAG-7, *F. myxofaciens* sp. P3G, and *Ferrovum* sp. JA12. Similarly, MAGs 4 and 7 and *Ferrovum* sp. JA12 may be able to colonize microaerobic environments that are not ideal for *F. myxofaciens* sp. P3G which lacks high affinity terminal oxidases. In Cabin Branch, these differences may lead to MAGs 4 and 7

preferentially inhabiting different portions of the biofilm with MAG-7 inhabiting the better oxygenated surface of the biofilms and MAG-4 residing in more microaerobic niches within the biofilm.

Nitrogen cycling

Ferroplasma myxofaciens sp. P3G can fix nitrogen, but this trait is not conserved within the genus. Neither *Ferroplasma* sp. Z-31 nor PN-J185 are capable of nitrogen fixation [15, 18]. At Cabin Branch, MAG-4 encodes the enzymes necessary for N-fixation (NifH, D, and K) and the hydrolysis of urea (Fig. 6) while MAG-7 does not. The ability to fix nitrogen would allow MAG-4 to colonize low NH_4 environments. Additionally, it may influence where MAG-4 resides in the biofilm community as nitrogenase proteins are oxygen sensitive [36]. Therefore, MAG-4 may preferentially reside in low-oxygen environments that provide enough oxygen to perform Fe(II) oxidation, but at low enough concentrations to protect nitrogenase from oxidative damage. This idea is supported by the presence of *cbb*₃ and *bd* type oxidases in MAG-4 that are preferentially expressed in microaerobic conditions as discussed above.

Motility

Ferroplasma spp. vary in their ability to be motile. *Ferroplasma* sp. JA-12 and the closely related *Ferroplasma* sp. PN-J185 lack genes both for flagella and for chemotaxis whereas *F. myxofaciens* sp. P3G contains the genes necessary for a functional flagellum [9, 15–17, 19]. The MAGs from Cabin Branch are more similar to *F. myxofaciens* sp. P3G in this respect, because they encode the genes necessary for the synthesis of a functional flagellum. MAG-7 and *F. myxofaciens* sp. P3G also contain homologs of *tar*, a chemotaxis related protein that responds to the presence of aspartate or maltose and *aer* which produces a protein that allow a chemotactic response to redox state [37, 38] potentially indicating that these taxa can respond to different environmental cues than MAG-4 which lacks these genes.

Conclusions

We recovered co-occurring *Ferroplasma* taxa with distinct metabolisms from the emergence and outflow of an AMD site. A DSV-based approach suggested multiple *Ferroplasma* taxa co-occur which contrasts to previous OTU-based studies [5]. These data and the *Ferroplasma* genomes recovered from metagenomic data highlight the limits of using OTU clustering approaches in identifying discrete species. We also identified the metabolic diversity among closely related *Ferroplasma* spp. that likely facilitates the co-occurrence of these taxa. Specifically, the differences in nutrient cycling, motility, and chemotaxis

may facilitate co-occurrence without direct competition for resources. These physiological differences may also help to explain why *Ferroplasma* spp. are nearly ubiquitous in AMD environments despite the geochemical diversity of these environments. These data help fill in gaps of missing diversity in the *Ferroplasma* clade which are ubiquitous in AMD environments but difficult to culture. Finally, our data highlight genetic diversity necessary for closely related co-occurring species.

Methods

Site location and sampling

Cabin Branch is a site in the Daniel Boone National Forest in Kentucky near the border with Tennessee. Limestone was added to the channel as a passive remediation strategy and groundwater flows out from an emergence and across a limestone-lined channel before entering a pond, the Rose Pool (Figs. 1 and 7). The microbial communities within Cabin Branch are predominantly the Fe (II)-oxidizing taxon *Ferroplasma myxofaciens* [5]. Samples were collected on March 16, 2017 between 10 am and 2 pm. Sample collection was approved by and performed in collaboration with the staff at Daniel Boone National Forest.

Geochemical analyses

Aqueous geochemistry

At the time of sample collection, temperature and pH were measured and recorded using a WTW 330i meter and probe (Xylem Analytics, Weilheim, Germany). A YSI 30 conductivity meter and probe (YSI Inc., Yellow Springs, OH, USA) was used to measure conductivity. Total ammonia ($\text{NH}_4(\text{T}) = \text{NH}_3 + \text{NH}_4^+$) and ferrous iron (Fe(II)) concentration were determined using Hach DR1900 Portable Spectrometers (Hach Company, Loveland, CO) using the salicylate method (Hach Method 8155) and the phenanthroline method (Hach Method 8146), respectively.

To examine water geochemistry including major ions, major cations, trace elements, and carbon concentration and isotopes, samples were collected and stored following previously described methods [39–41]. In brief, samples were collected in a 140-mL syringe, filtered through 0.8/0.2 μm polyethersulfone syringe filters, and dispensed into 15-mL polypropylene centrifuge tubes (for major anions), acid-washed 15-mL polypropylene centrifuge tubes (for major cations and trace elements), Labco Exetainers® (Labco Limited, Lampeter, UK) pre-flushed with He (for dissolved inorganic carbon (DIC)), and sterile 50-mL centrifuge tubes (for dissolved organic carbon (DOC)). Major anions were determined at the Analytical Geochemistry Laboratory in the Department of Earth and Environmental Sciences at the University of Minnesota using a Thermo Scientific Dionex ICS 5000+

ion chromatography system. Were stored in 15 mL polypropylene centrifuge tubes. Major cations and trace metals were determined at the Analytical Geochemistry Laboratory in the Department of Earth and Environmental Sciences using a Thermo Scientific iCAP 6000 series ICP-OES (inductively coupled plasma optical emission spectroscopy) or a Thermo Scientific X Series 2 ICP-MS (inductively coupled plasma mass spectrometry). DIC concentration and ^{13}C isotopic signal analyses were performed on a GasBench II system interfaced to a Delta V Plus isotope ratio mass spectrometer (IR-MS) (Thermo Scientific, Bremen, Germany) by the Stable Isotope Facility (SIF) at the University of California, Davis for analysis. DOC concentration and ^{13}C isotopic signal analyses were performed on a O.I. Analytical Model 1030 TOC Analyzer (O.I. Analytical, College Station, TX, USA) by the SIF at UC-Davis. The standard deviation for DOC was determined from replicate analyses of each sample. For each analysis, field blanks were taken using 18.2 M Ω /cm deionized water transported to the field in 1-l acid washed Nalgene bottles for comparison to samples.

Colorimetric tests are optimized for 20 °C and are known to have chemical interferences, thus the spectrophotometrically derived Fe(II) concentrations should be considered semi-quantitative whereas Fe_{total} is measured via ICP-MS or ICP-OES with samples that are acidified with concentrated HNO₃ to keep metals in solution. Our assumption is that the majority of Fe_{total} is Fe(II) and most Fe(III) present precipitates immediately as Fe(OH)_{3(S)} and is collected on 0.2 μm filters during sample collection.

CO₂ assimilation

In situ microcosms were performed to assess the potential for inorganic carbon uptake through the addition of NaH¹³CO₃. Sample collection and the microcosm set-up followed previously described methods [39–41]. In summary, samples (biofilms, filaments, etc.) were placed in sterile serum vials, overlaid with spring water, capped, and amended with NaH¹³CO₃ (100 μM final concentration) (Cambridge Isotope Laboratories, Inc., Andover, MA, USA). All assays were performed in triplicate between 10 AM and 2 PM. To assess the potential for photoautotrophic NaH¹³CO₃ uptake, microcosms were incubated in the light. In contrast, microcosms were wrapped in foil to assess the potential for chemoautotrophic (dark) NaH¹³CO₃ uptake. At the emergence, both green and white filaments were present. We performed assays on each by sampling predominantly green filaments and predominantly white filaments. We assumed the majority of uptake in the green biomass would be due to photoautotrophy and thus performed one set of microcosms on the bulk filaments/biofilm

(green + white) in the light and then separated the biofilm types (green or white) for microcosms performed in the dark. Following incubation, vials were flash frozen on dry ice and stored at –80 °C until processed (described below). Calculation of assimilation rates is described below.

C and N concentration and stable isotope signals

Samples for ^{13}C natural abundance were collected at the time of sampling using sterile spatulas. These were flash frozen on dry ice, stored in liquid N₂ for transport and stored at –80 °C until processed. To examine ^{13}C signals in natural abundance samples and microcosms, samples were acidified (for microcosms), dried, and weighed following previously described methods [39–41]. Samples were analyzed via an Elementar pyrocube elemental analyzer (EA) periphery connected to an Isoprime 100 continuous flow IRMS (IR-MS) at the University of Minnesota. NIST Standard 2710 was used for linearity corrections. $\delta^{13}\text{C}$ values were calibrated using reference standards USGS-40 and USGS-41 and checked with a laboratory working standard (glycine). The following precautions were taken to minimize cross contamination of natural abundance samples with samples that were amended with NaH¹³CO₃: natural abundance samples were processed, weighed, and analyzed separate from labeled samples and all laboratory processing and weighing equipment was cleaned with 80% ethanol between each sample. In addition, to check for memory effects or cross contamination of samples, standard checks and blanks were included with each analysis batch (no memory effects or cross contamination were detected). Assimilation rates were calculated as described previously [39–41] using difference between the total amount of ^{13}C in natural abundance samples and incubation assay replicates which represents total mass of ^{13}C -labelled DIC in biomass after the incubation period (1–2 h). We performed a one-way ANOVA followed by post hoc pairwise comparisons within the R software package (R version 3.3.2) to compare mean ^{13}C uptake rates and considered *p*-values < 0.05 as significantly different.

Molecular analyses

DNA extraction

Triplicate samples for DNA extraction were collected at the time of sampling using sterile spatulas, flash frozen onsite and stored at –80 °C until processed. DNA was extracted from each replicate sample (*n* = 3) using a DNeasy PowerSoil Kit (Qiagen, Carlsbad, CA, USA) according to the manufacturer's instructions and quantified using a Qubit 3.0 Fluorometer (Invitrogen, Burlington, ON, Canada). Equal volumes of each extraction were pooled and submitted to the University of Minnesota Genomics Center (UMGC) for amplicon

sequencing. As a negative DNA extraction control, we attempted to extract DNA from the filters used for the field blank water samples (described above). These extractions failed to yield detectable DNA and failed library preparation for amplicon sequencing (see below for amplicon sequencing details).

DNA sequencing

Amplicons were sequenced at the UMGC with MiSeq Illumina 2 × 300 bp chemistry targeting the V4 hypervariable region of bacterial and archaeal 16S SSU rRNA gene sequences (515F and 806r) [42] as described previously [40].

Preparation of amplicon libraries followed the UMGC's improved protocol to detect which taxonomic groups that often go undetected using existing methods [43]. Each sample was sequenced once. Total DNA was submitted to the UMGC for metagenomic sequencing and sequenced using HiSeq2500 High-Output 2 × 125 bp chemistry. Three samples were sequenced / lane.

16S rRNA analysis

16S rRNA sequences from Cabin Branch were used to examine community composition. Primers and unpaired sequences were removed using trimmomatic [44]) The surviving reads were processed in dada2 (v.1.4) following the pipeline tutorial [27]. Briefly, forward reads were trimmed to 210 bp and reverse reads were trimmed to 120 bp based on their quality profiles. Sequences with ambiguous bases and those with more than 2 expected errors were removed. Error rates were estimated using the learnErrors command. Sequences were dereplicated using the derepFastq command and the unique sequence variants were inferred using the dada command. Forward and reverse reads were merged using mergePairs. Contigs shorter than 250 or longer than 256 bp and chimeric sequences were removed. The surviving unique, denoised sequences are referred to as denoised sequence variants (DSVs). Taxonomy was assigned using the Silva training set [45] Eukaryotic sequences and those unclassified at the domain level were removed. The closest cultured and environmental relatives were identified using BLASTN [46].

We retrieved 1589 aligned, nearly-full length *Ferrovum* sequences from the Silva non-redundant database to serve as a reference alignment for the analyses described below. A 16S rRNA phylogeny was constructed by retrieving the sequences used by Ullrich et al. [18] with *Ferrotrophicum raditicola*, *Nitrospovibrio tenuis*, and *Nitrospira lenta* used as an outgroup. These sequences were aligned using MAFFT on the Cipres Science Gateway [47] using the aligned *Ferrovum* sequences from the Silva database as a reference alignment. A tree of the smaller subset of sequences was constructed in RAXML-

HPC2 on XSEDE [48] also on the Cipres Science Gateway. Sequences of DSVs classified as *Ferrovum* were added to the alignment using the -addfrags option in MAFFT [49]. Non-full-length sequences were added to the tree using the evolutionary placement algorithm. Trees were rooted and visualized in the interactive tree of life [50]. Raw sequences were uploaded to the NCBI Sequence Read Archive under BioProject PRJNA554371.

Metagenomic analysis

Individual metagenomes were assembled and a co-assembly of all metagenomes was constructed following the “tutorial on assembly-based metagenomics” [51]. Trimmed, quality-controlled sequences were assembled using MegaHit [52] using default parameters except minimum contig length, which was set at 1000 base pairs. Reads were mapped to the assembly using bowtie2 [53] and depth was calculated using the jgi_summarize_bam_contig_depths command in Anvi'o [54]. Contigs were binned using default parameters in meta-bat using [55]. Bin completeness was determined with CheckM [56]. Protein coding regions were identified with prodigal (within CheckM) [57] and GhostKoala was used to annotate protein coding sequences [58]. Pathways were considered to be absent from the taxon when none of the key genes associated with that pathway were present in the MAG.

Each bin was uploaded to KBASE [59] and annotated with Prokka using the “Annotate Assembly and Re-annotate Genomes with Prokka (v1.12)” app. A concatenated gene tree containing each bin and four published *Ferrovum* spp. genomes was constructed using the “Insert Set of Genomes Into Species Tree 2.1.10” app. This app uses up to 49 ribosomal and single-copy genes to construct a phylogenetic tree. MAGs that were more closely related to *Ferrovum* spp. than other taxa were selected for further analysis. The pairwise ANI between each bin and published *Ferrovum* genomes was calculated using anvi-compute-ani in Anvi'o. If MAGs shared >98% pairwise ANI and were phylogenetically cohesive, we considered them to be representing the same populations and the bin with the highest completeness was chosen for further analysis. Raw reads, assemblies, and MAGs were uploaded to the NCBI Sequence Read Archive under BioProject PRJNA554371. The abundance of *Ferrovum* in each metagenomes using One Codex [60].

16S rRNA A custom blast database containing the 16S rRNA gene from the *Ferrovum myxofaciens* type strain (NR_117782.1) using the makeblastdb command in BLAST+. Each of the *Ferrovum* bins were searched for 16S rRNA genes using BLAST and a cutoff value of E-30.

Cyc-2 A custom blast database of the high molecular weight Cyc-2 like proteins was constructed using the methods described above. Each of the *Ferrovum* bins were searched for the Cyc-2-like protein identified by Ullrich et al., [16] by blasting translated nucleotide sequences against the blast database with an E-value of E-120. The retrieved sequences, database sequences, and a sequence for Cyc-2 from an *Acidithiobacillus* sp. (AVK42811.1) were aligned using MAFFT and a maximum likelihood tree was constructed as described above. The tree was rooted on *Acidithiobacillus* and visualized in iTOL.

Comparison with other *Ferrovum* genomes

Existing *Ferrovum* genomes are located within either the Group I or Group IV *Ferrovum* clades [18]. The genome for *Ferrovum* sp. JA12 (Group I) and *Ferrovum myxofaciens* (Group IV) were downloaded from NCBI. CheckM was used to determine genome completeness, and within CheckM, prodigal was used to identify protein coding sequences. Ghost KOALA was used to annotate protein coding sequences using the genus_prokaryotes database. Protein coding sequences were also annotated with prokka. We then manually compared the gene content of MAGs to that of existing *Ferrovum* genomes.

Supplementary information

Supplementary information accompanies this paper at <https://doi.org/10.1186/s12866-020-01768-w>.

Additional file 1: Table S1. Carbon uptake experiment results.

Additional file 2: Table S2. Sequencing and assembly statistics for metagenomes used in this research.

Additional file 3: Table S3. Average nucleotide identity of MAGs retrieved from the Cabin Branch metagenomes.

Abbreviations

AMD: Acid mine drainage; ANI: Average nucleotide identity; APS: Ammonium persulfate 3-phosphoadenosine-5-phosphosulfate; DIC: Dissolved inorganic carbon; DSV: Denoised sequence variants; ICP-MS: Inductively coupled plasma mass spectrometry; ICP-OES: Inductively coupled plasma atomic emission spectroscopy; MAG: Metagenome assembled genomes; PAPS: 3-phosphoadenosine-5-phosphosulfate; TCA: Tricarboxylic acid cycle

Acknowledgements

We are grateful to the staff of the National Forest Service and Daniel Boone National Forest, especially Margueritte Wilson and Claudia Cotton, for the advice and insight regarding mine locations. We thank A. Gangidine, M. Berberich, R. Jain, and C. Schuler for assistance in field sampling and processing samples in the laboratory.

Authors' contributions

JRH, CLG, and TLH contributed to the conception of the project, data analysis interpretation of results, creation of figures, writing and editing the manuscript. JRH and TLH performed sample collection and sample analysis. The authors have read and approve of the manuscript.

Funding

This work was supported by the University of Cincinnati and the University of Minnesota. The authors acknowledge the Minnesota Supercomputing Institute (MSI) at the University of Minnesota for providing resources that contributed to the research results reported within this paper. The funders

had no role in the design of the study and collection, analysis, and interpretation of data.

Availability of data and materials

All analyses tools used in the study are publicly available. 16S rRNA and functional genes of closely related species and outgroups were downloaded from NCBI databases and accession numbers are provided for these sequences. Raw reads and assembled scaffolds for the metagenomes and MAG sequences as well as the amplicon libraries have been deposited in the NCBI under BioProject PRJNA554371.

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Author details

¹Department of Earth and Planetary Sciences, University of California, Davis, CA 95616, USA. ²Department of Earth and Environmental Sciences, University of Minnesota, Minneapolis, MN 55455, USA. ³Department of Plant and Microbial Biology, University of Minnesota, 218 Cargill Building, St. Paul, MN 55108, USA. ⁴The BioTechnology Institute, University of Minnesota, St. Paul, MN 55108, USA.

Received: 23 July 2019 Accepted: 29 March 2020

Published online: 18 May 2020

References

- Baker BJ, Banfield JF. Microbial communities in acid mine drainage. FEMS Microbiol Ecol. 2003;44:139–52.
- Johnson DB. Chemical and microbiological characteristics of mineral spoils and drainage waters at abandoned coal and metal mines. Water Air Soil Pollut Focus. 2003;3:47–66.
- Schippers A, Breuker A, Blazejak A, Bosecker K, Kock D, Wright TL. The biogeochemistry and microbiology of sulfidic mine waste and bioleaching dumps and heaps, and novel Fe (II)-oxidizing bacteria. Hydrometallurgy. 2010;104:342–50.
- Grettenberger CL, Pearce AR, Bibby KJ, Jones DS, Burgos WD, Macalady JL. Efficient low-pH iron removal by a microbial iron oxide mound ecosystem at scalp level run. Appl Environ Microbiol. 2017;83:e00015–7.
- Havig JR, Grettenberger C, Hamilton TL. Geochemistry and microbial community composition across a range of acid mine drainage impact and implications for the Neoproterozoic-Paleoproterozoic transition. J Geophys Res Biogeosciences. 2017;122:1404–22.
- Jones DS, Kohl C, Grettenberger C, Larson LN, Burgos WD, Macalady JL. Geochemical niches of iron-oxidizing acidophiles in acidic coal mine drainage. Appl Environ Microbiol. 2015;81:1242–50.
- Johnson DB, Hallberg KB, Hedrich S. Uncovering a microbial enigma: isolation and characterization of the streamer-generating, iron-oxidizing, acidophilic bacterium "*Ferrovum myxofaciens*". Appl Environ Microbiol. 2014; 80:672–80.
- Hallberg KB, Coupland K, Kimura S, Johnson DB. Macroscopic streamer growths in acidic, metal-rich mine waters in North Wales consist of novel and remarkably simple bacterial communities. Appl Environ Microbiol. 2006; 72:2022–30.
- Schlömann M, Kipy J, Mosler S, Poehlein A, Keller A, Janneck E, et al. Physiological, genomic, and proteomic characterization of new "*Ferrovum*" strains obtained from a pilot plant for mine-water treatment. Adv Mat Res. 2013;825:149–52.
- Johnson DB. Acidophilic microbial communities: candidates for bioremediation of acidic mine effluents. Int Biodeter Biodegr. 1995;35:41–58.
- Kipy J, Jwair RJ, Gelhaar N, Wiacek C, Janneck E, Schlömann M. Enrichment of "*Ferrovum*" spp. and *Gallionella* relatives using artificial mine water. Adv Mat Res. 2013;825:54–7.
- Tischler JS, Jwair RJ, Gelhaar N, Drechsel A, Skirl A-M, Wiacek C, et al. New cultivation medium for "*Ferrovum*" and *Gallionella*-related strains. J Microbiol Methods. 2013;95:138–44.

13. Kuang J-L, Huang L-N, Chen L-X, Hua Z-S, Li S-J, Hu M, et al. Contemporary environmental variation determines microbial diversity patterns in acid mine drainage. *ISME J.* 2013;7:1038.
14. Kay C, Rowe O, Rocchetti L, Coupland K, Hallberg K, Johnson D. Evolution of microbial "streamer" growths in an acidic, metal-contaminated stream draining an abandoned underground copper mine. *Life.* 2013;3:189–210.
15. Ullrich SR, González C, Poehlein A, Tischler JS, Daniel R, Schlömann M, et al. gene loss and horizontal gene transfer contributed to the genome evolution of the extreme acidophile "*Ferrovum*". *Front Microbiol.* 2016;7:797.
16. Ullrich SR, Poehlein A, Tischler JS, González C, Ossandon FJ, Daniel R, et al. Genome analysis of the biotechnologically relevant acidophilic iron oxidising strain JA12 indicates phylogenetic and metabolic diversity within the novel genus "*Ferrovum*". *PLoS One.* 2016;11:e0146832.
17. Mosler S, Poehlein A, Voget S, Daniel R, Kipry J, Schlömann M, et al. Predicting the metabolic potential of the novel iron oxidising bacterium "*Ferrovum*" sp. JA12 using comparative genomics. *Adv Mat Res.* 2013;825:153–6.
18. Ullrich SR, Poehlein A, Daniel R, Tischler JS, Vogel S, Schlömann M, et al. Comparative genomics underlines the functional and taxonomic diversity of novel "*Ferrovum*" related iron oxidizing bacteria. *Adv Mat Res.* 2015;1130:15–8.
19. Moya-Beltrán A, Cárdenas JP, Covarrubias PC, Issotta F, Ossandon FJ, Grail BM, et al. Draft genome sequence of the nominated type strain of "*Ferrovum myxofaciens*," an acidophilic, iron-oxidizing betaproteobacterium. *Genome Announc.* 2014;2:e00834–14.
20. Bowers RM, Kyrpides NC, Stepanauskas R, Harmon-Smith M, Doud D, Reddy TBK, et al. Minimum information about a single amplified genome (MISAG) and a metagenome-assembled genome (MIMAG) of bacteria and archaea. *Nat Biotechnol.* 2017;35:725.
21. Willis A, Woodhouse JN, Ongley SE, Jex AR, Burford MA, Neilan BA. Genome variation in nine co-occurring toxic *Cylindrospermopsis raciborskii* strains. *Harmful algae.* 2018;73. *J Bacteriol.* 2008;190:157–66.
22. Peña A, Teeling H, Huerta-Cepas J, Santos F, Yarza P, Brito-Echeverría J, et al. Fine-scale evolution: genomic, phenotypic and ecological differentiation in two coexisting *Salinibacter ruber* strains. *ISME J.* 2010;4:882.
23. Antón J, Lucio M, Peña A, Cifuentes A, Brito-Echeverría J, Moritz F, et al. High metabolomic microdiversity within co-occurring isolates of the extremely halophilic bacterium *Salinibacter ruber*. *PLoS One.* 2013;8:e64701.
24. Meyer JL, Huber JA. Strain-level genomic variation in natural populations of *Lebetimonas* from an erupting deep-sea volcano. *ISME J.* 2014;8:867.
25. Rodríguez-R LM, Castro JC, Kyrpides NC, Cole JR, Tiedje JM, Konstantinidis KT. How much do rRNA gene surveys underestimate extant bacterial diversity? *Appl Environ Microbiol.* 2018;84:e00014–8.
26. Callahan BJ, McMurdie PJ, Holmes SP. Exact sequence variants should replace operational taxonomic units in marker-gene data analysis. *ISME J.* 2017;11:jsmej2017119.
27. Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJA, Holmes SP. DADA2: High-resolution sample inference from Illumina amplicon data. *Nat Methods.* 2016;13:nmeth.3869.
28. Figueras MJ, Beaz-Hidalgo R, Hossain MJ, Liles MR. Taxonomic affiliation of new genomes should be verified using average nucleotide identity and multilocus phylogenetic analysis. *Genome Announc.* 2014;2:e00927–14.
29. MacArthur RH. Population ecology of some warblers of northeastern coniferous forests. *Ecology.* 1958;39:599–619.
30. Inouye DW. Resource partitioning in bumblebees: experimental studies of foraging behavior. *Ecology.* 1978;59:672–8.
31. Swanson BO, Gibb AC, Marks JC, Hendrickson DA. Trophic polymorphism and behavioral differences decrease intraspecific competition in a cichlid, *Herichthys minckleyi*. *Ecology.* 2003;84:1441–6.
32. Castelle C, Guiral M, Malarte G, Ledgham F, Leroy G, Brugna M, et al. A new iron-oxidizing/O₂-reducing supercomplex spanning both inner and outer membranes, isolated from the extreme acidophile *Acidithiobacillus ferrooxidans*. *J Biol Chem.* 2008;283:25803–11.
33. Jeans C, Singer SW, Chan CS, VerBerkmoes NC, Shah M, Hettich RL, et al. Cytochrome 572 is a conspicuous membrane protein with iron oxidation activity purified directly from a natural acidophilic microbial community. *ISME J.* 2008;2:542.
34. Cotter PA, Chepurí V, Gennis RB, Gunsalus RP. Cytochrome o (cyoABCDE) and d (cydAB) oxidase gene expression in *Escherichia coli* is regulated by oxygen, pH, and the fnr gene product. *J Bacteriol.* 1990;172:6333–8.
35. Chan C, McAllister SM, Garber A, Hallahan BJ, Rozovsky S. Fe oxidation by a fused cytochrome-porin common to diverse Fe-oxidizing bacteria. *Biorxiv.* 2018:228056.
36. Gallon JR. The oxygen sensitivity of nitrogenase: a problem for biochemists and micro-organisms. *Trends Biochem Sci.* 1981;6:19–23.
37. Rebbapragada A, Johnson MS, Harding GP, Zuccarelli AJ, Fletcher HM, Zhulin IB, et al. The Aer protein and the serine chemoreceptor Tsr independently sense intracellular energy levels and transduce oxygen, redox, and energy signals for *Escherichia coli* behavior. *Proc National Acad Sci.* 1997;94:10541–6.
38. Bibikov SI, Biran R, Rudd KE, Parkinson JS. A signal transducer for aerotaxis in *Escherichia coli*. *J Bacteriol.* 1997;179:4075–9.
39. Schuler CG, Havig JR, Hamilton TL. Hot spring microbial community composition, morphology, and carbon fixation: implications for interpreting the ancient rock record. *Front Earth Sci.* 2017;5:97.
40. Hamilton TL, Bennett AC, Murugapiran SK, Havig JR. Anoxygenic phototrophs span geochemical gradients and diverse morphologies in terrestrial geothermal springs. *Msystems.* 2019;4:e00498–19.
41. Havig JR, Hamilton TL. Productivity and community composition of low biomass/high silica precipitation hot springs: a possible window to earth's early biosphere? *Life Basel Switz.* 2019;9:64.
42. Caporaso JG, Lauber CL, Walters WA, Berg-Lyons D, Lozupone CA, Turnbaugh PJ, et al. Global patterns of 16S rRNA diversity at a depth of millions of sequences per sample. *Proc National Acad Sci.* 2011; 108(Supplement 1):4516–22.
43. Gohl DM, Vangay P, Garbe J, MacLean A, Hauge A, Becker A, et al. Systematic improvement of amplicon marker gene methods for increased accuracy in microbiome studies. *Nat Biotechnol.* 2016;34:nb.3601.
44. Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics.* 2014;30:2114–20.
45. Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, et al. The SILVA ribosomal RNA gene database project: improved data processing and web-based tools. *Nucleic Acids Res.* 2013;41:D590–6.
46. Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, et al. BLAST+: architecture and applications. *BMC Bioinformatics.* 2009;10:421.
47. Miller MA, Pfeiffer W, Schwartz T. Creating the CIPRES Science Gateway for Inference of Large Phylogenetic Trees. In: 2010 Gatew Comput Environ Work GCE; 2010. p. 1–8.
48. Stamatakis A. RAxML-VI-HPC: maximum likelihood-based phylogenetic analyses with thousands of taxa and mixed models. *Bioinformatics.* 2006;22:2688–90.
49. Katoh K, Standley DM. MAFFT Multiple Sequence Alignment Software Version 7: Improvements in Performance and Usability. *Mol Biol Evol.* 2013;30(4):772–80.
50. Letunic I, Bork P. Interactive tree of life (ITOL) v3: an online tool for the display and annotation of phylogenetic and other trees. *Nucleic Acids Res.* 2016;44(W1):W242–5.
51. Eren M. A tutorial on assembly-based metagenomics.
52. Li D, Liu C-M, Luo R, Sadakane K, Lam T-W. MEGAHIT: an ultra-fast single-node solution for large and complex metagenomics assembly via succinct de Bruijn graph. *Bioinformatics.* 2015;31:1674–6.
53. Langmead B, Salzberg SL. Fast gapped-read alignment with bowtie 2. *Nat Methods.* 2012;9:357.
54. Eren AM, Esen ÖC, Quince C, Vineis JH, Morrison HG, Sogin ML, et al. Anvi'o: an advanced analysis and visualization platform for 'omics data. *PeerJ.* 2015;3:e1319.
55. Kang DD, Froula J, Egan R, Wang Z. MetaBAT, an efficient tool for accurately reconstructing single genomes from complex microbial communities. *PeerJ.* 2015;3:e1165.
56. Parks DH, Imelfort M, Skennerton CT, Hugenholtz P, Tyson GW. CheckM: assessing the quality of microbial genomes recovered from isolates, single cells, and metagenomes. *Genome Res.* 2015;25:1043–55.
57. Hyatt D, Chen G-L, LoCascio PF, Land ML, Larimer FW, Hauser LJ. Prodigal: prokaryotic gene recognition and translation initiation site identification. *BMC Bioinformatics.* 2010;11:119.
58. Kanehisa M, Sato Y, Morishima K. BlastKOALA and GhostKOALA: KEGG tools for functional characterization of genome and metagenome sequences. *J Mol Biol.* 2016;428:726–31.
59. Arkin AP, Cottingham RW, Henry CS, Harris NL, Stevens RL, Maslov S, et al. KBase: the United States Department of Energy Systems Biology Knowledgebase. *Nat Biotechnol.* 2018;36:566.
60. S.S. Minot, N. Krumm, N.B. Greenfield One codex: a sensitive and accurate data platform for genomic microbial identification bioRxiv. 2015. <https://doi.org/10.1101/027607>.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.