

UC Davis

UC Davis Previously Published Works

Title

Comprehensive management of acute pulmonary embolism in primary care using telemedicine in the COVID-era

Permalink

<https://escholarship.org/uc/item/85v959s8>

Journal

BMJ Case Reports, 14(6)

ISSN

1757-790X

Authors

Chang, Joshua
Isaacs, Dayna J
Leung, Joseph
et al.

Publication Date

2021-06-01

DOI

10.1136/bcr-2021-243083

Peer reviewed



OPEN ACCESS

Comprehensive management of acute pulmonary embolism in primary care using telemedicine in the COVID-era

Joshua Chang ,¹ Dayna J Isaacs ,^{2,3} Joseph Leung,⁴ David R Vinson ^{5,6}

¹Internal Medicine Residency Program, Kaiser Foundation Health Plan Inc, Oakland, California, USA

²University of California Davis School of Medicine, Sacramento, California, USA

³Internal Medicine Residency Program, University of California Los Angeles Health, Los Angeles, California, USA

⁴Adult and Family Medicine, Kaiser Permanente San Francisco Medical Center, San Francisco, California, USA

⁵Emergency Medicine, Kaiser Permanente Roseville Medical Center, Roseville, California, USA

⁶Kaiser Permanente Division of Research and the CREST Network, Oakland, California, USA

Correspondence to

Dr David R Vinson;
drvinson@ucdavis.edu

Accepted 27 April 2021

SUMMARY

A healthy, active woman in her 70s reported intermittent exertional dyspnoea for 2 months, notable during frequent open-water swimming. Symptoms were similar to an episode of travel-provoked pulmonary embolism 3 years prior. She denied chest pain, cough, fever, extremity complaints and symptoms at rest. Due to the COVID-19 pandemic, her healthcare system was using secure telemedicine to evaluate non-critical complaints. During the initial video visit, she appeared well, conversing normally without laboured breathing. An elevated serum D-dimer prompted CT pulmonary angiography, which identified acute lobar pulmonary embolism. After haematology consultation and telephone conversation with the patient, her physician prescribed rivaroxaban. Her symptoms rapidly improved. She had an uneventful course and is continuing anticoagulation indefinitely. The pandemic has increased the application of telemedicine for acute care complaints. This case illustrates its safe and effective use for comprehensive management of acute pulmonary embolism in the primary care setting.

BACKGROUND

Since the early 2000s, telemedicine has seen increasing adoption in both primary and specialty care settings.¹ Telemedicine tools such as real-time patient–physician video or telephone visits and asynchronous secure messaging via a patient portal integrated within an electronic health record system can increase efficiency and healthcare access.^{1–3} These benefits come with challenges to implementation, including familiarity with technology, establishment of patient–physician relationship and availability of support infrastructure.^{3–5} The COVID-19 pandemic caused by SARS-CoV-2 has accelerated the adoption and reliance of telehealth to provide timely, low risk and accessible care.^{6,7} This may include triage, diagnosis and management of acute pulmonary embolism.

Initial care for acute, symptomatic pulmonary embolism is shifting away from hospital-based care to outpatient management for select low-risk patients.^{8–10} International consensus exists across major societies in recommending outpatient management for low-risk pulmonary embolism.^{11–15} The definition of ‘outpatient management’, however, varies widely in the literature.¹⁶ Most outpatient management studies of acute pulmonary embolism have been undertaken in the emergency department or secondary care settings. Little research attention has been directed on pulmonary embolism

management in the primary care setting.^{17–19} How the recent expansion of telemedicine in primary care can be applied to the diagnosis and treatment of patients with acute pulmonary embolism has not been well studied. We explore the utility of telemedicine in a case of comprehensive clinic-based outpatient pulmonary embolism management using video visit, telephone follow-up and secure messaging in a health system with an integrated electronic health record.

CASE PRESENTATION

A healthy, active woman in her 70s with a remote history of provoked pulmonary embolism presented to her primary care physician (PCP) with 2–3 months of dyspnoea on exertion during the COVID-19 pandemic. The patient is an active open-water swimmer and initially noticed reduction in exercise tolerance, both in the water and when walking up inclines, which she attributed to deconditioning. She also endorsed palpitations and occasional dizziness, but denied chest pain, cough, fevers, syncope and dyspnoea at rest. She had no recent surgery, periods of reduced mobility or known history of malignancy. However, she had a history of provoked pulmonary embolism in the setting of air travel 3 years prior, for which she had completed 3 months of dabigatran. The initial evaluation for this current episode was conducted through secure video visit as in-person visits were limited during the early COVID-19 pandemic. Through video, the PCP noted no acute distress and no laboured breathing.

INVESTIGATIONS

Initial outpatient laboratory work-up was ordered on day 1 and completed on day 3. It showed normal complete blood count, thyroid stimulating hormone and elevated D-dimer of 2.41 µg/mL (normal <0.49 µg/mL). Results were transmitted to the patient, who sent a secure message to her PCP reporting mild improvement in symptoms and inquiring about next steps. They discussed by messaging the likelihood of pulmonary embolism. He ordered advanced imaging by CT pulmonary angiography. It was performed the next day (4 days after presentation) and demonstrated new partial filling defects of the distal right upper lobe pulmonary artery. Prior hypercoagulable work-up was negative. COVID-19 testing was declined based on self-assessed low-risk exposure from strict social distancing adherence.



© BMJ Publishing Group Limited 2021. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Chang J, Isaacs DJ, Leung J, et al. *BMJ Case Rep* 2021;**14**:e243083. doi:10.1136/bcr-2021-243083

TREATMENT

At the end of the imaging study, the patient waited on campus while the radiologist called the PCP with the results. The PCP then spoke by phone with haematology, who advised indefinite anticoagulation therapy in the setting of recurrent pulmonary embolism. Age-appropriate cancer screening and COVID-19 testing were also recommended. The PCP called the patient, discussed the results and directed her to the pharmacy on campus to pick-up the rivaroxaban he ordered. He also submitted a referral for a haematology appointment.

OUTCOME AND FOLLOW-UP

The pharmacy-led anticoagulation management service provided next-day patient education by telephone and supplemental instructions sent by secure messaging. She had a secure video visit with haematology 1 week after her pulmonary embolism diagnosis. This second haematologist agreed with the earlier recommendation of indeterminate anticoagulation. After 1 month of rivaroxaban, the patient learnt that its reversal agent was less accessible in many hospitals than idarucizumab, the agent designed for dabigatran reversal. She then requested switching to dabigatran, with which she has continued treatment. She has experienced no bleeding complications nor recurrence of venous thromboembolism.

DISCUSSION

The COVID-19 pandemic has accelerated the utilisation of telemedicine as healthcare systems grappled with restrictions on in-person evaluation.²⁰ The Centers for Medicare and Medicaid Services and the US Department of Health and Human Services have also incentivised telehealth with waivers related to reimbursement and liability.^{21 22} One cross-sectional analysis found a 35.3% increase in telemedicine use along with a 21.4% decrease in office visits and fewer laboratory testing during the second quarter of 2020, suggesting both structural changes in healthcare delivery and associated challenges.²³

The diagnosis of pulmonary embolism in primary care

Many professional societies recommend evidence-based, high-value diagnostic strategies for the evaluation of patients with suspected acute pulmonary embolism using validated clinical prediction rules (eg, Wells criteria) and selective probability-driven D-dimer measurements.^{24–27} Such a risk-stratified approach has been externally validated in the primary care setting.^{28 29} The PCP in our case followed this internationally endorsed approach.

The time from initial presentation in primary care to diagnosis of acute pulmonary embolism varies widely. The evaluation can be prolonged if a pulmonary embolism diagnosis is not pursued until after diagnostic tests for other conditions (eg, chest radiographs for pneumonia) fail to explain the symptoms. Most in-person work-ups for acute pulmonary embolism (starting with the initial presentation) are completed within 7 days, though about 25% run longer.^{30 31} We would not expect a telemedicine approach to greatly differ in duration. Our patient's diagnostic evaluation was completed within 5 days, which is within the normal range of in-person work-ups described in the literature. The British Thoracic Society guideline recommends that 'patients with suspected (pulmonary embolism) should, where reasonably practical, undergo investigation on the same day of presentation to exclude a diagnosis of (pulmonary embolism).'¹⁴ US guidelines across multiple specialties do not include similar temporal parameters.^{24 26 32–34}

Eligibility for telemedical care of acute pulmonary embolism

Telemedicine for pulmonary embolism management in the primary care setting has not been well described in the literature. Our case sheds light on this novel approach. This patient was particularly well-suited to video-based outpatient diagnosis and treatment because of her familiarity with technology, established PCP relationship, access to advanced imaging and laboratory testing, as well as prior pulmonary embolism and anticoagulation experience. Additionally, the patient's ready access to close follow-up and her PCP's timely responses via secure messaging and telephone both facilitated outpatient management.

Although the PCP did not document any formal risk evaluation in the electronic health record, the patient appears to have met low-risk criteria for outpatient management from the American College of Chest Physicians: 'clinically stable with good cardiopulmonary reserve; no contraindications...; expected to be compliant with treatment; and the patient feels well enough to be treated at home.'¹³ She also likely met outpatient criteria of the Hestia clinical decision rule, if we assume that continued ocean swimming and absence of dyspnoea at rest suggest haemodynamic stability and pulse oximetry $\geq 90\%$, though this correlation has not been studied.³⁵

Home measurement of her vital signs would have helped complete her outpatient eligibility assessment. Obtaining heart rate, respiratory rate, blood pressure and temperature at home have been possible for many years. The ability to accurately measure pulse oximetry at home is a more recent development, following the advent of reliable portable devices.³⁶ Smart phones may also soon be used for pulse oximetry measurements.³⁷ The increasing availability of reliable home vital sign measurements may open the door for research on selective in-home use of risk stratification triage tools, like the Hestia decision rule and the Pulmonary Embolism Severity Index.^{9 35 38} Some insurance companies cover the costs of remote patient monitoring devices, such as pulse oximeters and blood pressure monitoring devices. Reimbursement may continue to expand with the modernisation of telehealth.

Comprehensive primary care-based pulmonary embolism management

This case report adds to the emerging literature describing comprehensive pulmonary embolism management in the primary care setting, including 3 in-person cases.^{17–19} All 3 patients did well and developed no recurrent venous thromboembolism or bleeding complications. These cases met Pulmonary Embolism Severity Index and Hestia eligibility criteria for outpatient management. As in our current case, their PCPs also had ready telephone access to on-call haematologists and pulmonologists for consultation. The patients received prompt anticoagulation therapy with supplemental education and long-term monitoring by pharmacy-led anticoagulation management services, a centralised model adopted by many institutions.³⁹

The promises and limitations of telemedicine

The scope of telemedicine has progressed since the American College of Physicians 2015 recommendation on telemedicine use in primary care citing the technology's potential to improve access to care and patient satisfaction and to reduce cost.³ As the COVID-19 pandemic catalysed telemedicine's rapid expansion from niche to necessity and revolutionised the structure of primary care,^{23 40} important limitations including access, quality and adoption must be considered. While telemedicine aims to increase access to care, particularly in rural and underserved

areas, persistent inequities exist, including age, socioeconomic status, disability, level of education, smart device literacy, bandwidth and broadband access.^{4 41–43} Systematic reviews and meta-analyses reveal that telemedicine is likely comparable with in-person visits for certain services, such as mental health and rehabilitation, although is likely less effective for others, such as dermatology.⁴⁴

With regard to acute pulmonary embolism management, the current literature is sparse on using telemedicine to evaluate acute complaints in general.^{45 46} The benefits of quick access via telemedicine are pitted against the absence of a complete physical examination. For this patient, missing critical objective components include a complete set of vital signs and detailed cardiopulmonary and extremity examinations. While additional testing including a 12-lead ECG and two-view chest radiograph may partly remedy initial examination shortcomings, these are predicated on timely access to diagnostic testing. As discussed above, the emergence of reliable tools for home vital sign measurements may help broaden the capacity of telemedicine. Elements of a proposed pathway for the telemedical evaluation of possible deep vein thrombosis may be adaptable for a similar pulmonary embolism pathway.⁴⁷

Telemedicine expanded in 2020 as a convenient means of providing healthcare without increasing the risk of SARS-CoV-2 exposure. Given an eligible patient in the right care setting supported by adequate resources and infrastructure, this case raises the possibility that telemedicine may be used safely and effectively to diagnose and manage select cases of acute pulmonary embolism without the need for transfer to a higher level of care. However, one case report cannot justify wholesale a new pattern of care. Research should be undertaken to establish the safety and effectiveness of this approach and to identify patients who are best suited for telemedicine management. How well triage tools like the Pulmonary Embolism Severity Index

and the Hestia criteria might perform to identify patients with acute pulmonary embolism eligible for telemedical care without complete physical examination data is unknown. Expanding these questions beyond acute pulmonary embolism to other acute care conditions raises questions about which patients with which complaints in which care settings are eligible for comprehensive telemedical care. More research is needed to help us understand the possibilities and limitations of telemedicine in the management of patients with high-acuity complaints, including pulmonary embolism, and to ensure equitable patient access and outcomes. Preference for using telemedicine will likely remain higher after the pandemic, but the degree to which it is used or useful in a non-pandemic period is not yet clear.

Acknowledgements We thank our patient for her enthusiastic participation. We are grateful to Kelli Peterman, MPH, Diane M Carpenter, MPH, and Mary E Reed, DrPH, Kaiser Permanente Division of Research, for their helpful assistance. We appreciate the generous support of the Kaiser Permanente Northern California Graduate Medical Education Program and the Delivery Science and Physician Researcher Programs, Oakland, California.

Contributors JC, DJI, JL and DRV acquired and analysed data, drafted the initial report, contributed to its critical revision and approved the final version. DRV is accountable for the work.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent for publication Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iDs

Joshua Chang <http://orcid.org/0000-0003-3591-602X>
Dayna J Isaacs <http://orcid.org/0000-0002-3302-774X>
David R Vinson <http://orcid.org/0000-0001-6559-1858>

REFERENCES

- 1 American Medical Association. Physicians' motivations and requirements for adopting digital health: adoption and attitudinal shifts from 2016 to 2019, 2020. Available: <https://www.ama-assn.org/system/files/2020-02/ama-digital-health-study.pdf> [Accessed 15 Apr 2021].
- 2 Reed ME, Parikh R, Huang J, et al. Real-Time patient-provider video telemedicine integrated with clinical care. *N Engl J Med* 2018;379:1478–9.
- 3 Daniel H, Sulmasy LS, Health and Public Policy Committee of the American College of Physicians. Policy recommendations to guide the use of telemedicine in primary care settings: an American College of Physicians position paper. *Ann Intern Med* 2015;163:787–9.
- 4 Nieman CL, Oh ES. Connecting with older adults via telemedicine. *Ann Intern Med* 2020;173:831–2.
- 5 Reed ME, Huang J, Parikh R, et al. Patient-Provider video telemedicine integrated with clinical care: patient experiences. *Ann Intern Med* 2019;171:222–4.
- 6 Lin S, Sattler A, Smith M. Retooling primary care in the COVID-19 era. *Mayo Clin Proc* 2020;95:1831–4.
- 7 The COVID-19 Healthcare Coalition Telehealth Impact Study Work Group. Telehealth impact: patient survey analysis, 2021. Available: <https://c19hcc.org/telehealth/patient-survey-analysis/> [Accessed 15 Apr 2021].
- 8 Vinson DR, Mark DG, Chettipally UK, et al. Increasing safe outpatient management of emergency department patients with pulmonary embolism: a controlled pragmatic trial. *Ann Intern Med* 2018;169:855–65.
- 9 Aujesky D, Roy P-M, Verschuren F, et al. Outpatient versus inpatient treatment for patients with acute pulmonary embolism: an international, open-label, randomised, non-inferiority trial. *Lancet* 2011;378:41–8.
- 10 Klil-Drori AJ, Coulombe J, Suissa S, et al. Temporal trends in outpatient management of incident pulmonary embolism and associated mortality. *Thromb Res* 2018;161:111–6.
- 11 Yussen RD, Jiménez D. Outpatient management eligibility criteria for patients who have acute symptomatic pulmonary embolism. *Ann Am Thorac Soc* 2015;12:623–4.

Patient's perspective

I loved the flexibility of speaking with my PCP in a video visit. Having the visit virtually was convenient and by far the safer approach in a pandemic, many months before COVID-19 vaccinations were available. I was especially glad to avoid a trip to the emergency room — at no point did I have that 'dread' feeling that compels one to hightail it to the ER.

Learning points

- ▶ Telemedicine, including real-time patient–physician video visits, and asynchronous secure message exchange, was used to successfully coordinate outpatient primary and specialty care for a patient with acute pulmonary embolism without transfer to the emergency department or hospital.
- ▶ This virtual approach to low-risk pulmonary embolism management is a particularly convenient means of providing healthcare without increasing the risk of SARS-CoV-2 exposure or the burden on the emergency department.
- ▶ This novel approach to the management of acute pulmonary embolism has never been studied and is not ready for broad clinical application. More research is needed to help us understand the possibilities and limitations of telemedicine in the management of patients with acute pulmonary embolism and other acute care complaints and diagnoses.

- 12 Ortel TL, Neumann I, Ageno W, *et al.* American Society of hematology 2020 guidelines for management of venous thromboembolism: treatment of deep vein thrombosis and pulmonary embolism. *Blood Adv* 2020;4:4693–738.
- 13 Kearon C, Akl EA, Ornelas J, *et al.* Antithrombotic therapy for VTE disease: CHEST guideline and expert panel report. *Chest* 2016;149:315–52.
- 14 Howard LSGE, Barden S, Condliffe R, *et al.* British Thoracic Society guideline for the initial outpatient management of pulmonary embolism (PE). *Thorax* 2018;73:ii1–29.
- 15 Konstantinides SV, Meyer G, Becattini C, *et al.* 2019 ESC guidelines for the diagnosis and management of acute pulmonary embolism developed in collaboration with the European Respiratory Society (ERS). *Eur Heart J* 2020;41:543–603.
- 16 Shan J, Isaacs DJ, Bath H, *et al.* "Outpatient management" of pulmonary embolism defined in the primary literature: a narrative review (In press). *Perm J* 2021.
- 17 Vinson DR, Aujesky D, Geersing GJ, *et al.* Comprehensive outpatient management of low-risk pulmonary embolism: can primary care do this? A narrative review. *Perm J* 2020;24:163.
- 18 Vinson DR, Isaacs DJ, Johnson EJ. Managing acute pulmonary embolism in primary care in a patient declining emergency department transfer: a case report. *Eur Heart J Case Rep* 2020;4:1–4.
- 19 Isaacs DJ, Johnson EJ, Hofmann ER, *et al.* Primary care physicians comprehensively manage acute pulmonary embolism without higher-level-of-care transfer: a report of two cases. *Medicine* 2020;99:e23031.
- 20 Mehrotra A, Cherner M, Linetsky D. *The impact of the COVID-19 pandemic on outpatient care: visits return to Prepandemic levels, but not for all providers and patients.* Commonwealth Fund, 2020. <https://www.commonwealthfund.org/publications/2020/oct/impact-covid-19-pandemic-outpatient-care-visits-return-prepandemic-levels>
- 21 U.S. Department of Health & Human Services Office for Civil Rights. Notification of enforcement discretion for telehealth remote communications during the COVID-19 nationwide public health emergency, 2020. Available: <https://www.hhs.gov/hipaa/for-professionals/special-topics/emergency-preparedness/notification-enforcement-discretion-telehealth/index.html#> [Accessed 15 Apr 2021].
- 22 Centers for Medicare & Medicaid Services. Emergency Preparedness & Response Operations. Coronavirus waivers & flexibilities, 2021. Available: <https://www.cms.gov/about-cms/emergency-preparedness-response-operations/current-emergencies/coronavirus-waivers> [Accessed 15 Apr 2021].
- 23 Alexander GC, Tajanlangit M, Heyward J, *et al.* Use and content of primary care office-based vs telemedicine care visits during the COVID-19 pandemic in the US. *JAMA Netw Open* 2020;3:e2021476.
- 24 Raja AS, Greenberg JO, Qaseem A, *et al.* Evaluation of patients with suspected acute pulmonary embolism: best practice advice from the clinical guidelines Committee of the American College of Physicians. *Ann Intern Med* 2015;163:701–11.
- 25 NICE. Venous thromboembolic diseases: diagnosis, management and thrombophilia testing, 2020. Available: <https://www.nice.org.uk/guidance/ng158> [Accessed 15 Apr 2021].
- 26 Lim W, Le Gal G, Bates SM, *et al.* American Society of Hematology 2018 guidelines for management of venous thromboembolism: diagnosis of venous thromboembolism. *Blood Adv* 2018;2:3226–56.
- 27 Tran HA, Gibbs H, Merriman E, *et al.* New guidelines from the Thrombosis and Haemostasis Society of Australia and New Zealand for the diagnosis and management of venous thromboembolism. *Med J Aust* 2019;210:227–35.
- 28 Hendriksen JMT, Geersing G-J, Lucassen WAM, *et al.* Diagnostic prediction models for suspected pulmonary embolism: systematic review and independent external validation in primary care. *BMJ* 2015;351:h4438.
- 29 Hendriksen JMT, Lucassen WAM, Erkens PMG, *et al.* Ruling Out Pulmonary Embolism in Primary Care: Comparison of the Diagnostic Performance of "Gestalt" and the Wells Rule. *Ann Fam Med* 2016;14:227–34.
- 30 Walen S, Damoiseaux RA, Uil SM, *et al.* Diagnostic delay of pulmonary embolism in primary and secondary care: a retrospective cohort study. *Br J Gen Pract* 2016;66:e444–50.
- 31 Hendriksen JMT, Koster-van Ree M, Morgenstern MJ, *et al.* Clinical characteristics associated with diagnostic delay of pulmonary embolism in primary care: a retrospective observational study. *BMJ Open* 2017;7:e012789.
- 32 Rosovsky RP, Grodzin C, Channick R, *et al.* Diagnosis and treatment of pulmonary embolism during the coronavirus disease 2019 pandemic. *Chest* 2020;158:2590–601.
- 33 American College of Emergency Physicians Clinical Policies Subcommittee (Writing Committee) on Thromboembolic Disease; Wolf SJ, Hahn SA, *et al.* Clinical Policy: Critical Issues in the Evaluation and Management of Adult Patients Presenting to the Emergency Department With Suspected Acute Venous Thromboembolic Disease. *Ann Emerg Med* 2018;71:e59–109.
- 34 Qaseem A, Snow V, Barry P, *et al.* Current diagnosis of venous thromboembolism in primary care: a clinical practice guideline from the American Academy of Family Physicians and the American College of Physicians. *Ann Fam Med* 2007;5:57–62.
- 35 Zondag W, Mos ICM, Creemers-Schild D, *et al.* Outpatient treatment in patients with acute pulmonary embolism: the Hestia study. *J Thromb Haemost* 2011;9:1500–7.
- 36 Schradung WA, McCafferty B, Grove J, *et al.* Portable, consumer-grade pulse oximeters are accurate for home and medical use: implications for use in the COVID-19 pandemic and other resource-limited environments. *J Am Coll Emerg Physicians Open* 2020;1:1450–8.
- 37 Browne SH, Bernstein M, Bickler PE. Accuracy of Samsung smartphone integrated pulse oximetry meets full FDA clearance standards for clinical use. *medRxiv* 2021. doi: 10.1101/2021.02.17.21249755. [Epub ahead of print: 18 Feb 2021].
- 38 Aujesky D, Obrosky DS, Stone RA, *et al.* Derivation and validation of a prognostic model for pulmonary embolism. *Am J Respir Crit Care Med* 2005;172:1041–6.
- 39 Sylvester KW, Ting C, Lewin A, *et al.* Expanding anticoagulation management services to include direct oral anticoagulants. *J Thromb Thrombolysis* 2018;45:274–80.
- 40 Patel SY, Mehrotra A, Huskamp HA, *et al.* Trends in outpatient care delivery and telemedicine during the COVID-19 pandemic in the US. *JAMA Intern Med* 2021;181:388–91.
- 41 Huang J, Graetz I, Millman A, *et al.* Primary care telemedicine during the COVID-19 pandemic: which patients choose a video vs. telephone visit (In press). *JAMIA Open* 2021.
- 42 Reed ME, Huang J, Graetz I, *et al.* Patient characteristics associated with choosing a telemedicine visit vs office visit with the same primary care clinicians. *JAMA Netw Open* 2020;3:e205873.
- 43 Scott Kruse C, Karem P, Shifflett K, *et al.* Evaluating barriers to adopting telemedicine worldwide: a systematic review. *J Telemed Telecare* 2018;24:4–12.
- 44 Shigekawa E, Fix M, Corbett G, *et al.* The current state of telehealth evidence: a rapid review. *Health Aff* 2018;37:1975–82.
- 45 Smith KM, Hunte HE, Graber ML. Telediagnosis for acute care: implications for the quality and safety of diagnosis, 2020. Available: <https://www.ahrq.gov/sites/default/files/wysiwyg/patient-safety/reports/issue-briefs/Telediagnosis-brief2.pdf> [Accessed 15 Apr 2021].
- 46 Porter P, Brisbane J, Abeyratne U, *et al.* Diagnosing community-acquired pneumonia via a smartphone-based algorithm: a prospective cohort study in primary and acute-care consultations. *Br J Gen Pract* 2021;71:e258–65.
- 47 Iyengar KP, Jain VK, Soni M, *et al.* Virtual risk assessment pathway for deep venous thrombosis: a preliminary model. *Postgrad Med J* 2020. doi:10.1136/postgradmedj-2020-138837. [Epub ahead of print: 28 Oct 2020].

Copyright 2021 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <https://www.bmj.com/company/products-services/rights-and-licensing/permissions/>
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

Customer Service

If you have any further queries about your subscription, please contact our customer services team on +44 (0) 207111 1105 or via email at support@bmj.com.

Visit casereports.bmj.com for more articles like this and to become a Fellow