

Rapidity dependence of proton cumulants and correlation functions

Adam Bzdak^{1,*} and Volker Koch^{2,†}¹*AGH University of Science and Technology, Faculty of Physics and Applied Computer Science, 30-059 Kraków, Poland*²*Nuclear Science Division, Lawrence Berkeley National Laboratory, Berkeley, California 94720, USA*

(Received 27 July 2017; published 13 November 2017)

The dependence of multiproton correlation functions and cumulants on the acceptance in rapidity and transverse momentum is studied. We find that the preliminary data of various cumulant ratios are consistent, within errors, with rapidity and transverse momentum-independent correlation functions. However, rapidity correlations which moderately increase with rapidity separation between protons are slightly favored. We propose to further explore the rapidity dependence of multiparticle correlation functions by measuring the dependence of the integrated reduced correlation functions as a function of the size of the rapidity window.

DOI: [10.1103/PhysRevC.96.054905](https://doi.org/10.1103/PhysRevC.96.054905)

I. INTRODUCTION

One of the central goals in strong interaction research is to explore the phase diagram of QCD. Of particular interest is the search for a possible first-order phase coexistence region and its associated critical point. A significant effort in this search, experimentally as well as theoretically, is concentrating on the measurements and calculations of correlations and cumulants of conserved charges. A particular emphasis has been put on the cumulants of the baryon number [1–6], see also Refs. [7–20] (see, e.g., Ref. [21] for an overview). Interpreting these higher-order cumulants and their measurements, however, is not a straightforward exercise as discussed, e.g., in Refs. [22–35]. Also, different, although related, ideas, based on an intermittency analysis in the transverse momentum phase space have been explored [36–38].

Recently, it has been pointed out [39,40] (see also Refs. [20,41]) that it may be more instructive to study (integrated) multiparticle correlations instead of cumulants. In the limit when antiparticles can be ignored, which is the case for antiprotons at low beam energies, the integrated multiparticle correlations are linear combinations of the various cumulants and thus can be extracted easily from the measured cumulants. This has been performed on the basis of preliminary data on proton cumulants from the STAR Collaboration [42]. It was found that the systems created at low beam energies (7.7–11.5 GeV) exhibit sizable three-proton and strong four-proton correlations [40,43]. Indeed, as pointed out in Ref. [44] in order to reproduce the observed magnitude of these correlations one has, for example, to assume a strong presence of eight-nucleon (or four-proton) clusters in the system. In addition to the sheer magnitude of the correlations, the centrality and rapidity dependence of these correlations give additional insight into properties of the systems created in these collisions [40].

In this paper we will explore the rapidity and to some extent transverse momentum dependence of multiparticle correlations in more detail. One of our motivations is a

recent preliminary observation by the STAR Collaboration [45,46] regarding the rapidity dependence of the two-proton correlation function. Within the rapidity window $|y| < 0.8$, the STAR Collaboration finds that across all Brookhaven National Laboratory's Relativistic Heavy Ion Collider energies the two-proton reduced correlation function (see the definition in Sec. II) in central Au + Au collisions is increasing with the rapidity separation $y_1 - y_2$ between the two protons. The shape of the correlation function can be approximately described by

$$c_2(y_1 - y_2) = c_2^0 + \gamma_2(y_1 - y_2)^2, \quad \gamma_2 > 0, \quad (1)$$

where c_2^0 is the value at $y_1 - y_2 = 0$ and γ_2 is a positive number with $\gamma_2 \sim 2 \times 10^{-2}$ at $\sqrt{s} = 7.7$ GeV [45].¹ Taking such a correlation at face value, one would conclude that protons prefer to be separated in rapidity, or, in other words, they seem to repel each other. The shape of the correlation function is roughly energy independent, which is rather surprising since protons at, say, 7.7 GeV originate almost exclusively from the target and projectile nuclei whereas at 200 GeV the protons at midrapidity mostly are produced.

The apparent anticorrelation between two protons was first observed in e^+e^- collisions at $\sqrt{s} = 29$ GeV [48]. Recently an analogous observation was made by the ALICE Collaboration in the context of the two-baryon azimuthal correlations [49]. This measurement also found similar anticorrelations between protons and λ 's, suggesting that the observed effects are not due to the Pauli exclusion principle or electromagnetic interactions. To our knowledge, the origin of this effect remains an open question, which is important to resolve. Formation of clusters, as suggested in Ref. [44] and as expected close to a critical point and a phase transition, would naively lead to attractive correlations in rapidity (i.e., protons would prefer to have similar rapidity) and not anticorrelations. However, we should keep in mind that these correlations are in rapidity and not

¹At the recent Critical Point and Onset of Deconfinement Conference, the STAR Collaboration reported [47] that the rapidity dependence of the two-proton correlation function depends considerably on the method employed to subtract the uncorrelated single-particle contribution from the data. Thus the value for γ_2 quoted here may still change and should be taken only as a rough guidance.

*bzdak@fis.agh.edu.pl

†vkoch@lbl.gov

in configuration space. Also, one should note that this effect, which, so far, is only observed for two-particle correlations, may not be inconsistent with the negative value for the integrated two-particle correlations extracted from the cumulant measurements [40,43]. In general, the sign of an integrated multiparticle correlation also is driven by a pedestal. For example, in the case of two protons, c_2^0 in Eq. (1) may depend on the fluctuations of the volume or rather the number of wounded nucleons [23,31,44,50] and is not necessarily related to a possible repulsion or attraction in rapidity between protons.

Clearly the rapidity dependence of the proton correlations need to be studied to gain further insight into the aforementioned sizable three-proton and strong four-proton correlations observed at low energies. It is the purpose of this paper to start exploring this issue. To this end we study the dependence of the multiproton correlation functions on rapidity and, to some extent, on the transverse momentum. We show that the preliminary STAR Collaboration data [42] are consistent with constant multiproton correlation functions and slightly favor multiproton anticorrelations in rapidity. We also demonstrate that these correlations can be constrained further by measuring integrated reduced or normalized correlation functions as a function of the rapidity window Δy .

This paper is organized as follows. In the next section we introduce the notation and discuss the behavior of cumulants and correlation functions in the limits of small and large acceptances. Next we analyze the preliminary STAR Collaboration data and extract some trends about the rapidity dependence of three-proton and four-proton correlations. We also will propose a means to extract more detailed information about the multiparticle correlations. In the last section we conclude with a discussion of the essential results.

II. NOTATION AND COMMENTS

In this paper we focus on protons only, and in the following we denote the proton number by N and its deviation from the mean by $\delta N = N - \langle N \rangle$. Here $\langle N \rangle$ is the mean number of protons at a given centrality. The cumulants of the proton distribution function as measured by the STAR Collaboration are then given by

$$\begin{aligned} K_1 &\equiv \langle N \rangle; & K_2 &\equiv \langle (\delta N)^2 \rangle; & K_3 &\equiv \langle (\delta N)^3 \rangle; \\ K_4 &\equiv \langle (\delta N)^4 \rangle - 3\langle (\delta N)^2 \rangle^2. \end{aligned} \quad (2)$$

As already eluded to in the Introduction, the cumulants can be expressed in terms of the multiparticle integrated correlation functions [40], which also are known as factorial cumulants [39],

$$K_2 = \langle N \rangle + C_2, \quad (3)$$

$$K_3 = \langle N \rangle + 3C_2 + C_3, \quad (4)$$

$$K_4 = \langle N \rangle + 7C_2 + 6C_3 + C_4, \quad (5)$$

where

$$\begin{aligned} C_2 &= \int dy_1 dy_2 C_2(y_1, y_2) \\ &= \int dy_1 dy_2 [\rho_2(y_1, y_2) - \rho(y_1)\rho(y_2)], \end{aligned} \quad (6)$$

and similar for higher-order correlation functions. See, e.g., Ref. [51] for explicit definitions of the correlation functions up to the sixth order. In Eq. (6) $C_2(y_1, y_2)$ is the two-particle rapidity correlation function, $\rho_2(y_1, y_2)$ is the two-particle rapidity density, and $\rho(y)$ is the single-particle rapidity distribution. The generalization of Eqs. (3)–(5) to two species of particles can be found in the Appendix of Ref. [40]. Here and in the following y_i denotes rapidity or, in general, a set of variables under consideration ($y_i, p_{t,i}, \phi_i$).

It is a convenient and common practice to define the reduced correlation function,

$$c_n(y_1, \dots, y_n) = \frac{C_n(y_1, \dots, y_n)}{\rho(y_1) \cdots \rho(y_n)}. \quad (7)$$

The integral of the reduced correlation function over some given acceptance range, we subsequently will call, for the lack of a better term, ‘‘coupling,’’

$$c_n = \frac{C_n}{\langle N \rangle^n} = \frac{\int \rho(y_1) \cdots \rho(y_n) c_n(y_1, \dots, y_n) dy_1 \cdots dy_n}{\int \rho(y_1) \cdots \rho(y_n) dy_1 \cdots dy_n}. \quad (8)$$

The cumulants K_n then may be expressed in terms of the couplings c_n ,

$$K_2 = \langle N \rangle + \langle N \rangle^2 c_2, \quad (9)$$

$$K_3 = \langle N \rangle + 3\langle N \rangle^2 c_2 + \langle N \rangle^3 c_3, \quad (10)$$

$$K_4 = \langle N \rangle + 7\langle N \rangle^2 c_2 + 6\langle N \rangle^3 c_3 + \langle N \rangle^4 c_4. \quad (11)$$

Of course, mathematically, the cumulants $K_1 = \langle N \rangle$, K_2 , K_3 , and K_4 carry exactly the same information as $[C_2, C_3, C_4]$ or $[c_2, c_3, c_4]$. However, as already discussed in Ref. [40], studying cumulants may not be the best way to extract information about the dynamics of the system since: (i) Cumulants mix the correlation functions of different orders, and (ii) they might be dominated by a trivial term $\langle N \rangle$ even in the presence of interesting dynamics.

One such example where the trivial term $\langle N \rangle$ dominates and thus hides the interesting physics is the limit of small acceptance as we will discuss next.

A. Effective Poisson limit

Before we discuss the rapidity and transverse momentum dependence of the various cumulants and correlations, let us briefly remind ourselves what happens if one considers the limit of small or vanishing acceptance. Here, we will restrict ourselves to correlations in rapidity, however, our arguments will be general and apply to any variables. Suppose that particles are measured in a rapidity interval $y_0 \leq y \leq y_0 + \Delta y$ and that $\Delta y \rightarrow 0$. Let us first consider two-particle correlations. For sufficiently small Δy any reasonable correlation function $c_2(y_1, y_2)$ may be approximated by a constant.² As

²For the extreme case of $c_2(y_1, y_2) \sim \delta(y_1 - y_2)$, c_2 , given by Eq. (8), depends on the acceptance window even for very small rapidity intervals, and our argument does not apply. However, a Dirac- δ correlation function is of no interest in any practical situation.

a consequence, for sufficiently small Δy , the coupling c_2 is independent of Δy as can be seen from Eq. (8). In other words, suppose that $c_2(y_1, y_2) \simeq c_2^0$ for very small Δy , then

$$c_2 = \frac{\int_{\Delta y} \rho(y_1)\rho(y_2)c_2(y_1, y_2)dy_1dy_2}{\int_{\Delta y} \rho(y_1)\rho(y_2)dy_1dy_2} \simeq c_2^0. \quad (12)$$

We emphasize that c_2^0 may assume any value. However, whatever the value of c_2^0 , in the limit of $\Delta y \rightarrow 0$, we have $\langle N \rangle \rightarrow 0$ and $K_2 \simeq \langle N \rangle$ [see Eq. (9)]. Exactly the same argument holds for any K_n , and we obtain $K_n \simeq \langle N \rangle$ and consequently all cumulant ratios equal to unity $K_n/K_m \simeq 1$.

Therefore, even in the presence of sizable correlations, their effects on the cumulants are suppressed for small acceptance. Actually, as can be seen from Eqs. (9)–(11), it is the number of particles which determines if the cumulants are dominated by $\langle N \rangle$ and, thus, their ratios are close to unity. For example, if $\langle N \rangle^4 c_4 \ll \langle N \rangle$, the fourth-order cumulant K_4 is practically not sensitive to four-proton correlations even if c_4 is different from zero and may carry some interesting information. Therefore, even for large acceptance, the cumulants are close to the Poisson limit if one is dealing with rare particles. This may very well be the reason that for low energies the STAR Collaboration observes a cumulant ratio of $K_4/K_2 \simeq 1$ for antiprotons and it would be interesting to measure the couplings c_n for antiprotons in order to see if antiprotons exhibit the same correlations as protons at low energies.

Clearly measuring cumulants and looking for the deviation from the Poisson limit is not the most optimal way to extract possible nontrivial correlations resulting from criticality, etc. Instead, one either should directly measure the differential multiparticle correlation [Eq. (7)] or, at the very least, extract the couplings c_n Eq. (8). Their dependence on the acceptance does reflect a change in physics and is not simply a consequence of a change in the number of particles.³

After having investigated the case of small acceptance let us next turn to the opposite limit of (nearly) full acceptance.

B. Full acceptance

Let us next study what happens in the situation when all baryons, including the spectators, are detected. In this case (again, we consider low energies and neglect antibaryons) $N = \langle N \rangle = B$, where B is the total baryon number of the entire system. Therefore, $\delta N = 0$ and obviously $K_n = 0$ for $n \geq 2$. Using Eqs. (3)–(5) and (9)–(11) we obtain

$$C_2 = -B, \quad C_3 = 2B, \quad C_4 = -6B, \quad (13)$$

and

$$c_2 = -\frac{1}{B}, \quad c_3 = \frac{2}{B^2}, \quad c_4 = -\frac{6}{B^3}. \quad (14)$$

We note that this is a general result and it is insensitive to the presence of any dynamics other than global baryon number conservation.

³An additional advantage of the couplings is that they are independent of the efficiency of the detector as long as the efficiency follows a binomial distribution and is phase-space independent [24,30,41].

Finally let us note that $K_3/K_2 \rightarrow -1$ and $K_4/K_2 \rightarrow 1$ when we approach the limit of full acceptance. To see this let us consider a region in phase space, denoted by (a) , and the remaining phase space, or complement, which we denote by (b) . Since the baryon number is conserved, having $N_{(a)}$ baryons in region (a) implies $N_{(b)} = B - N_{(a)}$ baryons in the complement (b) . Since $\delta B = 0$ we have

$$\delta N_{(b)} = \delta(B - N_{(a)}) = -\delta N_{(a)}, \quad (15)$$

and consequently,

$$\begin{aligned} K_{n,(a)} &= K_{n,(b)}, \quad n = 2, 4, 6, \dots, \\ K_{n,(a)} &= -K_{n,(b)}, \quad n = 3, 5, 7, \dots \end{aligned} \quad (16)$$

Here $K_{n,(a)}$ is the cumulant measured in region (a) , and $K_{n,(b)}$ is the cumulant in a remaining part of the full phase-space (b) . This is a rather nontrivial and general consequence of baryon conservation. A more rigorous derivation is presented in the Appendix.

In the previous subsection we argued that for very small acceptance the cumulant ratio goes to 1 and thus the cumulant ratio for the full acceptance goes to -1 for K_3/K_2 and to 1 for K_4/K_2 . The integrated correlation functions and the couplings, on the other hand, do not show such a symmetry between a given region of phase space and its complement. This is shown in detail in the Appendix but can already be inferred from the fact that in the limit of full acceptance the couplings are determined entirely by the total baryon number B . In the limit of vanishing acceptance, however, other physics also affects the value of the couplings as discussed in Sec. II A.

Having discussed the limits of small and full acceptances we now turn to the rapidity dependence of the cumulants and correlation functions.

III. RESULTS

In this section we discuss in detail the rapidity and, to some extent, the transverse momentum dependence of multiproton cumulants and correlation functions. First, we will explore the limit of rapidity and transverse momentum-independent correlations. Next we will discuss to which extent the present preliminary STAR Collaboration data allow us to set limits on the rapidity dependence of the underlying correlations.

A. Constant correlation

Let us start with the simplest assumption, namely, that the reduced correlation function does not depend on rapidity and transverse momentum, i.e.,

$$c_n(y_1, p_{t1}, \dots, y_n, p_{tn}) = \text{const} = c_n^0. \quad (17)$$

This rather extreme assumption, however, is, as we will show below, consistent with the preliminary STAR Collaboration data at 7.7 GeV (see also Ref. [40]). In addition, in this case




FIG. 1. The cumulant ratio K_4/K_2 in central 0–5% Au + Au collisions at $\sqrt{s} = 7.7$ GeV as a function of the number of measured protons $\langle N \rangle$ for different acceptance windows in rapidity and transverse momentum (in units of GeV). For all data points $p_t > 0.4$ GeV. The black solid line represents a prediction based on a constant correlation function, see Eq. (17). The shaded band is driven mostly by the large experimental uncertainty of K_4 . Based on the preliminary STAR Collaboration data [42].

the couplings c_n do not depend on rapidity and transverse momentum either as can be seen from Eq. (8),

$$c_n = c_n^0. \quad (18)$$

The multiparticle integrated correlation functions $C_n = \langle N \rangle^n c_n$ and cumulants K_n , in turn, depend on the acceptance only through their dependence on the number of protons $\langle N \rangle$, see Eqs. (9)–(11). Therefore, in Fig. 1 we plot K_4/K_2 as measured by the STAR Collaboration as a function of $\langle N \rangle$ for different rapidity and transverse momentum intervals.

The black solid line in Fig. 1 represents a prediction based on a constant correlation function. In this calculation we have three unknown parameters c_2^0 , c_3^0 , and c_4^0 . Since these numbers do not depend on acceptance, we determine them from the preliminary data for $|y| < 0.5$ ($\Delta y = 1$) and $0.4 < p_t < 2$ GeV, that is, from the maximal acceptance currently available. Here we use Eqs. (9)–(11) and the values for $\langle N \rangle$, K_2 , K_3 , and K_4 shown in Ref. [42].⁴ To determine $\langle N \rangle$ at a given acceptance region we assume the single-proton rapidity distribution to be flat as a function of rapidity, i.e., $\langle N \rangle = \langle N_{\Delta y=1} \rangle \Delta y$, and, for the transverse momentum single-proton distribution, we take $\rho(p_t) \sim p_t \exp(-m_t/T)$ with $T = 0.27$ GeV and $m_t = (m^2 + p_t^2)^{1/2}$ with $m = 0.94$ GeV. Both these assumptions are well supported by experimental data [52,53]. Having c_n^0 , we can predict the cumulants or the correlation functions for any acceptance charac-

terized by $\langle N \rangle$ whether in transverse momentum or in rapidity.⁵

Interestingly we find that, except for one point at $|y| < 0.5$ and $0.4 < p_t < 1.2$ GeV, all the points follow within the admittedly large experimental error bars one universal curve consistent with a constant correlation function. The fact that the rapidity dependence of the cumulant ratio K_4/K_2 is consistent with long-range rapidity correlations already has been found in Ref. [40]. That the transverse momentum dependence is also consistent with long-range correlations is new. If correct, it would, for example, imply that the cumulant ratio K_4/K_2 has roughly the same value (close to unity) for a transverse momentum range of $0.8 \text{ GeV} < p_t < 2 \text{ GeV}$ as the value for the range of $0.4 \text{ GeV} < p_t < 0.8 \text{ GeV}$ since, in both p_t windows, $\langle N \rangle$ is approximately the same. The result for the p_t range of $0.4 \text{ GeV} < p_t < 0.8 \text{ GeV}$ has been published by the STAR Collaboration in Ref. [5].

Of course, the error bars in the preliminary STAR Collaboration data are rather sizable and, therefore, a mild dependence of the correlation function on rapidity (and transverse momentum) cannot be ruled out. In addition, as already mentioned in the Introduction, the preliminary, explicit measurement of the two-proton correlation function [45,46] does exhibit an increase with increasing rapidity difference of a proton pair $y_1 - y_2$. To explore this further we next will allow for some mild rapidity dependence of the correlation function.

B. Rapidity-dependent correlation

In the previous subsection we demonstrated that the STAR Collaboration data for K_4/K_2 at 7.7 GeV are consistent with a constant multiproton correlation function. Here we study how sensitive the cumulant ratios and correlations are to a certain (weak) rapidity dependence. To this end we consider the leading correction to a constant correlation function, which should be even in $y_i - y_k$. Thus we explore the following Ansätze for the reduced correlation functions,

$$\begin{aligned} c_2(y_1, y_2) &= c_2^0 + \gamma_2 (y_1 - y_2)^2, \\ c_3(y_1, y_2, y_3) &= c_3^0 + \gamma_3 \frac{1}{3} [(y_1 - y_2)^2 + (y_1 - y_3)^2 \\ &\quad + (y_2 - y_3)^2], \\ c_4(y_1, y_2, y_3, y_4) &= c_4^0 + \gamma_4 \frac{1}{6} [(y_1 - y_2)^2 + (y_1 - y_3)^2 \\ &\quad + (y_1 - y_4)^2 + (y_2 - y_3)^2 \\ &\quad + (y_2 - y_4)^2 + (y_3 - y_4)^2], \end{aligned} \quad (19)$$

where γ_n measures the deviation from $c_n(y_1, \dots, y_n) = \text{const}$. Note that we have constructed the correlation function such that positive values of γ_n result in growing correlations with rapidity separation between particles. We further note that the above form for the two-proton reduced correlation function $c_2(y_1, y_2)$ is supported by the preliminary STAR Collaboration data [45,46] where $\gamma_2 > 0$, that is, two protons do not want

⁴We determine c_n^0 from the proton cumulants but compare to y and p_t dependences of the net-proton cumulants, which are the only data currently available. Although at 7.7 GeV the number of antiprotons is practically negligible, it results in a slight disagreement of the black solid line with the blue star in Fig. 1.

⁵Based on the preliminary STAR Collaboration data for the cumulants [42] we obtain $c_2^0 \approx -1.1 \times 10^{-3}$, $c_3^0 \approx -1.7 \times 10^{-4}$, and $c_4^0 \approx 7.3 \times 10^{-5}$.

to occupy the same rapidity. Our simple formulas for c_3 and c_4 are not supported by any known data, however, we believe they should serve as a reasonable representation for the correlation if the distance in rapidity between protons is not too long. Within the region of validity of our simple Ansatz, the coefficients γ_n have a clear physical interpretation, and here we will constrain their values or at least their signs. To this end we will use the preliminary STAR Collaboration data for K_3/K_2 and K_4/K_2 . Although, as already pointed out, the rapidity dependence of these cumulant ratios is consistent with constant correlations, we will see that the data allow for excluding certain values for γ_n and possibly even determine their sign.

Taking the above relations and integrating in Eq. (8) over $|y_i| < \Delta y/2$ we obtain for the couplings,

$$c_n(\Delta y) = \frac{C_n}{\langle N \rangle^n} = c_n^0 + \gamma_n \frac{1}{6} (\Delta y)^2. \quad (20)$$

The couplings $c_n(\Delta y)$, which depend on the region of acceptance Δy ($|y_i| < \Delta y/2$), should not be confused with the reduced correlation function $c_n(y_1, \dots, y_n)$, which depends on the rapidities of the individual particles. As before, for a given γ_n the constant term c_n^0 is extracted from the STAR Collaboration data at $\Delta y = 1$ ($|y| < 0.5$) and $0.4 < p_t < 2$ GeV. Consequently, c_n^0 will depend on the choice of γ_n .

In Fig. 2 we show K_3/K_2 for different values of γ_3 in panel (a) and γ_2 in panel (b). We observe that, as already discussed before, the preliminary STAR Collaboration data are consistent with a constant correlation function in rapidity ($\gamma_2 = \gamma_3 = 0$). However, a small positive value of $\gamma_2 \sim 10^{-2}$ or $\gamma_3 \sim 10^{-3}$ would actually improve the agreement slightly. The negative values for γ_2 and γ_3 , on the other hand, appear to be disfavored so are large positive values. The same is true for the comparison with the K_4/K_2 cumulant ratio, which we show in Fig. 3. Again, the data are consistent with constant rapidity correlation functions or perhaps slightly positive values for γ_2 , γ_3 , or γ_4 , whereas negative values for γ_n seem to be disfavored.⁶

Also, the overall picture of slightly “repulsive” corrections to the constant correlation functions, i.e., $\gamma_n \geq 0$ is consistent with the preliminary STAR Collaboration data on the two-proton rapidity correlation function, which, as discussed in the Introduction, indicates a peculiar repulsion between protons in rapidity. As these new STAR Collaboration measurements only address two-proton correlations, the most direct test would be a comparison of the rapidity dependence of the second-order cumulant or integrated correlation. This is shown in Fig. 4. Unfortunately, at present there are no data available for rapidity intervals other than $\Delta y = 1$, and since this point is used for the determination of the overall constant c_2^0 , no constraint can be made at this time. However, we wish to emphasize the strong dependence compared to the size of the error bar. Indeed, the increase in the correlation exhibited in the preliminary STAR Collaboration data for the differential correlation functions [45,46] is consistent with $\gamma_2 \sim 2 \times 10^{-2}$,




FIG. 2. The cumulant ratio K_3/K_2 in central Au + Au collisions at $\sqrt{s} = 7.7$ GeV as a function of the rapidity acceptance Δy , $|y| < \Delta y/2$ for (a) $\gamma_2 = 0$ and different values of γ_3 from Eq. (19) and (b) $\gamma_3 = 0$ and different values of γ_2 . Based on the preliminary STAR Collaboration data [42].

which would correspond to the red dashed curve in Fig. 4. Given the size of the error bar at $\Delta y = 1$, it should be possible to discriminate from a constant correlation function, shown by the green solid line. Needless to say, such a measurement of the rapidity dependence of K_2/K_1 would be very valuable to ensure the consistency of the cumulant measurement with that of the differential correlation function.⁷

Of course it would be even more valuable to have information about the differential three-particle and four-particle correlation functions. Therefore, we propose, as a first step, to measure the rapidity dependence of the couplings $c_n(\Delta y)$. This will allow for a direct determination of the coefficients γ_n as we demonstrate in Fig. 5 where we plot $c_n(\Delta y)/c_n^0 - 1$ for $\gamma_2 = 10^{-2}$, $\gamma_3 = 10^{-3}$, and $\gamma_4 = 2 \times 10^{-4}$. We note that $c_n(\Delta y)$ is rather sensitive to γ_n .

⁶Specifically we find the following values for c_n^0 and γ_n for the blue lines in Figs. 2 and 3: $\gamma_2 = 10^{-2}$, $c_2^0 \approx -2.8 \times 10^{-3}$, $\gamma_3 = 10^{-3}$, $c_3^0 \approx -3.4 \times 10^{-4}$, and $\gamma_4 = 2 \times 10^{-4}$, $c_4^0 \approx 3.9 \times 10^{-5}$.

⁷We note that the preliminary measurements of $c_2(y_1, y_2)$ and K_n use different centrality selections, which do affect the values of c_n^0 and possibly γ_n . Therefore, a direct comparison of the values for γ_2 needs to be performed with some care.




FIG. 3. The cumulant ratio K_4/K_2 in central Au + Au collisions at $\sqrt{s} = 7.7$ GeV as a function of the rapidity acceptance Δy , $|y| < \Delta y/2$ for (a) different values of γ_4 , (b) γ_3 , and (c) γ_2 from Eq. (19). Based on the preliminary STAR Collaboration data [42].

In principle it would also be interesting to measure $c_n(\Delta y)$ for higher n , such as $n = 5$ and 6 . In this case,

$$c_5(y_1, \dots, y_5) = c_5^0 + \gamma_5 \frac{1}{10} \sum_{i,k=1; i < k}^5 (y_i - y_k)^2,$$

$$c_6(y_1, \dots, y_6) = c_6^0 + \gamma_6 \frac{1}{15} \sum_{i,k=1; i < k}^6 (y_i - y_k)^2, \quad (21)$$

and $c_n(\Delta y)$ is given by Eq. (20).




FIG. 4. The cumulant ratio K_2/K_1 in central Au + Au collisions at $\sqrt{s} = 7.7$ GeV as a function of the rapidity acceptance Δy , $|y| < \Delta y/2$ for different values of γ_2 . Based on the preliminary STAR Collaboration data [42].

IV. DISCUSSION AND CONCLUSIONS

Before we conclude let us discuss the main findings of this paper.

- (1) The preliminary data for the proton cumulant ratio K_4/K_2 obtained by the STAR Collaboration at $\sqrt{s} = 7.7$ GeV are consistent with long-range correlations in




FIG. 5. The ratio of the couplings $c_n(\Delta y)/c_n^0 - 1$, see Eqs. (8) and (20) for $\gamma_2 = 10^{-2}$, $\gamma_3 = 10^{-3}$ and $\gamma_4 = 2 \times 10^{-4}$. Δy denotes the size of the rapidity window $|y| < \Delta y/2$, which the reduced correlation functions are integrated over.

both rapidity and transverse momentum. As a result the cumulants effectively depend only on the number of protons (N) in the acceptance. Therefore, we predict that new measurements with increased acceptance will lead to even larger values for K_4/K_2 . Naturally this increase will be limited eventually by global charge conservation as discussed in Ref. [22], and the Ansatz for the correlation function Eq. (17) will have its limitation for large Δy . Consequently our present prediction for large $\Delta y > 1$ needs to be taken with a grain of salt.

- (2) Allowing for small deviation from a constant value we find that a slightly repulsive rapidity dependence is favored by the data. By repulsive we mean that the correlation function increases with increasing rapidity separation between protons. Or in other words, we find that $\gamma_n > 0$ in Eq. (19) is favored. Perhaps this may be the first evidence for repulsive three-proton and four-proton correlations.
- (3) Clearly, as demonstrated in Fig. 5, a measurement of the couplings as a function of the rapidity and transverse momentum windows would be very valuable to shed more light on the ranges and detailed shapes of the correlation functions.
- (4) Finally we want to reiterate that the fact that cumulant ratios for small acceptance or, more precisely for a small number of particles, are close to unity does not necessarily imply the absence of correlations. This is demonstrated in Fig. 1 where we actually assume a *constant* correlation. In addition, this may also be the reason that antiprotons show a cumulant ratio of $K_4/K_2 \simeq 1$ at low energies, whereas the protons show a significant deviation from unity.

We further demonstrated that global baryon conservation fully determines the cumulant ratios, integrated correlation functions, and couplings close to the full acceptance regardless of any additional dynamics. In addition we showed that, as a result of baryon number conservation, the cumulants in a given phase-space window and their complements are closely related, see Eq. (16) and the Appendix.

To summarize, we have studied the rapidity dependence of cumulants, integrated correlation functions, and couplings based on the presently available preliminary STAR Collaboration data [42]. Although we found that, within the present experimental errors the data are consistent with rapidity independent correlations, a slightly repulsive component seems to be favored. This would be consistent with the preliminary measurement of two-particle differential proton correlations by the STAR Collaboration [45,46]. To gain further insight, in particular, into the three-proton and four-proton correlations, we proposed to measure the dependence of the couplings as a function of the rapidity window.

ACKNOWLEDGMENTS

We thank the STAR Collaboration for providing us with their preliminary data. We further thank A. Bialas and W.

Llope for stimulating discussions. A.B. was supported by the Ministry of Science and Higher Education (MNiSW) and by the National Science Centre, Grant No. DEC-2014/15/B/ST2/00175 and, in part, by Grant No. DEC-2013/09/B/ST2/00497. V.K. was supported by the U.S. Department of Energy, Office of Science, Office of Nuclear Physics, under Contract No. DE-AC02-05CH11231.

APPENDIX: FULL ACCEPTANCE

Suppose we divide the full phase space into the two not necessarily equal-sized regions denoted by the subscripts (a) and (b). Let $P_{(a)}(N_{(a)})$ be the probability to observe $N_{(a)}$ baryons in the phase-space region (a). The probability to have $N_{(b)}$ baryons in the remaining part of the entire phase-space $P_{(b)}(N_{(b)})$ is given by $P_{(b)}(N_{(b)}) = P_{(a)}(N_{(a)}) = P_{(a)}(B - N_{(b)})$ since $N_{(a)} = B - N_{(b)}$, where B is the conserved number of baryons. Here we assume that we can ignore antibaryons. The cumulant generating function for the phase-space region (a), $h_{(a)}(t)$ is given by

$$\begin{aligned} h_{(a)}(t) &= \log \left[\sum_{N_{(a)}} P_{(a)}(N_{(a)}) e^{N_{(a)}t} \right] \\ &= \log \left[\sum_{N_{(b)}} P_{(a)}(B - N_{(b)}) e^{(B - N_{(b)})t} \right] \\ &= \log \left[\sum_{N_{(b)}} P_{(b)}(N_{(b)}) e^{(B - N_{(b)})t} \right] \\ &= h_{(b)}(-t) + Bt, \end{aligned} \quad (\text{A1})$$

where $h_{(b)}(t)$ is the cumulant generating function for phase-space region (b). The cumulants in the two regions (a) and (b) are given by the derivatives at $t = 0$,

$$K_{n,(a)} = \left. \frac{d^n}{dt^n} h_{(a)}(t) \right|_{t=0}, \quad K_{n,(b)} = \left. \frac{d^n}{dt^n} h_{(b)}(t) \right|_{t=0}. \quad (\text{A2})$$

Thus we get for $n = 1$,

$$\langle N_{(a)} \rangle = K_{1,(a)} = B - K_{1,(b)} = B - \langle N_{(b)} \rangle, \quad (\text{A3})$$

and for $n \geq 2$,

$$K_{n,(a)} = (-1)^n K_{n,(b)}. \quad (\text{A4})$$

Given this relation between the cumulants of the two regions and using Eqs. (2)–(5) we also can find the relation between the integrated correlation functions $C_{n,(a)}$ and $C_{n,(b)}$ in regions (a) and (b), respectively,

$$\begin{aligned} C_{2,(a)} &= -B + 2\langle N_{(b)} \rangle + C_{2,(b)}, \\ C_{3,(a)} &= 2B - 6\langle N_{(b)} \rangle - 6C_{2,(b)} - C_{3,(b)}, \\ C_{4,(a)} &= -6B + 24\langle N_{(b)} \rangle + 36C_{2,(b)} + 12C_{3,(b)} + C_{4,(b)}. \end{aligned} \quad (\text{A5})$$

Clearly, the integrated correlation functions do not show any symmetry between the two complementary regions of the phase space. The same is also true for the couplings c_n . In the limit where $\langle N_{(a)} \rangle \rightarrow B$ and thus $\langle N_{(b)} \rangle \rightarrow 0$ we find,

following the above equations, that $C_{2,(a)} \rightarrow -B$, $C_{3,(a)} \rightarrow 2B$, and $C_{4,(a)} \rightarrow -6B$. In this case, the couplings become $c_{2,(a)} \rightarrow -\frac{1}{B}$, $c_{3,(a)} \rightarrow \frac{2}{B^2}$, and $c_{4,(a)} \rightarrow -\frac{6}{B^3}$ and again are determined entirely by the total baryon number B . For the

complementary region (b), on the other hand, we have the limit of $\langle N_b \rangle \rightarrow 0$ in which case as discussed in Sec. II A, dynamics beyond baryon number conservation also affects the couplings.

-
- [1] M. A. Stephanov, K. Rajagopal, and E. V. Shuryak, *Phys. Rev. Lett.* **81**, 4816 (1998).
- [2] M. A. Stephanov, *Phys. Rev. Lett.* **102**, 032301 (2009).
- [3] V. Skokov, B. Friman, and K. Redlich, *Phys. Rev. C* **83**, 054904 (2011).
- [4] B. Friman, F. Karsch, K. Redlich, and V. Skokov, *Eur. Phys. J. C* **71**, 1694 (2011).
- [5] L. Adamczyk *et al.* (STAR Collaboration), *Phys. Rev. Lett.* **112**, 032302 (2014).
- [6] L. Adamczyk *et al.* (STAR Collaboration), *Phys. Rev. Lett.* **113**, 092301 (2014).
- [7] S. Borsanyi, Z. Fodor, S. D. Katz, S. Krieg, C. Ratti *et al.*, *J. High Energy Phys.* **01** (2012) 138.
- [8] A. Bazavov *et al.* (HotQCD Collaboration), *Phys. Rev. D* **86**, 034509 (2012).
- [9] R. Bellwied, S. Borsanyi, Z. Fodor, S. D. Katz, A. Pasztor, C. Ratti, and K. K. Szabo, *Phys. Rev. D* **92**, 114505 (2015).
- [10] R. V. Gavai and S. Gupta, *Phys. Lett. B* **696**, 459 (2011).
- [11] M. Kitazawa, M. Asakawa, and H. Ono, *Phys. Lett. B* **728**, 386 (2014).
- [12] M. Asakawa and M. Kitazawa, *Prog. Part. Nucl. Phys.* **90**, 299 (2016).
- [13] A. Mukherjee, J. Steinheimer, and S. Schramm, *Phys. Rev. C* **96**, 025205 (2017).
- [14] C. Herold, M. Nahrgang, Y. Yan, and C. Kobdaj, *Phys. Rev. C* **93**, 021902(R) (2016).
- [15] A. Chatterjee, S. Chatterjee, T. K. Nayak, and N. R. Sahoo, *J. Phys. G* **43**, 125103 (2016).
- [16] R. A. Lacey, P. Liu, N. Magdy, B. Schweid, and N. N. Ajitanand, [arXiv:1606.08071](https://arxiv.org/abs/1606.08071).
- [17] M. Hippert and E. S. Fraga, *Phys. Rev. D* **96**, 034011 (2017).
- [18] R. Rougemont, R. Critelli, J. Noronha-Hostler, J. Noronha, and C. Ratti, *Phys. Rev. D* **96**, 014032 (2017).
- [19] G. A. Almási, B. Friman, and K. Redlich, *Phys. Rev. D* **96**, 014027 (2017).
- [20] S. He and X. Luo, *Phys. Lett. B* **774**, 623 (2017).
- [21] V. Koch, in *Relativistic Heavy Ion Physics*, edited by R. Stock, Landolt-Börnstein New Series I Vol. 23 (Springer, Heidelberg, 2010), pp. 626–652.
- [22] A. Bzdak, V. Koch, and V. Skokov, *Phys. Rev. C* **87**, 014901 (2013).
- [23] V. Skokov, B. Friman, and K. Redlich, *Phys. Rev. C* **88**, 034911 (2013).
- [24] A. Bzdak and V. Koch, *Phys. Rev. C* **86**, 044904 (2012).
- [25] A. Bzdak and V. Koch, *Phys. Rev. C* **91**, 027901 (2015).
- [26] X. Luo, *Phys. Rev. C* **91**, 034907 (2015).
- [27] T. Nonaka, T. Sugiura, S. Esumi, H. Masui, and X. Luo, *Phys. Rev. C* **94**, 034909 (2016).
- [28] G. D. Westfall, *Phys. Rev. C* **92**, 024902 (2015).
- [29] Z. Fecková, J. Steinheimer, B. Tomášik, and M. Bleicher, *Phys. Rev. C* **92**, 064908 (2015).
- [30] A. Bzdak, R. Holzmann, and V. Koch, *Phys. Rev. C* **94**, 064907 (2016).
- [31] P. Braun-Munzinger, A. Rustamov, and J. Stachel, *Nucl. Phys. A* **960**, 114 (2017).
- [32] M. Bluhm, M. Nahrgang, S. A. Bass, and T. Schäfer, *Eur. Phys. J. C* **77**, 210 (2017).
- [33] H.-J. Xu, *Phys. Lett. B* **765**, 188 (2017).
- [34] T. Nonaka, M. Kitazawa, and S. Esumi, *Phys. Rev. C* **95**, 064912 (2017).
- [35] P. Garg and D. K. Mishra, *Phys. Rev. C* **96**, 044908 (2017).
- [36] T. Anticic *et al.* (NA49 Collaboration), *Phys. Rev. C* **81**, 064907 (2010).
- [37] T. Anticic *et al.* (NA49 Collaboration), *Eur. Phys. J. C* **75**, 587 (2015).
- [38] N. G. Antoniou, N. Davis, and F. K. Diakonou, *Phys. Rev. C* **93**, 014908 (2016).
- [39] B. Ling and M. A. Stephanov, *Phys. Rev. C* **93**, 034915 (2016).
- [40] A. Bzdak, V. Koch, and N. Strodthoff, *Phys. Rev. C* **95**, 054906 (2017).
- [41] M. Kitazawa and X. Luo, *Phys. Rev. C* **96**, 024910 (2017).
- [42] X. Luo (STAR Collaboration), *Proceedings, 9th International Workshop on Critical Point and Onset of Deconfinement (CPOD 2014): Bielefeld, Germany, 2014* (SISSA, Trieste, Italy, 2015).
- [43] X. Luo and N. Xu, *Nucl. Sci. Tech.* **28**, 112 (2017).
- [44] A. Bzdak, V. Koch, and V. Skokov, *Eur. Phys. J. C* **77**, 288 (2017).
- [45] S. Jowzaee (STAR Collaboration), in *the XXVI International Conference on Ultrarelativistic Heavy-Ion Collisions* (Quark Matter 2017), Chicago, 2017, edited by O. Evdokimov, R. Betts, and U. Heinz (CERN, Meyrin, Switzerland, 2017).
- [46] A. Lipiec (STAR Collaboration), *XII Workshop on Particle Correlations and Femtoscopy, Amsterdam, 2017* (Nikhef, Amsterdam, 2017).
- [47] W. Llope (STAR Collaboration), *the Conference on Critical Point and Onset of Deconfinement (CPOD 2017), Stony Brook, New York, 2017* (Brookhaven National Laboratory, Upton, NY, 2017).
- [48] H. Aihara *et al.* (TPC/Two Gamma Collaboration), *Phys. Rev. Lett.* **57**, 3140 (1986).
- [49] J. Adam *et al.* (ALICE Collaboration), *Eur. Phys. J. C* **77**, 569 (2017).
- [50] A. Bialas, M. Bleszynski, and W. Czyz, *Nucl. Phys. B* **111**, 461 (1976).
- [51] A. Bzdak and P. Bozek, *Phys. Rev. C* **93**, 024903 (2016).
- [52] L. Adamczyk *et al.* (STAR Collaboration), *Phys. Rev. C* **96**, 044904 (2017).
- [53] T. Anticic *et al.* (NA49 Collaboration), *Phys. Rev. C* **83**, 014901 (2011).