



Lawrence Berkeley Laboratory

UNIVERSITY OF CALIFORNIA

Materials Sciences Division

To be presented at Quantum Electronics and Laser Science, Anaheim, CA,
June 2-7, 1996, and to be published in the Proceedings

Room Temperature Observation of Quantum Jumps of Single Molecule into Dark States

T. Ha, Th. Enderle, D.F. Ogletree, P.R. Selvin, S. Weiss,
and D.S. Chemla

November 1995



REFERENCE COPY |
Does Not |
Circulate |

Bldg. 50 Library.

Copy 1

LBL-38015

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

LBL-38015
UC-404

**ROOM TEMPERATURE OBSERVATION OF QUANTUM JUMPS OF SINGLE
MOLECULE INTO DARK STATES**

T. Ha, Th. Enderle, D. F. Ogletree, P.R. Selvin, S. Weiss, and D.S. Chemla

Molecular Design Institute
Materials Sciences Division
Lawrence Berkeley National Laboratory
University of California
Berkeley, CA 94720

November 1995

This work was supported by the Office of Energy Research, Office of Basic Energy Sciences, Division of Materials Sciences of the U.S. Department of Energy, Contract No. DE-AC03-76SF00098.

Room temperature observation of quantum jumps of single molecule into dark states

*T. Ha, Th. Enderle, D.F. Ogletree,
P.R. Selvin, S. Weiss and D.S. Chemla*

Molecular Design Institute, Materials Sciences Division

Lawrence Berkeley National Laboratory

1 Cyclotron Road, Berkeley, CA 94720

TEL: 510-486-5202

FAX: 510-486-5530

e-mail: sweiss@ux5.lbl.gov

Abstract:

Fluctuations in the room temperature emission rate from single dye molecules which are excited with the near field scanning optical microscope reveal long (~ seconds) and short (~ milliseconds) lived dark states.

Room temperature observation of quantum jumps of single molecule into dark states

*T. Ha, Th. Enderle, D.F. Ogletree,
P.R. Selvin, S. Weiss and D.S. Chemla*

Molecular Design Institute, Materials Sciences Division
Lawrence Berkeley National Laboratory
1 Cyclotron Road, Berkeley, CA 94720
TEL: 510-486-5202
FAX: 510-486-5530
e-mail: sweiss@ux5.lbl.gov

Summary:

We monitored the fluorescence emission rate of single dye molecules on dry surface with the near field scanning optical microscope (NSOM) [1]. A Texas Red fluorophore was covalently attached to a short strand of a DNA molecule and bound to an amino-propyl-silanized glass coverslip [2]. After imaging, the NSOM probe was stationed over a single molecule and its emission rate was recorded as function of time until its photodestruction. Counts were integrated for 5 - 20 ms and typical count rates were 5 - 30 kHz. A typical emission time trace is shown in Fig. 1.

In addition to the irreversible photodestruction event at 26s, reversible jumps to background level around 4s and 9s are noticeable. These long lived dark states last few seconds. The two curves are polarization resolved emission time traces recorded simultaneously with circular polarization excitation. As can be seen, the jumps to the background level are correlated on both channels, eliminating sudden rotation as the explanation. Spectral diffusion which explains similar jumps at low temperatures [3] is unlikely for our case, since the absorption spectrum at room temperature is very broad. Similar data, taken from many molecules, reveal transitions to dark states which last from 50ms to 5s. The origin of these transitions is yet to be understood.

The noise in the "on" state does not follow Poisson statistic. With improved time resolution (5ms vs. 20ms), the fluctuations in emission rate get bigger, suggesting transitions to short lived dark states (few ms).

Fig. 2a contrasts a molecule with shot noise limited emission (no transition to dark states) with molecules which are very "noisy" (Fig. 2b and 2c). Fig. 2a represents the typical emission from many molecules. It also suggests that systematic noise such as mechanical drift, vibration and laser coupling fluctuation are minimal at the relevant time scale. In some cases, however, large fluctuations are observed (Fig. 2b and 2c). In particular, Fig. 2c displays almost digital signal fluctuations.

To check the hypothesis of dark states, we modeled the molecular emission with Poissonic distribution of photon arrival at the "on" state, with $1/t_2$ transition probability to a dark state and with t_1 dark state lifetime. Noise distribution histograms were generated from simulated data and compared to histograms obtained from the measurements. In this way t_1 and t_2 could be extracted with better than 0.5ms accuracy. The traces in Fig. 2b and 2c resulted in $t_1 = .075 (\pm 0.25)$ ms, $t_2 = 1.75 (\pm 0.5)$ ms and $t_1 = 2.0 (\pm 0.5)$ ms, $t_2 = 2.0 (\pm 0.5)$ respectively. We found that t_1 ranges from below 0.5ms to 2ms and t_2 from 1.5ms to 2ms. Assuming 5% collection efficiency of fluorescence photons, we estimate that the transfer efficiency to the dark state is about 0.1%. Both the lifetime of dark state (< 2 ms) and the transfer efficiency (or transfer rate of $\sim \mu$ s) are consistent with triplet state lifetime and intersystem crossing rate of dye molecules. Recently, similar quantum jumps to a triplet state were observed from single terrylene molecules in *p*-terphenyl at cryogenic temperatures where triplet state lifetime is the same for every molecule [4]. The distribution in t_1 and t_2 in our measurements, is probably a manifestation of different local environments and surface interactions.

References:

[1] E. Betzig and J.K. Trautman, *Science*, **257**, 189 (1992).

[2] T. Ha, Th. Enderle, D.F. Ogletree, D.S. Chemla, P.R. Selvin, and S. Weiss, to be published.

[3] W.E. Moerner, *Science*, **265**, 46 (1994)

[4] Th. Basche, S. Kummer and C. Brauchle, *Nature*, **373**, 132 (1995).

Figure captions:

Figure1: Simultaneously acquired time traces of single molecular emission at two orthogonal polarizations. Integration time is 20ms.

Figure2: Time traces of single molecular emission from three different molecules. The mean value and the standard deviation of the "on" portion of the signal are given.




Fig. 1, Ha et.al.






Fig. 2, Ha et.al.

LAWRENCE BERKELEY LABORATORY
UNIVERSITY OF CALIFORNIA
TECHNICAL INFORMATION DEPARTMENT
BERKELEY, CALIFORNIA 94720