Lawrence Berkeley National Laboratory

Recent Work

Title

L3-EDGE ANOMALOUS SCATTERING OF X-RAYS BY PRASEODYMIUM AND SAMARIUM

Permalink

<https://escholarship.org/uc/item/8746169b>

Author

Templeton, L.K.

Publication Date

1979-10-01

Lawrence Berkeley Laboratory

UNIVERSITY OF CALIFORNIA

Materials & Molecular Research Division

Submitted to the Journal of the American Chemical Society

L_3 -EDGE ANOMALOUS SCATTERING OF X-RAYS BY PRASEODYMIUM
AND SAMARIUM

Lieselotte K. Templeton, David H. Templeton and
R. Paul Phizackerley

October 1979

RECEIVED
LAWRENCE
BERKELEY LABORATORY

NOV 16 1979

LIBRARY AND
DOCUMENTS SECTION

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 6782.*

LBL-9865 c.2

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

L_3 -Edge Anomalous Scattering of X-rays by Praseodymium and Samarium

Lieselotte K. Templeton and David H. Templeton

Materials and Molecular Research Division
Lawrence Berkeley Laboratory
and
Department of Chemistry
University of California
Berkeley, California 94720

R. Paul Phizackerley

Stanford Synchrotron Radiation Laboratory
and
Department of Structural Biology
Stanford University, Stanford, California 94305

OCTOBER 1979

ABSTRACT

Anomalous scattering terms f' and f'' for Pr and Sm were derived from diffraction experiments with synchrotron radiation and crystals of $\text{NaPrEDTA} \cdot 8\text{H}_2\text{O}$ and $\text{NaSmEDTA} \cdot 8\text{H}_2\text{O}$ at wavelengths near the L_3 absorption edges. Changes in f' ranging from -7 to -26 e for Pr and from -6 to -21 e for Sm, and f'' values up to 28 e for Pr and 23 e for Sm occur at wavelengths within 0.003 Å of the L_3 edges. These large effects can be used to solve the phase problem for macromolecular crystallography by diffraction measurements with one crystal at two or more wavelengths. It is estimated that one atom of praseodymium in a molecule of 800,000 daltons could be enough.

L_3 -Edge Anomalous Scattering of X-rays by Praseodymium and Samarium

Sir:

We report measurements of the anomalous scattering of X-rays by atoms of praseodymium and samarium at wavelengths through their L_3 absorption edges. Values of both f' and f'' , the real and imaginary components of the anomalous scattering, were derived from diffraction experiments using synchrotron radiation. Both components exhibit exceptionally large changes in a narrow interval of wavelength, changes which offer a powerful tool for solving the phase problem in crystal structure analysis.

Synchrotron radiation provides a continuous X-ray spectrum intense enough for single-crystal diffraction experiments at arbitrary wavelengths selected by a narrow-bandpass crystal monochromator. Experiments with cesium hydrogen tartrate^{1,2} showed that f' for cesium dips to -26.7 ± 0.3 electrons and f'' rises to 16.1 ± 0.8 electrons near the L_3 edge ($2.474 \text{ \AA}$). These effects are large enough to be useful in solving macromolecular crystal structures, but cesium is not very convenient for this purpose because of strong absorption of X-rays by light atoms at this wavelength. The rare earth elements have L_3 edges ($2.26 \text{ \AA}$ for lanthanum to $1.34 \text{ \AA}$ for lutetium) which span the wavelength range normally used to study large molecules. X-ray absorption spectra³ for several rare earth compounds show sharp and intense lines at the respective L_3 edges, indicating anomalous scattering effects as large or larger than those for cesium. This indication is confirmed by the present results.

The method⁴ is to use least-squares adjustment to derive f' , f'' and a scale factor from diffraction intensities measured at the wavelength of interest with a crystal whose structure is known from a conventional experiment at another wavelength. We used crystals of sodium praseodymium ethylenediaminetetraacetate octahydrate ($\text{NaPrC}_{10}\text{H}_{12}\text{N}_2\text{O}_8 \cdot 8\text{H}_2\text{O}$) and the isomorphous samarium salt which, like the same salts of most of the other rare earth elements, crystallize in the non-centric space group $Fdd2$.⁵⁻⁷ The atomic coordinates and thermal parameters were determined using Mo $K\alpha$ radiation.⁸ In one praseodymium experiment we measured 970 reflections (without any repetitions) in the range $0.08 < \lambda^{-1} \sin \theta < 0.42$. At each other wavelength we measured 48 reflections in the range of $\lambda^{-1} \sin \theta$ from 0.37 to 0.41, including 23 Bijvoet pairs and two centric reflections, from one to three times each. One reflection was repeated after each 10 or 12 measurements to serve as an intensity standard. Subsequent study of the standard-reflection profiles indicate changes in wavelength during some of the praseodymium experiments, probably caused by movement of the electron orbits in the storage ring. Corrections as large as 0.0011 Å based on the observed changes in Bragg angle are included in the wavelengths listed in Table 1. Absorption corrections were calculated by analytical integration⁹ with the crystal shapes described by 9 or 10 plane faces and absorption coefficients ranging from 100 to 460 cm^{-1} . Other details of the method and apparatus are described elsewhere.^{1,2,10,11}

The values found for the anomalous scattering terms are plotted in Fig. 1 and listed, with the respective R values for each data set, in Tables 1 and 2. For both elements, f'' exhibits the striking resonance line which is observed in the absorption curves. The peak values are substantially larger than we observed for cesium, and in the praseodymium case more than double the values for any element near K or L edges, calculated by Cromer and Liberman¹² with a model which neglects fine structure lines.

The curves for f' show the characteristic shape demanded by the dispersion relation for a sharp absorption line. In Fig. 1 the broken lines show f' calculated from f'' by a Kramers-Kronig integration;² for this purpose the experimental f'' curve was extended to higher and lower wavelengths by estimated values. Errors in these estimates affect the calculated curve by an additive constant, but contribute little to its shape in this narrow wavelength interval. The agreement of the curves is excellent.

The wavelength spread of the X-ray beam from the monochromator, estimated as $\Delta\lambda/\lambda = 10^{-3}$, is comparable to the widths observed for the peaks in f' and f'' . Thus it may be that even larger variations can be observed with more nearly monochromatic X-rays. We intend to test this possibility.

The use of f'' to help solve the phase problem in X-ray crystallography, by means of the intensity difference it causes for reflections hkl and $\bar{k}\bar{h}\bar{l}$, is well established. Larger values of f''

make the method more powerful. The effect of a change in f' is similar to that of a change in atomic number. Thus experiments at two or more wavelengths where f' is different can be analyzed like isomorphous replacement data. Again the utility of the method is sensitive to the magnitude of the changes in f' . Large anomalous scattering terms also facilitate the determination of heavy atom positions. Once the positions are known for the anomalously scattering atoms, one can derive phase angles without ambiguity from diffraction data measured for one crystal at three or more wavelengths, or at two or more wavelengths if both members of each Bijvoet pair are measured. Mathematical procedures for this method have been described by Hoppe and Jakubowski.¹³ They showed that phases could be determined with an accuracy of about 50° for many of the reflections of a protein (erythrocrucorin, M.W. ~16,000) using two wavelengths near the K edge of iron, where the anomalous scattering effects are about 7 times smaller than those reported here for praseodymium. An atom of praseodymium in a molecule of 800,000 daltons would be a somewhat similar case, since according to Wilson statistics the percentage effects of anomalous scattering depend on the ratio of changes in f to the square root of the number of atoms.¹⁴

Acknowledgement. The crystals for this experiment were prepared by Helena W. Ruben. We thank her and Dr. Allan Zalkin for cooperation in the determination of the structural parameters needed for this work, Dr. James C. Phillips and Professor Keith O. Hodgson for

development of the diffractometer facility and its mode of operation at SSRL and for many stimulating discussions, and the staff members of SSRL, too numerous to list here, who helped make it possible for us to work with synchrotron radiation. This work was supported in part by the Division of Nuclear Sciences, Office of Basic Energy Sciences, U.S. Department of Energy under contract No. W-7405-Eng-48. Some of the materials incorporated in this work were developed at the Stanford Synchrotron Radiation Laboratory which is supported by the National Science Foundation (under contract DMR77-27489), in cooperation with SLAC and the Department of Energy.

Lieselotte K. Templeton, David H. Templeton*

Materials and Molecular Research Division,
Lawrence Berkeley Laboratory and Department
of Chemistry, University of California,
Berkeley, California 94720

R. Paul Phizackerley

Stanford Synchrotron Radiation Laboratory
and Department of Structural Biology,
Stanford University, Stanford, California 94305

References and Notes

- (1) Phillips, J.C.; Templeton, D.H.; Templeton L.K.; Hodgson, K.O.
Science 1978, 201, 257-259.
- (2) Templeton, D.H.; Templeton L.K.; Phillips, J.C.; Hodgson, K.O.
Submitted to Acta Crystallographica.
- (3) Phillips, J.C. Ph.D. Thesis 1978, Stanford University.
- (4) Templeton, L.K.; Templeton, D.H. Acta Crystallogr. 1978, A34,
368-371.
- (5) Hoard, J.L.; Lee, B.; Lind, M.D. J. Am. Chem. Soc. 1965, 87,
1612-1613.
- (6) Koetzle, T.F.; Hamilton, W.C. "Anomalous Scattering", Ramaseshan,
S.; Abrahams, S.C.; Ed.; Munksgaard: Copenhagen, 1975; pp. 489-
502.
- (7) Nassimbeni, L.R.; Wright, M.R.W.; van Niekerk, J.C.; McCallum,
P.A. Acta Crystallogr. 1979, B35, 1341-1345.
- (8) Zalkin, A.; Ruben, H.W.; Templeton D.H. unpublished. For
 $\text{NaPr(EDTA)} \cdot 8\text{H}_2\text{O}$, $a = 19.589(3)$, $b = 35.763(5)$, $c = 12.121(2)$ Å,
 $R = 0.037$ for 3449 reflections ($I > \sigma$). For $\text{NaSm(EDTA)} \cdot 8\text{H}_2\text{O}$,
 $a = 19.503(11)$, $b = 35.596(17)$, $c = 12.119(5)$ Å; $R = 0.030$ for
3368 reflections ($I > \sigma$).
- (9) Templeton, L.K.; Templeton, D.H. Abstracts, American Crystallo-
graphic Association Proceedings 1973, Ser. 2, 1, 143.
- (10) Hastings, J.B.; Kincaid, B.M.; Eisenberger, P. Nucl Instr. and
Meth. 1978, 152, 167-171.

- (11) Phillips, J.C.; Cerino, J.A.; Hodgson, K.O. J. Appl. Cryst. 1979, in press.
- (12) Cromer, D.T.; Liberman, D. J. Chem. Phys. 1970, 53, 1891-1898.
- (13) Hoppe, W.; Jakubowski, U. "Anomalous Scattering", Ramaseshan, S.; Abrahams, S.C.; Ed.; Munksgaard: Copenhagen, 1975; pp. 437-461.
- (14) Crick, F.H.C.; Magdoff, B.S. Acta Crystallogr. 1956, 9, 901-908.

Table 1. Praseodimium Anomalous Scattering.

$\lambda(\text{\AA})$	f'	f''	R^a
2.0833	-18.5(6)	4.3(6)	0.045
2.0817	-21.6(5)	5.1(4)	0.050
2.0810	-22.8(5)	5.4(4)	0.077
2.0805	-25.0(7)	7.9(6)	0.083
2.0796	-25.7(6)	8.1(4)	0.068
2.0790	-26.5(3)	19.7(2)	0.084 ^b
2.0788	-25.8(11)	22.3(9)	0.056
2.0784	-24.0(13)	25.6(11)	0.053
2.0776	-17.2(13)	27.8(11)	0.041
2.0772	-14.8(12)	26.9(11)	0.038
2.0767	-6.6(10)	18.7(13)	0.036
2.0745	-10.3(7)	10.5(9)	0.036

$$^a R = \Sigma |\Delta F| / \Sigma |F_o|.$$

^b970 reflections.

Table 2. Samarium Anomalous Scattering.

$\lambda(\text{\AA})$	f'	f''	R
1.8512	-15.6(5)	4.1(6)	0.035
1.8490	-17.2(5)	4.5(5)	0.037
1.8479	-18.6(5)	6.5(5)	0.057
1.8468	-21.3(10)	7.7(10)	0.082
1.8463	-19.5(10)	14.0(13)	0.072
1.8457	-18.6(8)	22.0(7)	0.033
1.8452	-8.5(16)	23.3(18)	0.050
1.8441	-6.3(18)	14.0(24)	0.078
1.8425	-10.0(9)	9.9(12)	0.047
1.8397	-12.7(10)	10.7(12)	0.042

FIGURE CAPTION

Fig. 1. Anomalous scattering terms f' and f'' near the L_3 edges for samarium (left) and praseodymium (right). The broken lines are calculated from f'' by a dispersion relation.

XBL 799-11731

Fig. 1

This report was done with support from the Department of Energy. Any conclusions or opinions expressed in this report represent solely those of the author(s) and not necessarily those of The Regents of the University of California, the Lawrence Berkeley Laboratory or the Department of Energy.

Reference to a company or product name does not imply approval or recommendation of the product by the University of California or the U.S. Department of Energy to the exclusion of others that may be suitable.

TECHNICAL INFORMATION DEPARTMENT
LAWRENCE BERKELEY LABORATORY
UNIVERSITY OF CALIFORNIA
BERKELEY, CALIFORNIA 94720