

UC Irvine

UC Irvine Previously Published Works

Title

Identification and Functional Characterization of Brainstem Cannabinoid CB2 Receptors

Permalink

<https://escholarship.org/uc/item/88q0m818>

Journal

Science, 310(5746)

ISSN

0036-8075

Authors

Van Sickle, Marja D
Duncan, Marnie
Kingsley, Philip J
et al.

Publication Date

2005-10-14

DOI

10.1126/science.1115740

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

Identification and Functional Characterization of Brainstem Cannabinoid CB₂ Receptors

Marja D. Van Sickle,^{1*} Marnie Duncan,^{1*} Philip J. Kingsley,²
Abdeslam Mouihate,¹ Paolo Urbani,³ Ken Mackie,⁴ Nephi Stella,⁵
Alexandros Makriyannis,⁶ Daniele Piomelli,⁷ Joseph S. Davison,¹
Lawrence J. Marnett,² Vincenzo Di Marzo,³ Quentin J. Pittman,¹
Kamala D. Patel,¹ Keith A. Sharkey^{1†}

The presence and function of CB₂ receptors in central nervous system (CNS) neurons are controversial. We report the expression of CB₂ receptor messenger RNA and protein localization on brainstem neurons. These functional CB₂ receptors in the brainstem were activated by a CB₂ receptor agonist, 2-arachidonoylglycerol, and by elevated endogenous levels of endocannabinoids, which also act at CB₁ receptors. CB₂ receptors represent an alternative site of action of endocannabinoids that opens the possibility of nonpsychotropic therapeutic interventions using enhanced endocannabinoid levels in localized brain areas.

Endocannabinoids, anandamide, and 2-arachidonoylglycerol (2-AG) are lipid mediators that act at CB₁ and CB₂ receptors (1, 2). Their actions are terminated through cellular uptake facilitated by a putative endocannabinoid transporter, followed by intracellular enzymatic hydrolysis. Two degradative enzymes for endocannabinoid metabolism are known; fatty acid amide hydrolase (FAAH) preferentially degrades anandamide, and monoacylglycerol lipase preferentially degrades 2-AG (1, 3, 4). The CB₁ receptor is highly expressed in the CNS, where cannabinoids act at presynaptic CB₁ receptors to elicit changes in the synaptic efficacy of neuronal circuits (5). The CB₂ receptor has been found outside the CNS and is particularly associated with immune tissues, such as the spleen and thymus, as well as in various circulating immune cell populations (6). In the CNS, CB₂ receptor mRNA has been reported in cerebel-

lar granule cells (7), and CB₂ receptors have been described on perivascular microglial cells and in cultured cerebrovascular endothelium (8, 9). CB₂ receptor expression is enhanced on glia in neuritic plaques and on immune cells in simian immunodeficiency virus encephalitis (10, 11). To date, however, the CB₂ receptor protein has not been localized on central neurons, and the effects of endocannabinoids in the brain have always been attributed to an action at CB₁ receptors.

We found CB₂ receptor mRNA expression in the brain (cerebellum, cortex, and brainstem) and spleen of the rat using reverse transcription polymerase chain reaction (RT-PCR) (Fig. 1). Sequencing of the 472-base pair PCR product showed that the products amplified from the spleen, cortex, and brainstem were identical to the rat CB₂ receptor sequence (12, 13). Using quantitative real-time RT-PCR, we determined that the brainstem contained 1.5 ± 0.9% of the CB₂ receptor mRNA found in the spleen (fig. S1) (14). Amplification of CB₂ receptor from the brainstem was not due to genomic contamination of our sample, because amplification of RNA that was not reverse-transcribed did not lead to the generation of a product. Furthermore, our real-time PCR primers spanned intron-exon borders, which ensured that the only product that could be amplified was the spliced mRNA.

We next investigated whether CB₂ receptor protein could be detected by Western blotting and/or immunohistochemistry (14). Western blotting for the CB₂ receptor in the brainstem and cerebellum revealed three bands at about 45, 55, and 60 kD (Fig. 1), similar to previous reports in spleen (15). In the brainstem, CB₂ receptor immunoreactivity was found in neurons within the dorsal motor nucleus of the vagus (DMNX), the nucleus ambiguus, and

the spinal trigeminal nucleus; glial cells and blood vessels did not express detectable CB₂ receptor immunoreactivity (16). Preincubation with the cognate peptide of the CB₂ receptor antibody completely abolished cellular staining. These results are in contrast to the conclusions drawn by Derbenev *et al.* (17), who reported no CB₂ mRNA or receptor protein in similar regions of the rat brainstem. However, evaluation of their figures reveals a faint band of immunoreactivity in Western blots, consistent with our observations, and their RT-PCR primers were directed against different regions of the CB₂ receptor mRNA. The differences in the conclusions drawn likely reflect the low abundance of CB₂ receptor in the brain relative to the spleen and the choice of RT-PCR primers.

We next pursued the functional significance of this observation. The major psychoactive cannabinoid, Δ⁹-tetrahydrocannabinol (THC), is effective in the treatment of nausea and vomiting (emesis) (18). THC acts on neurons in the dorsal vagal complex of the brainstem, the site of integration of emetic reflexes that includes the nucleus of the solitary tract (NTS), the area postrema, and DMNX (18, 19), where we found CB₂ receptor expression. These well-characterized actions of CBs have been demonstrated in the ferret. Because of the inability of rodents to vomit, we verified our observations of the receptor distribution in this species (Fig. 1). Indeed, as in the rat, we observed major bands of immunoreactivity in Western blots at about the same relative molecular weights and a similar distribution of CB₂ receptor expression in neurons of the DMNX.

These results led us to investigate whether endocannabinoids could act at the CB₂ receptor in the brainstem to inhibit emesis. Using morphine-6-glucuronide (M6G) as an emetic stimulus, we found that both anandamide and 2-AG dose-dependently reduced emesis in the ferret (Fig. 2) (14). Using selective CB receptor antagonists, we attempted to reverse the actions of these endocannabinoids. The antiemetic effect of anandamide was almost completely reversed by the selective CB₁ receptor antagonist AM251 but was not significantly altered by the CB₂ antagonist AM630, which is consistent with the fact that the anandamide is not very efficacious at CB₂ receptors (2) and indicates that the dose of AM630 used does not antagonize CB₁ receptors. In contrast, the antiemetic effects of 2-AG were reversed by both AM251 and AM630 (Fig. 2). Thus, the CB₂ receptor may be functionally expressed in the ferret brainstem and could be targeted by an endocannabinoid. As we observed the ferrets during our studies, we noted that 2-AG administration was far less sedating than anandamide (Fig. 2). This action is consistent with a preferential effect at a CB₂ receptor, because CB₁ receptor activation *in vivo* is associated with sedation (2).

¹Institute of Infection, Immunity, and Inflammation and Hotchkiss Brain Institute, Department of Physiology and Biophysics, University of Calgary, Calgary, AB, Canada T2N 4N1. ²Department of Biochemistry, Vanderbilt Institute of Chemical Biology, Center in Molecular Toxicology, Vanderbilt-Ingram Cancer Center, Vanderbilt University School of Medicine, Nashville, TN 37232-0146, USA. ³Endocannabinoid Research Group, Institute of Biomolecular Chemistry, Consiglio Nazionale delle Ricerche, Pozzuoli, Naples, Italy. ⁴Department of Anesthesiology and Department of Physiology and Biophysics, University of Washington, Seattle, WA 98195-6540, USA. ⁵Department of Pharmacology, Psychiatry, and Behavioral Sciences, University of Washington, Seattle, WA 98195-7280, USA. ⁶Center for Drug Discovery, Department of Pharmaceutical Sciences, Northeastern University, Boston, MA 02115, USA. ⁷Department of Pharmacology, Gillespie Neuroscience Facility, University of California, Irvine, CA 92697-6180, USA.

*These authors contributed equally to this work.

†To whom correspondence should be addressed. E-mail: ksharkey@ucalgary.ca





Fig. 1. CB₂ receptor is expressed in the rat, mouse, and ferret CNS. (A) RNA was isolated from the spleen (SPL), cerebellum (CER), cortex (COR), and brainstem (BS) of rats. RT-PCR was performed using primers for CB₂ receptor (CB₂) or β -actin, and the expected amplicons were 472 and 277 base pairs, respectively. No band was detected in the no template control (NT). RNA from rat brainstem was reverse-transcribed or mock-treated before PCR. No bands in the RNA sample indicates amplification was not due to genomic DNA in the RNA sample. (B) Western blot containing protein homogenates of ferret or rat brain as indicated, beside Western blot of the same homogenates incubated with antibody preabsorbed with control peptide. Major bands were observed at about 45, 55, and 60 kD in rat and ferret brainstem (arrows; $n = 6$). In the ferret spleen, higher molecular weight bands were also observed (arrow). In the rat brain and spleen and ferret brain, we completely preabsorbed all the immunoreactive bands. However, in the ferret spleen, bands were substantially reduced, but not completely preabsorbed, with the concentration of peptide used. (C) CB₂ receptor immunoreactivity in the dorsal motor nucleus of the vagus of wild-type (left) and CB₂^{-/-} (right) mice (30).

Note the lack of immunoreactive cell bodies in the knockout mouse ($n = 3$ per group). Scale bar, 50 μ m. (D) Immunoreactivity for the neuronal nuclear marker NeuN (left, green) and CB₂ receptor immunoreactivity (center, red) in neurons of the dorsal motor nucleus of the vagus of the ferret and rat, as indicated. Overlay (right) of NeuN and CB₂ receptor illustrates that CB₂ receptor immunoreactivity is present in neurons of both the ferret and rat ($n = 4$), where it is localized on the cell membrane and in the cytoplasm of the neurons. Scale bars, 20 μ m.

Fig. 2. Endocannabinoids reduce emesis induced by M6G. (A) Episodes of emesis after treatments of emetic agent alone, M6G (0.05 mg/kg, subcutaneously; $n = 10$) or the following treatments ($n = 3$ to 7) preceding M6G; anandamide (AEA-1, 1 mg/kg; and AEA-2, 2 mg/kg); and 2-arachidonoyl glycerol (2-AG-0.5, 0.5 mg/kg; and 2-AG-2, 2 mg/kg). AEA and 2-AG had no emetic actions when given alone. (B) Episodes of emesis after M6G alone; or with the transport inhibitor (VDM11-0.5, 0.5 mg/kg; and VDM11-2, 2 mg/kg); paired ineffective doses of VDM11 and 2-AG (0.5 mg/kg); an FAAH inhibitor URB 597 (3 mg/kg). (C) Activity during the 60-min observation after the treatments indicated. (D) CB₁ antagonist AM251 (5 mg/kg) reversed the effects of anandamide (AEA, 2 mg/kg), 2-AG (2 mg/kg), and VDM11 (2 mg/kg). In contrast, the CB₂ antagonist AM630 (5 mg/kg) did not reverse the effect of AEA (2 mg/kg), but effectively reversed the action of 2-AG (2 mg/kg) and VDM11 (2 mg/kg). In the presence of AM630 (5 mg/kg), there were 3.4 ± 0.7 emetic episodes ($n = 5$) in animals given a lower dose of AEA (1 mg/kg), which is not significantly different from the 2.8 ± 1.3 emetic episodes ($n = 5$) in the absence of AM630. AM251 and AM630 had no emetic actions when given alone. AM630 did not potentiate the effects of M6G, whereas AM251 enhanced emesis as previously described (17). Results are expressed as percentage of number of episodes of emesis induced by M6G. *Significant differences compared with M6G alone, $P < 0.05$. Bars represent mean \pm SEM.



Although these data were strongly indicative of the actions of 2-AG at CB₂ receptors, it was important to evaluate whether endogenously released endocannabinoids reduce emesis through an action at CB₂ receptors. We used two approaches that would raise endocannabinoid levels in the brain. First, we tested whether the selective endocannabinoid reuptake inhibitor VDM11 (20) inhibited emesis. Second, we tested whether an FAAH inhibitor (URB 597) (21) reduced emesis. In both cases, we blocked or reduced the extent of emesis induced by M6G (Fig. 2). We extended these studies to find whether VDM11 would potentiate the effects of 2-AG. We used doses of VDM11 and 2-AG that alone did not significantly reduce emesis. When VDM11 was given together with 2-AG, we found a significant attenuation in the number of emetic episodes (Fig. 2). URB 597 strongly sedated animals, which suggested that this compound may

selectively enhance anandamide levels. Conversely, VDM11 had no sedative effects, and the antiemetic effects were reversed by both AM251 and AM630, which suggested that VDM11 may preferentially affect 2-AG levels.

If the arguments made above were correct and if endocannabinoids undergo increased turnover in response to an emetic stimulus, treatment with VDM11 or URB 597 would be expected to increase the levels of endocannabinoids in the brainstem. We investigated this using a model of emesis under anesthetized conditions, so that the brains could be rapidly removed and frozen in order to limit the inherent instability of endocannabinoids in tissue samples (14). Ferrets were given hypertonic saline as an emetic stimulus because M6G is an inconsistent emetic in anesthetized animals. Levels of anandamide and total arachidonoyl glycerol [which reflect tissue 2-AG levels (14)] were measured in the brainstem and, for com-

parison, the cerebellum. Levels of endocannabinoids in the ferret under basal conditions or after emesis were comparable to levels previously found in rodents (Fig. 3). As expected from the pharmacological experiments described above, VDM11 specifically increased levels of total arachidonoyl glycerol in the brainstem and cerebellum, whereas pretreatment with URB 597 only increased the levels of anandamide in the brainstem.

These results led us to investigate whether selective CB₂ receptor agonists reduced emesis. We observed no statistically significant reductions in emesis in animals given the CB₂ receptor agonists AM1241 (1 or 2 mg/kg) or JWH 133 (1 or 5 mg/kg) before M6G (22). This finding was not completely surprising, as inhibitors of endocannabinoid inactivation can be more efficacious than "direct" agonists (23–25). Furthermore, these data suggest that CB₂ receptor activation alone is not sufficient to inhibit emesis and that, under appropriate conditions, for example, those produced by inhibiting endocannabinoid inactivation, the CB₂ receptor can be activated in local brain regions together with CB₁ receptors and can inhibit emesis. This hypothesis was supported by a significant reduction in episodes of emesis (7.1 ± 0.5 to 5.0 ± 0.7 ; $n = 6$ to 10 ; $P < 0.05$) when anandamide (0.5 mg/kg) and AM1241 (1 mg/kg) were administered together at doses that were not antiemetic when either compound was given alone with M6G.

The behavioral evidence cited above is not a direct measure of neuronal activation and does not directly show functionally active CB₂ receptors in the brainstem. To determine whether CB₂ receptor agonists can activate neurons of the DMNX, we investigated the expression of phosphorylated extracellular signal-regulated kinase 1/2 (pERK) in rat DMNX neurons by immunohistochemistry (14), because phosphorylation of this enzyme is enhanced by cannabinoid agonists in other regions of the brain and in cell lines (26, 27). Administration of the CB₂ receptor agonist AM1241 increased pERK in DMNX neurons when compared with vehicle-treated controls (Fig. 4).





Fig. 3. Endocannabinoid levels in the ferret brainstem (left) and cerebellum (right). Control ferrets received intragastric normal saline ($n = 4$ to 7). After the emetic stimulus intragastric hypertonic saline (HS), levels were not significantly increased. URB597 (5 mg/kg) increased anandamide (AEA) levels in the brainstem but not in the cerebellum. VDM11 (3 mg/kg) increased total arachidonoyl glycerol (1-AG and 2-AG, 1AG accounts for about 20% and 2-AG about 80% of total) in both the brainstem and cerebellum when compared with the HS group ($P < 0.05$).

Fig. 4. A CB₂ receptor agonist activates neurons in the dorsal motor nucleus of the vagus (DMNX) of the rat. Immunoreactivity for pERK in rats treated with vehicle (A) ($n = 3$) and (B) the selective CB₂ receptor agonist, AM1241 (1 mg/kg; $n = 7$). AM1241 stimulated the expression of 5.6 ± 1.2 pERK immunoreactive cells per section in the DMNX compared with 1.2 ± 0.2 pERK immunoreactive cells in vehicle-treated animals. pERK immunoreactive cells were also observed in the nucleus of the solitary tract (not shown on figure). pERK immunoreactivity (red) was observed in nuclei and cytoplasm of activated cells. (C) CB₂ receptor immunoreactivity (green) was observed on the cell membrane and



in the cytoplasm of DMNX neurons. (D) Overlay of pERK and CB₂ receptor illustrate the presence of pERK in neurons that express the CB₂ receptor (arrows). pERK immunoreactivity was observed in about 15 to 20% of the CB₂ immunoreactive neurons. Scale bar, 50 μ m.

We have shown that CB₂ receptors are present in the brainstem and also in the cortex and cerebellum. As inferred by the use of a selective CB₂ antagonist, the brainstem receptors are functionally coupled to inhibition of emesis when costimulated with CB₁ receptors by an endogenous cannabinoid capable of activating both receptors. The extent of participation of CB₂ receptors in this effect is sufficient to reduce the widespread behavioral actions associated with the administration of CB₁ agonists. However, generalized activation of CB₂ receptors leads to immunosuppression (28) and is potentially deleterious if used as a therapy. Others have suggested that modulating the endocannabinoid system in the CNS represents a promising strategy for therapies for CNS disorders (29). Our observations suggest that targeting specific local populations of cannabinoid receptors (both CB₁ and CB₂) by enhancing endocannabinoid levels where they are released represents a therapeutic strategy that may be useful in disorders where either CB₁ or CB₂ receptor activation alone would not be desirable. This approach would circumvent the psychotropic and immunosuppressive side effects of exogenously administered cannabinoids and would provide an alternative approach for the therapeutic utilization of this unique neuroregulatory system.

peutic utilization of this unique neuroregulatory system.

References and Notes

1. V. Di Marzo, M. Bifulco, L. De Petrocellis, *Nat. Rev. Drug Discov.* **3**, 771 (2004).
2. A. C. Howlett et al., *Pharmacol. Rev.* **54**, 161 (2002).
3. L. De Petrocellis, M. G. Cascio, V. Di Marzo, *Br. J. Pharmacol.* **141**, 765 (2004).
4. T. P. Dinh et al., *Proc. Natl. Acad. Sci. U.S.A.* **99**, 10819 (2002).
5. T. F. Freund, I. Katona, D. Piomelli, *Physiol. Rev.* **83**, 1017 (2003).
6. T. W. Klein et al., *J. Leukoc. Biol.* **74**, 486 (2003).
7. S. D. Skaper et al., *Proc. Natl. Acad. Sci. U.S.A.* **93**, 3984 (1996).
8. E. Nunez et al., *Synapse* **53**, 208 (2004).
9. S. A. Golech et al., *Brain Res. Mol. Brain Res.* **132**, 87 (2004).
10. C. Benito et al., *J. Neurosci.* **25**, 2530 (2005).
11. C. Benito et al., *J. Neurosci.* **23**, 11136 (2003).
12. G. Griffin, Q. Tao, M. E. Abood, *J. Pharmacol. Exp. Ther.* **292**, 886 (2000).
13. S. M. Brown, J. Wager-Miller, K. Mackie, *Biochim. Biophys. Acta* **1576**, 255 (2002).
14. Materials and methods are available as supporting material on Science Online.
15. I. Matias et al., *Eur. J. Biochem.* **269**, 3771 (2002).
16. M. D. Van Sickle, Q. J. Pittman, K. A. Sharkey, unpublished observations.
17. A. V. Derbenev, T. C. Stuart, B. N. Smith, *J. Physiol.* **559**, 923 (2004).
18. M. D. Van Sickle et al., *Gastroenterology* **121**, 767 (2001).
19. P. J. Hornby, *Am. J. Med.* **111** (suppl. 8A), 106S (2001).
20. L. De Petrocellis, T. Bisogno, J. B. Davis, R. G. Pertwee, V. Di Marzo, *FEBS Lett.* **483**, 52 (2000).

21. D. Fegley et al., *J. Pharmacol. Exp. Ther.* **313**, 352 (2005).
22. M. D. Van Sickle, J. S. Davison, K. A. Sharkey, unpublished observations.
23. G. Marsicano et al., *Science* **302**, 84 (2003).
24. S. Gaetani, V. Cuomo, D. Piomelli, *Trends Mol. Med.* **9**, 474 (2003).
25. J. P. Chhatwal, M. Davis, K. A. Maguschak, K. J. Ressler, *Neuropsychopharmacology* **30**, 516 (2005).
26. E. Valjent et al., *Proc. Natl. Acad. Sci. U.S.A.* **102**, 491 (2005).
27. M. Bouaboula et al., *Eur. J. Biochem.* **237**, 704 (1996).
28. T. W. Klein et al., *J. Leukoc. Biol.* **74**, 486 (2003).
29. B. F. Cravatt, A. H. Lichtman, *Curr. Opin. Chem. Biol.* **7**, 469 (2003).
30. N. E. Buckley et al., *Eur. J. Pharmacol.* **396**, 141 (2000).
31. We thank L. Oland, W. Ho, and N. Shariat for excellent technical support and animal care. Q.J.P., J.S.D., and K.A.S. were supported by the Canadian Institutes of Health Research (CIHR); K.D.P. by a CIHR group grant in leukocyte trafficking; L.J.M. by GM15431; and K.M., N.S., A.M., and D.P. by U.S. National Institutes of Health. M.D.V.S. is an Alberta Heritage Foundation for Medical Research (AHFMR) graduate student; M.D. is an AHFMR and Canadian Association of Gastroenterology/CIHR/Janssen-Ortho postdoctoral fellow; K.D.P. holds a Canada research chair and is an AHFMR senior scholar; and Q.J.P. and K.A.S. are AHFMR medical scientists.

Supporting Online Material

www.sciencemag.org/cgi/content/full/310/5746/329/DC1
Materials and Methods
Fig. S1
References

6 June 2005; accepted 9 September 2005
10.1126/science.1115740

Observing Others: Multiple Action Representation in the Frontal Lobe

Koen Nelissen,¹ Giuseppe Luppino,² Wim Vanduffel,^{1,3}
Giacomo Rizzolatti,² Guy A Orban^{1*}

Observation of actions performed by others activates monkey ventral premotor cortex, where action meaning, but not object identity, is coded. In a functional MRI (fMRI) study, we investigated whether other monkey frontal areas respond to actions performed by others. Observation of a hand grasping objects activated four frontal areas: rostral F5 and areas 45B, 45A, and 46. Observation of an individual grasping an object also activated caudal F5, which indicates different degrees of action abstraction in F5. Observation of shapes activated area 45, but not premotor F5. Convergence of object and action information in area 45 may be important for full comprehension of actions.

Understanding actions performed by others is a fundamental social ability. There is now wide consensus that the activation of the motor

system is a necessary requisite for this ability. A mere visual representation, without involvement of the motor system, provides a description of the visible aspects of the movements of the agent, but does not give information critical for understanding action semantics, i.e., what the action is about, what its goal is, and how it is related to other actions (1, 2). Action information, however, without knowledge about the identity of the object acted upon, is not sufficient to provide a full understanding of the observed action. Only when information about the object identity is added to the se-

mantic information about the action can the actions of other individuals be completely understood (3).

The functional properties of a set of neurons in monkey ventral premotor cortex (area F5) provide evidence for the involvement of the motor system in action understanding. These "mirror" neurons discharge both when the individual performs an action and when the individual observes another person performing the same action (4, 5). They therefore match the observed action with its internal motor representation. F5 neurons responding to the observation of grasping respond equally well when a piece of food or a solid object of similar size and shape is being grasped. The object's identity appears to be ignored in F5 (4, 5).

We used fMRI in five awake monkeys (M1, M3 to M6) (6–9) to test how actions performed by others are represented in the monkey frontal lobe. In experiment 1, we intended to localize the frontal lobe regions involved in action observation. Monkeys saw video clips showing a full view of a person grasping an object ("acting person"), or an isolated hand grasping objects ("hand action") and static single frames or scrambled videos as controls. The acting person movies approximate the visual stimulation used in F5 single-cell studies (4, 5) and provide context information that is lacking in the hand action movies, which has been used in most human

¹Laboratorium voor Neuro-en Psychofysiologie, Katholieke Universiteit Leuven Medical School, Leuven, Belgium. ²Dipartimento di Neuroscienze, Università di Parma, via Volturno 39, 43100 Parma, Italy. ³Massachusetts General Hospital, Massachusetts Institute of Technology, Harvard Medical School, Athinoula A. Martino's Center for Biomedical Imaging, Charlestown, MA 02129, USA.

*To whom correspondence should be addressed.
E-mail: guy.orban@med.kuleuven.ac.be