UC Irvine

Journal of Education and Teaching in Emergency Medicine

Title

Bedside Ultrasound for the Diagnosis of Small Bowel Obstruction

Permalink

<https://escholarship.org/uc/item/88t2d13c>

Journal

Journal of Education and Teaching in Emergency Medicine, 2(4)

Authors

Anshus, Alexander
Alvarado, Maili

Publication Date

2017

DOI

10.5070/M524036786

Copyright Information

Copyright 2017 by the author(s). This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

Bedside Ultrasound for the Diagnosis of Small Bowel Obstruction

Alexander Anshus, BS* and Maili Alvarado, MD^

*University of California, Irvine, Department of Emergency Medicine, Orange, CA

^University of Arizona, University Medical Center Tucson, Department of Emergency Medicine, Tucson, AZ

Correspondence should be addressed to Maili Alvarado, MD at maili.alvarado@gmail.com

Submitted: May 31, 2017; Accepted: September 19, 2017; Electronically Published: October 15, 2017; <https://doi.org/10.21980/J86W6P>

Copyright: © 2017 Anshus, et al. This is an open access article distributed in accordance with the terms of the Creative Commons Attribution (CC BY 4.0) License. See: <http://creativecommons.org/licenses/by/4.0/>

Video Link: <https://youtu.be/VNsSjVsV9vI>

History of present illness: An elderly female with no history of prior abdominal surgeries presented to the emergency department (ED) with acute onset of abdominal pain and distention. Upon arrival, she began having large volume bilious emesis. While waiting for a computed tomography (CT) scan of her abdomen and pelvis, a point of care ultrasound (POCUS) was performed which showed evidence of a small bowel obstruction (SBO). The patient had a nasogastric tube placed that put out over two liters of bilious contents. A subsequent CT scan confirmed the diagnosis of SBO from a left inguinal hernia and the patient was admitted to the surgical service.

Significant findings: The POCUS utilizing the low frequency curvilinear probe demonstrates fluid-filled, dilated bowel loops greater than 2.5cm with to-and-fro peristalsis, and thickened bowel walls greater than 3mm, concerning for SBO.

Discussion: Gastrointestinal obstruction is a common diagnosis in the ED, accounting for approximately 15% of all ED visits for acute abdominal pain.¹ SBO accounts for approximately 80% of all obstructions.² In the diagnosis of SBO, studies show that abdominal x-rays have a sensitivity of 66-77% and specificity of 50-57%,³ CT scans have a sensitivity of 92% and specificity of 93%,⁴ and ultrasound has a sensitivity of 88% and specificity of 96%.⁵

While CT scan remains a widely accepted modality for diagnosing SBO, ultrasound is more cost effective, well tolerated, does not involve ionizing radiation, and can be done in a timely manner at the patient's bedside. Ultrasound can also identify transition points as well as distinguish between functional and mechanical obstruction.⁶ In addition to SBO, ultrasound can be used to diagnose external hernias, intussusception, tumors, superior mesenteric artery (SMA) syndrome, foreign bodies, bezoars, and ascariasis.⁷

Topics: Small bowel obstruction, ultrasound, point of care ultrasound, POCUS.

References:

1. Irvin TT. Abdominal pain: a surgical audit of 1190 emergency admissions. *Br J Surg*. 1989;76(11):1121-1125. doi: 10.1002/bjs.1800761105
2. Markogiannakis H, Messaris E, Dardamanis D, et al. Acute mechanical bowel obstruction: clinical presentation, etiology, management and outcome. *World J Gastroenterol*. 2007;13(3):432-437. doi: 10.3748/wjg.v13.i3.432
3. Shrake PD, Rex DK, Lappas JC, Maglinte DD. Radiographic evaluation of suspected small bowel obstruction. *Am J Gastroenterol*. 1991;86(2):175-178.
4. Mallo RD, Salem L, Lalani T, Flum DR. Computed tomography diagnosis of ischemia and complete obstruction in small bowel obstruction: A systematic review. *J Gastrointest Surg*. 2005;9(5):690-694. doi: 10.1016/j.gassur.2004.10.006
5. Ogata M, Mateer JR, Condon RE. Prospective evaluation of abdominal sonography for the diagnosis of bowel obstruction. *Ann Surg*. 1996;223(3):237-241.
6. Truong S, Arlt G, Pfingsten F, Schumpelick V. Importance of sonography in diagnosis of ileus. A retrospective study of 459 patients. *Chirurg*. 1992;63(8):634-640.
7. Wale A and Pilcher J. Current role of ultrasound in small bowel imaging. *Semin Ultrasound CT MR*. 2016;37(4):301-312. doi: 10.1053/j.sult.2016.03.001