

Lawrence Berkeley National Laboratory

LBL Publications

Title

Summary of the Research Progress Meeting

Permalink

<https://escholarship.org/uc/item/88w380qs>

Author

Kramer, H P

Publication Date

1949-05-01

UNIVERSITY OF
CALIFORNIA

*Radiation
Laboratory*

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 5545*

BERKELEY, CALIFORNIA

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

UNIVERSITY OF CALIFORNIA

Radiation Laboratory

Contract No. W-7405-eng-48

UNCLASSIFIED

SUMMARY OF THE RESEARCH PROGRESS MEETING

May 12, 1949

H. P. Kramer

Some of the results reported in this document may be of a preliminary or incomplete nature. It is the request of the Radiation Laboratory that the document not be circulated off the project nor the results quoted without permission.

Berkeley, California

Installation	No. Copies
Argonne National Laboratory	10
Armed Forces Special Weapons Project	1
Atomic Energy Commission, Washington	2
Battelle Memorial Institute	1
Brookhaven National Laboratory	10
Bureau of Ships	1
Carbide & Carbon Chemicals Corporation (K-25)	4
Carbide & Carbon Chemicals Corporation (Y-12)	4
Chicago Operations Office	1
Columbia University (Dunning)	1
Columbia University (Failla)	1
General Electric Company, Richland	7
Hanford Operations Office	1
Iowa State College	1
Kellex Corporation	2
Knolls Atomic Power Laboratory	4
Los Alamos	6
Massachusetts Institute of Technology (Gaudin)	1
Massachusetts Institute of Technology (Kaufmann)	1
Mound Laboratory	8
National Advisory Committee for Aeronautics	2
National Bureau of Standards	2
Naval Radiological Defense Laboratory	2
NEPA-Project	1
New York Operations Office	8
North American Aviation, Inc.	2
Oak Ridge National Laboratory	12
Patent Advisor, Washington	1
Sandia Base	2
Technical Information Branch, ORE	15
UCLA Medical Research Laboratory (Warren)	1
University of California Radiation Laboratory	6
University of Rochester	2
Western Reserve University (Friedell)	4
Sylvania Electric Products, Inc.	1

INFORMATION DIVISION
 Radiation Laboratory
 Univ. of California
 Berkeley, California

SUMMARY OF THE RESEARCH PROGRESS MEETING

May 12, 1949

H. P. Kramer

Controlled Atmosphere for a Spectrographic Source. J. Conway.

The spectrographic analysis of the rare earth elements by means of the conventional apparatus is impeded by the blanketing of the region from 3200-4200 Å by heavy cyanogen bands. The search for a method of shielding the plates from these bands without removing the desired spectral lines has resulted in a successful apparatus.

A gas-tight arc stand has been developed that is equipped with a variable speed motor that automatically adjusts the separation of the electrodes. The enclosed space is filled with oxygen gas at 2 lbs. per square inch above atmospheric pressure. It was found best to retain the conventional copper electrodes taking proper care to protect them from corrosion when a spectrographic analysis of elements contained in solution is undertaken.

The Use of a Magnet in Scattering Experiments. J. Hadley.

An apparatus has been set up for the magnetic analysis of charged particles emanating from a target exposed to the 200 Mev neutron beam that is produced in the 184-inch cyclotron on a 2 inch beryllium target. The arrangement is shown in Fig. 1. Charged particles are ejected from the target, pass through a counter into a magnetic field of 15000 gauss and after being deflected into a circular path appropriate to their charge and momentum again register on a counter that is in coincidence with the first tube.

Very narrow counting tubes are used instead of collimators that might introduce uncertainties due to slit penetration. It is desired to avoid the use of absorbers as much as possible since the manner of interaction of highly energetic

charged particles with the atoms of absorbers is not sufficiently well known. However, because of the small ionization that fast particles produce, it would be necessary to increase the sensitivity of the counting circuits correspondingly and if this is done, one conjures the likelihood of spurious counts.

Another difficulty that is encountered in the use of a magnetic field for energy discrimination is that at high energies the resolving power of the magnet is very poor. It was suggested from the audience that this trouble might be cured by using a number of counters placed side by side rather than employing one counter and thus making the measurements depend completely on the resolving power of the magnet.

Such a scheme it was felt would also contribute toward the solution of a further problem in connection with the apparatus: the extremely low counting rate and consequent poor statistics. It was found that at a scatter angle of 30° , at the poorest magnet resolution, and with the full intensity beam, the counting rate was only one count every two seconds.

If the performance of the apparatus can be improved sufficiently it is contemplated to extend the measurements made by H. York, et al. on secondary particles produced in various materials to the higher neutron energy.

Some Meson Work with the Deflected Proton Beam. C. Richman.

An experiment is being run to measure the distribution as to angle and energy of mesons produced in a carbon target by the 350 Mev proton beam from the 184-inch cyclotron. As can be seen in Fig. 2, the mesons that are produced enter wells sunk into a block of Pb along discrete radii from the target. Each one of these wells contains a photographic plate that records the mesons impinging on it. If a complete count is made and the beam is integrated the equipment also serves to furnish data for a calculation of the cross section for meson production. Ultimately it would be desirable for the advancement of theory to perform the experiment with a hydrogen

target in order to obtain the meson cross section and distributions for proton-proton interaction.

The figure indicates the complete shielding of the apparatus. That the shielding is successful was borne out by the results of a 2 hour run made without the target. The plates came out very clean.

Next a seven hour run with the target in position was performed. The density of meson tracks was about one track per mm^2 . However, the run was unsatisfactory because the background from scattered protons was so considerable that only the plates placed at angles greater than 90° could be read without difficulty and the plates at 10° and 20° were completely useless. Upon examination of the beam by means of a photographic plate directly exposed to the incoming protons, it was found that its outline was not sharply defined. This fuzziness was thought to be due to the fact that the beam was striking the collimating block. The geometry was changed and plates that were exposed in the forward direction after this alteration had been made were much clearer and can be read with some work.

The plate exposed at 115° was hurriedly counted and a total of 98 meson tracks were counted. 27 of these culminated in a star, 17 indicated the decay of a π meson into a μ meson, 41 were virgin tracks, and 13 stopped only after they had left the emulsion. A calculation based on the empirical relation that the number of negative π mesons is to the number of star producing mesons as 1.37 is to 1, yields the value 1.5 for the ratio of negative to positive π mesons. The average energy of the mesons that were observed is about 30 Mev.




Fig. 1




Fig. 2