

UC San Diego

UC San Diego Previously Published Works

Title

Alpine soil carbon is vulnerable to rapid microbial decomposition under climate cooling

Permalink

<https://escholarship.org/uc/item/89h1g6rc>

Journal

The ISME Journal: Multidisciplinary Journal of Microbial Ecology, 11(9)

ISSN

1751-7362

Authors

Wu, Linwei
Yang, Yunfeng
Wang, Shiping
[et al.](#)

Publication Date

2017-09-01

DOI

10.1038/ismej.2017.75

Peer reviewed

ORIGINAL ARTICLE

Alpine soil carbon is vulnerable to rapid microbial decomposition under climate cooling

Linwei Wu^{1,2,11}, Yunfeng Yang^{1,11}, Shiping Wang^{3,4}, Haowei Yue¹, Qiaoyan Lin⁵, Yigang Hu^{5,6}, Zhili He², Joy D Van Nostrand², Lauren Hale², Xiangzhen Li⁷, Jack A Gilbert^{8,9} and Jizhong Zhou^{1,2,10}

¹State Key Joint Laboratory of Environment Simulation and Pollution Control, School of Environment, Tsinghua University, Beijing, China; ²Institute for Environmental Genomics, Department of Microbiology and Plant Biology, and School of Civil Engineering and Environmental Sciences, University of Oklahoma, Norman, OK, USA; ³Key Laboratory of Alpine Ecology and Biodiversity, Institute of Tibetan Plateau Research, Chinese Academy of Sciences, Beijing, China; ⁴CAS Center for Excellence in Tibetan Plateau Earth Science, Beijing, China; ⁵Key Laboratory of Adaption and Evolution of Plateau Biota, Northwest Institute of Plateau Biology, Chinese Academy of Sciences, Xining, China; ⁶Shapotou Desert Experiment and Research Station, Cold and Arid Regions and Environmental & Engineering Research Institute, Chinese Academy of Sciences, Lanzhou, China; ⁷Chengdu Institute of Biology, Chinese Academy of Sciences, Chengdu, China; ⁸The Microbiome Center, BioScience Division, Argonne National Laboratory, Argonne, IL, USA; ⁹Department of Surgery, University of Chicago, Chicago, IL, USA and ¹⁰Earth and Environmental Sciences, Lawrence Berkeley National Laboratory, Berkeley, CA, USA

As climate cooling is increasingly regarded as important natural variability of long-term global warming trends, there is a resurging interest in understanding its impact on biodiversity and ecosystem functioning. Here, we report a soil transplant experiment from lower to higher elevations in a Tibetan alpine grassland to simulate the impact of cooling on ecosystem community structure and function. Three years of cooling resulted in reduced plant productivity and microbial functional potential (for example, carbon respiration and nutrient cycling). Microbial genetic markers associated with chemically recalcitrant carbon decomposition remained unchanged despite a decrease in genes associated with chemically labile carbon decomposition. As a consequence, cooling-associated changes correlated with a decrease in soil organic carbon (SOC). Extrapolation of these results suggests that for every 1 °C decrease in annual average air temperature, 0.1 Pg (0.3%) of SOC would be lost from the Tibetan plateau. These results demonstrate that microbial feedbacks to cooling have the potential to differentially impact chemically labile and recalcitrant carbon turnover, which could lead to strong, adverse consequences on soil C storage. Our findings are alarming, considering the frequency of short-term cooling and its scale to disrupt ecosystems and biogeochemical cycling.

The ISME Journal (2017) 11, 2102–2111; doi:10.1038/ismej.2017.75; published online 23 May 2017

Introduction

Global climate change science has predominantly targeted climate warming (Luo, 2007; Frey *et al.*, 2013; Nie *et al.*, 2013), but this focus is changing, as it is increasingly recognized that temporary, local cooling events are common amidst long-term global warming trends (Alley *et al.*, 2003; Ji *et al.*, 2014). There has been significant cooling in the Antarctic

Peninsula as the late 1990s, arising from natural variability of the regional atmospheric circulation (Turner *et al.*, 2016). It has been projected that the climate in the 21st century is likely to produce periods as long as one or two decade(s) of cooling (Easterling and Wehner, 2009; Lyubushin and Klyashtorin, 2012), which is alarming because historic evidence shows that cooling may perturb ecosystems and biogeochemical cycling at a scale comparable to what is known for warming (McAnena *et al.*, 2013). For instance, cooling on the Antarctic continent between 1966 and 2000 led to a rapid decrease in the primary productivity of lakes (6–9% per year) as well as the number of soil invertebrates (more than 10% loss per year) (Doran *et al.*, 2002). Therefore, further understanding of the full range of possible climate change scenarios and

Correspondence: Y Yang, State Key Joint Laboratory of Environment Simulation and Pollution Control, School of Environment, Tsinghua University, 1 Tsinghua Garden Road, Haidian District, Beijing 100084, China.
E-mail: yangyf@tsinghua.edu.cn

¹¹These authors contributed equally to this work.

Received 27 September 2016; revised 14 March 2017; accepted 23 March 2017; published online 23 May 2017

their potential impacts is of considerable value for policy makers and citizens to promulgate effective responses. Despite the impact, we still lack a fundamental understanding of how ecosystems respond to climate cooling.

To date, the effects of cooling are estimated indirectly by historical records (Campbell and McAndrews, 1993; Doran *et al.*, 2002) or model simulation (Lucht *et al.*, 2002), owing to the difficulty of carrying out *in situ* studies. Soil transplant experiments provide an opportunity to quantify the direct influence of substantial climate changes on the plant and soil microbial community structure and function (Breeuwer *et al.*, 2010; Vanhala *et al.*, 2011; Luan *et al.*, 2014). For example, soil transplants into warmer climates have shown comparable results to long-term *in situ* artificial warming (Petchey *et al.*, 1999; Vanhala *et al.*, 2011; Zhou *et al.*, 2012; Luan *et al.*, 2014; Yue *et al.*, 2015). In this study, we conducted a soil transplant study in a Tibetan alpine grassland, which is the largest grassland on the Eurasian continent (Yang *et al.*, 2008). Soils on the Tibetan alpine store a large amount of organic C (about 10% of the terrestrial C in China) and are especially vulnerable to global climate change (Qiu, 2008; Yang *et al.*, 2008), making it an ideal region to study the feedback of the grassland ecosystem to climate change. In this study, soils with attached vegetation from lower elevations were moved sequentially to higher elevations, along an elevation gradient of 3200, 3400, 3600 and 3800 m above sea level (see Materials and Methods for details). Soils mock transplanted, that is, soils removed from and then reinstalled to the same place, were used as controls. Various theoretical and empirical studies have suggested that climate warming increases both plant productivity and soil respiration (Luo, 2007; Zhou *et al.*, 2012), resulting in an increase in organic C loss through increased soil respiration, which is not offset by increased net primary production (Frey *et al.*, 2013). Therefore, we hypothesize that, in contrast to warming, cooling would suppress associated ecosystem functional processes, such as C uptake and respiration, and consequently cause a net increase in SOC.

Materials and Methods

Experimental design and soil sampling

The transplant experiments were carried out in Haibei Alpine Meadow Ecosystem Research Station (37°37'N, 101°12'E) of the Northeastern Tibet Plateau, Qinghai, China. Experimental plots were set up in May 2007. Along an elevation gradient of 3200, 3400, 3600 and 3800 m above sea level, triplicate soils with sizes of 1 m length × 1 m width × 0.3 m depth were dug out from the ground, with soil and aboveground vegetation intact, and transplanted upwards to plots at higher elevations. This strategy resulted in 18 transplanted samples (namely,

3200T3400, 3200T3600, 3200T3800, 3400T3600, 3400T3800 and 3600T3800, which means 3200 m plots to 3400 m plots, and so on). To simulate the disturbance effect due to soil extraction, triplicate plots at the elevations of 3200, 3400 and 3600 m were mock transplanted and reestablished in place to serve as controls, that is, they were dug out from the ground but put back to the original plots. All plots were randomized in block design, and surrounded by plastic to minimize exchange with neighboring soils.

Soil samples were collected in August 2009 and used for 454 pyrosequencing of 16S rRNA gene amplicon, GeoChip 4.0 (Tu *et al.*, 2014) and environmental variable measurements. Three soil cores with a diameter of 1.5 cm at the depth of 0–20 cm were taken randomly from each plot. Then soil samples were transported back to the laboratory at 4 °C in cooler boxes and sieved with a 2 mm mesh to remove visible grassroots and stones. Soil samples for pyrosequencing and GeoChip experiments were kept at –80 °C until DNA extraction, and soil samples for environmental variable measurements were kept at 4 °C.

Environmental variable measurements

Vegetation variables were measured for a sub-plot within each plot. Vegetation species, density, biomass and average height were recorded using established protocols (Klein *et al.*, 2007) when soils were sampled. Soil temperature was measured at depths of 5 and 20 cm using type-K thermocouples (Campbell Scientific, Logan, Utah, USA) coupled to a CR1000 datalogger, while soil moisture was recorded every 30 min at depths of 5 and 10 cm with time domain reflectometry (TDR) (Model Diviner-2000, Sentek Pty Ltd., Australia). Soil biogeochemical variables were measured as previously described (Yang *et al.*, 2014). In brief, total organic C (TOC) and total N (TN) were measured by a TOC-5000 A analyzer (Shimadzu Corp., Kyoto, Japan) and a Vario EL III Elemental Analyzer (Elementar, Hanau, Germany) using standard protocols (Ryba and Burgess, 2002). Soil NH₄⁺-N and NO₃⁻-N were analyzed with a FIAstar 5000 Analyzer (FOSS, Hillerd, Danmark). The soil C/N ratio was calculated as the TOC to TN ratio.

SOC density can be estimated using TOC and soil bulk density values (Yang *et al.*, 2009). SOC density in the top 20 cm was calculated as follows:

$$\text{SOC} = \sum_{i=1}^n T_i \times \text{BD}_i \times \text{TOC}_i \times (1 - C_i) / 100$$

where SOC, T_i , BD_i , TOC_i and C_i represent SOC density (kg C m^{-2}), layer thickness (cm), bulk density (g cm^{-3}), TOC (g kg^{-1}) and percentage of the fraction > 2 mm of the i th soil layer, respectively.

Ecosystem respiration (Re) and CH₄, N₂O flux estimation

During the growing seasons, the Re was measured by opaque, static, manual stainless steel chambers (Lin *et al.*, 2011) every 7–10 days at 9:00 to 11:00 am from May to September, depending on weather conditions. Based on previous experiments, the measurements obtained between 9:00 and 11:00 am best represented the average daily CO₂ flux (Lin *et al.*, 2011). The chambers were of the same architecture and dimension (40 cm × 40 cm × 40 cm) as previously described (Ma *et al.*, 2006). Chambers were closed for half an hour and gas samples (100 ml) were collected every 10 min using plastic syringes. The CO₂, CH₄ and N₂O concentrations of gas samples were analyzed with a gas chromatograph (HP Series 4890D, Hewlett Packard, USA) within 24 h after gas sampling. The gas chromatograph configurations for analyzing gas concentrations and the methods for calculating gas flux were previously described (Song *et al.*, 2003). The average gas flux in August 2009, when soil samples were collected, was used for statistical analyses in this study.

DNA extraction, 454 pyrosequencing and GeoChip 4.0 experiments

DNA extraction, purification, pyrosequencing and GeoChip 4.0 experiments, and raw data processing methods were described in recent studies (Rui *et al.*, 2015; Yue *et al.*, 2015). While GeoChip experiments were conducted for all samples, pyrosequencing experiments were conducted in most samples except for 3600 and 3600T3800 samples, which cannot be made up due to the lack of DNA from them for sequencing. However, the missing samples did not considerably affect our results. In brief, the Fast DNA Spin kit (MP Biomedical, Carlsbad, CA, USA) was used to extract DNA from 0.5 g soil, following the manufacturer's instructions. The V4-V5 hypervariable regions of 16S rRNA genes were PCR amplified with primers 515F (5'-GTGYCAGCMGCCGCGGTA-3') and 909R (5'-CCCGYCAATTCMTTTRAGT-3'). The pyrosequencing experiments were conducted with a GS FLX system (454 Life Sciences, Branford, CT, USA). The raw sequences were trimmed for quality control using the RDP Pipeline Initial Process (<http://pyro.cme.msu.edu/>). Sequences with low quality (length <300 bp, with ambiguous base 'N', or average base quality score <20) were removed, resulting in 170 161 high-quality and chimera-free reads with an average length of 408 bp. Then sequences were aligned using the Aware Infernal Aligner in the RDP pyrosequencing pipeline, and subjected to chimera check using the Uchime algorithm, and resampling to 2291 sequences per sample. Sequences were clustered by the complete-linkage clustering method incorporated in the RDP platform. Operational taxonomic units were classified using a 97% sequence identity cutoff and singletons were removed. Natural logarithmic

transformation was used before statistical analyses. The GeoChip experiments were performed as described previously (Yang *et al.*, 2013). In brief, DNA was labeled with the fluorescent dye Cy-5 using a random priming method and then purified with the QIA quick purification kit (Qiagen, Valencia, CA, USA) according to the manufacturer's instructions. After checking the dye incorporation on a NanoDrop ND-1000 spectrophotometer (NanoDrop Technologies, Wilmington, DE, USA), DNA was dried in a SpeedVac (ThermoSavant, Milford, MA, USA) at 45 °C for 45 min. Subsequently, labeled DNA was resuspended in 120 µl hybridization solution containing 40% formamide, 3 × SSC, 10 µg of unlabeled herring sperm DNA (Promega, Madison, WI, USA), and 0.1% SDS. The hybridizations were performed with a MAUI hybridization station (Bio-Micro, Salt Lake City, UT, USA) according to the manufacturer's instructions. After washing and drying, microarray was scanned by a NimbleGen MS200 scanner (Roche, Madison, WI, USA) at 633 nm using a laser power of 100% and a photomultiplier tube (PMT) gain of 75%. Signal intensities were subsequently quantified. Briefly, raw data were processed as following: (i) poor-quality spots, those flagged or with a signal to noise ratio (SNR) <2.0, were removed; (ii) at least two valid values of three biological replicates were required for each probe; (iii) relative abundance normalization was applied to all data; and (iv) natural logarithmic transformation was used before statistical analyses.

Statistical analyses

We used a pure random-effects model to estimate the overall weighted mean effect size of cooling on environmental variables and functional genes, based on the method of meta-analysis of response ratios (Hedges *et al.*, 1999). Briefly, for each of the six comparison pairs (3200T3400 vs 3200, 3200T3600 vs 3200, 3200T3800 vs 3200, 3400T3600 vs 3400, 3400T3800 vs 3400 and 3600T3800 vs 3600), the effect size, represented by the response ratio (RR) metric, was calculated as log-proportional change between the means of the transplant and control group. To correct potential bias introduced by small sample sizes, we used the bias-corrected metric RR^A and validated the accuracy of RR^A metric with Geary's test for each comparison pair prior to pooling outcomes (Lajeunesse, 2015). For each pair of transplant-control comparison, the effect size RR^A was calculated as:

$$RR^A = RR + \frac{1}{2} \left[\frac{(s.d._T)^2}{N_T \bar{X}_T^2} - \frac{(s.d._C)^2}{N_C \bar{X}_C^2} \right]$$

$$= \ln \frac{\bar{X}_T}{\bar{X}_C} + \frac{1}{2} \left[\frac{(s.d._T)^2}{N_T \bar{X}_T^2} - \frac{(s.d._C)^2}{N_C \bar{X}_C^2} \right]$$

where \bar{X}_T and \bar{X}_C represent the mean of the variable in transplant and control group; s.d._T and s.d._C

represent standard deviations of transplant and control group; N_T and N_C represent the number of replicates of transplant and control group.

The sample sizes and standard deviations were used to help quantify the sampling variability in the effect size RR^A within each comparison. For each comparison, the variance of RR^A (denoted by v) was calculated as:

$$v = \frac{(\text{s.d.}_T)^2}{N_T \bar{X}_T^2} + \frac{(\text{s.d.}_C)^2}{N_C \bar{X}_C^2} + \frac{1}{2} \left[\frac{(\text{s.d.}_T)^4}{N_T^2 \bar{X}_T^4} + \frac{(\text{s.d.}_C)^4}{N_C^2 \bar{X}_C^4} \right]$$

The random between-comparisons variance component was calculated as:

$$\hat{\sigma}_\lambda^2 = \frac{Q - (k - 1)}{\sum_{i=1}^k w_i - \frac{\sum_{i=1}^k w_i^2}{\sum_{i=1}^k w_i}}$$

Where

$$Q = \sum_{i=1}^k w_i (RR_i^A)^2 - \frac{(\sum_{i=1}^k w_i RR_i^A)^2}{\sum_{i=1}^k w_i}$$

and $w_i = 1/v_i$, k is the number of comparisons. In this study, $k = 6$.

The within-comparison variance and the between-comparisons variance component were converted into weights to help minimize the influence of comparisons with low statistical power when analyzing and pooling multiple study outcomes. The weight of each comparison was determined as:

$$w_i^* = \frac{1}{v_i + \hat{\sigma}_\lambda^2}$$

The weighted mean effect size (\overline{RR}^*) and its standard error were then determined based on the effect size and the weight of each comparison:

$$\overline{RR}^* = \frac{\sum_{i=1}^k w_i^* RR_i^A}{\sum_{i=1}^k w_i^*}$$

$$\text{s.e.}(\overline{RR}^*) = \sqrt{\frac{1}{\sum_{i=1}^k w_i^*}}$$

In addition, we examined the Pearson correlation between the effect size and the cooling degree, that is, the air temperature difference. For variables whose effect sizes were positively and significantly ($P < 0.05$) correlated with the cooling degree, the average change per °C cooling was calculated as the mean of the absolute change divided by the cooling degree in each comparison.

To test the cooling effects on the microbial community structure, three different complementary non-parametric dissimilarity analyses for multivariate data were used: analysis of similarity (ANOSIM) (Clarke, 1993), non-parametric multivariate analysis of variance (adonis) (Anderson, 2001) using distance matrices, and multi-response permutation procedure (MRPP) (McCune *et al.*, 2002). The Detrended Correspondence Analysis (DCA) (Hill and H.G.

Gauch, 1980) was used to visualize the difference of microbial community structure between cooling sample and its corresponding control.

The Canonical Correspondence Analysis (CCA) was performed to determine the most significant environmental variables linking to microbial functional gene structure (Ramette and Tiedje, 2007). Based on the variance inflation factors (VIF) values, redundant variables with $VIF > 20$ were removed from the CCA model. A partial-CCA based Variation Partitioning Analysis (VPA) was then performed to determine the contribution of different groups of variables. To determine the effects of microbial functional gene or environmental variables on greenhouse gas flux, partial Mantel tests (Smouse *et al.*, 1986) were performed in which the co-varying effects between microbial functional gene structure and environmental variables were controlled. Bray-Curtis coefficient and Euclidean distance were used to construct dissimilarity matrices of microbial communities and environmental variables, respectively.

Results and Discussion

For the six comparison pairs; 3200T3400 vs 3200, 3200T3600 vs 3200, 3200T3800 vs 3200, 3400T3600 vs 3400, 3400T3800 vs 3400 and 3600T3800 vs 3600, air temperature decreased by 0.26, 1.00, 2.09, 0.74, 1.83 and 1.09 °C, respectively (Supplementary Table 1). We calculated the weighted mean effect sizes of cooling on environmental variables. The temperature of the top 5 cm and lower 20 cm of the soil profile significantly decreased, with mean effect size of -0.22 (± 0.05 s.e.) and -0.23 (± 0.05). The effect size of soil temperature of the top 5 cm profile was significantly correlated with the cooling degree (Pearson's $r = 0.82$, $P = 0.04$) across the six comparisons. Further analysis showed that soil temperature of the top 5 cm profile decreased by 2.41 (± 0.48) °C per °C decrease in air temperature. Soil pH marginally significantly decreased under cooling, with an effect size of -0.02 (± 0.01), while soil moisture remained largely unchanged. In addition, plant species richness remained unchanged, but the plant biomass and total plant coverage (TCOV) responses were negative, with effect sizes of -0.53 (± 0.17) and -0.07 (± 0.04), respectively (Figure 1). These results were consistent with known cooling effects (Campbell and McAndrews, 1993), demonstrating that the field simulation we performed was reliable. Also, they directly contrasted observations from warming-based studies, in which plant productivity increased in tundras, grasslands and forests (Lal, 2005; Zhou *et al.*, 2012; Natali *et al.*, 2014).

Amplicon sequencing of the 16S rRNA gene was combined with GeoChip 4.0 (Tu *et al.*, 2014) to examine soil microbial communities. GeoChip 4.0 is a microarray containing probes to detect ~141 995 gene sequences from 410 gene families derived from




Figure 1 Effects of climate cooling on various environmental variables. The weighted mean effect size was calculated by the method of meta-analysis of response ratio. Error bars are at the 95% confidence level. The significance level is indicated by asterisk: *** $P < 0.001$, ** $P < 0.050$, and * $P < 0.100$. For the environmental variables, Biomass—plant biomass; TCOV—total plant coverage; Species—plant species richness; TOC_10 and TOC_20—total soil organic C at the 0–10 cm depth and 10–20 cm depth; TN_10 and TN_20—total soil N at the 0–10 cm depth and 10–20 cm depth; CN_10 and CN_20—the ratio of TOC to TN ratio at the 0–10 cm depth and 10–20 cm depth; NO₃_10—soil nitrate at the 0–10 cm depth; and NH₃_10—soil ammonia at the 0–10 cm depth.

bacteria, archaea or fungi, including genes associated with nitrogen (N), carbon (C), sulfur (S) and phosphorus (P) cycling, metal resistance, organic remediation, and other processes. DCA and dissimilarity tests showed that, for each transplant-control comparison, both taxonomic composition and functional gene structure of microbial communities in transplanted soils differed compared with in-place controls (Table 1; Supplementary Figure 1; $P < 0.05$). In addition, the Mantel test showed a very weak but significant correlation ($r = 0.18$, $P = 0.03$) between the variations of 16S rRNA gene data and GeoChip data across samples, unveiling some consistency between microbial taxonomic composition and functional gene structure. Furthermore, CCA indicated that microbial functional gene structure correlated ($P < 0.01$) with changes in soil temperature, pH, moisture, NH₃-N, TOC, TN, C/N ratio, plant biomass, plant species richness and TCOV (Supplementary Figure 2). A total of 63.6% of the variation in soil microbial functional gene structure could be explained by soil climatic variables (13.1%), soil geochemical variables (26.9%), plant variables (8.8%) and their interactions (Supplementary Figure 2), as indicated by VPA. It is noted that microbial functional gene structure of 3200T3400 was more similar to that of 3200T3800 rather than 3200T3600 (Supplementary Figure 1). A possible explanation is that some environmental variables were more similar between 3400 m and 3800 m sites than with the 3600 m site, such as plant biomass and pH, which may have important roles in shaping microbial functional gene structure. In fact, the

Table 1 Effects of cooling on microbial community compositions and functional genes with three different statistical approaches

| Data sets | adonis | | ANOSIM | | MRPP | |
|-------------------|--------|---------|--------|---------|----------|---------|
| | F | P-value | R | P-value | δ | P-value |
| 16S sequencing | 1.763 | 0.005 | 0.363 | 0.008 | 0.695 | 0.009 |
| All GeoChip genes | 3.521 | 0.018 | 0.160 | 0.020 | 0.232 | 0.014 |
| C cycling genes | 3.537 | 0.015 | 0.166 | 0.024 | 0.239 | 0.012 |
| N cycling genes | 3.557 | 0.019 | 0.174 | 0.012 | 0.233 | 0.010 |

changes of most environmental variables and microbial functional structure were not related to the degree of cooling.

Carbon decomposition and nutrient cycling were both impacted by cooling. Both the relative abundance and α -diversity (Simpson index) of genes associated with chemically labile C decomposition were markedly decreased in the transplanted plots, as indicated by the weighted mean effect sizes. These included *apu*, *cda* and *glucoamylase* genes for starch decomposition, *xylA* and *mannanase* genes for hemi-cellulose decomposition, *exochitinase* genes for chitin decomposition, and *pectinase* genes for pectin decomposition (Figure 2; Supplementary Figure 3), which have previously been shown to increase in response to warming (Zhou *et al.*, 2012). The most notably decreased C decomposition genes included *cda* genes derived from *Vibrio harveyi* HY01 and *Roseburia intestinalis* L1-82, *mannanase* genes derived from *Cellvibrio japonicus* Ueda107 and *Streptomyces sviceps* ATCC 29083 (Supplementary Figure 4). Consistently, the 16S rRNA gene amplicon data showed significant decreases in the relative abundances of the *Streptomyces* genus. In fact, the relative abundance of *Gammaproteobacteria*, *Firmicutes* and *Actinobacteria* were significantly decreased under cooling (Supplementary Figure 5). Interestingly, the functional genes responsible for the decomposition of chemically recalcitrant C, such as lignin and aromatics, which account for about 95% of topsoil organic C in Tibetan grasslands (Genxu *et al.*, 2002), were unchanged. Chemically recalcitrant C was previously thought to be stable in soil and used by microorganisms only when labile C substrates are exhausted (Lützow *et al.*, 2006). However, there is accumulating evidence that microorganisms decompose chemically recalcitrant C under suitable conditions, and that not all old and stable C compounds are as persistent as once considered (Kleber *et al.*, 2011; Lehmann and Kleber, 2015). Soil heterotrophic respiration was consistently steady at our study site (Hu *et al.*, 2008), which would be expected under consistent rates of C decomposition.

The abundance of most nutrient cycling genes either decreased or remained unchanged except *ureC*. For N cycling genes, the abundances of *amoA* genes decreased (Supplementary Figure 6a), which




Figure 2 Effects of climate cooling on relative abundances of C decomposition genes. The weighted mean effect size was calculated by the method of meta-analysis of response ratio. Error bars are at the 95% confidence level. The significance level is indicated by asterisk: *** $P < 0.001$, ** $P < 0.050$ and * $P < 0.100$.

Table 2 The influence of environmental variables and microbial functional genes on greenhouse gas emission by partial Mantel test

| Effects of | Functional gene groups ^a | | Soil variables | | Plant variables | |
|-----------------------|-------------------------------------|------------------------------|-----------------|-----------------|-----------------|----------------|
| | Soil variables ^b | Plant variables ^c | Functional gene | Plant variables | Functional gene | Soil variables |
| CO ₂ flux | -0.310 ^d | 0.045 | 0.615*** | 0.458*** | 0.406*** | 0.198** |
| CH ₄ flux | 0.159* | 0.143* | -0.002 | -0.031 | 0.213* | 0.226* |
| N ₂ O flux | 0.331** | 0.345** | -0.039 | 0.116 | 0.039 | 0.050 |

^aFunctional gene groups were carbon decomposition genes for CO₂ flux, *mmoX* genes for CH₄ flux and *norB* genes for N₂O flux. ^bSoil variables included for testing the relationship with CO₂ or CH₄ flux: soil temperature, soil moisture, pH, TOC, C/N ratio. Soil variables included for testing the relationship with N₂O flux: soil temperature, soil moisture, pH, NO₃-N, NH₃-N, TN, C/N ratio. ^cPlant variables were plant biomass, total coverage of plants (TCOV) and plant species richness. ^dThe *r*-value of Partial Mantel test is shown. Significance: *** $P < 0.001$, ** $P < 0.050$, * $P < 0.100$.

were derived from both ammonia-oxidizing archaea (AOA) and ammonia-oxidizing bacteria (AOB) (Wessén *et al.*, 2011). The most notably decreased *amoA* genes included those derived from uncultured archaea, *Edwardsiella tarda* EIB202 and *Pseudomonas putida* KT2440 (Supplementary Figure 6b), which was consistent with the amplicon sequencing data showing that the relative abundance of *Gammaproteobacteria* significantly decreased under cooling (Supplementary Figure 5). The abundance of *norB* genes encoding an enzyme to convert NO to N₂O in both nitrification and denitrification processes decreased, too (Supplementary Figure 6a, $P < 0.05$). However, we could not determine which process the *norB* genes belong to.

The ecosystem respiration (Re, represented by the measured CO₂ flux) response was negative, with an effect size of -0.30 (±0.15) (Figure 1), similar to a recent cooling experiment in forests (Luan *et al.*, 2014). However, partial Mantel tests demonstrated an insignificant correlation between C decomposition gene abundance and CO₂ flux when soil or plant variables were controlled (Table 2). Rather, soil and plant variables showed a correlation ($P < 0.001$) with CO₂ flux when controlled for C decomposition genes.

Specifically, plant biomass, TCOV, plant species richness, soil temperature, pH and TOC at a depth of 0–10 cm positively correlated with CO₂ flux, whereas soil moisture negatively correlated (Figure 3a; Supplementary Figure 7). The forward stepwise regression method was applied to select variables for predicting CO₂ flux based on Akaike information criterion (AIC). The final linear model included two variables of air temperature and TCOV, which was able to explain ~90% ($P < 0.001$) of the variation in CO₂ flux (Supplementary Table 2). Furthermore, partial correlation analysis showed that air temperature positively and significantly correlated with CO₂ flux ($r = 0.75$, $P = 0.05$) when soil and plant variables were controlled. The decrease in air temperature (that is, cooling) could be the major driver of Re decrease, since often biological activities are controlled by temperature (Brown *et al.*, 2004). The positive correlation ($r^2 = 0.57$, $P < 0.001$) between plant biomass and CO₂ flux (Figure 3a) together with the partial Mantel test results (Table 2) suggested that a reduction in ecosystem respiration might be primarily attributed to decreased autotrophic respiration under cooling. Consistently, plant autotrophic respiration was




Figure 3 (a) Correlation between CO₂ efflux and plant biomass and (b) correlation between the change of SOC density and temperature difference. Linear regression lines and their 95% confidence limits (shaded) are shown. SOC Change was calculated as the difference in SOC density between transplant soils (cooling) and their corresponding controls. Negative values of SOC changes indicate the decrease of SOC under cooling.

shown to be the main explanatory factor of variations in ecosystem respiration on the Tibetan alpine grasslands (Hu *et al.*, 2008). Further, soil TOC at depths of 0–10 and 10–20 cm were decreased, with effect sizes of $-0.05 (\pm 0.02)$ and $-0.06 (\pm 0.01)$, respectively (Figure 1). The effect size of soil TOC at depths of 10–20 cm strongly and significantly correlated with the cooling degree (Pearson's $r = 0.91$, $P = 0.01$), and soil TOC at depths of 10–20 cm decreased by $2.55 (\pm 0.44)$ mg g⁻¹ per °C decrease in air temperature.

TN contents at the depths of 0–10 cm decreased, with an effect size of $-0.03 (\pm 0.01)$, though the C/N ratio remained largely unchanged (Figure 1). The N₂O fluxes were either decreased (*t*-test, $P < 0.05$) or unchanged by cooling (Supplementary Table 1), which was consistent with previous studies showing that the effect of temperature on N₂O emission is generally positive in non-wetland soils (Smith, 1997). N₂O flux was significantly correlated ($P < 0.006$) with the abundance of *norB* genes when soil or plant variables were controlled (Table 2), suggesting that N₂O flux in these sites was microbially-driven (Singh *et al.*, 2010). The grassland appeared to be a CH₄ sink rather than a source (Supplementary Table 1), implying that CH₄ oxidation could have a more important role in CH₄ flux than CH₄ production. In support of this, *mcrA* abundance, associated with CH₄ production, was not correlated to CH₄ flux. In contrast, the abundance of *mmoX*, associated with CH₄ oxidation, including those derived from methanotrophs possessing the soluble methane monooxygenase such as *Methylocella spp.* and *Methylomonas spp.*, was correlated with CH₄ flux ($P < 0.100$) (Table 2). Methane flux remained unchanged under cooling (Supplementary Table 1; Figure 1). In comparison, previous studies showed that soil CH₄ uptake increased (Peterjohn *et al.*, 1994; Hart, 2006) or remained unchanged (Torn and Harte, 1996; Rustad and Fernandez, 1998)

under warming. As CH₄ production is an anaerobic process, soil moisture might have stronger influence on soil CH₄ fluxes than soil temperature (Bowden *et al.*, 1998; Wei *et al.*, 2015).

To determine whether the lower soil C was related to cooling, we calculated the correlation between the decrease of SOC density (kg C m⁻²) under cooling and the temperature difference. SOC loss in the top 20 cm was correlated ($r^2 = 0.67$, $P < 0.001$) with the degree of cooling (Figure 3b). Although there were considerable variations in the SOC changes at certain temperature differences, all of the SOC changes are exclusively negative. The SOC decrease for the extreme of the altitudinal transplant experiment (from 3200 to 3800 m) was the largest. When the data of this pair was removed, the linear and significant correlation between SOC change and temperature difference still holds ($r^2 = 0.44$, $P < 0.001$). The decrease of SOC was estimated to be 0.088 kg C m⁻² °C⁻¹, ranging from 0.071 to 0.105 kg C m⁻² °C⁻¹ at the 95% confidence level. By extrapolating these observations to the entire Tibetan plateau, it is estimated that a 1°C decrease in temperature would result in a roughly 0.1 Pg SOC loss plateau-wide, which accounts for about 0.3% of the SOC stored in the Tibetan plateau (Genxu *et al.*, 2002). However, obviously, this extrapolation should be considered a rough estimation due to the heterogeneity of the plateau, which will likely result in substantial variance in responses of the soil microbial community.

Various studies have suggested that climate warming would result in soil C loss through increased soil respiration (Frey *et al.*, 2013). Therefore, warming will feedback positively on climate change. However, recent studies showed contradictory results. For example, long-term warming has increased net ecosystem C storage on Arctic tundra, while the soil carbon storage remained unchanged (Sistla *et al.*, 2013). In a downward

transplant warming experiment, we also found that SOC remained unchanged under warming (Yue *et al.*, 2015). In the current study, however, we found that SOC was significantly decreased under cooling (Figure 3b; Supplementary Table 1). A possible explanation is that the decrease in soil C inputs through litter and root exudates was not offset by the decrease in soil C consumption through respiration. Plant biomass decreased under cooling, suggesting a significant decrease in soil C inputs. In contrast, the abundance and presence of chemically recalcitrant C decomposition genes were largely unchanged. Persistent microbially-driven turnover of chemically recalcitrant carbon accelerates soil C loss, since the chemically recalcitrant C constitutes the majority of topsoil C in Tibetan grasslands (Genxu *et al.*, 2002). Even though soil C loss by climate cooling was similar to that observed under warming (Cox *et al.*, 2000; Nie *et al.*, 2013), we propose that the underlying mechanisms are fundamentally different.

Our results highlight the importance of understanding the role of plant and microbial communities in providing ecosystem feedbacks to climate change. To improve the prediction of ecosystem feedbacks to climate cooling, it is important to consider various types of feedback mechanisms resulting from the changes in plant and microbial community ecology. Our findings also provide insights into ecological consequences of numerous cooling events in earth's history.

Data accessibility

GeoChip data are available online (www.ncbi.nlm.nih.gov/geo/) with the accession number GSE52425. 454 sequencing data are available from the MG-RAST database (accession no. 4565936.3 to 4565973.3).

Conflict of Interest

The authors declare no conflict of interest.

Acknowledgements

We thank Haibei Research Station staff for sampling, Hao Yu for GeoChip assistance. This research was supported by grants to Yunfeng Yang from the National Key Basic Research Program of China (2013CB956601), the Strategic Priority Research Program of the Chinese Academy of Sciences (XDB15010102), and National Science Foundation of China (41471202), to Shiping Wang from the National Basic Research Program (2013CB956000) and National Science Foundation of China (41230750), and to Jizhong Zhou from the National Science Foundation of China (41430856), and the United States Department of Energy, Biological Systems Research on the Role of Microbial Communities in Carbon Cycling Program (DE-SC0010715).

Author contributions

This study was conceived and led by SW, JZ and YY; YH, QL, XL and YY carried out GeoChip experiments and environmental measurements; LW and HY performed the analytical work; LW, YY and JZ wrote the manuscript with the help from SW, ZH, JVN, LH and JG.

References

- Alley RB, Marotzke J, Nordhaus W, Overpeck J, Peteet D, Pielke R *et al.* (2003). Abrupt climate change. *Science* **299**: 2005–2010.
- Anderson MJ. (2001). A new method for non-parametric multivariate analysis of variance. *Austral Ecol* **26**: 32–46.
- Bowden RD, Newkirk KM, Rullo GM. (1998). Carbon dioxide and methane fluxes by a forest soil under laboratory-controlled moisture and temperature conditions. *Soil Biol Biochem* **30**: 1591–1597.
- Breeuwer A, Heijmans MM, Robroek BJ, Berendse F. (2010). Field simulation of global change: transplanting northern bog mesocosms southward. *Ecosystems* **13**: 712–726.
- Brown JH, Gillooly JF, Allen AP, Savage VM, West GB. (2004). Toward a metabolic theory of ecology. *Ecology* **85**: 1771–1789.
- Campbell ID, McAndrews JH. (1993). Forest disequilibrium caused by rapid Little Ice Age cooling. *Nature* **366**: 336–338.
- Clarke KR. (1993). Non-parametric multivariate analyses of changes in community structure. *Austr J Ecol* **18**: 117–117.
- Cox PM, Betts RA, Jones CD, Spall SA, Totterdell IJ. (2000). Acceleration of global warming due to carbon-cycle feedbacks in a coupled climate model. *Nature* **408**: 184–187.
- Doran PT, Prisco JC, Lyons WB, Walsh JE, Fountain AG, McKnight DM *et al.* (2002). Antarctic climate cooling and terrestrial ecosystem response. *Nature* **415**: 517–520.
- Easterling DR, Wehner MF. (2009). Is the climate warming or cooling. *Geophys Res Lett* **36**: 8.
- Frey SD, Lee J, Melillo JM, Six J. (2013). The temperature response of soil microbial efficiency and its feedback to climate. *Nat Clim Change* **3**: 395–398.
- Genxu W, Ju Q, Guodong C, Yuanmin L. (2002). Soil organic carbon pool of grassland soils on the Qinghai-Tibetan Plateau and its global implication. *Sci Total Environ* **291**: 207–217.
- Hart SC. (2006). Potential impacts of climate change on nitrogen transformations and greenhouse gas fluxes in forests: a soil transfer study. *Global Change Biol* **12**: 1032–1046.
- Hedges LV, Gurevitch J, Curtis PS. (1999). The meta-analysis of response ratios in experimental ecology. *Ecology* **80**: 1150–1156.
- Hill MO, H.G. Gauch J. (1980). Detrended correspondence analysis: an improved ordination technique. *Vegetatio* **42**: 47–58.
- Hu QW, Wu Q, Cao GM, Li D, Long RJ, Wang YS. (2008). Growing season ecosystem respirations and associated

- component fluxes in two alpine meadows on the Tibetan Plateau. *J Integr Plant Biol* **50**: 271–279.
- Ji F, Wu Z, Huang J, Chassignet EP. (2014). Evolution of land surface air temperature trend. *Nat Clim Change* **4**: 462–466.
- Kleber M, Nico PS, Plante A, Filley T, Kramer M, Swanston C *et al.* (2011). Old and stable soil organic matter is not necessarily chemically recalcitrant: implications for modeling concepts and temperature sensitivity. *Global Change Biol* **17**: 1097–1107.
- Klein JA, Harte J, Zhao X-Q. (2007). Experimental warming, not grazing, decreases rangeland quality on the Tibetan Plateau. *Ecol Appl* **17**: 541–557.
- Lajeunesse MJ. (2015). Bias and correction for the log response ratio in ecological meta-analysis. *Ecology* **96**: 2056–2063.
- Lal R. (2005). Forest soils and carbon sequestration. *Forest Ecol Manage* **220**: 242–258.
- Lehmann J, Kleber M. (2015). The contentious nature of soil organic matter. *Nature* **528**: 60–68.
- Lin X, Zhang Z, Wang S, Hu Y, Xu G, Luo C *et al.* (2011). Response of ecosystem respiration to warming and grazing during the growing seasons in the alpine meadow on the Tibetan plateau. *Agric Forest Meteorol* **151**: 792–802.
- Luan J, Liu S, Chang SX, Wang J, Zhu X, Liu K *et al.* (2014). Different effects of warming and cooling on the decomposition of soil organic matter in warm-temperate oak forests: a reciprocal translocation experiment. *Biogeochemistry* **121**: 551–564.
- Lucht W, Prentice IC, Myneni RB, Sitch S, Friedlingstein P, Cramer W *et al.* (2002). Climatic control of the high-latitude vegetation greening trend and Pinatubo effect. *Science* **296**: 1687–1689.
- Luo Y. (2007). Terrestrial carbon-cycle feedback to climate warming. *Annu Rev Ecol Evol Syst* **38**: 683–712.
- Lützw Mv, Kögel-Knabner I, Ekschmitt K, Matzner E, Guggenberger G, Marschner B *et al.* (2006). Stabilization of organic matter in temperate soils: mechanisms and their relevance under different soil conditions—a review. *Eur J Soil Sci* **57**: 426–445.
- Lyubushin AA, Klyashtorin LB. (2012). Short term global dT prediction using (60-70)-years periodicity. *Energy Environ* **23**: 75–86.
- Ma X, Wang S, Wang Y, Jiang G, Nyren P. (2006). Short-term effects of sheep excrement on carbon dioxide, nitrous oxide and methane fluxes in typical grassland of Inner Mongolia. *N Zeal J Agric Res* **49**: 285–297.
- McAnena A, Flögel S, Hofmann P, Herrle J, Griesand A, Pross J *et al.* (2013). Atlantic cooling associated with a marine biotic crisis during the mid-Cretaceous period. *Nat Geosci* **6**: 558–561.
- McCune B, Grace JB, Urban DL. (2002). *Analysis of ecological communities*, vol. 28. MjM software design Glenden Beach, OR.
- Natali SM, Schuur EAG, Webb EE, Pries CEH, Crummer KG. (2014). Permafrost degradation stimulates carbon loss from experimentally warmed tundra. *Ecology* **95**: 602–608.
- Nie M, Pendall E, Bell C, Gasch CK, Raut S, Tamang S *et al.* (2013). Positive climate feedbacks of soil microbial communities in a semi-arid grassland. *Ecol Lett* **16**: 234–241.
- Petchey OL, McPhearson PT, Casey TM, Morin PJ. (1999). Environmental warming alters food-web structure and ecosystem function. *Nature* **402**: 69–72.
- Peterjohn WT, Melillo JM, Steudler PA, Newkirk KM, Bowles FP, Aber JD. (1994). Responses of trace gas fluxes and N availability to experimentally elevated soil temperatures. *Ecol Appl* **4**: 617–625.
- Qiu J. (2008). China: the third pole. *Nat News* **454**: 393–396.
- Ramette A, Tiedje JM. (2007). Multiscale responses of microbial life to spatial distance and environmental heterogeneity in a patchy ecosystem. *Proc Natl Acad Sci USA* **104**: 2761–2766.
- Rui J, Li J, Wang S, An J, Liu W-t, Lin Q *et al.* (2015). Responses of bacterial communities to simulated climate changes in alpine meadow soil of the Qinghai-Tibet Plateau. *Appl Environ Microbiol* **81**: 6070–6077.
- Rustad LE, Fernandez IJ. (1998). Experimental soil warming effects on CO₂ and CH₄ flux from a low elevation spruce–fir forest soil in Maine, USA. *Global Change Biol* **4**: 597–605.
- Ryba SA, Burgess RM. (2002). Effects of sample preparation on the measurement of organic carbon, hydrogen, nitrogen, sulfur, and oxygen concentrations in marine sediments. *Chemosphere* **48**: 139–147.
- Singh BK, Bardgett RD, Smith P, Reay DS. (2010). Microorganisms and climate change: terrestrial feedbacks and mitigation options. *Nat Rev Microbiol* **8**: 779–790.
- Sistla SA, Moore JC, Simpson RT, Gough L, Shaver GR, Schimel JP. (2013). Long-term warming restructures Arctic tundra without changing net soil carbon storage. *Nature* **497**: 615–618.
- Smith K. (1997). The potential for feedback effects induced by global warming on emissions of nitrous oxide by soils. *Global Change Biol* **3**: 327–338.
- Smouse PE, Long JC, Sokal RR. (1986). Multiple regression and correlation extensions of the Mantel test of matrix correspondence. *Syst Zool* **35**: 627–632.
- Song C, Yan B, Wang Y, Wang Y, Lou Y, Zhao Z. (2003). Fluxes of carbon dioxide and methane from swamp and impact factors in Sanjiang Plain, China. *Chin Sci Bull* **48**: 2749–2753.
- Torn MS, Harte J. (1996). Methane consumption by montane soils: implications for positive and negative feedback with climatic change. *Biogeochemistry* **32**: 53–67.
- Tu Q, Yu H, He Z, Deng Y, Wu L, Nostrand JD *et al.* (2014). GeoChip 4: a functional gene-array-based high-throughput environmental technology for microbial community analysis. *Mol Ecol Resour* **14**: 914–928.
- Turner J, Lu H, White I, King JC, Phillips T, Hosking JS *et al.* (2016). Absence of 21st century warming on Antarctic Peninsula consistent with natural variability. *Nature* **535**: 411–415.
- Vanhala P, Karhu K, Tuomi M, Björklöf K, Fritze H, Hyvärinen H *et al.* (2011). Transplantation of organic surface horizons of boreal soils into warmer regions alters microbiology but not the temperature sensitivity of decomposition. *Global Change Biol* **17**: 538–550.
- Wei D, Wang Y, Wang Y. (2015). Considerable methane uptake by alpine grasslands despite the cold climate: in situ measurements on the central Tibetan plateau, 2008–2013. *Global Change Biol* **21**: 777–788.
- Wessén E, Söderström M, Stenberg M, Bru D, Hellman M, Welsh A *et al.* (2011). Spatial distribution of ammonia-oxidizing bacteria and archaea across a 44-hectare farm related to ecosystem functioning. *ISME J* **5**: 1213–1225.

- Yang Y, Fang J, Tang Y, Ji C, Zheng C, He J *et al.* (2008). Storage, patterns and controls of soil organic carbon in the Tibetan grasslands. *Global Change Biol* **14**: 1592–1599.
- Yang Y, Fang J, Smith P, Tang Y, Chen A, Ji C *et al.* (2009). Changes in topsoil carbon stock in the Tibetan grasslands between the 1980s and 2004. *Global Change Biol* **15**: 2723–2729.
- Yang Y, Wu L, Lin Q, Yuan M, Xu D, Yu H *et al.* (2013). Responses of the functional structure of soil microbial community to livestock grazing in the Tibetan alpine grassland. *Global Change Biol* **19**: 637–648.
- Yang Y, Gao Y, Wang S, Xu D, Yu H, Wu L *et al.* (2014). The microbial gene diversity along an elevation gradient of the Tibetan grassland. *ISME J* **8**: 430–440.
- Yue H, Wang M, Wang S, Gilbert JA, Sun X, Wu L *et al.* (2015). The microbe-mediated mechanisms affecting topsoil carbon stock in Tibetan grasslands. *ISME J* **9**: 2012–2020.
- Zhou JZ, Xue K, Xie JP, Deng Y, Wu LY, Cheng XH *et al.* (2012). Microbial mediation of carbon-cycle feedbacks to climate warming. *Nat Clim Change* **2**: 106–110.

Supplementary Information accompanies this paper on The ISME Journal website (<http://www.nature.com/ismej>)