UC Davis

UC Davis Previously Published Works

Title

Review of the Pathways Involved in the Osteogenic Differentiation of Adipose-Derived Stem Cells.

Permalink

<https://escholarship.org/uc/item/8b876731>

Journal

The Journal of craniofacial surgery, 30(3)

ISSN

1049-2275

Authors

Asserson, Derek B

Orbay, Hakan

Sahar, David E

Publication Date

2019-05-01

DOI

10.1097/scs.00000000000005447

Peer reviewed

Review of the Pathways Involved in the Osteogenic Differentiation of Adipose-Derived Stem Cells

Derek B. Asserson, BS, Hakan Orbay, MD, PhD, and David E. Sahar, MD

Abstract: Grafts and prosthetic materials used for the repair of bone defects are often accompanied by comorbidity and rejection. Therefore, there is an immense need for novel approaches to combating the issues surrounding such defects. Because of their accessibility, substantial proportion, and osteogenic differentiation potential, adipose-derived stem cells (ASCs) make for an ideal source of bone tissue in regenerative medicine. However, efficient induction of ASCs toward an osteoblastic lineage in vivo is met with challenges, and many signaling pathways must come together to secure osteoblastogenesis. Among them are bone morphogenic protein, wingless-related integration site protein, Notch, Hedgehog, fibroblast growth factor, vascular endothelial growth factor, and extracellular regulated-signal kinase. The goal of this literature review is to conglomerate the present research on these pathways to formulate a better understanding of how ASCs are most effectively transformed into bone in the context of tissue engineering.

Key Words: Adipose-derived stem cells, bone regeneration, osteogenic differentiation, tissue engineering

(*J Craniofac Surg* 2019;30: 703–708)

In 2001, 12,700 bone grafts were performed to repair craniofacial defects at a cost of over \$549 million in children alone.¹ Another 1.3–1.5 million grafts were constructed in 2010 to fix defects of the cranium, sternum, ribs, and extremities.² It is said that bone grafting is also required in 1 out of every 4 dental implants.³ This reach seems to be a transnational occurrence, as graft materials are approved all over the world, with some countries placing great importance on the procedures in training programs.^{2,4}

Current treatment methods using autologous bone, allogenic transplants, and prosthetic materials all carry significant drawbacks, such as donor-site morbidity, immune rejection, and extrusion and infection, while failing to meet the enormous and ever-increasing demand for reconstruction of skeletal deficiencies.⁵ This gap has provided impetus to develop regenerative medicine that is being relied on more often to deal with disease and trauma. Engineering bone tissue was only made possible with the discovery and utilization of stem cells that are capable of self-renewal and differentiation

into a bevy of lineages.⁶ The implementation of stem cells for the purpose of bone regeneration potentially eliminates the aforementioned problems associated with autologous and allogenic grafts.⁵

Because of their high differentiation potential and low morbidity during harvesting, bone marrow stem cells (BMSCs) have been considered the gold standard in bone tissue engineering.^{7–9} However, low cellular yields, as well as extremely painful extraction, have pushed research in a novel direction, one that places fat at the core of regenerative medicine.^{10–13} Adipose-derived stem cells (ASCs) have similar transcription profiles to BMSCs for genes induced in stem-cell phenotypes, and are identified for their ability to differentiate into a vast array of lineages, including adipogenic, chondrogenic, myogenic, neurogenic, and osteogenic forms.^{14,15} Roughly 5000 fibroblast colony-forming units (CFU-F) are obtained per gram of adipose tissue, and ASCs constitute 2% of the nucleated cells in processed lipoaspirate.¹⁶ Compared with only 100 to 1000 CFU-F per milliliter of bone marrow, adipose tissue makes for an excellent source of mesenchymal stem cells (MSCs).¹⁶

Secondary to ASC harvesting, scaffolds are employed to ensure proper behavior of the stem cells by generating controlled niches and delivering appropriate biomaterials.^{17–19} Optimal bone regeneration is achieved with certain physical and biological characteristics of the scaffold, especially cell attachment sites, in which the ideal osteogenic scaffold is a porous, biodegradable, three-dimensional structure.^{20–23} The culture medium of ASCs also largely helps determine differentiation potential toward certain lineages.²⁴

Although several papers demonstrate the osteogenic differentiation of ASCs, the molecular mechanisms by which the ASCs differentiate into osteoprogenitor cells are not as easily understood and no systematic review of them exists in the literature. This review serves to elucidate those mechanisms through research endeavors that have successfully turned ASCs into bone. The pertinent papers are classified according to main signaling pathways in Table 1.

METHODS

A search of the PubMed/Medline database was performed with the search terms “adipose-derived stem cells” OR “ASCs” AND “osteogenic differentiation.” Initial results yielded 256 papers, which was reduced to 179 after title screening. When subsequently screened for content, 14 backgrounds (some of which were not used) and 29 experimental papers were left. An additional 62 investigations found via connected articles were incorporated to further substantiate the literature review.

RESULTS

Osteoblastogenesis or osteoblast differentiation during development is controlled by a complex network at both the transcriptional level and extracellular signaling pathways.⁵ The key players in the transcriptional control of osteoblast differentiation are Runt-related family 2 (RUNX2), formerly called Cbfa1,^{25–29} and the late

From the Department of Surgery, Surgical Bioengineering Laboratory, University of California Davis Medical Center, Sacramento, CA.

Received November 2, 2018.

Accepted for publication February 4, 2019.

Address correspondence and reprint requests to Hakan Orbay, MD, PhD, 4625 2nd Avenue, Research Building II, Room 3004 Sacramento, CA 95817; E-mail: hakanorbay78@hotmail.com; Derek B. Asserson, BS, University of California Davis, Sacramento, CA;

E-mail: derek.asserson2615@cnsu.edu

The authors report no conflicts of interest.

Copyright © 2019 by Mutaz B. Habal, MD

ISSN: 1049-2275

DOI: 10.1097/SCS.0000000000005447

TABLE 1. Classification of Signaling Pathways by Paper

Author	Year	Signaling Pathway
Arnsford et al	2009	Wnt
Bandyopadhyay et al	2006	BMP
Behr et al	2012	VEGF
Bergwitz et al	2001	Wnt
Boyden et al	2002	Wnt
Burke et al	1998	FGF
Cadigan et al	1997	Wnt
Chang et al	2007	Wnt
Chen et al	2004	BMP
Chillakuri et al	2012	Notch
Cowan et al	2004	BMP
Daluiski et al	2001	BMP
Deregowski et al	2006	Notch
Ding et al	2014	VEGF
Dragoo et al	2003	BMP
Dragoo et al	2005	BMP
Eswarakumar et al	2002	FGF
Fan et al	2016	Notch
Fan et al	2013	BMP
Fan et al	2014	BMP
Fan et al	2016	BMP
Fischer et al	2002	Wnt
Gilboa et al	2000	BMP
Gong et al	2001	Wnt
Grottkau et al	2013	BMP, Wnt, Hedgehog
Heldin et al	1997	BMP
Hilton et al	2008	Notch
Hung et al	2015	ERK
Jacob et al	2006	FGF
Kadesch	2004	Notch
Kokabu et al	2012	BMP
Kwan et al	2011	FGF
Levi et al	2010	BMP
Levi et al	2011	BMP
Li et al	2016	VEGF
Lin et al	2008	BMP
Liu et al	2002	FGF
Long et al	2013	Hedgehog
Lu et al	2012	BMP
Lu et al	2013	BMP
Marie	2003	FGF
Marie	2012	FGF
Mie et al	2000	BMP
Montero et al	2000	FGF
Nakamura et al	1997	Hedgehog
Ohbayashi et al	2002	FGF
Quarto et al	2006	FGF
Quarto et al	2008	FGF
Rice	2008	BMP, Wnt
Santos et al	2010	VEGF
Sarkar et al	2001	FGF
Schmuhl et al	2014	ERK
Spinella-Jaegle et al	2001	Hedgehog
St-Jacques et al	1999	Hedgehog
Stevens et al	2010	Wnt
Tu et al	2007	Wnt
Vanhatupa et al	2015	BMP
Westendorf et al	2004	Wnt
Wodarz et al	1998	Wnt
Yuasa et al	2002	Hedgehog

transcription factor osterix (OSX).^{30,31} Extracellular signaling pathways converge on these transcription factors to orchestrate and regulate osteoblastogenesis. The pathways are bone morphogenic protein (BMP), wingless-related integration site protein (Wnt), Notch, Hedgehog, fibroblast growth factor (FGF), vascular endothelial growth factor (VEGF), and extracellular regulated-signal kinase (ERK) (Fig. 1). The same signaling pathways play a role in the direction of putative adult stem cells to an osteoblastic lineage, demonstrated by the bone markers alkaline phosphatase (ALP), type I collagen, osteopontin (OPN), and osteocalcin (OCN), as well as matrix mineralization,³² for the purpose of tissue engineering.^{33,34}

Bone Morphogenic Protein

The BMPs belong to the transcription growth factor beta superfamily.^{35,36} This group is perhaps the most well-known of the cytokines involved in osteogenesis, and as such, the majority of studies related to ASCs and bone formation has revolved around the BMP pathway.

The BMPs operate through signaling mothers against decapentaplegic (Smad) 1 phosphorylation.^{35,37} The phosphorylated Smads promote osteogenic differentiation by forming a complex with Smad4.³⁴ The BMPs also increase the transcription of RUNX2.³⁸ Subtypes BMP2 and BMP4 are particularly important, governing crucial steps of differentiation by managing the transition from RUNX2- to OSX-positive cells.^{39–41} The ASCs increase bone regeneration significantly when loaded onto a scaffold with BMP2 or when transfected with the BMP2 gene.^{42–47} On the contrary, BMP3 is thought to have a negative regulatory effect on osteogenesis⁴⁸ through interaction with the BMP receptor type II to inhibit BMP2 and BMP4 signaling.⁴⁹

Downregulation of the osteogenic inhibitor Noggin in ASCs triggered more appreciable mandibular regeneration in rats, and this effect became more pronounced when Noggin^{-/-} ASCs were supplemented with BMP2.^{50–52} Furthermore, tumor necrosis factor alpha (TNF-α), a major inflammatory factor peaking after bone fracture, promoted BMP2 establishment in human primary osteoblasts and inhibited osteogenic differentiation when the pathway was contacted with Noggin.⁵³ Preconditioning with TNF-α augmented proliferation, mobilization, and osteogenic differentiation of ASCs and upregulated BMP2, upon silencing of BMP2 by siRNA.⁵⁴

FIGURE 1. Schematic of osteogenic differentiation induced by adipose-derived stem cells. ALP, alkaline phosphatase; ASC, adipose-derived stem cell; BMP, bone morphogenic protein; ERK, extracellular regulated-signal kinase; FGF, fibroblast growth factor; MSC, mesenchymal stem cell; OCN, osteocalcin; OPN, osteopontin; OSX, osterix; RUNX2, runt-related family 2; VEGF, vascular endothelial growth factor; Wnt, wingless-related integration site protein.

Wingless-Related Integration Site Protein

The Wnt family consists of a large number of secreted glycol proteins that are involved in embryonic development, tissue induction, and axis polarity.^{55,56} Wnt ligands bind to frizzled receptors at the cell surface and to LRP5 and LRP6 coreceptors.³⁴ LRP5 gain of function results in increased bone mass,⁵⁷ whereas mutations in the gene cause osteoporosis-pseudoglioma syndrome.⁵⁸

Canonical Wnt signaling results in stabilization and translocation of β -catenin to the nucleus, where it binds to T-cell factor/lymphoid enhancer factor (TCF/Lef) transcription factors.³⁸ β -catenin-TCF/Lef complexes activate transcription of multiple Wnt-responsive genes, including those implicated in proliferation, osteoblast differentiation, and osteogenesis.^{59–61}

Noncanonical Wnt10b is identified for its unique requirement for maintenance of mesenchymal progenitor activity in adult bone after it was distinguished as the only Wnt ligand linking to mesenchymal progenitor function in humans and mice.⁶² Some studies revealed that noncanonical Wnt signaling, namely that of Wnt5a with Ror2 and RhoA as counterparts, plus *N*-cadherin-mediated β -catenin signaling, are necessary for mechanically induced osteogenesis.⁶³ Furthermore, Wnt4 may have potentiality in improving bone regeneration and repair of craniofacial defects.⁶⁴ Wnt3a and Wnt7b, also of the noncanonical variety, signal through G-proteins to activate phosphatidylinositol signaling and PKC δ , the latter of which is necessary for osteoblastogenesis.⁶⁵

Notch

The Notch network is known to be part of an evolutionarily conserved mechanism that balances proliferation and differentiation of stem cells.⁶⁶ Such action begins with the Jagged Delta protein's attachment to the receptor³⁴ prior to cleavage by γ -secretase to release the Notch intracellular domain.⁶⁷

Notch inhibition in the embryonic limb leads to increased bone mass and a reciprocal decrease in bone marrow mesenchymal progenitors, thereby suggesting Notch's function in suppressing osteogenic differentiation and maintaining a sizable progenitor pool.⁶⁸ Notch is thought to act by quelling RUNX2 transcriptional activity by induction of the transcription factors HEY1 and HEYL, which physically interact with RUNX2.⁶⁹

The miR-34a was found to upregulate *RUNX2* by inhibiting retinoblastoma binding protein 2 via the *NOTCH1/CYCLIN D1* coregulatory network.⁷⁰ Another report, however, observed that Notch-1 overexpression inhibited osteogenesis by interrupting canonical Wnt signaling, but failed to do so with interrupted BMP.⁷¹

Hedgehog

Pattern arrangement of bone tissue is greatly influenced by Hedgehog signaling, albeit little work on it has been executed.³⁴ In particular, Indian Hedgehog (IHH) plays a decisive job in endochondral development, as *Ihh*^{-/-} mice lack osteoblasts within the endochondral tissue, but not in intramembranous tissue.⁷² Without IHH, mesenchymal progenitors in the perichondrium of the cartilaginous anlagen do not express *Runx2* and hence fail to undergo differentiation.⁷³ The IHH receptor Smoothened (Smo) has been deemed necessary to build trabecular bone.⁷²

On the contrary to IHH, Sonic Hedgehog (Shh) induces ALP expression^{74,75} and augments selectivity in the differentiation of MSCs into the osteoblast lineage.⁷⁶

Fibroblast Growth Factor

The FGF2, FGF9, FGF18, and their corresponding receptors, FGFR1, FGFR2, and FGFR3, are all linked to skeletal development of the long bones and calvarium.^{77,78} Embryos of genotype *Fgf18*^{-/-}

had defects in osteoblast maturation despite normal *Runx2* expression.^{79,80} In contrast, FGFR1 activity has a tendency to launch osteogenic differentiation at an early stage, only to inhibit mineralization capabilities in mature osteoblasts.⁸¹ Gain of function mutations in FGFR1, FGFR2, and FGFR3 caused craniosynostosis, while implantation of beads soaked with FGF2 and FGF4 around sutures caused osteogenesis and later suture closure.^{82,83}

FGF2 serves as the quintessential example of how FGF is connected to bone formation.⁸⁴ Although the promotion of bone repair via ASC mediation had been previously reported in vivo, FGF2 was additionally found to inhibit terminal osteogenic differentiation by antagonizing the retinoic acid-mediated upregulation of BMP receptor type IB.^{85–87} This paradox can be explained when one considers the principal function of FGF2, to promote the proliferation and expansion of osteoprogenitor cells to maximize the osteoprogenitor pool for future differentiation.⁸⁵ Still, other discussions are rooted in the fact that FGF2 can activate different signaling pathways, including ERK, PI3K, and PKC, of which stage and context would resolve the proliferation-versus-differentiation debate.^{84,88} Given the complexity of FGF, as well as other signaling pathways, the ability to guide a specific one in the context of ASC-based therapy for bone formation would be a sweeping step in regenerative medicine.⁵

Vascular Endothelial Growth Factor

Neurovascularization and angiogenesis are paramount to proper bone foundation.⁸⁹ It has been exhibited that VEGFA had a more potent effect in precipitating ASC-mediated calvarial regeneration than either BMP2 or FGF2, through a combination of osteogenesis and angiogenesis.⁹⁰

Dimethylloxalglycine was tested for a dose-dependent effect by levels of *RUNX2*, *OCN*, and *ALP* expression, plus VEGF, which is produced as a product of *HIF-1 α* overexpression.⁹¹ The ASCs expressing *Runx2* in combination with a vascularized flap led to more effective bone repair than either facet acting alone, VEGF and collagen type I were indicators.⁹²

Extracellular Signal-Regulated Kinase

Another pathway that deals with osteogenic differentiation is ERK, which is sometimes referred to as mitogen-activating protein kinase, and where platelet-derived growth factor (PDGF) comes into play.⁹³ In 1 study, in the presence of PDGF-BB, ASCs, not BMSCs, heightened osteogenic differentiation by increasing *Runx2* and *OCN* output.⁹⁴

Other Pathways Involved in Osteogenic Differentiation

Erythropoietin has been reported to give rise to osteogenesis by inhibiting PPAR γ , while IGF1 does the same through TAZ, and when joined, resulted in even higher levels of *Runx2*, *OPN*, *OCN*, *ALP*, and matrix mineralization.⁹⁵ The ASCs magnofected with *Bcl2*, an inhibitor of apoptosis, prompted greater ALP, extracellular matrix mineralization, and expression of *Ocn*, *Opn*, and *Runx2* than nucleofected cells.⁹⁶ miR-135 was shown to positively regulate osteogenic differentiation based on bone markers and extracellular matrix decomposition through a potentially new miR135/Hoxa2/*Runx2* pathway.⁹⁷ The role of the transcription factor δ FosB has been explored, and the results obtained pointed to increased bone mass and decreased adipocyte origination upon overexpression.^{98,99} A final biomolecule that has been scrutinized is growth and differentiation factor 5, which appears to be more effective than BMP2 in inducing onset of osteogenic differentiation while simultaneously prompting vascularization through VEGF.^{99,100}

DISCUSSION

Taken together, these limited numbers of studies have assisted in mapping out the signaling pathways that ASCs go through to become bone. Most work has looked at the function that the BMP course fulfills, followed by those of Wnt, Notch, Hedgehog, FGF, VEGF, and ERK. By differing scaffold type and culture medium, molecular composition, mechanical stress, chemical amalgamation, or bioengineering, the effectiveness of osteogenic differentiation can be evaluated. Activity of ALP, emergence of collagen type I, expression of *OCN* and *OPN*, and mineralization through calcium precipitation were quantified in most cases to analyze potential.

It has become increasingly clear that with proper control of outside inputs, ASCs make an excellent choice for bone tissue engineering in clinical practice. The ideal mixture of favorable inducers is not as straightforward, but the depth of research insists that multiple manipulation strategies exist, as long as the stem cells are exposed to at least one inducer of osteogenesis early on. Additionally, there is a trend in combining inducers as to achieve optimal outcomes.

New methods are necessitated to not only reduce the culture period and quantity of growth factors, but also to enhance the efficiency of osteogenesis and thus bone regeneration. One approach is delivery of cytokines incorporating these molecules into scaffolds as basic as liposomes and microspheres.¹⁷ This makes the growth factor retainable at the site of interest for an extended period while maintaining its biologic activity. Engineered ASCs with gene transfection by virus vectors have evolved to be an attractive option to ameliorate bone repair, especially in large bone defects.³⁴ As bone formation by implantation of ASCs must be preceded by the in vitro osteogenic differentiation of these cells, it is compulsory to design procedures that ensure a well-characterized and consistent cell population following the differentiation process.³⁴

REFERENCES

- Steiner C, Elixhauser A, Schnaier J. The healthcare cost and utilization project: an overview. *Eff Clin Pract* 2002;5:143–151
- Deev RV, Drobyshev AY, Bozo IY, et al. Ordinary and activated bone grafts: applied classification and the main features. *BioMed Res Int* 2015;2015:365050
- Chiapasco M, Casentini P, Zaniboni M. Bone augmentation procedures in implant dentistry. *Int J Oral Maxillofac Implants* 2009;24:237–259
- Kaing L, Grubor D, Chandu A. Assessment of bone grafts placed within an oral and maxillofacial training programme for implant rehabilitation. *Aust Dent J* 2011;56:406–411
- Senarath-Yapa K, McArdle A, Renda A, et al. Adipose-derived stem cells: a review of signaling networks governing cell fate and regenerative potential in the context of craniofacial and long bone skeletal repair. *Int J Mol Sci* 2014;15:9314–9330
- Zech NH, Preisegger KH, Hollands P. Stem cell therapeutics—reality versus hype and hope. *J Assist Reprod Genet* 2011;28:287–290
- Johnstone B, Hering TM, Caplan AI, et al. In vitro chondrogenesis of bone marrow-derived mesenchymal progenitor cells. *Exp Cell Res* 1998;238:265–272
- Pittenger MF, Mackay AM, Beck SC, et al. Multilineage potential of adult human mesenchymal stem cells. *Science* 1999;284:143–147
- Jaiswal RK, Jaiswal N, Bruder SP, et al. Adult human mesenchymal stem cell differentiation to the osteogenic or adipogenic lineage is regulated by mitogen-activated protein kinase. *J Biol Chem* 2000;275:9645–9652
- Zuk PA, Zhu M, Mizuno H, et al. Multilineage cells from human adipose tissue: implications for cell-based therapies. *Tissue Eng* 2001;7:211–228
- Zuk PA, Zhu M, Ashjian P, et al. Human adipose tissue is a source of multipotent stem cells. *Mol Biol Cell* 2002;13:4279–4295
- Gimble JM, Katz AJ, Bunnell BA. Adipose-derived stem cells for regenerative medicine. *Circ Res* 2007;100:1249–1260
- Zuk PA. Tissue engineering craniofacial defects with adult stem cells? Are we ready yet? *Pediatr Res* 2008;63:478–486
- Guilak F, Estes BT, Diekman BO, et al. 2010 Nicolas Andry Award: multipotent adult stem cells from adipose tissue for musculoskeletal tissue engineering. *Clin Orthop Relat Res* 2010;468:2530–2540
- Saulnier N, Puglisi MA, Lattanzi W, et al. Gene profiling of bone marrow- and adipose tissue-derived stromal cells: a key role of Kruppel-like factor 4 in cell fate regulation. *Cytotherapy* 2011;13:329–340
- Strem BM, Hicok KC, Zhu M, et al. Multipotential differentiation of adipose tissue-derived stem cells. *Keio J Med* 2005;54:132–141
- Basmanav FB, Kose GT, Hasirci V. Sequential growth factor delivery from complexed microspheres for bone tissue engineering. *Biomaterials* 2008;29:4195–4204
- Jeon O, Song SJ, Yang HS, et al. Long-term delivery enhances in vivo osteogenic efficacy of bone morphogenetic protein-2 compared to short-term delivery. *Biochem Biophys Res Commun* 2008;369:774–780
- Zhang S, Doschak MR, Uludag H. Pharmacokinetics and bone formation by BMP-2 entrapped in polyethylenimine-coated albumin nanoparticles. *Biomaterials* 2009;30:5143–5155
- Engler AJ, Sen S, Sweeney HL, et al. Matrix elasticity directs stem cell lineage specification. *Cell* 2006;126:677–689
- Comisar WA, Kazmers NH, Mooney DJ, et al. Engineering RGD nanopatterned hydrogels to control preosteoblast behavior: a combined computational and experimental approach. *Biomaterials* 2007;28:4409–4417
- Dalby MJ, Gadegaard N, Tare R, et al. The control of human mesenchymal cell differentiation using nanoscale symmetry and disorder. *Nat Mater* 2007;6:997–1003
- Hofmann S, Hagenmuller H, Koch AM, et al. Control of in vitro tissue-engineered bone-like structures using human mesenchymal stem cells and porous silk scaffolds. *Biomaterials* 2007;28:1152–1162
- Lindroos B, Suuronen R, Miettinen S. The potential of adipose stem cells in regenerative medicine. *Stem Cell Rev* 2011;7:269–291
- Ducy P, Zhang R, Geoffroy V, et al. *Osf2/Cbfa1*: a transcriptional activator of osteoblast differentiation. *Cell* 1997;89:747–754
- Otto F, Thornell AP, Crompton T, et al. *Cbfa1*, a candidate gene for cleidocranial dysplasia syndrome, is essential for osteoblast differentiation and bone development. *Cell* 1997;89:765–771
- Ducy P, Starbuck M, Priemel M, et al. A *Cbfa1*-dependent genetic pathway controls bone formation beyond embryonic development. *Genes Dev* 1999;13:1025–1036
- Choi JY, Pratap J, Javed A, et al. Subnuclear targeting of Runx/Cbfa/AML factors is essential for tissue-specific differentiation during embryonic development. *Proc Natl Acad Sci U S A* 2001;98:8650–8655
- Li J, Zhao Q, Wang E, et al. Transplantation of *Cbfa1*-overexpressing adipose stem cells together with vascularized periosteal flaps repair segmental bone defects. *J Surg Res* 2012;176:e13–e20
- Nakashima K, Zhou X, Kunkel G, et al. The novel zinc finger-containing transcription factor osterix is required for osteoblast differentiation and bone formation. *Cell* 2002;108:17–29
- Zhou X, Zhang Z, Feng JQ, et al. Multiple functions of Osterix are required for bone growth and homeostasis in postnatal mice. *Proc Natl Acad Sci U S A* 2010;107:12919–12924
- Hicok KC, Du Laney TV, Zhou YS, et al. Human adipose-derived adult stem cells produce osteoid in vivo. *Tissue Eng* 2004;10:371–380
- Levi B, Longaker MT. Concise review: adipose-derived stromal cells for skeletal regenerative medicine. *Stem Cells* 2011;29:576–582
- Grottkau BE, Lin Y. Osteogenesis of adipose-derived stem cells. *Bone Res* 2013;1:133–145
- Heldin CH, Miyazono K, Dijke P. TGF-beta signalling from cell membrane to nucleus through SMAD proteins. *Nature* 1997;390:465–471
- Chen D, Zhao M, Mundy GR. Bone morphogenetic proteins. *Growth Factors* 2004;22:233–241
- Gilboa L, Nohe A, Geissendorfer T, et al. Bone morphogenetic protein receptor complexes on the surface of live cells: a new oligomerization mode for serine/threonine kinase receptors. *Mol Biol Cell* 2000;11:1023–1035
- Rice DP. Craniofacial sutures. Development, disease and treatment. *Preface Front Oral Biol* 2008;12:xi

39. Mie M, Ohgushi H, Yanagida Y, et al. Osteogenesis coordinated in C3H10T1/2 cells by adipogenesis-dependent BMP-2 expression system. *Tissue Eng* 2000;6:9–18
40. Bandyopadhyay A, Tsuji K, Cox K, et al. Genetic analysis of the roles of BMP2, BMP4, and BMP7 in limb patterning and skeletogenesis. *PLoS Genet* 2006;2:e216
41. Vanhatupa S, Ojansivu M, Autio R, et al. Bone morphogenetic protein-2 induces donor-dependent osteogenic and adipogenic differentiation in human adipose stem cells. *Stem Cells Transl Med* 2015;4:1391–1402
42. Dragoo JL, Choi JY, Lieberman JR, et al. Bone induction by BMP-2 transduced stem cells derived from human fat. *J Orthop Res* 2003;21:622–629
43. Cowan CM, Shi YY, Aalami OO, et al. Adipose-derived adult stromal cells heal critical-size mouse calvarial defects. *Nat Biotechnol* 2004;22:560–567
44. Dragoo JL, Lieberman JR, Lee RS, et al. Tissue-engineered bone from BMP-2-transduced stem cells derived from human fat. *Plast Reconstr Surg* 2005;115:1665–1673
45. Lin Y, Tang W, Wu L, et al. Bone regeneration by BMP-2 enhanced adipose stem cells loading on alginate gel. *Histochem Cell Biol* 2008;129:203–210
46. Levi B, James AW, Nelson ER, et al. Human adipose derived stromal cells heal critical size mouse calvarial defects. *PLoS One* 2010;5:e11177
47. Levi B, Hyun JS, Nelson ER, et al. Nonintegrating knockdown and customized scaffold design enhances human adipose-derived stem cells in skeletal repair. *Stem Cells* 2011;29:2018–2029
48. Daluiski A, Engstrand T, Bahamonde ME, et al. Bone morphogenetic protein-3 is a negative regulator of bone density. *Nat Genet* 2001;27:84–88
49. Kokabu S, Gamer L, Cox K, et al. BMP3 suppresses osteoblast differentiation of bone marrow stromal cells via interaction with Aevr2b. *Mol Endocrinol* 2012;26:87–94
50. Fan J, Park H, Tan S, et al. Enhanced osteogenesis of adipose derived stem cells with Noggin suppression and delivery of BMP-2. *PLoS One* 2013;8:e72474
51. Fan J, Park H, Lee MK, et al. Adipose-derived stem cells and BMP-2 delivery in chitosan-based 3D constructs to enhance bone regeneration in a rat mandibular defect model. *Tissue Eng Part A* 2014;20:2169–2179
52. Fan J, Im CS, Guo M, et al. Enhanced osteogenesis of adipose-derived stem cells by regulating bone morphogenetic protein signaling antagonists and agonists. *Stem Cells Transl Med* 2016;5:539–551
53. Lu Z, Wang G, Dunstan CR, et al. Short-term exposure to tumor necrosis factor- α enables human osteoblasts to direct adipose tissue-derived mesenchymal stem cells into osteogenic differentiation. *Stem Cells Dev* 2012;21:2420–2429
54. Lu Z, Wang G, Dunstan CR, et al. Activation and promotion of adipose stem cells by tumor necrosis factor- α preconditioning for bone regeneration. *J Cell Physiol* 2013;228:1737–1744
55. Cadigan KM, Nusse R. Wnt signaling: a common theme in animal development. *Genes Dev* 1997;11:3286–3305
56. Wodarz A, Nusse R. Mechanisms of Wnt signaling in development. *Annu Rev Cell Dev Biol* 1998;14:59–88
57. Boyden LM, Mao J, Belsky J, et al. High bone density due to a mutation in LDL-receptor-related protein 5. *N Engl J Med* 2002;346:1513–1521
58. Gong Y, Slee RB, Fukui N, et al. LDL receptor-related protein 5 (LRP5) affects bone accrual and eye development. *Cell* 2001;107:513–523
59. Bergwitz C, Wendlandt T, Kispert A, et al. Wnts differentially regulate colony growth and differentiation of chondrogenic rat calvaria cells. *Biochim Biophys Acta* 2001;1538:129–140
60. Fischer L, Boland G, Tuan RS. Wnt signaling during BMP-2 stimulation of mesenchymal chondrogenesis. *J Cell Biochem* 2002;84:816–831
61. Westendorf JJ, Kahler RA, Schroeder TM. Wnt signaling in osteoblasts and bone diseases. *Gene* 2004;341:19–39
62. Stevens JR, Miranda-Carboni GA, Singer MA, et al. Wnt10b deficiency results in age-dependent loss of bone mass and progressive reduction of mesenchymal progenitor cells. *J Bone Miner Res* 2010;25:2138–2147
63. Arnsdorf EJ, Tummala P, Jacobs CR. Non-canonical Wnt signaling and N-cadherin related beta-catenin signaling play a role in mechanically induced osteogenic cell fate. *PLoS One* 2009;4:e5388
64. Chang J, Sonoyama W, Wang Z, et al. Noncanonical Wnt-4 signaling enhances bone regeneration of mesenchymal stem cells in craniofacial defects through activation of p38 MAPK. *J Biol Chem* 2007;282:30938–30948
65. Tu X, Joeng KS, Nakayama KI, et al. Noncanonical Wnt signaling through G protein-linked PKC δ activation promotes bone formation. *Dev Cell* 2007;12:113–127
66. Kadesch T. Notch signaling: the demise of elegant simplicity. *Curr Opin Genet Dev* 2004;14:506–512
67. Chillakuri CR, Sheppard D, Lea SM, et al. Notch receptor-ligand binding and activation: insights from molecular studies. *Semin Cell Dev Biol* 2012;23:421–428
68. Tu X, Chen J, Lim J, et al. Physiological notch signaling maintains bone homeostasis via RBPjk and Hey upstream of NFATc1. *PLoS Genet* 2012;8:e1002577
69. Hilton MJ, Tu X, Wu X, et al. Notch signaling maintains bone marrow mesenchymal progenitors by suppressing osteoblast differentiation. *Nat Med* 2008;14:306–314
70. Fan C, Jia L, Zheng Y, et al. MiR-34a promotes osteogenic differentiation of human adipose-derived stem cells via the RBP2/NOTCH1/CYCLIN D1 coregulatory network. *Stem Cell Reports* 2016;7:236–248
71. Derogowski V, Gazzero E, Priest L, et al. Notch 1 overexpression inhibits osteoblastogenesis by suppressing Wnt/beta-catenin but not bone morphogenetic protein signaling. *J Biol Chem* 2006;281:6203–6210
72. St-Jacques B, Hammerschmidt M, McMahon AP. Indian hedgehog signaling regulates proliferation and differentiation of chondrocytes and is essential for bone formation. *Genes Dev* 1999;13:2072–2086
73. Long F, Ornitz DM. Development of the endochondral skeleton. *Cold Spring Harb Perspect Biol* 2013;5:a008334
74. Nakamura T, Aikawa T, Iwamoto-Enomoto M, et al. Induction of osteogenic differentiation by hedgehog proteins. *Biochem Biophys Res Commun* 1997;237:465–469
75. Yuasa T, Kataoka H, Kinto N, et al. Sonic hedgehog is involved in osteoblast differentiation by cooperating with BMP-2. *J Cell Physiol* 2002;193:225–232
76. Spinella-Jaegle S, Rawadi G, Kawai S, et al. Sonic hedgehog increases the commitment of pluripotent mesenchymal cells into the osteoblastic lineage and abolishes adipocytic differentiation. *J Cell Sci* 2001;114:2085–2094
77. Montero A, Okada Y, Tomita M, et al. Disruption of the fibroblast growth factor-2 gene results in decreased bone mass and bone formation. *J Clin Invest* 2000;105:1085–1093
78. Eswarakumar VP, Monsonego-Ornan E, Pines M, et al. The IIIc alternative of Fgfr2 is a positive regulator of bone formation. *Development* 2002;129:3783–3793
79. Liu Z, Xu J, Colvin JS, et al. Coordination of chondrogenesis and osteogenesis by fibroblast growth factor 18. *Genes Dev* 2002;16:859–869
80. Ohbayashi N, Shibayama M, Kurotaki Y, et al. FGF18 is required for normal cell proliferation and differentiation during osteogenesis and chondrogenesis. *Genes Dev* 2002;16:870–879
81. Jacob AL, Smith C, Partanen J, et al. Fibroblast growth factor receptor 1 signaling in the osteo-chondrogenic cell lineage regulates sequential steps of osteoblast maturation. *Dev Biol* 2006;296:315–328
82. Burke D, Wilkes D, Blundell TL, et al. Fibroblast growth factor receptors: lessons from the genes. *Trends Biochem Sci* 1998;23:59–62
83. Sarkar S, Petiot A, Copp A, et al. FGF2 promotes skeletogenic differentiation of cranial neural crest cells. *Development* 2001;128:2143–2152
84. Marie PJ. Fibroblast growth factor signaling controlling osteoblast differentiation. *Gene* 2003;316:23–32
85. Quarto N, Longaker MT. FGF-2 inhibits osteogenesis in mouse adipose tissue-derived stromal cells and sustains their proliferative and osteogenic potential state. *Tissue Eng* 2006;12:1405–1418

86. Quarto N, Wan DC, Longaker MT. Molecular mechanisms of FGF-2 inhibitory activity in the osteogenic context of mouse adipose-derived stem cells (mASCs). *Bone* 2008;42:1040–1052
87. Kwan MD, Sellmyer MA, Quarto N, et al. Chemical control of FGF-2 release for promoting calvarial healing with adipose stem cells. *J Biol Chem* 2011;286:11307–11313
88. Marie PJ. Fibroblast growth factor signaling controlling bone formation: an update. *Gene* 2012;498:1–4
89. Santos MI, Reis RL. Vascularization in bone tissue engineering: physiology, current strategies, major hurdles and future challenges. *Macromol Biosci* 2010;10:12–27
90. Behr B, Sorkin M, Lehnhardt M, et al. A comparative analysis of the osteogenic effects of BMP-2, FGF-2, and VEGFA in a calvarial defect model. *Tissue Eng Part A* 2012;18:1079–1086
91. Ding H, Gao YS, Wang Y, et al. Dimethylloxaloylglycine increases the bone healing capacity of adipose-derived stem cells by promoting osteogenic differentiation and angiogenic potential. *Stem Cells Dev* 2014;23:990–1000
92. Li Q, Wang T, Zhang GF, et al. A comparative evaluation of the mechanical properties of two calcium phosphate/collagen composite materials and their osteogenic effects on adipose-derived stem cells. *Stem Cells Int* 2016;2016:6409546
93. Schmuhl E, Ramer R, Salamon A, et al. Increase of mesenchymal stem cell migration by cannabidiol via activation of p42/44 MAPK. *Biochem Pharmacol* 2014;87:489–501
94. Hung BP, Hutton DL, Kozielski KL, et al. Platelet-derived growth factor BB enhances osteogenesis of adipose-derived but not bone marrow-derived mesenchymal stromal/stem cells. *Stem Cells* 2015;33:2773–2784
95. Zhou J, Wei F, Ma Y. Inhibiting PPARgamma by erythropoietin while upregulating TAZ by IGF1 synergistically promote osteogenic differentiation of mesenchymal stem cells. *Biochem Biophys Res Commun* 2016;478:349–355
96. Brett E, Zielins ER, Luan A, et al. Magnetic nanoparticle-based upregulation of B-cell lymphoma 2 enhances bone regeneration. *Stem Cells Transl Med* 2017;6:151–160
97. Xie Q, Wang Z, Zhou H, et al. The role of miR-135-modified adipose-derived mesenchymal stem cells in bone regeneration. *Biomaterials* 2016;75:279–294
98. Sabatakis G, Sims NA, Chen J, et al. Overexpression of DeltaFosB transcription factor(s) increases bone formation and inhibits adipogenesis. *Nat Med* 2000;6:985–990
99. Rada T, Reis RL, Gomes ME. Adipose tissue-derived stem cells and their application in bone and cartilage tissue engineering. *Tissue Eng Part B Rev* 2009;15:113–125
100. Lough DM, Chambers C, Germann G, et al. Regulation of ADSC osteoinductive potential using notch pathway inhibition and gene rescue: a potential on/off switch for clinical applications in bone formation and reconstructive efforts. *Plast Reconstr Surg* 2016;138:642e–652e