UC Berkeley

UC Berkeley Previously Published Works

Title

Multiplicity dependence of light (anti-)nuclei production in p-Pb collisions at s NN = 5.02 TeV

Permalink

https://escholarship.org/uc/item/8bb1r9q5

Authors

Acharya, S Adamová, D Adhya, SP <u>et al.</u>

Publication Date

2020

DOI

10.1016/j.physletb.2019.135043

Peer reviewed

Physics Letters B 800 (2020) 135043

Contents lists available at ScienceDirect

Physics Letters B

www.elsevier.com/locate/physletb

Multiplicity dependence of light (anti-)nuclei production in p–Pb collisions at $\sqrt{s_{\text{NN}}} = 5.02$ TeV



ARTICLE INFO

Article history: Received 9 July 2019 Received in revised form 10 October 2019 Accepted 18 October 2019 Available online 23 October 2019 Editor: L. Rolandi

ABSTRACT

The measurement of the deuteron and anti-deuteron production in the rapidity range -1 < y < 0 as a function of transverse momentum and event multiplicity in p–Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV is presented. (Anti-)deuterons are identified via their specific energy loss dE/dx and via their time-of-flight. Their production in p–Pb collisions is compared to pp and Pb–Pb collisions and is discussed within the context of thermal and coalescence models. The ratio of integrated yields of deuterons to protons (d/p) shows a significant increase as a function of the charged-particle multiplicity of the event starting from values similar to those observed in pp collisions at low multiplicities and approaching those observed in Pb–Pb collisions at high multiplicities. The mean transverse particle momenta are extracted from the deuteron spectra and the values are similar to those obtained for p and Λ particles. Thus, deuteron spectra do not follow mass ordering. This behaviour is in contrast to the trend observed for non-composite particles in p–Pb collisions. In addition, the production of the rare ³He and ³He nuclei has been studied. The spectrum corresponding to all non-single diffractive p-Pb collisions is obtained in the rapidity window -1 < y < 0 and the p_{T} -integrated yield dN/dy is extracted. It is found that the yields of protons, deuterons, and ³He, normalised by the spin degeneracy factor, follow an exponential decrease with mass number.

© 2019 Published by Elsevier B.V. This is an open access article under the CC BY license (http://creativecommons.org/licenses/by/4.0/). Funded by SCOAP³.

1. Introduction

The energy densities reached in the collisions of ultrarelativistic particles lead to a significant production of complex (anti-)(hyper-)nuclei. The high yield of anti-quarks produced in these reactions has led to the first observation of the anti-alpha particle [1] as well as of the anti-hyper-triton [2] by the STAR collaboration, and to detailed measurements by the ALICE collaboration [3–6] at energies reached at the CERN LHC. However, the production mechanism is not fully understood. In a more general context, these measurements also provide input for the background determination in searches for anti-nuclei in space. Such an observation of anti-deuterons or ${}^{3}\overline{\text{He}}$ of cosmic origin could carry information on the existence of large amounts of anti-matter in our universe or provide a signature of the annihilation of dark matter particles [7–11].

Recent data in pp and in heavy-ion collisions provide evidence for an interesting observation regarding the production mechanism of (anti-)nuclei [3,5,6,12,13]: in Pb–Pb interactions, the d/p ratio does not vary with the collision centrality and the value agrees with expectations from thermal-statistical models which feature a common chemical freeze-out temperature of all hadrons around 156 MeV [3,14,15]. In inelastic pp collisions, the corresponding ratio is a factor 2.2 lower than in Pb–Pb collisions [3,12]. With respect to these measurements, the results of d and ³He produced in p–Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV, being a system in between the two extremes of pp and Pb–Pb collisions, are of prominent interest and they are the subject of this letter. While deuterons have been measured differentially in multiplicity, the ³He (³He) spectrum was only obtained inclusively for all non-single diffractive events because of their low production rate.

In addition to the evolution of the integrated d/p ratio for various multiplicity classes, the question whether the transverse momentum distribution of deuterons is consistent with a collective radial expansion together with the non-composite hadrons is of particular interest. Such behaviour has been observed for light nuclei in Pb–Pb collisions [3,5]. The presence of collective effects in p–Pb collisions at LHC energies has recently been supported by several experimental findings (see for instance [16–22] and recent reviews in [23,24]). These include a clear mass ordering of the mean transverse momenta of light flavoured hadrons in p–Pb collisions as expected from hydrodynamical models [18].

https://doi.org/10.1016/j.physletb.2019.135043

0370-2693/© 2019 Published by Elsevier B.V. This is an open access article under the CC BY license (http://creativecommons.org/licenses/by/4.0/). Funded by SCOAP³.





^{*} E-mail address: alice-publications@cern.ch.

2. Analysis

The results presented here are based on a low pile-up p-Pb data sample collected with the ALICE detector during the LHC running campaign at $\sqrt{s_{NN}}$ = 5.02 TeV in 2013. A detailed description of the detector is available in [25–29]. The main detectors used in this analysis are the Inner Tracking System (ITS) [30], the Time Projection Chamber (TPC) [31], and the Time-Of-Flight detector (TOF) [32,33]. The two innermost layers of the ITS consist of Silicon Pixel Detectors (SPD), followed by two layers of Silicon Drift Detectors (SDD), and two layers of Silicon Strip Detectors (SSD). As the main tracking device, the TPC provides full azimuthal acceptance for tracks in the pseudo-rapidity region $|\eta_{lab}| < 0.8$. In addition, it provides particle identification via the measurement of the specific energy loss dE/dx. The TOF array is located at about 3.7 m from the beam line and provides particle identification by measuring the particle speed with the time-of-flight technique. In p-Pb collisions, the overall time resolution is about 85 ps for high multiplicity events. In peripheral events, where multiplicities are similar to pp, it decreases to about 120 ps due to a worse starttime (collision-time) resolution [34]. All detectors are positioned in a solenoidal magnetic field of B = 0.5 T.

The event sample used for the analysis presented in this letter was collected exclusively in the beam configuration where the proton travels towards negative η_{lab} . The minimum-bias trigger signal and the definition of the multiplicity classes was provided by the V0 detector consisting of two arrays of 32 scintillator tiles each covering the full azimuth within 2.8 $<\eta_{lab}<$ 5.1 (V0A, Pb-beam direction) and $-3.7 < \eta_{lab} < -1.7$ (VOC, p-beam direction). The event selection was performed in a similar way to that described in Ref. [18]. A coincidence of signals in both VOA and VOC was required online in order to remove background from single diffractive and electromagnetic events. In the offline analysis, further background suppression was achieved by requiring that the arrival time of the signals in the two neutron Zero Degree Calorimeters (ZDC), which are located ± 112.5 m from the interaction point, is compatible with a nominal p-Pb collision. The contamination from pile-up events was reduced to a negligible level (< 1%) by rejecting events in which more than one primary vertex was reconstructed either from SPD tracklets or from tracks reconstructed in the whole central barrel. The position of the reconstructed primary vertex was required to be located within ± 10 cm of the nominal interaction point in the longitudinal direction. In total, an event sample of about 100 million minimum-bias (MB) events after all selections was analysed. The corresponding integrated luminosity, $L_{\text{int}} = N_{\text{MB}}/\sigma_{\text{MB}}$, where σ_{MB} is the MB trigger cross-section measured with van-der-Meer scans, amounts to 47.8 μ b⁻¹ with a relative uncertainty of 3.7% [35].

The final results are given normalised to the total number of non-single diffractive (NSD) events. Therefore, a correction of $3.6\% \pm 3.1\%$ [36] is applied to the minimum-bias results, which corresponds to the trigger and vertex reconstruction inefficiency for this selection. For the study of d and \overline{d} , the sample is divided into five multiplicity classes, which are defined as percentiles of the VOA signal. This signal is proportional to the charged-particle multiplicity in the corresponding pseudo-rapidity region in the direction of the Pb-beam. Following the approach in [37], the multiplicity dependent results are normalized to the number of events N_{ev} corresponding to the visible (triggered) cross-section. The event sample is corrected for the vertex reconstruction efficiency. This correction is of the order of 4% for the lowest VOA multiplicity class (60-100%) and negligible (<1%) for the other multiplicity classes. The chosen selection and the corresponding charged-particle multiplicity at mid-rapidity are summarized in Table 1.

Т	blo	1
Id	Die	

Multiplicity intervals and the corresponding charged-particle multiplicities at midrapidity. The uncertainties reported for the $\langle dN_{ch}/d\eta_{lab}\rangle|_{|\eta_{lab}|<0.5}$ are the systematic ones, statistical uncertainties are negligible. Values are taken from [18].

VOA Class	$\langle dN_{ch}/d\eta_{lab}\rangle _{ \eta_{lab} <0.5}$
0-10%	40.6 ± 0.9
10-20%	30.5 ± 0.7
20-40%	23.2 ± 0.5
40-60%	16.1 ± 0.4
60-100%	7.1 ± 0.2

In this analysis, the production of primary deuterons and ³He-nuclei and that of their respective anti-particles are measured in a rapidity window -1 < y < 0 in the centre-of-mass system. Since the energy per nucleon of the proton beam is higher than that of the Pb beam, the nucleon-nucleon system moves in the laboratory frame with a rapidity of -0.465. Potential differences of the spectral shape or normalisation due to the larger *y*-range with respect to the measurement of π , K, and p [18] are found to be negligible for the (anti-)deuteron and ³He minimum-bias spectra with respect to the overall statistical and systematic uncertainties. In order to select primary tracks of suitable guality, various track selection criteria are applied. At least 70 clusters in the TPC and two hits in the ITS (out of which at least one in the SPD) are required. These selections guarantee a track momentum resolution of 2% in the relevant $p_{\rm T}$ -range and a dE/dx resolution of about 6% for minimum ionising particles. The maximum allowed Distance-of-Closest-Approach (DCA) to the primary collision vertex is 0.12 cm in the transverse (DCA_{xv}) and 1.0 cm in the longitudinal (DCA_z) plane. Furthermore, it is required that the χ^2 per TPC cluster is less than 4 and tracks of weak-decay products with kink topology are rejected [29], as they cannot originate from the tracks of primary nuclei.

The particle identification performance of the TPC and TOF detectors in p-Pb collisions is shown in Fig. 1. For the mass determination with the TOF detector, the contribution of tracks with a wrongly assigned TOF cluster is largely reduced by a 3σ preselection in the TPC dE/dx, where σ corresponds to the TPC dE/dx resolution. Nevertheless, due to the small abundance of deuterons the background is still significant and it is removed using a fit to the squared mass distribution. An example of a fit for antideuterons with transverse momenta 2.2 GeV/ $c < p_T < 2.4$ GeV/cis shown in the right panel of Fig. 1. The squared rest mass of the deuteron has been subtracted to simplify the fitting function. The signal has a Gaussian shape with an exponential tail on the right side. This tail is necessary to describe the time-signal shape of the TOF detector [33]. For the background, the sum of two exponential functions is used. One of the exponential functions accounts for the mismatched tracks and the other accounts for the tail of the proton peak. For $(anti-)^3$ He nuclei, the dE/dx is sufficient for a clean identification using only this technique over the entire momentum range 1.5 GeV/ $c < p_T < 5$ GeV/c as the atomic number Z = 2 for ³He leads to a clear separation from other particles.

The tracking acceptance \times efficiency determination is based on a Monte-Carlo simulation using the DPMJET event generator [38] and a full detector description in GEANT3 [39]. As discussed in [3], the hadronic interaction of (anti-)nuclei with detector material is not fully described in GEANT3, therefore two additional correction factors are applied. Firstly, in order to account for the material between the collision vertex and the TPC, the track reconstruction efficiencies extracted from GEANT3 are scaled to match those from GEANT4 [40,41]. Secondly, for tracks which cross in addition the material between the TPC and the TOF detectors, a data-driven



Fig. 1. Energy loss d*E*/d*x* in the TPC and the corresponding expected energy loss from a parametrization of the Bethe-Bloch curve (left). Example of the fit to the squared TOF mass difference which shows separately the signal and the background from the exponential tail of protons and from mismatched tracks (right).



Fig. 2. Tracking acceptance \times efficiency correction for (anti-)deuterons (left) and for ³He and ³He (right) in the minimum-bias class. The efficiencies for anti-nuclei are lower due to the larger cross-section for hadronic interactions.

correction factor has been evaluated by comparing the matching efficiency of tracks to TOF hits in data and Monte Carlo simulation. Since the TRD was not fully installed in 2013, this study was repeated for regions in azimuth with and without installed TRD modules. The matching efficiencies for tracks crossing the TRD material were then scaled such that the corrected yield agrees with the one obtained for tracks that are not crossing any TRD material. This procedure results in a further reduction of the acceptance × efficiency of 6% for deuterons and 11% for anti-deuterons. The acceptance and efficiency corrections are found to be independent of the event multiplicity and are shown in Fig. 2 for primary deuterons and anti-deuterons, with and without requiring a TOF match, as well as for ³He and ³He.

The raw yields of deuterons and ³He also include secondary particles which stem from the interactions of primary particles with the detector material. To subtract this contribution, a data-driven approach as in [3,18] is used. The distribution of the DCA_{xy} is fitted with two distributions (called "templates" in the following) obtained from Monte-Carlo simulations describing primary and secondary deuterons, respectively. The fit is performed in the range $|DCA_{xy}| < 0.5$ cm which allows the contribution from material to be constrained by the plateau of the distribution at larger distances ($|DCA_{xy}| > 0.15$ cm). The contamination of secondaries amounts to about 45% to 55% in the lowest p_T -interval and decreases exponentially towards higher p_T until it becomes negligible (<1%) above 2 GeV/*c*. The limited number of ³He candidate tracks does not allow a background subtraction based on templates, in-

stead a bin counting procedure in the aforementioned DCA_{xy} signal and background regions is used.

The systematic uncertainties of the measurement are summarised for deuterons and ³He as well as for their antiparticles in Table 2. For deuterons, the uncertainty related to the secondary correction is estimated by repeating the template fit procedure under a variation of the DCA_z cut. The corresponding uncertainty for ³He nuclei is determined by varying the ranges in DCA_{xv} for the signal and background regions in the bin counting procedure. For d and ³He the systematic uncertainty on the cross-section for hadronic interaction is determined by a systematic comparison of different propagation codes (GEANT3 and GEANT4). The material between TPC and TOF needs to be considered only for the (anti-)deuteron spectrum and increases the uncertainty by additional 3% and 5% for deuterons and anti-deuterons, respectively. This corresponds to the half of the observed discrepancy in the TPC-TOF matching efficiencies evaluated in data and Monte Carlo. For both deuterons and anti-deuterons, the particle identification procedure introduces only a small uncertainty which slightly increases at high $p_{\rm T}$ and is estimated based on the variation of the $n\sigma$ -cuts in the TPC dE/dx as well as on a variation of the signal extraction in the TOF with different fit functions. The PID related uncertainties for 3 He and 3 He remain negligible over the entire $p_{\rm T}$ -range due to the background-free identification based on the TPC dE/dx. Feed-down from weakly decaying hyper-tritons $\begin{pmatrix} 3 \\ A \end{pmatrix}$ H) is negligible for deuterons [3,4]. Since only about 4-8% of all $^{3}_{\Lambda}$ H deTransport code

Total

TPC-TOF matching

<i>A</i> ain sources of systematic uncertainties for deuterons and ³ He as well as their anti-particles for low and high $p_{\rm T}$.							
	(d		1	3	He	
$p_{\rm T}~({\rm GeV}/c)$	0.9	2.9	0.9	2.9	2.2	5.0	
Tracking (ITS-TPC matching)	5%	5%	5%	5%	6%	4%	
Secondaries material	1%	negl.	negl.	negl.	20%	1%	
Secondaries weak decay	negl.	negl.	negl.	negl.	5%	negl.	
Material budget	3%	3%	3%	3%	3%	1%	
Particle identification	1%	3%	1%	3%	3%	3%	

3%

5%

8%

3%

5%

9%

6%

23%

6%

8%

Table 2			
Main sources of systematic uncertainties for deuterons	; and ³ He as well as	their anti-particles for low and	high $p_{\rm T}$.
	1	d	

3%

3%

8%

3%

3%

7%



Fig. 3. Transverse momentum distributions of deuterons (left) and anti-deuterons (right) for various multiplicity classes. The multiplicity class definition is based on the signal amplitude observed in the VOA detector located on the Pb-side. The vertical bars represent the statistical errors, the empty boxes show the systematic uncertainty. The lines represent individual fits using a $m_{\rm T}$ -exponential function.

caying into ³He pass the track selection criteria for primary ³He, the remaining contamination has not been subtracted and the uncertainty related to it was further investigated by a variation of the DCA_{xv}-cut in data and a final uncertainty of 5% is assigned. The influence of uncertainties in the material budget on the reconstruction efficiency has been studied by simulating events varying the amount of material by $\pm 10\%$. The estimates of the uncertainties related to the tracking and ITS-TPC matching are based on a variation of the track cuts and are found to be approximately 5%. The uncertainties related to tracking, transport code, material budget and TPC-TOF matching are fully correlated across different multiplicity intervals.

3. Results and discussion

3.1. Spectra and yields

The transverse momentum spectra of deuterons and antideuterons in the rapidity range -1 < y < 0 are presented in Fig. 3 for several multiplicity classes. The spectra show a hardening with increasing event multiplicity. This behaviour was already observed for lower mass particles in p-Pb collisions [18]. For the extraction of $\langle p_{\rm T} \rangle$ and $p_{\rm T}$ -integrated yields dN/dy, the spectra are fitted individually using a $m_{\rm T}$ -exponential function [42].

The values obtained for dN/dy for (anti-)deuterons are summarized in Table 3. They have been calculated by summing up

Table 3

Integrated yields dN/dy of (anti-)deuterons. The first value is the statistical and the second is the total systematic uncertainty which includes both the systematic uncertainty on the measured spectra and the uncertainty of the extrapolation to low and high $p_{\rm T}$.

³He

50

4%

negl

negl.

1%

3%

11%

12%

1.8

6%

5%

3%

3%

18%

20%

negl.

Multiplicity classes	dN/dy (d)	$dN/dy \ (\overline{d})$
0-10%	$(2.86\pm0.03\pm0.30)\times10^{-3}$	$(2.83\pm0.03\pm0.35)\times10^{-3}$
10-20%	$(2.08 \pm 0.02 \pm 0.22) \times 10^{-3}$	$(1.94 \pm 0.03 \pm 0.24) imes 10^{-3}$
20-40%	$(1.43 \pm 0.01 \pm 0.15) \times 10^{-3}$	$(1.43 \pm 0.02 \pm 0.17) \times 10^{-3}$
40-60%	$(8.93\pm0.08\pm0.93)\times10^{-4}$	$(9.06\pm0.15\pm1.09)\times10^{-4}$
60-100%	$(2.89\pm0.05\pm0.30)\times10^{-4}$	$(3.02\pm0.07\pm0.36)\times10^{-4}$

the p_{T} -differential yield in the region where the spectrum is measured and by integrating the fit result in the unmeasured region at low and high transverse momenta. While the fraction of the extrapolated yield at high $p_{\rm T}$ is negligible, the fraction at low $p_{\rm T}$ ranges from 23% at high to 38% at low multiplicities. The uncertainty introduced by this extrapolation is estimated by comparing the result obtained with the $m_{\rm T}$ -exponential fit to fit results from several alternative functional forms (Boltzmann, Blast-wave [43], and $p_{\rm T}$ -exponential).

Fig. 4 shows the \overline{d}/d ratios as a function of $p_{\rm T}$ for all multiplicity intervals. The ratios are found to be consistent with unity within uncertainties. This behaviour is expected, since thermal and coalescence models predict that the \overline{d}/d ratio is given by $(\overline{p}/p)^2$



Fig. 4. Anti-deuteron to deuteron production ratio for the five multiplicity classes. All ratios are compatible with unity, indicated as a dashed grey line. The vertical bars represent the statistical errors while the empty boxes show the total systematic uncertainty.

(see for instance [15]) and the \bar{p}/p ratio measured in p–Pb collisions is consistent with unity for all multiplicity intervals [18].

The rare production of A > 2 nuclei only allows the extraction of minimum-bias spectra for ³He and ³He with the available statistics and thus the result is normalised to all non-single diffractive (NSD) events. In total, 40 ³He nuclei are observed, while about 29400 tracks from \overline{d} are reconstructed in the same data sample. The corresponding spectra are shown in Fig. 5 together with a $m_{\rm T}$ -exponential fit which is used for the extraction of the dN/dy and $\langle p_{\rm T} \rangle$ of the spectra. The fit is performed such that the residuals to both the ³He and ³He spectrum are minimised simultaneously. The fraction of the extrapolated yield corresponds to about 58%. The uncertainty introduced by this extrapolation is also estimated by comparing the result obtained with the $m_{\rm T}$ -exponential fit to fit results from several alternative functional forms (Boltzmann, Blastwave [43], and p_T -exponential). A p_T -integrated yield of dN/dy = $(1.36\pm0.16(stat)\pm0.52(syst))\times10^{-6}$ and an average transverse momentum of $\langle p_T \rangle = (1.78 \pm 0.11(\text{stat}) \pm 0.77(\text{syst}))$ GeV/c are obtained.

The yields of p, d and ³He for NSD p–Pb events and normalised to their spin degeneracy are shown in Fig. 6 as a function of the mass number A together with results for inelastic pp collisions and central Pb-Pb collisions. An exponential decrease with increasing A is observed in all cases, yet with different slopes. The penalty factor, i.e. the reduction of the yield for each additional nucleon, is obtained from a fit to the data and a value of 635 ± 90 in p-Pb collisions is found which is significantly larger than the factor of 359 ± 41 which was observed for central Pb–Pb collisions [3]. The penalty factor obtained for the inelastic pp collisions [12] is found to be 942 ± 107 . Such an exponential decrease of the (anti-)nuclei yield with mass number has also been observed at lower incident energies in heavy-ion [1,44–46] as well as in p–A collisions [47].

3.2. Coalescence parameter

In the traditional coalescence model, deuterons and other light nuclei are formed by protons and neutrons, which are close in



Fig. 5. Transverse momentum distribution of ³He and ³He for all NSD collisions (N_{NSD}). The vertical bars represent the statistical errors while the empty boxes show the total systematic uncertainty. The line represents a χ^2 fit with a m_{T} -exponential function (see text for details).



Fig. 6. Production yield dN/dy normalised by the spin degeneracy as a function of the mass number for inelastic pp collisions, minimum-bias p-Pb and central Pb-Pb collisions [12,13,18,48,49]. The empty boxes represent the total systematic uncertainty while the statistical errors are shown by the vertical bars. The lines represent fits with an exponential function.

phase space. In this picture, the deuteron momentum spectra are related to those of its constituent nucleons via [50,51]

$$E_{\rm d} \, \frac{{\rm d}^3 N_{\rm d}}{{\rm d} p_{\rm d}^3} = B_2 \, \left(E_{\rm p} \, \frac{{\rm d}^3 N_{\rm p}}{{\rm d} p_{\rm p}^3} \right)^2, \tag{1}$$

where the momentum of the deuteron is given by $p_d = 2p_p$. Since the neutron spectra are experimentally not accessible, they are approximated by the proton spectra. The value of B_2 is computed as a function of event multiplicity and transverse momentum as the ratio between the deuteron yield measured at $p_{T} = p_{T,d}$ and the square of the proton yield at $p_{T,p} = 0.5 p_{T,d}$. The obtained B_2 -values are shown in Fig. 7. In its simplest implementation, the coalescence model for uncorrelated particle emission from a pointlike source predicts that the observed B_2 -values are independent of $p_{\rm T}$ and of event multiplicity (called "simple coalescence" in the following). Within uncertainties and given the current width of the multiplicity classes, the observed $p_{\rm T}$ dependence is still compatible with the expected flat behaviour (for a detailed discussion see [6]). Moreover, a decrease of the measured B_2 parameter with increasing event multiplicity for a fixed p_{T} is observed. This effect is even more pronounced in Pb-Pb collisions [3] and a possible explana-



Fig. 7. Coalescence parameter B_2 as a function of p_T for different VOA multiplicity classes. The vertical lines represent the statistical errors and the empty boxes show the total systematic uncertainty.

tion is an increasing source volume, which can effectively reduce the coalescence probability [7,51].

3.3. Mean transverse momenta

In Fig. 8 (left), the mean values of the transverse momenta of deuterons are compared with the corresponding results for π^{\pm} , K^{\pm} , $p(\bar{p})$, and $\Lambda(\overline{\Lambda})$ [18]. As for all other particles, the $\langle p_T \rangle$ of deuterons shows an increase with increasing event multiplicity, which reflects the observed hardening of the spectra. However, it is striking that deuterons violate the mass ordering which was observed for non-composite particles [18,52]: despite their much larger mass, the $\langle p_T \rangle$ values are similar to those of $\Lambda(\overline{\Lambda})$ and only slightly higher than those of $p(\bar{p})$.

Note that simple coalescence models give a significantly different prediction for the $\langle p_T \rangle$ of deuterons with respect to hydrodynamical models. This can be best illustrated with two simplifying requirements which are approximately fulfilled in data. Firstly, the coalescence parameter is assumed flat in p_T and secondly the proton spectrum can be described by an exponential shape, i.e. $C \exp(-p_T/T)$ with two parameters *C* and *T*. In this case, the shape of the deuteron spectrum can be analytically calculated based on the definition of B_2 . Due to the self-similarity feature of the exponential function, $(\exp(x/a))^a = \exp(x)$, the spectral shape of the proton and the deuteron are then found to be identical:

$$\frac{1}{2\pi p_{\rm T}^d} \frac{{\rm d}^2 N^d}{{\rm d}y \, {\rm d}p_{\rm T}^d} = B_2 \Big(\frac{1}{2\pi p_{\rm T}^p} \frac{{\rm d}^2 N^p}{{\rm d}y \, {\rm d}p_{\rm T}^p} \Big)^2 = B_2 \Big(C \exp(-\frac{p_{\rm T}^p}{T}) \Big)^2$$
$$= B_2 \Big(C \exp(-\frac{p_{\rm T}^d}{2T}) \Big)^2 = B_2 C^2 \exp(-\frac{p_{\rm T}^d}{T}) \,. \tag{2}$$

Thus, the same $\langle p_T \rangle$ for both particles is expected and the behaviour observed in p–Pb collisions is well described by simple coalescence models. This finding can be even further substantiated by directly calculating the $\langle p_T \rangle$ of deuterons assuming a constant value of B_2 and using the measured proton spectrum as input. As shown in Fig. 8 (right), in this case, a good agreement with the data is found considering that a large fraction of the systematic uncertainty is correlated among different multiplicity bins. The Blast-Wave model [43] fails to describe the $\langle p_T \rangle$ values for deuterons using the common kinetic freeze-out parameters from [18], which describe simultaneously the spectra of pions, kaons, and protons.



Fig. 8. Mean p_T of various particle species as a function of the mean chargedparticle density at mid-rapidity for different V0A multiplicity classes. The empty boxes show the total systematic uncertainty while the shaded boxes indicate the contribution which is uncorrelated across multiplicity intervals (left). Comparison of $\langle p_T \rangle$ of protons and deuterons with the simple coalescence and the Blast-Wave model expectations. The shaded areas show the expected $\langle p_T \rangle$ for deuterons from a simple coalescence model assuming a p_T -independent B_2 as well as the calculated $\langle p_T \rangle$ for protons and deuterons from the Blast-Wave model [43] using the kinetic freeze-out parameters for pions, kaons, protons and Λ from [18] (right).



Fig. 9. Deuteron-over-proton ratio as a function of charged-particle multiplicity at mid-rapidity for pp, p–Pb and Pb–Pb collisions [3,6,12]. The empty boxes show the systematic uncertainty while the vertical lines represent the statistical uncertainty.

3.4. Deuteron-over-proton ratio

The deuteron-over-proton ratio is shown in Fig. 9 for three collision systems as a function of the charged-particle density at mid-rapidity. In Pb–Pb collisions it has been observed that the d/p ratio does not vary with centrality within uncertainties (red symbols). Such a trend is consistent with a thermal-statistical approach and the magnitude of the measured values agree with freeze-out temperatures in the range of 150-160 MeV [3]. The d/p ratio obtained in inelastic pp collisions increases with multiplicity [6]. The results in p–Pb collisions bridge the two measurements in terms of multiplicity and system size and show an increase of the d/p ratio with multiplicity. Here, the low (high) multiplicity value is compatible with the result from pp (Pb–Pb) collisions. Note that the experimental significance of this enhancement is further substantiated by considering only the part of the systematic uncertainty which is uncorrelated across multiplicity intervals.

A similar rise with multiplicity is observed for the ratios of the yields of multi-strange particles to that of pions in p–Pb collisions [53]. In this case the canonical suppression due to exact strangeness conservation in smaller systems gives a qualitative explanation [54]. An interpretation of the d/p ratio within thermal

7

models is difficult, since the measured p/π ratio in these three systems is about the same [18]. Therefore, the available parameter space for a change in the freeze-out temperature or a suppression due to exact conservation of baryon number is limited [55]. Coalescence models are able to explain such an observation. The probability of forming a deuteron increases with the nucleon density and thus also with the charged-particle density. The results from pp and p–Pb collisions at low charged-particle density fit with this concept.

4. Conclusions

The production of deuterons and ³He and their antiparticles in p-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV has been studied at midrapidity. The results on deuteron production in p-Pb collisions exhibit a continuous evolution with multiplicity between pp and Pb-Pb collisions. The production of complex nuclei shows an exponential decrease with mass (number). The penalty factor (decrease of yield for each additional nucleon) is larger than the one observed in central Pb-Pb collisions and smaller than the one measured in pp collisions. The transverse momentum distributions of deuterons become harder with increasing multiplicity. Two intriguing observations that have been recently reported by ALICE [6] in high multiplicity pp collisions are confirmed in the present paper. Firstly, the $\langle p_T \rangle$ values of deuterons are comparable to those of the much lighter Λ baryons and thus do not follow a mass ordering. This behaviour is observed for all multiplicity intervals and it is in contrast to the expectation from simple hydrodynamical models. These observations made in p-Pb collisions support a coalescence mechanism, while in Pb-Pb collisions the deuteron seems to follow the collective expansion of the fireball. Secondly, the d/p ratio rises strongly with multiplicity, while this ratio remains approximately constant as a function of multiplicity in Pb-Pb collisions, where its value agrees with thermal-model predictions.

Acknowledgements

The ALICE Collaboration would like to thank all its engineers and technicians for their invaluable contributions to the construction of the experiment and the CERN accelerator teams for the outstanding performance of the LHC complex. The ALICE Collaboration gratefully acknowledges the resources and support provided by all Grid centres and the Worldwide LHC Computing Grid (WLCG) collaboration. The ALICE Collaboration acknowledges the following funding agencies for their support in building and running the ALICE detector: A. I. Alikhanyan National Science Laboratory (Yerevan Physics Institute) Foundation (ANSL), State Committee of Science and World Federation of Scientists (WFS), Armenia; Austrian Academy of Sciences, Austrian Science Fund (FWF): [M 2467-N361 and Nationalstiftung für Forschung. Technologie und Entwicklung, Austria; Ministry of Communications and High Technologies, National Nuclear Research Center, Azerbaijan; Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq), Universidade Federal do Rio Grande do Sul (UFRGS), Financiadora de Estudos e Projetos (Finep) and Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP), Brazil; Ministry of Science & Technology of China (MSTC), National Natural Science Foundation of China (NSFC) and Ministry of Education of China (MOEC), China; Croatian Science Foundation and Ministry of Science and Education, Croatia; Centro de Aplicaciones Tecnológicas y Desarrollo Nuclear (CEADEN), Cubaenergía, Cuba; Ministry of Education, Youth and Sports of the Czech Republic, Czech Republic; The Danish Council for Independent Research | Natural Sciences, the Carlsberg Foundation and Danish National Research Foundation (DNRF), Denmark; Helsinki Institute of Physics (HIP), Finland; Commissariat à

l'Énergie Atomique (CEA), Institut National de Physique Nucléaire et de Physique des Particules (IN2P3) and Centre National de la Recherche Scientifique (CNRS) and Région des Pays de la Loire, France; Bundesministerium für Bildung und Forschung (BMBF) and GSI Helmholtzzentrum für Schwerionenforschung GmbH, Germany; General Secretariat for Research and Technology, Ministry of Education, Research and Religions, Greece: National Research Development and Innovation Office, Hungary; Department of Atomic Energy, Government of India (DAE), Department of Science and Technology, Government of India (DST), University Grants Commission, Government of India (UGC) and Council of Scientific and Industrial Research (CSIR), India; Indonesian Institute of Sciences, Indonesia; Centro Fermi - Museo Storico della Fisica e Centro Studi e Ricerche Enrico Fermi and Istituto Nazionale di Fisica Nucleare (INFN). Italy: Institute for Innovative Science and Technology. Nagasaki Institute of Applied Science (IIST), Japan Society for the Promotion of Science (JSPS) KAKENHI and Japanese Ministry of Education, Culture, Sports, Science and Technology (MEXT), Japan; Consejo Nacional de Ciencia (CONACYT) y Tecnología, through Fondo de Cooperación Internacional en Ciencia y Tecnología (FON-CICYT) and Dirección General de Asuntos del Personal Academico (DGAPA), Mexico; Nederlandse Organisatie voor Wetenschappelijk Onderzoek (NWO), Netherlands; The Research Council of Norway, Norway; Commission on Science and Technology for Sustainable Development in the South (COMSATS), Pakistan; Pontificia Universidad Católica del Perú, Peru; Ministry of Science and Higher Education and National Science Centre, Poland; Korea Institute of Science and Technology Information and National Research Foundation of Korea (NRF), Republic of Korea; Ministry of Education and Scientific Research, Institute of Atomic Physics and Ministry of Research and Innovation and Institute of Atomic Physics, Romania; Joint Institute for Nuclear Research (JINR), Ministry of Education and Science of the Russian Federation, National Research Centre Kurchatov Institute, Russian Science Foundation and Russian Foundation for Basic Research, Russia; Ministry of Education, Science, Research and Sport of the Slovak Republic, Slovakia; National Research Foundation of South Africa, South Africa; Swedish Research Council (VR) and Knut & Alice Wallenberg Foundation (KAW), Sweden; European Organization for Nuclear Research, Switzerland; National Science and Technology Development Agency (NSDTA), Suranaree University of Technology (SUT) and Office of the Higher Education Commission under NRU project of Thailand, Thailand; Turkish Atomic Energy Agency (TAEK), Turkey; National Academy of Sciences of Ukraine, Ukraine; Science and Technology Facilities Council (STFC), United Kingdom; National Science Foundation of the United States of America (NSF) and United States Department of Energy, Office of Nuclear Physics (DOE NP), United States of America.

References

- STAR Collaboration, H. Agakishiev, et al., Observation of the antimatter helium-4 nucleus, Nature 473 (2011) 353, arXiv:1103.3312 [nucl-ex].
- [2] STAR Collaboration, B.I. Abelev, et al., Observation of an antimatter hypernucleus, Science 328 (2010) 58–62, arXiv:1003.2030 [nucl-ex].
- [3] ALICE Collaboration, J. Adam, et al., Production of light nuclei and anti-nuclei in pp and Pb-Pb collisions at energies available at the CERN Large Hadron Collider, Phys. Rev. C 93 (2) (2016) 024917, arXiv:1506.08951 [nucl-ex].
- [4] ALICE Collaboration, J. Adam, et al., ${}^{3}_{\Lambda}$ H and ${}^{3}_{\Lambda}\overline{H}$ production in Pb-Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV, Phys. Lett. B 754 (2016) 360–372, arXiv:1506.08453 [nucl-ex].
- [5] ALICE Collaboration, S. Acharya, et al., Measurement of deuteron spectra and elliptic flow in Pb–Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV at the LHC, Eur. Phys. J. C77 (10) (2017) 658, arXiv:1707.07304 [nucl-ex].
- [6] ALICE Collaboration, S. Acharya, et al., Multiplicity dependence of (anti-) deuteron production in pp collisions at $\sqrt{s} = 7$ TeV, Phys. Lett. B 794 (2019) 50–63, arXiv:1902.09290 [nucl-ex].

- [7] K. Blum, K.C.Y. Ng, R. Sato, M. Takimoto, Cosmic rays, antihelium, and an old navy spotlight, Phys. Rev. D96 (10) (2017) 103021, arXiv:1704.05431 [astro-ph. HE1.
- [8] V. Poulin, P. Salati, I. Cholis, M. Kamionkowski, J. Silk, Where do the AMS-02 antihelium events come from?, Phys. Rev. D 99 (2) (2019) 023016, arXiv:1808, 08961 [astro-ph.HE].
- [9] N. Tomassetti, A. Oliva, Production of cosmic-ray antinuclei in the Galaxy and background for dark matter searches, PoS EPS-HEP2017 (2017) 620, arXiv:1712. 03177 [astro-ph.HE].
- [10] M. Korsmeier, F. Donato, N. Fornengo, Prospects to verify a possible dark matter hint in cosmic antiprotons with antideuterons and antihelium, Phys. Rev. D97 (10) (2018) 103011, arXiv:1711.08465 [astro-ph.HE].
- [11] Y. Cui, I.D. Mason, L. Randall, General analysis of antideuteron searches for dark matter, J. High Energy Phys. 1011 (2010) 017, arXiv:1006.0983 [hep-ph].
- [12] ALICE Collaboration, S. Acharya, et al., Production of deuterons, tritons, ³He nuclei and their antinuclei in pp collisions at $\sqrt{s} = 0.9$, 2.76 and 7 TeV, Phys. Rev. C 97 (2) (2018) 024615, arXiv:1709.08522 [nucl-ex].
- [13] ALICE Collaboration, S. Acharya, et al., Production of ⁴He and ⁴He in Pb-Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV at the LHC, Nucl. Phys. A 971 (2018) 1–20, arXiv: 1710.07531 [nucl-ex].
- [14] A. Andronic, P. Braun-Munzinger, J. Stachel, H. Stoecker, Production of light nuclei, hypernuclei and their antiparticles in relativistic nuclear collisions, Phys. Lett. B 697 (2011) 203-207, arXiv:1010.2995 [nucl-th].
- [15] J. Cleymans, S. Kabana, I. Kraus, H. Oeschler, K. Redlich, et al., Antimatter production in proton-proton and heavy-ion collisions at ultrarelativistic energies, Phys. Rev. C 84 (2011) 054916, arXiv:1105.3719 [hep-ph].
- [16] ALICE Collaboration, B. Abelev, et al., Long-range angular correlations on the near and away side in *p*-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV, Phys. Lett. B 719 (2013) 29-41, arXiv:1212.2001 [nucl-ex].
- [17] ALICE Collaboration, B. Abelev, et al., Long-range angular correlations of π , K and p in p-Pb collisions at $\sqrt{s_{NN}}$ = 5.02 TeV, Phys. Lett. B 726 (2013) 164–177, arXiv:1307.3237 [nucl-ex].
- [18] ALICE Collaboration, B. Abelev, et al., Multiplicity dependence of pion, kaon, proton and lambda production in p-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV, Phys. Lett. B 728 (2014) 25-38, arXiv:1307.6796 [nucl-ex].
- [19] CMS Collaboration, S. Chatrchyan, et al., Observation of long-range near-side angular correlations in proton-lead collisions at the LHC, Phys. Lett. B 718 (2013) 795-814, arXiv:1210.5482 [nucl-ex].
- [20] ATLAS Collaboration, G. Aad, et al., Observation of associated near-side and away-side long-range correlations in $\sqrt{s_{NN}} = 5.02$ TeV proton-lead collisions with the ATLAS detector, Phys. Rev. Lett. 110 (18) (2013) 182302, arXiv:1212. 5198 [hep-ex].
- [21] ATLAS Collaboration, G. Aad, et al., Measurement with the ATLAS detector of multi-particle azimuthal correlations in p+Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV, Phys. Lett. B 725 (2013) 60-78, arXiv:1303.2084 [hep-ex].
- [22] CMS Collaboration, S. Chatrchyan, et al., Multiplicity and transverse momentum dependence of two- and four-particle correlations in pPb and PbPb collisions. Phys. Lett. B 724 (2013) 213-240, arXiv:1305.0609 [nucl-ex].
- [23] C. Loizides, Experimental overview on small collision systems at the LHC, Nucl. Phys. A 956 (2016) 200-207, arXiv:1602.09138 [nucl-ex].
- [24] J.L. Nagle, W.A. Zajc, Small system collectivity in relativistic hadronic and nuclear collisions, Annu, Rev. Nucl. Part, Sci. 68 (2018) 211-235, arXiv:1801.03477 [nucl-ex].
- [25] ALICE Collaboration, B. Alessandro, et al., ALICE: physics performance report, volume II, J. Phys. G 32 (2006) 1295-2040.
- [26] ALICE Collaboration, P. Cortese, et al., ALICE: physics performance report, volume I. I. Phys. G 30 (2004) 1517-1763.
- [27] ALICE Collaboration, K. Aamodt, et al., The ALICE experiment at the CERN LHC, JINST 3 (2008) S08002.
- [28] ALICE Collaboration, K. Aamodt, et al., Midrapidity antiproton-to-proton ratio in pp collisions at $\sqrt{s} = 0.9$ and 7 TeV measured by the ALICE experiment, Phys. Rev. Lett. 105 (2010) 072002, arXiv:1006.5432 [hep-ex].
- [29] ALICE Collaboration, B. Abelev, et al., Performance of the ALICE experiment at the CERN LHC, Int. J. Mod. Phys. A 29 (2014) 1430044, arXiv:1402.4476 [nuclexl
- [30] ALICE Collaboration, K. Aamodt, et al., Alignment of the ALICE Inner Tracking System with cosmic-ray tracks, JINST 5 (2010) P03003, arXiv:1001.0502 [physics.ins-det].

- [31] J. Alme, et al., The ALICE TPC, a large 3-dimensional tracking device with fast readout for ultra-high multiplicity events, Nucl. Instrum. Meth. A 622 (2010) 316-367, arXiv:1001.1950 [physics.ins-det].
- [32] A. Akindinov, A. Alici, P. Antonioli, S. Arcelli, M. Basile, et al., Results of the ALICE time-of-flight detector from the 2009 cosmic-ray data taking, Eur. Phys. I. C 68 (2010) 601-607.
- [33] A. Akindinov, et al., Performance of the ALICE Time-Of-Flight detector at the LHC, Eur. Phys. J. Plus 128 (2013) 44.
- [34] ALICE Collaboration, J. Adam, et al., Determination of the event collision time with the ALICE detector at the LHC, Eur. Phys. J. Plus 132 (2) (2017) 99, arXiv: 1610.03055 [physics.ins-det].
- [35] ALICE Collaboration, B. Abelev, et al., Measurement of visible cross sections in proton-lead collisions at $\sqrt{s_{\rm NN}}$ = 5.02 TeV in van der Meer scans with the ALICE detector, IINST 9 (11) (2014) P11003, arXiv:1405.1849 [nucl-ex].
- [36] ALICE Collaboration, B. Abelev, et al., Pseudorapidity density of charged particles in p + Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV, Phys. Rev. Lett. 110 (3) (2013) 032301, arXiv:1210.3615 [nucl-ex].
- [37] ALICE Collaboration, J. Adam, et al., Multiplicity dependence of charged pion, kaon, and (anti)proton production at large transverse momentum in p-Pb collisions at $\sqrt{s_{\rm NN}}$ = 5.02 TeV, Phys. Lett. B 760 (2016) 720–735, arXiv:1601.03658 [nucl-ex].
- [38] S. Roesler, R. Engel, J. Ranft, The Monte Carlo event generator DPMJET-III, arXiv: hep-ph/0012252 [hep-ph].
- [39] R. Brun, F. Bruyant, F. Carminati, S. Giani, M. Maire, A. McPherson, G. Patrick, L. Urban, GEANT Detector Description and Simulation Tool, CERN Report W5013, 1994
- [40] GEANT4 Collaboration, S. Agostinelli, et al., GEANT4: a simulation toolkit, Nucl. Instrum. Meth. A 506 (2003) 250-303.
- [41]V. Uzhinsky, J. Apostolakis, A. Galoyan, G. Folger, V. Grichine, et al., Antinucleus-nucleus cross sections implemented in Geant4, Phys. Lett. B 705 (2011) 235-239.
- [42] R. Hagedorn, Statistical thermodynamics of strong interactions at highenergies, Nuovo Cim. Suppl. 3 (1965) 147-186.
- [43] E. Schnedermann, J. Sollfrank, U.W. Heinz, Thermal phenomenology of hadrons from 200-A/GeV S+S collisions, Phys. Rev. C 48 (1993) 2462-2475, arXiv:nuclth/9307020
- [44] E814 Collaboration, J. Barrette, et al., Production of light nuclei in relativistic heavy ion collisions, Phys. Rev. C 50 (1994) 1077-1084.
- [45] P. Braun-Munzinger, J. Stachel, Production of strange clusters and strange matter in nucleus-nucleus collisions at the AGS, J. Phys. G 21 (1995) L17-L20, arXiv:nucl-th/9412035 [nucl-th].
- [46] NA52 Collaboration, R. Arsenescu, et al., An investigation of the anti-nuclei and nuclei production mechanism in Pb + Pb collisions at 158-A-GeV, New J. Phys. 5 (2003) 150.
- [47] Yu.M. Antipov, et al., Observation of anti-he-3, Yad. Fiz. 12 (1970) 311-322.
- [48] ALICE Collaboration, B. Abelev, et al., Centrality dependence of π , K, p production in Pb-Pb collisions at $\sqrt{s_{NN}}$ = 2.76 TeV, Phys. Rev. C 88 (2013) 044910, arXiv:1303.0737 [hep-ex].
- [49] ALICE Collaboration, J. Adam, et al., Measurement of pion, kaon and proton production in proton–proton collisions at $\sqrt{s} = 7$ TeV, Eur. Phys. J. C 75 (5) (2015) 226, arXiv:1504.00024 [nucl-ex].
- [50] S. Butler, C. Pearson, Deuterons from high-energy proton bombardment of matter, Phys. Rev. 129 (1963) 836-842.
- [51] R. Scheibl, U.W. Heinz, Coalescence and flow in ultrarelativistic heavy ion collisions, Phys. Rev. C 59 (1999) 1585-1602, arXiv:nucl-th/9809092 [nucl-th].
- [52] ALICE Collaboration, J. Adam, et al., Production of K^{*} (892)⁰ and ϕ (1020) in p-Pb collisions at $\sqrt{s_{NN}}$ = 5.02 TeV, Eur. Phys. J. C 76 (5) (2016) 245, arXiv: 1601.07868 [nucl-ex].
- [53] ALICE Collaboration, J. Adam, et al., Multi-strange baryon production in p-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV, Phys. Lett. B 758 (2016) 389-401, arXiv:1512. 07227 [nucl-ex].
- [54] ALICE Collaboration, S. Acharya, et al., Multiplicity dependence of light-flavor hadron production in pp collisions at $\sqrt{s} = 7$ TeV, Phys. Rev. C 99 (2) (2019) 024906, arXiv:1807.11321 [nucl-ex].
- [55] V. Vovchenko, B. Dönigus, H. Stoecker, Multiplicity dependence of light nuclei production at LHC energies in the canonical statistical model, Phys. Lett. B 785 (2018) 171-174, arXiv:1808.05245 [hep-ph].

ALICE Collaboration

- S. Acharya¹⁴¹, D. Adamová⁹³, S.P. Adhya¹⁴¹, A. Adler⁷³, J. Adolfsson⁷⁹, M.M. Aggarwal⁹⁸,
- G. Aglieri Rinella³⁴, M. Agnello³¹, N. Agrawal^{48,10}, Z. Ahammed¹⁴¹, S. Ahmad¹⁷, S.U. Ahn⁷⁵, A. Akindinov⁹⁰, M. Al-Turany¹⁰⁵, S.N. Alam¹⁴¹, D.S.D. Albuquerque¹²², D. Aleksandrov⁸⁶, B. Alessandro⁵⁸, H.M. Alfanda⁶, R. Alfaro Molina⁷¹, B. Ali¹⁷, Y. Ali¹⁵, A. Alici^{10,53,27}, A. Alkin²,

9

J. Alme²², T. Alt⁶⁸, L. Altenkamper²², I. Altsybeev¹¹², M.N. Anaam⁶, C. Andrei⁴⁷, D. Andreou³⁴, H.A. Andrews ¹⁰⁹, A. Andronic ¹⁴⁴, M. Angeletti ³⁴, V. Anguelov ¹⁰², C. Anson ¹⁶, T. Antičić ¹⁰⁶, H.A. Andrews ¹⁰⁵, A. Andronic ¹⁴⁴, M. Angeletti ³⁴, V. Anguelov ¹⁰², C. Anson ¹⁶, T. Antičić ¹⁰⁶, F. Antinori ⁵⁶, P. Antonioli ⁵³, R. Anwar ¹²⁵, N. Apadula ⁷⁸, L. Aphecetche ¹¹⁴, H. Appelshäuser ⁶⁸, S. Arcelli ²⁷, R. Arnaldi ⁵⁸, M. Arratia ⁷⁸, I.C. Arsene ²¹, M. Arslandok ¹⁰², A. Augustinus ³⁴, R. Averbeck ¹⁰⁵, S. Aziz ⁶¹, M.D. Azmi ¹⁷, A. Badalà ⁵⁵, Y.W. Baek ⁴⁰, S. Bagnasco ⁵⁸, X. Bai ¹⁰⁵, R. Bailhache ⁶⁸, R. Bala ⁹⁹, A. Baldisseri ¹³⁷, M. Ball ⁴², S. Balouza ¹⁰³, R.C. Baral ⁸⁴, R. Barbera ²⁸, L. Barioglio ²⁶, G.G. Barnaföldi ¹⁴⁵, L.S. Barnby ⁹², V. Barret ¹³⁴, P. Bartalini ⁶, K. Barth ³⁴, E. Bartsch ⁶⁸, F. Baruffaldi ²⁹, N. Bastid ¹³⁴, S. Basu ¹⁴³, G. Batigne ¹¹⁴, B. Batyunya ⁷⁴, P.C. Batzing ²¹, D. Bauri ⁴⁸, J.L. Bazo Alba ¹¹⁰, I.G. Bearden ⁸⁷, C. Bedda ⁶³, N.K. Behera ⁶⁰, I. Belikov ¹³⁶, F. Bellini ³⁴, R. Bellwied ¹²⁵, V. Belyaev ⁹¹, G. Bencedi ¹⁴⁵, S. Beole ²⁶, A. Bercuci ⁴⁷, Y. Berdnikov ⁹⁶, D. Berenyi ¹⁴⁵, R.A. Bertens ¹³⁰, D. Berzano ⁵⁸, M.G. Besoiu ⁶⁷, L. Betev ³⁴, A. Bhasin ⁹⁹, I.R. Bhat ⁹⁹, M.A. Bhat ³, H. Bhatt ⁴⁸, B. Bhattacharjee ⁴¹, A. Bianchi ²⁶, L. Bianchi ^{125,26}, N. Bianchi ⁵¹, I. Bielčík ³⁷, I. Bielčíková ⁹³, A. Bilandzic ^{117,103}, C. Biro ¹⁴⁵, R. Biswas ³ L. Belev ¹⁷, A. Blashi¹⁰, I.K. Blat¹⁰, M.A. Blat¹⁰, H. Blat¹⁰, B. Blatt¹⁰, B. Blattacharjee¹¹, A. Blanchi¹⁰, L. Bianchi^{125,26}, N. Bianchi⁵¹, J. Bielčík³⁷, J. Bielčíková⁹³, A. Bilandzic^{117,103}, G. Biro¹⁴⁵, R. Biswas³, S. Biswas³, J.T. Blair¹¹⁹, D. Blau⁸⁶, C. Blume⁶⁸, G. Boca¹³⁹, F. Bock^{94,34}, A. Bogdanov⁹¹, L. Boldizsár¹⁴⁵, A. Bolozdynya⁹¹, M. Bombara³⁸, G. Bonomi¹⁴⁰, H. Borel¹³⁷, A. Borissov^{144,91}, M. Borri¹²⁷, H. Bossi¹⁴⁶, E. Botta²⁶, L. Bratrud⁶⁸, P. Braun-Munzinger¹⁰⁵, M. Bregant¹²¹, T.A. Broker⁶⁸, M. Broz³⁷, E.J. Brucken⁴³, E. Bruna⁵⁸, G.E. Bruno^{33,104}, M.D. Buckland¹²⁷, D. Budnikov¹⁰⁷, H. Buesching⁶⁸, S. Bufalino³¹, O. Bugnon¹¹⁴, P. Buhler¹¹³, P. Buncic³⁴, Z. Buthelezi⁷², J.B. Butt¹⁵, J.T. Buxton⁹⁵, D. Caffarri⁸⁸, A. Caliva¹⁰⁵, E. Calvo Villar¹¹⁰, R.S. Camacho⁴⁴, P. Camerini²⁵, A.A. Capon¹¹³, F. Carnesecchi¹⁰, J. Castillo Castellanos¹³⁷, A.J. Castro¹³⁰, E.A.R. Casula⁵⁴, F. Catalano³¹, C. Ceballos Sanchez⁵², P. Chakraborty⁴⁸, S. Chandra¹⁴¹, B. Chang¹²⁶, W. Chang⁶, S. Chapeland³⁴, M. Chartier¹²⁷, S. Chattopadhyay¹⁰⁸, A. Chauvin²⁴, C. Cheshkov¹³⁵, B. Cheynis¹³⁵, V. Chibante Barroso³⁴, D.D. Chinellato¹²², S. Cho⁶⁰, P. Chochula³⁴, T. Chowdhury¹³⁴, P. Christakoglou⁸⁸, C.H. Christianaen⁸⁷, P. Christianaen⁷⁹, T. Chesi, ¹³³, C. Cir, J. ⁵⁴, J. ⁵⁴, J. ⁵⁴, J. ⁵⁴, J. ⁵⁵, S. ⁵⁴, S. ⁵⁵, S. ⁵⁶, P. ⁵⁶, J. ⁵⁶, S. ⁵⁷, S. ⁵⁶, S. ⁵⁷, S. ⁵⁶, S. ⁵⁶, S. ⁵⁶, S. ⁵⁶, S. ⁵⁶, S. ⁵⁷, S. ⁵⁷, S. ⁵⁶, S. ⁵⁷, S. ⁵⁷, S. ⁵⁶, S. ⁵⁶, S. ⁵⁶, S. ⁵⁶, S. ⁵⁶, S. ⁵⁶, S. ⁵⁷, S. ⁵⁷, S. ⁵⁶, S. ⁵⁷, S. ⁵⁷, S. ⁵⁶, S. ⁵⁷, S. ⁵⁷, S. ⁵⁷, S. ⁵⁷, S. ⁵⁶, S. ⁵⁷, S. ⁵⁶, S. ⁵⁶, S. ⁵⁶, S. ⁵⁶, S. ⁵⁶, S. ⁵⁷, S. ⁵⁷, S. ⁵⁷, S. ⁵⁷, S. ⁵⁷, S. ⁵⁷, S. ⁵⁶, S. ⁵⁶, S. ⁵⁶, S. ⁵⁶, S. ⁵⁶, S. ⁵⁷, S. ⁵⁶, S. ⁵⁶, S. ⁵⁶, S. ⁵⁶, S. ⁵⁶, S. ⁵⁷, C.H. Christensen⁸⁷, P. Christiansen⁷⁹, T. Chujo¹³³, C. Cicalo⁵⁴, L. Cifarelli^{10,27}, F. Cindolo⁵³, J. Cleymans¹²⁴, F. Colamaria⁵², D. Colella⁵², A. Collu⁷⁸, M. Colocci²⁷, M. Concas^{58,ii}, G. Conesa Balbastre⁷⁷, Z. Conesa del Valle⁶¹, G. Contin^{59,127}, J.G. Contreras³⁷, T.M. Cormier⁹⁴, Y. Corrales Morales ^{58,26}, P. Cortese ³², M.R. Cosentino ¹²³, F. Costa ³⁴, S. Costanza ¹³⁹, J. Crkovská ⁶¹, Y. Corrales Morales ^{58,26}, P. Cortese ³², M.R. Cosentino ¹²³, F. Costa ³⁴, S. Costanza ¹³⁹, J. Crkovská ⁶¹, P. Crochet ¹³⁴, E. Cuautle ⁶⁹, L. Cunqueiro ⁹⁴, D. Dabrowski ¹⁴², T. Dahms ^{103,117}, A. Dainese ⁵⁶, F.P.A. Damas ^{137,114}, S. Dani ⁶⁵, M.C. Danisch ¹⁰², A. Danu ⁶⁷, D. Das ¹⁰⁸, I. Das ¹⁰⁸, P. Das ³, S. Das ³, A. Dash ⁸⁴, S. Dash ⁴⁸, A. Dashi ¹⁰³, S. De ^{84,49}, A. De Caro ³⁰, G. de Cataldo ⁵², C. de Conti ¹²¹, J. de Cuveland ³⁹, A. De Falco ²⁴, D. De Gruttola ¹⁰, N. De Marco ⁵⁸, S. De Pasquale ³⁰, R.D. De Souza ¹²², S. Deb ⁴⁹, H.F. Degenhardt ¹²¹, K.R. Deja ¹⁴², A. Deloff ⁸³, S. Delsanto ^{131,26}, P. Dhankher ⁴⁸, D. Di Bari ³³, A. Di Mauro ³⁴, R.A. Diaz ⁸, T. Dietel ¹²⁴, P. Dillenseger ⁶⁸, Y. Ding ⁶, R. Divià ³⁴, Ø. Djuvsland ²², U. Dmitrieva ⁶², A. Dobrin ^{34,67}, B. Dönigus ⁶⁸, O. Dordic ²¹, A.K. Dubey ¹⁴¹, A. Dubla ¹⁰⁵, S. Dudi ⁹⁸, M. Dukhishyam ⁸⁴, P. Dupieux ¹³⁴, R.J. Ehlers ¹⁴⁶, D. Elia ⁵², H. Engel ⁷³, E. Epple ¹⁴⁶, B. Erazmus ¹¹⁴, F. Erhardt ⁹⁷, A. Erokhin ¹¹², M.R. Ersdal ²², B. Espagnon ⁶¹, G. Eulisse ³⁴, J. Eum ¹⁸, D. Evans ¹⁰⁹, S. Evdokimov ⁸⁹, L. Fabbietti ^{117,103}, M. Faggin ²⁹, J. Faivre ⁷⁷, A. Fantoni ⁵¹, M. Fasel ⁹⁴, P. Fecchio ³¹, A. Feliciello ⁵⁸, G. Feofilov ¹¹², A. Fernández Téllez ⁴⁴, A. Ferrero ¹³⁷, A. Ferretti ²⁶, A. Festanti ³⁴, V.I.G. Feuillard ¹⁰², I. Figiel ¹¹⁸, S. Filchagin ¹⁰⁷, D. Finogeev ⁶², E.M. Fionda ²², G. Fiorenza ⁵², E. Flor ¹²⁵. A. Ferrero ¹⁰⁵, G. Feorliov¹¹², A. Ferrandez Tellez¹¹, A. Ferrero¹⁰⁵, A. Ferretti¹⁰⁵, A. Festanti¹⁰⁷,
V.J.G. Feuillard¹⁰², J. Figiel¹¹⁸, S. Filchagin¹⁰⁷, D. Finogeev⁶², F.M. Fionda²², G. Fiorenza⁵², F. Flor¹²⁵,
S. Foertsch⁷², P. Foka¹⁰⁵, S. Fokin⁸⁶, E. Fragiacomo⁵⁹, U. Frankenfeld¹⁰⁵, G.G. Fronze²⁶, U. Fuchs³⁴,
C. Furget⁷⁷, A. Furs⁶², M. Fusco Girard³⁰, J.J. Gaardhøje⁸⁷, M. Gagliardi²⁶, A.M. Gago¹¹⁰, A. Gal¹³⁶,
C.D. Galvan¹²⁰, P. Ganoti⁸², C. Garabatos¹⁰⁵, E. Garcia-Solis¹¹, K. Garg²⁸, C. Gargiulo³⁴, A. Garibli⁸⁵,
K. Garner¹⁴⁴, P. Gasik^{103,117}, E.F. Gauger¹¹⁹, M.B. Gay Ducati⁷⁰, M. Germain¹¹⁴, J. Ghosh¹⁰⁸,
P. Ghosh¹⁴¹, S.K. Ghosh³, P. Gianotti⁵¹, P. Giubellino^{105,58}, P. Giubilato²⁹, P. Glässel¹⁰²,
D.M. Coméz Corzi⁷¹, A. Comez Pamirez⁷³, V. Conzalez¹⁰⁵, P. Conzález ⁷⁴⁴, S. Corbunov³⁹ D.M. Goméz Coral ⁷¹, A. Gomez Ramirez ⁷³, V. Gonzalez ¹⁰⁵, P. González-Zamora ⁴⁴, S. Gorbunov ³⁹, L. Görlich ¹¹⁸, S. Gotovac ³⁵, V. Grabski ⁷¹, L.K. Graczykowski ¹⁴², K.L. Graham ¹⁰⁹, L. Greiner ⁷⁸, A. Grelli ⁶³, C. Grigoras³⁴, V. Grigoriev⁹¹, A. Grigoryan¹, S. Grigoryan⁷⁴, O.S. Groettvik²², J.M. Gronefeld¹⁰⁵, F. Grosa³¹, J.F. Grosse-Oetringhaus³⁴, R. Grosso¹⁰⁵, R. Guernane⁷⁷, B. Guerzoni²⁷, M. Guittiere¹¹⁴, K. Gulbrandsen⁸⁷, T. Gunji¹³², A. Gupta⁹⁹, R. Gupta⁹⁹, I.B. Guzman⁴⁴, R. Haake^{34,146}, M.K. Habib¹⁰⁵, C. Hadjidakis⁶¹, H. Hamagaki⁸⁰, G. Hamar¹⁴⁵, M. Hamid⁶, R. Hannigan¹¹⁹, M.R. Haque⁶³, A. Harlenderova ¹⁰⁵, J.W. Harris ¹⁴⁶, A. Harton ¹¹, J.A. Hasenbichler ³⁴, H. Hassan ⁷⁷, D. Hatzifotiadou ^{10,53}, P. Hauer ⁴², S. Hayashi ¹³², S.T. Heckel ⁶⁸, E. Hellbär ⁶⁸, H. Helstrup ³⁶, A. Herghelegiu ⁴⁷, E.G. Hernandez ⁴⁴, G. Herrera Corral ⁹, F. Herrmann ¹⁴⁴, K.F. Hetland ³⁶, T.E. Hilden ⁴³, H. Hillemanns ³⁴,

 C. Hills ¹²⁷, B. Hippolyte ¹³⁶, B. Hohlweger ¹⁰³, D. Horak ³⁷, S. Hornung ¹⁰⁵, R. Hosokawa ¹³³, P. Hristov ³⁴, C. Huang ⁶¹, C. Hughes ¹³⁰, P. Huhn ⁶⁸, T.J. Humanic ⁹⁵, H. Hushnud ¹⁰⁸, L.A. Husova ¹⁴⁴, N. Hussain ⁴¹, S.A. Hussain ¹⁵, T. Hussain ¹⁷, D. Hutter ³⁹, D.S. Hwang ¹⁹, J.P. Iddon ^{127,34}, R. Ilkaev ¹⁰⁷, M. Inaba ¹³³, M. Ippolitov ⁸⁶, M.S. Islam ¹⁰⁸, M. Ivanov ¹⁰⁵, V. Ivanov ⁹⁶, V. Izucheev ⁸⁹, B. Jacak ⁷⁸, N. Jacazio ²⁷, P.M. Jacobs ⁷⁸, M.B. Jadhav ⁴⁸, S. Jadlovska ¹¹⁶, J. Jadlovsky ¹¹⁶, S. Jaelani ⁶³, C. Jahnke ¹²¹, M.J. Jakubowska ¹⁴², M.A. Janik ¹⁴², M. Jercic ⁹⁷, O. Jevons ¹⁰⁹, R.T. Jimenez Bustamante ¹⁰⁵, M. Jin ¹²⁵, F. Jonas ^{144,94}, P.G. Jones ¹⁰⁹, A. Jusko ¹⁰⁹, P. Kalinak ⁶⁴, A. Kalweit ³⁴, J.H. Kang ¹⁴⁷, V. Kaplin ⁹¹, S. Kar ⁶, A. Karasu Uysal ⁷⁶, O. Karavichev ⁶², T. Karavicheva ⁶², P. Karczmarczyk ³⁴, E. Karpechev ⁶², U. Kebschull ⁷³, R. Keidel ⁴⁶, M. Keil ³⁴, B. Ketzer ⁴², Z. Khabanova ⁸⁸, A.M. Khan ⁶, S. Khan ¹⁷, S.A. Khan ¹⁴¹, A. Khanzadeev ⁹⁶, Y. Kharlov ⁸⁹, A. Khatun ¹⁷, J. Kim ¹⁴⁷, J.S. Kim ⁴⁰, J. Kim ¹⁰², J. Kim ¹²⁶, E.J. Kim ¹⁴⁷, T. Kim ¹⁴⁷, J. Kim ¹⁴⁷, J.S. Kim ⁴⁰, J. Kim ¹⁰², J. Kim ¹⁴⁷, J. Kim ¹⁴⁷, S. Kirsch ³⁹, I. Kisel ³⁹, S. Kiselev ⁹⁰, A. Kisiel ¹⁴², J.L. Klay ⁵, C. Klein ⁶⁸, J. Klein ⁷⁸, C. Klein ⁷⁸, C. Klein ^{-Bö}sing ¹⁴⁴, S. Klewin ¹⁰², A. Kluge ³⁴, M.L. Knichel ³⁴, A.G. Knospe ¹²⁵, C. Kobdaj ¹¹⁵, M.K. Köhler ¹⁰², T. Kollegger ¹⁰⁵, A. Kondratyev ⁷⁴, N. Kondratyeva ⁹¹, E. Kondratyk ⁸⁹, P.J. Konopka ³⁴, L. Koska ¹¹⁶, O. Kovalenko ⁸³, V. Kovalenko ¹¹², M. Kowalski ¹¹⁸, I. Králik ⁶⁴, A. Kravčáková ³⁸, L. Kreis ¹⁰⁵, M. Krivda ^{109,64}, F. Krizek ⁹³, K. Krizkova Gajdosova ³⁷, M. Krüger ⁶⁸, E. Kryshen ⁹⁶, M. Krzewicki ³⁹, A.M. Kubera ⁹⁵, V. Kučera ⁶⁰, C. Kuhn ¹³⁶, P.G. Kuijer ⁸⁸, L. Kumar ⁹⁸, S. Kumar ⁴⁸, S. Kundu ⁴⁸, P. Kurashvili ⁸³, A. Kurepin N. Klijkker, K. Kližkova G., E. Kreis G., M. Klivda G., K. Kližkova G., Kuhi 136, P.G. Kuijer 88, M. Krüger 68, E. Kryshen 96, M. Krzewicki 39, A.M. Kubera 95, V. Kučera 60, C. Kuhn 136, P.G. Kuijer 88, L. Kumar 98, S. Kundar 48, S. Kundu 84, P. Kurashvili 83, A. Kurepin 62, A.B. Kurepin 62, S. Kushpil 93, J. Kvapil 109, M.J. Kweon 60, J.Y. Kwon 60, Y. Kwon 147, S.L. La Pointe 39, P. La Rocca 28, Y.S. Lai 78, R. Langoy 129, K. Lapidus 34, 146, A. Lardeux 21, P. Larionov 51, E. Laudi 34, R. Lavicka 37, T. Lazareva 112, R. Lea 25, L. Leardini 102, S. Lee 147, F. Lehas 88, S. Lehner 113, J. Lehrbach 39, R.C. Lemmon 92, I. León Monzón 120, E.D. Lesser 20, M. Lettrich 34, P. Lévai 145, X. Li 12, X.L Li 6, J. Lien 129, R. Lietava 109, B. Lim 18, S. Lindal 21, V. Lindenstruth 39, S.W. Lindsay 127, C. Lippmann 105, M.A. Lisa 95, V. Litichevskyi 43, A. Liu 78, S. Liu 95, W.J. Llope 143, I.M. Lofnes 22, V. Loginov 91, C. Loizides 94, P. Loncar 35, X. Lopez 134, E. López Torres 8, P. Luettig 68, J.R. Luhder 144, M. Lunardon 29, G. Luparello 59, M. Lupi 73, A. Maevskaya 62, M. Mager 34, S.M. Mahmood 21, T. Mahmoud 42, A. Maire 136, R.D. Majka 146, M. Malaev 96, O.W. Malik 21, L. Malinina 74,iii, D. Mal'Kevich 90, P. Malzacher 105, A. Maronov 107, V. Manko 86, F. Manso 134, V. Manzari 52, Y. Mao 6, M. Marchisone 135, J. Mareš 66, G.V. Margagliotti 25, A. Margotti 53, J. Margutti 63, A. Marín 105, C. Markert 119, M. Marquard 68, N.A. Martin 102, P. Martinengo 34, J.L. Martinez 125, M.I. Martínez 44, G. Martínez García 114, M. Martinez Pedreira 34, S. Masciocchi 105, M. Masera 26, A. Masoni 54, L. Massacrier 61, E. Meason 114, A. Mastroserio 138, A.M. Mathis ^{103,117}, P.F.T. Matuoka 121, A. Mastori 54, L. Massacrier 61, E. Masson 114, A. Mastroserio 138, A.M. Mathis ^{103,117}, P.F.T. Matuoka 121, A. Matyia 118, C. Mayer 118, M. Mazzilli 33, M.A. Mazzoni 57, A.F. Mechler 68, F. Meddi 23, Y. Melikyan 91, A. Mastoserio 57, A.F. Mechler 68, F. Meddi 23, Y. Melikyan 91, A. Mastoserio 54, L. Massacrier 61, E. Meninno 30, M. Mohisin Khan ^{17,1v}, M. Mondal ¹⁴¹, M.M. Mondal ⁶⁵, C. Mordasini ¹⁰³, D.A. Moreira De Godoy ¹⁴⁴, LA.P. Moreno ⁴⁴, S. Moretto ²⁹, A. Morreale ¹¹⁴, A. Morsch ³⁴, T. Mrnjavac ³⁴, V. Muccifora ⁵¹, E. Mudnic ³⁵, D. Mühlheim ¹⁴⁴, S. Muhuri ¹⁴¹, J.D. Mulligan ^{78,146}, M.G. Munhoz ¹²¹, K. Münning ⁴², R.H. Munzer ⁶⁸, H. Murakami ¹³², S. Murray ⁷², L. Musa ³⁴, J. Musinsky ⁶⁴, C.J. Myers ¹²⁵, J.W. Myrcha ¹⁴², B. Naik ⁴⁸, R. Nair ⁸³, B.K. Nandi ⁴⁸, R. Nania ^{53,10}, E. Nappi ⁵², M.U. Naru ¹⁵, A.F. Nassirpour ⁷⁹, H. Natal da Luz ¹²¹, C. Nattrass ¹³⁰, R. Nayak ⁴⁸, T.K. Nayak ^{141,84}, S. Nazarenko ¹⁰⁷, R.A. Negrao De Oliveira ⁶⁸, L. Nellen ⁶⁹, S.V. Nesbo ³⁶, G. Neskovic ³⁹, B.S. Nielsen ⁸⁷, S. Nikolaev ⁸⁶, S. Nikulin ⁸⁶, V. Nikulin ⁹⁶, F. Noferini ^{10,53}, P. Nomokonov ⁷⁴, G. Nooren ⁶³, J. Norman ⁷⁷, P. Nowakowski ¹⁴², A. Nyanin ⁸⁶, J. Nystrand ²², M. Ogino ⁸⁰, A. Ohlson ¹⁰², J. Oleniacz ¹⁴², A.C. Oliveira Da Silva ¹²¹, M.H. Oliver ¹⁴⁶, C. Oppedisano ⁵⁸, R. Orava ⁴³, A. Ortiz Velasquez ⁶⁹, A. Oskarsson ⁷⁹, J. Otwinowski ¹¹⁸, K. Oyama ⁸⁰, Y. Pachmayer ¹⁰², V. Pacik ⁸⁷, D. Pagano ¹⁴⁰, G. Paić ⁶⁹, P. Palni ⁶, J. Pan ¹⁴³, A.K. Pandey ⁴⁸, S. Panebianco ¹³⁷, V. Papikyan ¹, P. Pareek ⁴⁹, J. Park ⁶⁰, J.E. Parkkila ¹²⁶, S. Parmar ⁹⁸, A. Passfeld ¹⁴⁴, S.P. Pathak ¹²⁵, R.N. Patra ¹⁴¹, B. Paul ^{24,58}, H. Pei ⁶, T. Peitzmann ⁶³, X. Peng ⁶, L.G. Pereira ⁷⁰, H. Pereira Da Costa ¹³⁷, D. Peresunko ⁸⁶, G.M. Perez ⁸, E. Perez Lezama ⁶⁸, V. Peskov ⁶⁸, Y. Pestov ⁴, V. Petráček ³⁷, M. Petrovici ⁴⁷, R.P. Pezzi ⁷⁰, S. Piano ⁵⁹, M. Pikna ¹⁴, P. Pillot ¹¹⁴, L.O.D.L. Pimentel ⁸⁷, O. Pinazza ^{53,34}, L. Pinsky ¹²⁵, S. Pisano ⁵¹, D.B. Piyarathna ¹²⁵, M. Płoskoń ⁷⁸, M. Planinic ⁹⁷, F. Pliquett ⁶⁸, J. Pluta ¹⁴², S. Pochybova ¹⁴⁵, M.G. Poghosyan ⁹⁴, B. Polichtchouk ⁸⁹, N. Poljak ⁹⁷, W. Poonsawat ¹¹⁵, A. Pop ⁴⁷, H. Poppenborg ¹⁴⁴,

S. Porteboeuf-Houssais ¹³⁴, V. Pozdniakov ⁷⁴, S.K. Prasad ³, R. Preghenella ⁵³, F. Prino ⁵⁸, C.A. Pruneau ¹⁴³, I. Pshenichnov ⁶², M. Puccio ^{34,26}, V. Punin ¹⁰⁷, K. Puranapanda ¹⁴¹, J. Putschke ¹⁴³, R.E. Quishpe ¹²⁵, S. Ragoni ¹⁰⁹, S. Raha ³, S. Rajput ⁹⁹, J. Rak ¹²⁶, A. Rakotozafindrabe ¹³⁷, L. Ramello ³², F. Rami ¹³⁶, R. Raniwala ¹⁰⁰, S. Raniwala ¹⁰⁰, S.S. Räsänen ⁴³, B.T. Rascanu ⁶⁸, R. Rath ⁴⁹, V. Ratza ⁴², I. Ravasenga ³¹, K.F. Read ^{130,94}, K. Redlich ^{83,v}, A. Rehman ²², P. Reichelt ⁶⁸, F. Reidt ³⁴, X. Ren ⁶, R. Renfordt ⁶⁸, A. Reshetin ⁶², J.-P. Revol ¹⁰, K. Reygers ¹⁰², V. Riabov ⁹⁶, T. Richert ^{79,87}, M. Richter ²¹, P. Riedler ³⁴, W. P. Let ³⁴, J. P. K. P. K. P. Revol ¹⁰, K. Reygers ¹⁰², V. Riabov ⁹⁶, T. Richert ^{79,87}, M. Richter ²¹, P. Riedler ³⁴, M. R. Reshetin ⁶², J.-P. Revol ¹⁰, K. Reygers ¹⁰², V. Riabov ⁹⁶, T. Richert ^{79,87}, M. Richter ²¹, P. Riedler ³⁴, M. R. Reshetin ⁶², J.-P. Revol ¹⁰, K. Reygers ¹⁰², V. Riabov ⁹⁶, T. Richert ^{79,87}, M. Richter ²¹, P. Riedler ³⁴, M. R. Reshetin ⁶², J.-P. Revol ¹⁰, K. Reygers ¹⁰², V. Riabov ⁹⁶, T. Richert ^{79,87}, M. Richter ²¹, P. Riedler ³⁴, M. Richter ¹¹, P. Riedler ³⁴, M. Richter ¹¹, P. Riedler ³⁴, M. Richter ¹², P. Riedler ³⁴, M. Richter ¹², P. Riedler ³⁴, M. Richter ¹², P. Riedler ³⁴, M. Richter ¹³, M. Richter ¹³, M. Richter ¹³, M. Richter ¹⁴, M. Rich W. Riegler ³⁴, F. Riggi ²⁸, C. Ristea ⁶⁷, S.P. Rode ⁴⁹, M. Rodríguez Cahuantzi ⁴⁴, K. Røed ²¹, R. Rogalev ⁸⁹, E. Rogochaya ⁷⁴, D. Rohr ³⁴, D. Röhrich ²², P.S. Rokita ¹⁴², F. Ronchetti ⁵¹, E.D. Rosas ⁶⁹, K. Roslon ¹⁴², P. Rosnet ¹³⁴, A. Rossi ²⁹, A. Rotondi ¹³⁹, F. Roukoutakis ⁸², A. Roy ⁴⁹, P. Roy ¹⁰⁸, O.V. Rueda ⁷⁹, R. Rui ²⁵, B. Rumyantsev ⁷⁴, A. Rustamov ⁸⁵, E. Ryabinkin ⁸⁶, Y. Ryabov ⁹⁶, A. Rybicki ¹¹⁸, H. Rytkonen ¹²⁶, S. Sadhu ¹⁴¹, S. Sadovsky ⁸⁹, K. Šafařík ^{37,34}, S.K. Saha ¹⁴¹, B. Sahoo ⁴⁸, P. Sahoo ⁴⁹, R. Sahoo ⁴⁹, S. Sahoo ⁶⁵, P.K. Sahu ⁶⁵, J. Saini ¹⁴¹, S. Sakai ¹³³, S. Sambyal ⁹⁹, V. Samsonov ^{91,96}, A. Sandoval ⁷¹, A. Sarkar ⁷², D. Sarkar ¹⁴³, N. Sarkar ¹⁴¹, P. Sarma ⁴¹, V.M. Sarti ¹⁰³, M.H.P. Sas ⁶³, E. Scapparone ⁵³, B. Schaefer ⁹⁴, D. Sarkar ¹⁴³, N. Sarkar ¹⁴¹, P. Sarma ⁴¹, V.M. Sarti ¹⁰³, M.H.P. Sas ⁶³, E. Scapparone ⁵³, B. Schaefer ⁹⁴, J. Schambach ¹¹⁹, H.S. Scheid ⁶⁸, C. Schiaua ⁴⁷, R. Schicker ¹⁰², A. Schmah ¹⁰², C. Schmidt ¹⁰⁵, H.R. Schmidt ¹⁰¹, M.O. Schmidt ¹⁰², M. Schmidt ¹⁰¹, N.V. Schmidt ^{94,68}, A.R. Schmier ¹³⁰, J. Schukraft ^{34,87}, Y. Schutz ^{34,136}, K. Schwarz ¹⁰⁵, K. Schweda ¹⁰⁵, G. Scioli ²⁷, E. Scomparin ⁵⁸, M. Šefčík ³⁸, J.E. Seger ¹⁶, Y. Sekiguchi ¹³², D. Sekihata ^{132,45}, I. Selyuzhenkov ^{105,91}, S. Senyukov ¹³⁶, D. Serebryakov ⁶², E. Serradilla ⁷¹, P. Sett ⁴⁸, A. Sevcenco ⁶⁷, A. Shabanov ⁶², A. Shabetai ¹¹⁴, R. Shahoyan ³⁴, W. Shaikh ¹⁰⁸, A. Shangaraev ⁸⁹, A. Sharma ⁹⁸, A. Sharma ⁹⁹, M. Sharma ⁹⁹, N. Sharma ⁹⁸, A.I. Sheikh ¹⁴¹, K. Shigaki ⁴⁵, M. Shimomura ⁸¹, S. Shirinkin ⁹⁰, Q. Shou ¹¹¹, Y. Sibiriak ⁸⁶, S. Siddhanta ⁵⁴, T. Siemiarczuk ⁸³, D. Silvermyr ⁷⁹, C. Silvestre ⁷⁷, G. Simatovic ⁸⁸, G. Simonetti ^{103,34}, R. Singh ⁸⁴, R. Singh ⁹⁹, V.K. Singh ¹⁴¹, V. Singhal ¹⁴¹, T. Sinha ¹⁰⁸, B. Sitar ¹⁴, M. Sitta ³², T.B. Skaali ²¹, M. Slupecki ¹²⁶, N. Smirnov ¹⁴⁶, R.J.M. Snellings ⁶³, T.W. Snellman ¹²⁶, J. Sochan ¹¹⁶, C. Soncco ¹¹⁰, J. Song ^{60,125}, A. Songmoolnak ¹¹⁵, F. Soramel ²⁹, S. Sorensen ¹³⁰, I. Sputowska ¹¹⁸, J. Stachel ¹⁰², I. Stan ⁶⁷, P. Stankus ⁹⁴, P.J. Steffanic ¹³⁰, E. Stenlund ⁷⁹, D. Stocco ¹¹⁴, M.M. Storetvedt ³⁶, P. Strmen ¹⁴, A.A.P. Suaide ¹²¹, T. Sugitate ⁴⁵, C. Suire ⁶¹, M. Suleymanov ¹⁵, M. Suljic ³⁴, R. Sultanov ⁹⁰, M. Šumbera ⁹³, S. Sumowidagdo ⁵⁰, K. Suzuki ¹¹³, S. Swain ⁶⁵, A. Szabo ¹⁴, I. Szarka ¹⁴, U. Tabassam ¹⁵, G. Taillepied ¹³⁴, J. Takahashi ¹²², G.J. Tambave ²², M. Suleymanov ¹⁵, M. Suljic ³⁴, K. Sultanov ³⁵, M. Sumbera ³⁵, S. Sumowidagdo ³⁰, K. Suzuki ¹¹³, S. Swain ⁶⁵, A. Szabo ¹⁴, I. Szarka ¹⁴, U. Tabassam ¹⁵, G. Taillepied ¹³⁴, J. Takahashi ¹²², G.J. Tambave ²², S. Tang ^{134,6}, M. Tarhini ¹¹⁴, M.G. Tarzila ⁴⁷, A. Tauro ³⁴, G. Tejeda Muñoz ⁴⁴, A. Telesca ³⁴, C. Terrevoli ^{125,29}, D. Thakur ⁴⁹, S. Thakur ¹⁴¹, D. Thomas ¹¹⁹, F. Thoresen ⁸⁷, R. Tieulent ¹³⁵, A. Tikhonov ⁶², A.R. Timmins ¹²⁵, A. Toia ⁶⁸, N. Topilskaya ⁶², M. Toppi ⁵¹, F. Torales-Acosta ²⁰, S.R. Torres ¹²⁰, S. Tripathy ⁴⁹, T. Tripathy ⁴⁸, S. Trogolo ^{26,29}, G. Trombetta ³³, L. Tropp ³⁸, V. Trubnikov ², W.H. Trzaska ¹²⁶, T.P. Trzcinski ¹⁴², B.A. Trzeciak ⁶³, T. Tsuji ¹³², A. Tumkin ¹⁰⁷, R. Turrisi ⁵⁶, T.S. Tveter ²¹, K. Ullaland ²², E.N. Umaka ¹²⁵, A. Uras ¹³⁵, G.L. Usai ²⁴, A. Utrobicic ⁹⁷, M. Vala ^{116,38}, N. Valle ¹³⁹, S. Vallero ⁵⁸, N. van der Kolk ⁶³, I.V.R. van Doremalen ⁶³, M. van Leeuwen ⁶³, P. Vando Vanro ³⁴ S. Vallero⁵⁸, N. van der Kolk⁶³, L.V.R. van Doremalen⁶³, M. van Leeuwen⁶³, P. Vande Vyvre³⁴, D. Varga¹⁴⁵, Z. Varga¹⁴⁵, M. Varga-Kofarago¹⁴⁵, A. Vargas⁴⁴, M. Vargyas¹²⁶, R. Varma⁴⁸, M. Vasileiou⁸², A. Vasiliev⁸⁶, O. Vázquez Doce^{117,103}, V. Vechernin¹¹², A.M. Veen⁶³, E. Vercellin²⁶, S. Vergara Limón⁴⁴, L. Vermunt⁶³, R. Vernet⁷, R. Vértesi¹⁴⁵, M.G.D.L.C. Vicencio⁹, L. Vickovic³⁵, J. Viinikainen¹²⁶, Z. Vilakazi¹³¹, O. Villalobos Baillie¹⁰⁹, A. Villatoro Tello⁴⁴, G. Vino⁵², A. Vinogradov⁸⁶, T. Virgili³⁰, Z. Vilakazi ¹³¹, O. Villalobos Baillie ¹⁰⁹, A. Villatoro Tello ⁴⁴, G. Vino ⁵², A. Vinogradov ⁸⁶, T. Virgili ³⁰, V. Vislavicius ⁸⁷, A. Vodopyanov ⁷⁴, B. Volkel ³⁴, M.A. Völkl ¹⁰¹, K. Voloshin ⁹⁰, S.A. Voloshin ¹⁴³, G. Volpe ³³, B. von Haller ³⁴, I. Vorobyev ¹⁰³, D. Voscek ¹¹⁶, J. Vrláková ³⁸, B. Wagner ²², Y. Watanabe ¹³³, M. Weber ¹¹³, S.G. Weber ^{144,105}, A. Wegrzynek ³⁴, D.F. Weiser ¹⁰², S.C. Wenzel ³⁴, J.P. Wessels ¹⁴⁴, E. Widmann ¹¹³, J. Wiechula ⁶⁸, J. Wikne ²¹, G. Wilk ⁸³, J. Wilkinson ⁵³, G.A. Willems ³⁴, E. Willsher ¹⁰⁹, B. Windelband ¹⁰², W.E. Witt ¹³⁰, Y. Wu ¹²⁸, R. Xu ⁶, S. Yalcin ⁷⁶, K. Yamakawa ⁴⁵, S. Yang ²², S. Yano ¹³⁷, Z. Yasin ^{vi}, Z. Yin ⁶, H. Yokoyama ⁶³, I.-K. Yoo ¹⁸, J.H. Yoon ⁶⁰, S. Yuan ²², A. Yuncu ¹⁰², V. Yurchenko ², V. Zaccolo ^{58,25}, A. Zaman ¹⁵, C. Zampolli ³⁴, H.J.C. Zanoli ¹²¹, N. Zardoshti ³⁴, A. Zarochentsev ¹¹², P. Závada ⁶⁶, N. Zaviyalov ¹⁰⁷, H. Zbroszczyk ¹⁴², M. Zhalov ⁹⁶, X. Zhang ⁶, Z. Zhang ^{6,134}, C. Zhao ²¹, V. Zherebchevskii ¹¹², N. Zhigareva ⁹⁰, D. Zhou ⁶, Y. Zhou ⁸⁷, Z. Zhou ²², J. Zhu ⁶, Y. Zhu ⁶, A. Zichichi ^{27,10}, M.B. Zimmermann ³⁴, G. Zinovjev ², N. Zurlo ¹⁴⁰

- ² Bogolyubov Institute for Theoretical Physics, National Academy of Sciences of Ukraine, Kiev, Ukraine
- ³ Bose Institute, Department of Physics and Centre for Astroparticle Physics and Space Science (CAPSS), Kolkata, India

- ⁵ California Polytechnic State University, San Luis Obispo, CA, United States
- ⁶ Central China Normal University, Wuhan, China

¹ A.I. Alikhanyan National Science Laboratory (Yerevan Physics Institute) Foundation, Yerevan, Armenia

⁴ Budker Institute for Nuclear Physics, Novosibirsk, Russia

- ⁸ Centro de Aplicaciones Tecnológicas y Desarrollo Nuclear (CEADEN), Havana, Cuba
- ⁹ Centro de Investigación y de Estudios Avanzados (CINVESTAV), Mexico City and Mérida, Mexico
- ¹⁰ Centro Fermi Museo Storico della Fisica e Centro Studi e Ricerche "Enrico Fermi', Rome, Italy
- ¹¹ Chicago State University, Chicago, IL, United States
- ¹² China Institute of Atomic Energy, Beijing, China
- ¹³ Chonbuk National University, Jeonju, Republic of Korea
- ¹⁴ Comenius University Bratislava, Faculty of Mathematics, Physics and Informatics, Bratislava, Slovakia
- ¹⁵ COMSATS University Islamabad, Islamabad, Pakistan
- ¹⁶ Creighton University, Omaha, NE, United States
- ¹⁷ Department of Physics, Aligarh Muslim University, Aligarh, India
- ¹⁸ Department of Physics, Pusan National University, Pusan, Republic of Korea
- ¹⁹ Department of Physics, Sejong University, Seoul, Republic of Korea
- ²⁰ Department of Physics, University of California, Berkeley, CA, United States
- ²¹ Department of Physics, University of Oslo, Oslo, Norway
- ²² Department of Physics and Technology, University of Bergen, Bergen, Norway
- ²³ Dipartimento di Fisica dell'Università 'La Sapienza' and Sezione INFN, Rome, Italy
- ²⁴ Dipartimento di Fisica dell'Università and Sezione INFN, Cagliari, Italy
- ²⁵ Dipartimento di Fisica dell'Università and Sezione INFN, Trieste, Italy
- ²⁶ Dipartimento di Fisica dell'Università and Sezione INFN, Turin, Italy
- ²⁷ Dipartimento di Fisica e Astronomia dell'Università and Sezione INFN, Bologna, Italy
- ²⁸ Dipartimento di Fisica e Astronomia dell'Università and Sezione INFN, Catania, Italy
- ²⁹ Dipartimento di Fisica e Astronomia dell'Università and Sezione INFN, Padova, Italy
- ³⁰ Dipartimento di Fisica 'E.R. Caianiello' dell'Università and Gruppo Collegato INFN, Salerno, Italy
- ³¹ Dipartimento DISAT del Politecnico and Sezione INFN, Turin, Italy
- ³² Dipartimento di Scienze e Innovazione Tecnologica dell'Università del Piemonte Orientale and INFN Sezione di Torino, Alessandria, Italy
- ³³ Dipartimento Interateneo di Fisica 'M. Merlin' and Sezione INFN, Bari, Italy
- ³⁴ European Organization for Nuclear Research (CERN), Geneva, Switzerland
- ³⁵ Faculty of Electrical Engineering, Mechanical Engineering and Naval Architecture, University of Split, Split, Croatia
- ³⁶ Faculty of Engineering and Science, Western Norway University of Applied Sciences, Bergen, Norway
- ³⁷ Faculty of Nuclear Sciences and Physical Engineering, Czech Technical University in Prague, Prague, Czech Republic
- ³⁸ Faculty of Science, P.J. Šafárik University, Košice, Slovakia
- ³⁹ Frankfurt Institute for Advanced Studies, Johann Wolfgang Goethe-Universität Frankfurt, Frankfurt, Germany
- ⁴⁰ Gangneung-Wonju National University, Gangneung, Republic of Korea
- ⁴¹ Gauhati University, Department of Physics, Guwahati, India
- ⁴² Helmholtz-Institut für Strahlen- und Kernphysik, Rheinische Friedrich-Wilhelms-Universität Bonn, Bonn, Germany
- ⁴³ Helsinki Institute of Physics (HIP), Helsinki, Finland
- ⁴⁴ High Energy Physics Group, Universidad Autónoma de Puebla, Puebla, Mexico
- ⁴⁵ Hiroshima University, Hiroshima, Japan
- ⁴⁶ Hochschule Worms, Zentrum für Technologietransfer und Telekommunikation (ZTT), Worms, Germany
- ⁴⁷ Horia Hulubei National Institute of Physics and Nuclear Engineering, Bucharest, Romania
- ⁴⁸ Indian Institute of Technology Bombay (IIT), Mumbai, India
- ⁴⁹ Indian Institute of Technology Indore, Indore, India
- ⁵⁰ Indonesian Institute of Sciences, Jakarta, Indonesia
- ⁵¹ INFN, Laboratori Nazionali di Frascati, Frascati, Italy
- ⁵² INFN, Sezione di Bari, Bari, Italy
- ⁵³ INFN, Sezione di Bologna, Bologna, Italy
- ⁵⁴ INFN, Sezione di Cagliari, Cagliari, Italy
- ⁵⁵ INFN, Sezione di Catania, Catania, Italy
- ⁵⁶ INFN. Sezione di Padova, Padova, Italy
- ⁵⁷ INFN, Sezione di Roma, Rome, Italy
- ⁵⁸ INFN, Sezione di Torino, Turin, Italy
- ⁵⁹ INFN, Sezione di Trieste, Trieste, Italy
- ⁶⁰ Inha University, Incheon, Republic of Korea

61 Institut de Physique Nucléaire d'Orsay (IPNO), Institut National de Physique Nucléaire et de Physique des Particules (IN2P3/CNRS), Université de Paris-Sud, Université Paris-Saclay, Orsay, France

- ⁶² Institute for Nuclear Research, Academy of Sciences, Moscow, Russia
- ⁶³ Institute for Subatomic Physics, Utrecht University/Nikhef, Utrecht, Netherlands
- ⁶⁴ Institute of Experimental Physics, Slovak Academy of Sciences, Košice, Slovakia
- ⁶⁵ Institute of Physics, Homi Bhabha National Institute, Bhubaneswar, India
- ⁶⁶ Institute of Physics of the Czech Academy of Sciences, Prague, Czech Republic
- ⁶⁷ Institute of Space Science (ISS), Bucharest, Romania
- ⁶⁸ Institut für Kernphysik, Johann Wolfgang Goethe-Universität Frankfurt, Frankfurt, Germany
- ⁶⁹ Instituto de Ciencias Nucleares, Universidad Nacional Autónoma de México, Mexico City, Mexico
- ⁷⁰ Instituto de Física, Universidade Federal do Rio Grande do Sul (UFRGS), Porto Alegre, Brazil
- ⁷¹ Instituto de Física, Universidad Nacional Autónoma de México, Mexico City, Mexico
- ⁷² iThemba LABS, National Research Foundation, Somerset West, South Africa
- 73 Johann-Wolfgang-Goethe Universität Frankfurt Institut für Informatik, Fachbereich Informatik und Mathematik, Frankfurt, Germany
- ⁷⁴ Joint Institute for Nuclear Research (JINR), Dubna, Russia
- ⁷⁵ Korea Institute of Science and Technology Information, Daejeon, Republic of Korea
- ⁷⁶ KTO Karatay University, Konya, Turkey
- ⁷⁷ Laboratoire de Physique Subatomique et de Cosmologie, Université Grenoble-Alpes, CNRS-IN2P3, Grenoble, France
- ⁷⁸ Lawrence Berkeley National Laboratory, Berkeley, CA, United States
- 79 Lund University Department of Physics, Division of Particle Physics, Lund, Sweden
- ⁸⁰ Nagasaki Institute of Applied Science, Nagasaki, Japan
- ⁸¹ Nara Women's University (NWU), Nara, Japan
- ⁸² National and Kapodistrian University of Athens, School of Science, Department of Physics, Athens, Greece
- ⁸³ National Centre for Nuclear Research, Warsaw, Poland
- ⁸⁴ National Institute of Science Education and Research, Homi Bhabha National Institute, Jatni, India

⁸⁵ National Nuclear Research Center, Baku, Azerbaijan

- ⁸⁶ National Research Centre Kurchatov Institute, Moscow, Russia
- ⁸⁷ Niels Bohr Institute. University of Copenhagen, Copenhagen, Denmark
- ⁸⁸ Nikhef, National institute for subatomic physics, Amsterdam, Netherlands
- 89 NRC Kurchatov Institute IHEP, Protvino, Russia
- 90 NRC Kurchatov Institute ITEP, Moscow, Russia
- ⁹¹ NRNU Moscow Engineering Physics Institute, Moscow, Russia
- ⁹² Nuclear Physics Group, STFC Daresbury Laboratory, Daresbury, United Kingdom
- ⁹³ Nuclear Physics Institute of the Czech Academy of Sciences, Řež u Prahy, Czech Republic
- ⁹⁴ Oak Ridge National Laboratory, Oak Ridge, TN, United States
- 95 Ohio State University, Columbus, OH, United States
- 96 Petersburg Nuclear Physics Institute, Gatchina, Russia
- ⁹⁷ Physics department, Faculty of science, University of Zagreb, Zagreb, Croatia
- 98 Physics Department, Panjab University, Chandigarh, India
- ⁹⁹ Physics Department, University of Jammu, Jammu, India
- ¹⁰⁰ Physics Department, University of Raiasthan, Jaipur, India
- ¹⁰¹ Physikalisches Institut, Eberhard-Karls-Universität Tübingen, Tübingen, Germany
- ¹⁰² Physikalisches Institut, Ruprecht-Karls-Universität Heidelberg, Heidelberg, Germany
- ¹⁰³ Physik Department, Technische Universität München, Munich, Germany
- ¹⁰⁴ Politecnico di Bari, Bari, Italy
- ¹⁰⁵ Research Division and ExtreMe Matter Institute EMMI, GSI Helmholtzzentrum für Schwerionenforschung GmbH, Darmstadt, Germany
- ¹⁰⁶ Rudjer Bošković Institute, Zagreb, Croatia
- ¹⁰⁷ Russian Federal Nuclear Center (VNIIEF), Sarov, Russia
- ¹⁰⁸ Saha Institute of Nuclear Physics, Homi Bhabha National Institute, Kolkata, India
- ¹⁰⁹ School of Physics and Astronomy, University of Birmingham, Birmingham, United Kingdom
- ¹¹⁰ Sección Física, Departamento de Ciencias, Pontificia Universidad Católica del Perú, Lima, Peru
- ¹¹¹ Shanghai Institute of Applied Physics, Shanghai, China
- ¹¹² St. Petersburg State University, St. Petersburg, Russia
- ¹¹³ Stefan Mever Institut für Subatomare Physik (SMI), Vienna, Austria
- ¹¹⁴ SUBATECH, IMT Atlantique, Université de Nantes, CNRS-IN2P3, Nantes, France
- ¹¹⁵ Suranaree University of Technology, Nakhon Ratchasima, Thailand
- ¹¹⁶ Technical University of Košice, Košice, Slovakia
- ¹¹⁷ Technische Universität München, Excellence Cluster 'Universe', Munich, Germany
- ¹¹⁸ The Henryk Niewodniczanski Institute of Nuclear Physics, Polish Academy of Sciences, Cracow, Poland
- ¹¹⁹ The University of Texas at Austin, Austin, TX, United States
- ¹²⁰ Universidad Autónoma de Sinaloa, Culiacán, Mexico
- ¹²¹ Universidade de São Paulo (USP), São Paulo, Brazil
- ¹²² Universidade Estadual de Campinas (UNICAMP), Campinas, Brazil
- 123 Universidade Federal do ABC, Santo Andre, Brazil
- ¹²⁴ University of Cape Town, Cape Town, South Africa ¹²⁵ University of Houston, Houston, TX, United States
- 126 University of Jyväskylä, Jyväskylä, Finland
- ¹²⁷ University of Liverpool, Liverpool, United Kingdom ¹²⁸ University of Science and Techonology of China, Hefei, China
- 129 University of South-Eastern Norway, Tonsberg, Norway
- ¹³⁰ University of Tennessee, Knoxville, TN, United States
- ¹³¹ University of the Witwatersrand, Johannesburg, South Africa
- ¹³² University of Tokyo, Tokyo, Japan
- ¹³³ University of Tsukuba, Tsukuba, Japan
- ¹³⁴ Université Clermont Auvergne, CNRS/IN2P3, LPC, Clermont-Ferrand, France
- ¹³⁵ Université de Lyon, Université Lyon 1, CNRS/IN2P3, IPN-Lyon, Villeurbanne, Lyon, France
- ¹³⁶ Université de Strasbourg, CNRS, IPHC UMR 7178, F-67000 Strasbourg, France, Strasbourg, France
- 137 Université Paris-Saclay Centre d'Etudes de Saclay (CEA), IRFU, Départment de Physique Nucléaire (DPhN), Saclay, France
- ¹³⁸ Università degli Studi di Foggia, Foggia, Italy
- ¹³⁹ Università degli Studi di Pavia, Pavia, Italy
- ¹⁴⁰ Università di Brescia, Brescia, Italy
- ¹⁴¹ Variable Energy Cyclotron Centre, Homi Bhabha National Institute, Kolkata, India
- 142 Warsaw University of Technology, Warsaw, Poland
- ¹⁴³ Wayne State University, Detroit, MI, United States
- ¹⁴⁴ Westfälische Wilhelms-Universität Münster, Institut für Kernphysik, Münster, Germany
- ¹⁴⁵ Wigner Research Centre for Physics, Hungarian Academy of Sciences, Budapest, Hungary
- ¹⁴⁶ Yale University, New Haven, CT, United States
- ¹⁴⁷ Yonsei University, Seoul, Republic of Korea
- i Deceased.
- Dipartimento DET del Politecnico di Torino, Turin, Italy.
- iii M.V. Lomonosov Moscow State University, D.V. Skobeltsyn Institute of Nuclear, Physics, Moscow, Russia.
- iv Department of Applied Physics, Aligarh Muslim University, Aligarh, India.
- Institute of Theoretical Physics, University of Wroclaw, Poland.
- vi PINSTECH, Islamabad, Pakistan.